

This is the peer reviewed version of the following article:

Surgery of Moderate Coronary Artery Stenosis / Mattioli, A. V.; Farinetti, A.; Manenti, A.; Roncati, L.. - In: ANNALS OF THORACIC SURGERY. - ISSN 0003-4975. - 106:5(2018), pp. 1594-1595. [10.1016/j.athoracsur.2018.05.035]

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.


16/05/2026 00:23

(Article begins on next page)

# Accepted Manuscript

Surgery of Moderate Coronary Artery Stenosis

Anna Vittoria Mattioli, MD, Alberto Farinetti, MD, Antonio Manenti, MD, Luca Roncati, MD



PII: S0003-4975(18)30833-6

DOI: [10.1016/j.athoracsur.2018.05.035](https://doi.org/10.1016/j.athoracsur.2018.05.035)

Reference: ATS 31688

To appear in: *The Annals of Thoracic Surgery*

Received Date: 24 April 2018

Accepted Date: 14 May 2018

Please cite this article as: Mattioli AV, Farinetti A, Manenti A, Roncati L, Surgery of Moderate Coronary Artery Stenosis, *The Annals of Thoracic Surgery* (2018), doi: 10.1016/j.athoracsur.2018.05.035.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Surgery of Moderate Coronary Artery Stenosis

*To the Editor:*

The interesting paper by Raza et al. demonstrates that progression to severe stenosis is faster in grafted than in non-grafted moderate coronary artery stenosis (MCAS), especially when performed with saphenous vein (SV) rather than with internal thoracic artery (ITA) [1]. We correlate this to some particular pathophysiological and surgical aspects of ITA-grafts.

Coronary artery hemodynamics greatly changes after a bypass, because the new flow largely overwhelms the pre-existing, inducing a subsequent competition [2]. This is more striking in case of bypass with a SV than with an ITA. In the first case the blood flow, directly correlated with that of the ascending aorta, cancels or largely decreases any other downstream from a MCAS, favoring a subsequent thrombosis. In the second case, the new flow is modulated by ITA, a second-order elastic branch of the aortic arch. This artery behaves as a “feeding” elastic vessel, naturally predisposed to satisfy increased demands of blood. In coronary surgery this occurs especially when fashioned in a Y-shape on a single pedicle. Moreover, ITA benefits from a low susceptibility to atherosclerosis, and of a particular power of repairing elastolysis of the tunica media, secondary to age, diabetes, hypertension, etc., generating unspecialized smooth muscle-like cells not of vascular origin [3]. On contrary, SV-conduits remodel with deposition of extracellular collagen material, predisposing to fibrosis, with consequent loss of elasticity and increased risk of progressive obstruction. All this could explain the better results of ITA-grafts and assure surgeons about ITA as preferred conduits also in revascularizing symptomatic MCAS [1]. Besides, in multi-segmental lesions, typically of the left anterior descending coronary artery, an associated endarterectomy downstream to a MCAS, increases the graft run-off, through new-open septal perforator collaterals [4].

Beyond these surgical considerations, a basic diagnostic problem still remains concerning MCAS. After a first angiographic approach, further hemodynamic studies would demonstrate conditions of real vascular insufficiency, with an impending risk of myocardial severe ischemia, which represents an unquestionable indication to surgery [5].

Anna Vittoria Mattioli, MD

Department of Cardiology

Modena University

Modena, Italy

Alberto Farinetti, MD

Antonio Manenti, MD

Department of Surgery

Modena University

41124 Modena, Italy

Email: [antonio.manenti@unimore.it](mailto:antonio.manenti@unimore.it)

Luca Roncati, MD

Department of Pathology

Modena, Italy

**References**

1. Raza S, Blackstone EH, Houghtaling PL, Olivares G, Ravichandren K, Koprivanac M, et al. Natural history of moderate coronary artery stenosis after surgical revascularization. *Ann Thorac Surg* 2018;105:815-21.
2. Ferguson TB Jr. Physiology of in-situ arterial revascularization in coronary artery bypass grafting: preoperative, intraoperative and postoperative factors and influences. *World J Cardiol* 2016;8:623-37.
3. Roncati L, Manenti A, Caprili L, Fedeli C. Internal thoracic artery histologic characteristics clarify the high performance in coronary bypass. *Ann Thorac Surg* 2016; 101:2429-30.
4. Nishigawa K, Fukui T, Yamazaki Y, et al. Ten-year experience of coronary endarterectomy for the diffusely diseased left anterior descending artery. *Ann Thorac Surg* 2017; 103: 710-6.
5. Jeremias A, Kirtane AJ, Stone GW. A test in context: fractional flow reserve: accuracy, prognostic implications, and limitations. *JACC* 2017;69:2748–58.