

This is the peer reviewed version of the following article:

Chondrocyte expression of apoptotic and pro-inflammatory factors in the development of post- traumatic arthritis in humans / Sena, P., Benincasa, M., Cavani, F., Ferretti, M., Smargiassi, A., Manfredini, G., Palumbo, C.. - In: ITALIAN JOURNAL OF ANATOMY AND EMBRYOLOGY. - ISSN 1122-6714. - STAMPA. - 118 - n. 2 (suppl.):(2013), pp. 172-172. (67th Meeting of the Italian Society of Anatomy and Histology Brescia (Italy) 20-22 September, 2013).

Firenze University Press

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

30/10/2026 04:58

(Article begins on next page)

Chondrocyte expression of apoptotic and pro-inflammatory factors in the development of post-traumatic arthritis in humans

Paola Sena, Marta Benincasa, Francesco Cavani, Marzia Ferretti, Alberto Smargiassi, Giuseppe Manfredini and Carla Palumbo

Dip. Scienze Biomediche, Metaboliche e Neuroscienze - Sez. Morfologia umana, Università degli Studi di Modena e RE, Italy

The development of post-traumatic arthritis following intra-articular fracture remains an important unsolved clinical problem. The possibility that extensive chondrocyte apoptosis occurs following intra-articular fracture, thus contributing to the development of post-traumatic arthritis, has received increasing attention [1]. It has been demonstrated the existence of a direct correlation between the rate of apoptosis and the severity of osteoarthritis [2]. Pharmacologic inhibitors of enzymes involved in apoptosis have been explored as potential therapeutic agents [3]. In the present study we aimed to deepen the characterization of apoptotic mediators, expressed by chondrocytes, involved in human post-traumatic arthritis following intra-articular fracture and the possible implication of pro-inflammatory receptors in arthritis. The expression of a panel of pro/anti apoptotic factors (Caspase-3, PARP-1, BCL2) and inflammation-related receptors (ChemR23) were analysed in chondrocytes from patients undergoing surgery for intra-articular calcaneal fractures. The factors were investigated by immunofluorescence coupled with confocal analysis and western blotting, followed by densitometric evaluation of chondrocyte cultures harvested from patients with intra-articular fractures compared with control ones. The results clearly demonstrated that a statistically significant difference exists in the expression of pro/anti apoptotic factors and ChemR23 between fractured and control patients. In conclusion our data suggest that increased chondrocyte death, occurring after cartilage injury together with inflammatory process, could play a pivotal role in the onset of arthritic disease.

References

- [1] Hembree W.C. et al. (2007) Viability and apoptosis of human chondrocytes in osteochondral fragments following joint trauma. *J Bone Joint Surg Br* 89(10): 1388-95.
- [2] Kim H.A. et al. (2000) Apoptotic chondrocyte death in human osteoarthritis. *J Rheumatol* 27: 455-462.
- [3] D'Lima D. et al. (2006) Caspase inhibitors reduce severity of cartilage lesions in experimental osteoarthritis. *Arthritis Rheum* 54(6): 1814-1821.

Key words

Articular fracture, post-traumatic arthritis, chondrocyte culture, apoptosis, inflammation.