

This is the peer reviewed version of the following article:

Recurrence of hypertensive disorders of pregnancy: an individual patient data metaanalysis / van Oostwaard, Miriam F.; Langenveld, Josje; Schuit, Ewoud; Papatsonis, Dimitri N. M.; Brown, Mark A.; Byaruhanga, Romano N.; Bhattacharya, Sohinee; Campbell, Doris M.; Chappell, Lucy C.; Chiaffarino, Francesca; Crippa, Isabella; Facchinetti, Fabio; Ferrazzani, Sergio; Ferrazzi, Enrico; Figueiró Filho, Ernesto A.; Gaugler Senden, Ingrid P. M.; Haavaldsen, Camilla; Lykke, Jacob A.; Mbah, Alfred K.; Oliveira, Vanessa M.; Poston, Lucilla; Redman, Christopher W. G.; Salim, Raed; Thilaganathan, Baskaran; Vergani, Patrizia; Zhang, Jun; Steegers, Eric A. P.; Mol, Ben Willem J.; Ganzevoort, Wessel. - In: AMERICAN JOURNAL OF OBSTETRICS AND GYNECOLOGY. - ISSN 0002-9378. - STAMPA. - 212:5(2015), pp. 1-17. [10.1016/j.ajog.2015.01.009]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

19/12/2025 00:27

19/12/2025 00:27

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/270967359>

Recurrence of hypertensive disorders of pregnancy, an Individual Patient Data Meta-Analysis.

Article in *American Journal of Obstetrics and Gynecology* · January 2015

DOI: 10.1016/j.ajog.2015.01.009 · Source: PubMed

CITATIONS

6

READS

189

29 authors, including:



Sohinee Bhattacharya

University of Aberdeen

122 PUBLICATIONS **1,473** CITATIONS

[SEE PROFILE](#)



Enrico Ferrazzi

University of Milan

386 PUBLICATIONS **5,641** CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



THINC. - The Healthcare Innovation Center [View project](#)



Opioid pseudopeptides [View project](#)

All content following this page was uploaded by [Ernesto Antonio Figueiró-Filho](#) on 07 February 2015.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.

Accepted Manuscript



Recurrence of hypertensive disorders of pregnancy, an Individual Patient Data Meta-Analysis

Miriam F. Van Oostwaard, MD, Josje Langenveld, MD, PhD, Ewoud Schuit, MSc, PhD, Dimitri N.M. Papatsonis, MD, PhD, Mark A. Brown, MD, PhD, Romano N. Byaruhanga, MD, PhD, Sohinee Bhattacharya, MD, PhD, Doris M. Campbell, MD, PhD, Lucy C. Chappell, MD, PhD, Francesca Chiaffarino, ScD, Isabella Crippa, MD, PhD, Fabio Facchinetti, MD, PhD, Sergio Ferrazzani, MD, PhD, Enrico Ferrazzi, MD, PhD, Ernesto A. Figueiró-Filho, MD, PhD, Ingrid P.M. Gaugler-Senden, MD, PhD, Camilla Haavaldsen, MD, PhD, Jacob A. Lykke, MD, PhD, Alfred K. Mbah, PhD, Vanessa M. Oliveira, MD, PhD, Lucilla Poston, MD, PhD, Christopher W.G. Redman, MD, PhD, Raed Salim, MD, PhD, Baskaran Thilaganathan, MD, PhD, Patrizia Vergani, MD, PhD, Jun Zhang, MD, PhD, Eric A.P. Steegers, MD, PhD, Ben Willem J. Mol, MD, PhD, Wessel Ganzevoort, MD, PhD.

PII: S0002-9378(15)00010-1

DOI: [10.1016/j.ajog.2015.01.009](https://doi.org/10.1016/j.ajog.2015.01.009)

Reference: YMOB 10210

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 22 June 2014

Revised Date: 10 November 2014

Accepted Date: 7 January 2015

Please cite this article as: Van Oostwaard MF, Langenveld J, Schuit E, Papatsonis DNM, Brown MA, Byaruhanga RN, Bhattacharya S, Campbell DM, Chappell LC, Chiaffarino F, Crippa I, Facchinetti F, Ferrazzani S, Ferrazzi E, Figueiró-Filho EA, Gaugler-Senden IPM, Haavaldsen C, Lykke JA, Mbah AK, Oliveira VM, Poston L, Redman CWG, Salim R, Thilaganathan B, Vergani P, Zhang J, Steegers EAP, Mol BWJ, Ganzevoort W, Recurrence of hypertensive disorders of pregnancy, an Individual Patient Data Meta-Analysis, *American Journal of Obstetrics and Gynecology* (2015), doi: 10.1016/j.ajog.2015.01.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please

note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Recurrence of hypertensive disorders of pregnancy, an Individual Patient Data Meta-Analysis

Miriam F. VAN OOSTWAARD¹, MD, Josje LANGENVELD², MD, PhD, Ewoud SCHUIT^{3,4,5}, MSc, PhD Dimitri N.M. PAPATSONIS⁶, MD, PhD, Mark A. BROWN⁷, MD, PhD, Romano N. BYARUHANGA⁸, MD, PhD, Sohinee BHATTACHARYA⁹, MD, PhD, Doris M. CAMPBELL¹⁰, MD, PhD, Lucy C. CHAPPELL¹¹, MD, PhD, Francesca CHIAFFARINO¹², ScD, Isabella CRIPPA¹³, MD, PhD, Fabio FACCHINETTI¹⁴, MD, PhD, Sergio FERRAZZANI¹⁵, MD, PhD, Enrico FERRAZZI¹⁶, MD, PhD, Ernesto A. FIGUEIRO-FILHO¹⁷, MD, PhD, Ingrid P.M. GAUGLER-SENDEN¹⁸, MD, PhD, Camilla HAAVALDSEN¹⁹, MD, PhD, Jacob A. LYKKE²⁰, MD, PhD, Alfred K. MBAH²¹, PhD, Vanessa M. OLIVEIRA²², MD, PhD, Lucilla POSTON¹¹, MD, PhD, Christopher W.G. REDMAN²³, MD, PhD, Raed SALIM²⁴, MD, PhD, Baskaran THILAGANATHAN²⁵, MD, PhD, Patrizia VERGANI¹³, MD, PhD, Jun ZHANG²⁶, MD, PhD, Eric A.P. STEEGERS¹, MD, PhD, Ben Willem J. MOL²⁷, MD, PhD, Wessel GANZEVOORT⁴, MD, PhD.

1. Department of Obstetrics and Gynecology Erasmus Medical Centre, Rotterdam, the Netherlands
2. Department of Obstetrics and Gynecology Atrium Medical Centre, Heerlen, the Netherlands
3. Julius Centre for Health Sciences and Primary Care, Universitair Medisch Centrum, Utrecht, the Netherlands
4. Department of Obstetrics and Gynecology Academic Medical Centre, Amsterdam, the Netherlands
5. Stanford Prevention Research Center, Stanford University, Stanford, CA, USA
6. Department of Obstetrics and Gynecology Amphia Hospital, Breda, the Netherlands
7. Department of Renal Medicine, St George Hospital, Sydney, Australia
8. Department Obstetrics and Gynecology St. Raphael of St. Francis Hospital, Nsambya Kampala, Uganda
9. Dugald Baird Centre for Research on Women's Health, Aberdeen Maternity Hospital, Aberdeen, United Kingdom
10. Department of Obstetrics and Gynecology, Aberdeen Maternity Hospital, Aberdeen, United Kingdom
11. Women's Health Academic Centre, King's College London, United Kingdom
12. Department of Obstetrics, Gynecology and Neonatology, IRCSS Fondazione Ca' Granda, Ospedale Maggiore Policlinico, Milan, Italy
13. Department of Obstetrics and Gynecology, Ospedale San Gerardo, Università degli Studi di Milano-Bicocca, Monza, Italy
14. Department of Obstetrics and Gynecology, Università degli Studi di Modena e Reggio Emilia, Modena, Italy
15. Department of Obstetrics and Gynecology, Università Cattolica del S. Cuore, Roma, Italy
16. Department of Woman, Mother and Neonate, Buzzi Children's Hospital, Istituti Clinici di Perfezionamento, Biomedical and Clinical School of Medicine University of Milan, Italy
17. Faculty of Medicine (FAMED), Federal University of Mato Grosso do Sul (UFMS), Campo Grande, Brazil
18. Department of Obstetrics and Gynecology, Jeroen Bosch Ziekenhuis, 's Hertogenbosch, The Netherlands
19. Department of Gynecology and Obstetrics, and Institute of Clinical Medicine, Akershus University Hospital, University of Oslo, Lørenskog, Norway
20. Department of Obstetrics and Gynecology, Hvidovre Hospital, Copenhagen, Denmark and Faculty of Health Science, University of Copenhagen, Denmark
21. Department of Epidemiology and Biostatistics, University of South Florida, Tampa, USA
22. Center for Biological and Health Sciences, Federal University of Mato Grosso do Sul, Campo Grande, Brazil
23. Nuffield Department of Obstetrics and Gynaecology, John Radcliffe Hospital, University of Oxford, United Kingdom
24. Department of Obstetrics and Gynecology, Ha'Emek Medical Center, Afula, Israel
25. Lanesborough Wing, St George's, University of London, United Kingdom
26. Xinhua Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, China
27. The Robinson Institute, School of Paediatrics and Reproductive Health, University of Adelaide, Australia

Disclosure of Interests:

The authors report no conflict of interest

Sources of financial support:

No financial support has been received for realization of this research

Corresponding author:Reprints:

Miriam F. van Oostwaard
Erasmus MC - Sophia
Department of Obstetrics and Gynecology
PO BOX 2060
3000 CB Rotterdam
The Netherlands
E-mail: miriamvanoostwaard@gmail.com

For Editorial use only:

Miriam F. van Oostwaard
Bremstraat 15
3053DC Rotterdam
The Netherlands
miriamvanoostwaard@gmail.com
Tel: +31-6-16432022

Word count:

Abstract: 336
Manuscript: 3814

Top Heading: Obstetrics**Condensation:**

The recurrence rates of hypertensive disorders of pregnancy, preeclampsia, gestational hypertension, and HELLP syndrome were 20.7%, 13.8%, 8.6%, and 0.2% respectively.

Short version of title:

Individual participant data meta-analysis on the recurrence of hypertensive disorders of pregnancy

This research was presented as a poster at the 19th Annual Scientific Meeting of the International Society for the Study of Hypertension in Pregnancy, ISSHP, New Orleans, Louisiana, USA, October 26-29 2014.

ABSTRACT:**Objective**

We performed an Individual Participant Data (IPD) meta-analysis to calculate the recurrence risk of hypertensive disorders of pregnancy (HDP) and recurrence of individual hypertensive syndromes.

Study design

We performed an electronic literature search for cohort studies that reported on women suffering from HDP and who had a subsequent pregnancy. The principal investigators were contacted, informed and requested for their original study data. The obtained data were merged to form one combined database. The results will be presented as % with 95% confidence interval (CI) and odds ratios (OR) with 95% CI.

Results

Out of 94 eligible cohort studies, we obtained IPD of 22 studies, including a total of 99,415 women. Pooled data of 64 studies using published data (IPD where available) showed a recurrence rate of 18.1% (N=152,213, 95% CI: 17.9 – 18.3). In the 22 studies included in our IPD, the recurrence rate of a HDP was 20.7% (95%CI: 20.4 – 20.9). Recurrence manifested as preeclampsia (PE) in 13.8% (95%CI: 13.6 – 14.1), gestational hypertension (GH) in 8.6% (95%CI: 8.4 – 8.8) and Hemolysis, Elevated Liver enzymes and Low Platelets (HELLP) syndrome in 0.2% (95%CI: 0.16 – 0.25). The delivery of a small for gestational age (SGA) child accompanied the recurrent HDP in 3.4% (95%CI: 3.2 – 3.6). Concomitant HELLP syndrome or delivery of a SGA child increased the risk of recurrence of HDP. Recurrence increased with decreasing

gestational age at delivery in the index pregnancy. If the HDP recurred, it was in general milder, regarding maximum diastolic blood pressure, proteinuria, use of oral antihypertensive and anticonvulsive medication, delivery of a SGA child, premature delivery and perinatal mortality. Normotensive women developed chronic hypertension after pregnancy more often after experiencing recurrence (OR 3.7 95% CI: 2.3 – 6.1).

Conclusion

Among women that suffer hypertension in pregnancy, the recurrence rate in a next pregnancy is relatively low and the course of disease is milder for most women with recurrent disease. These reassuring data should be used for shared decision making in women who consider a new pregnancy after a pregnancy complicated by hypertension.

Key words

Gestational hypertension, HELLP syndrome, IPD, preeclampsia, pregnancy, recurrence

INTRODUCTION:

Hypertensive disorders of pregnancy (HDP) are a common health problem and are the second most common cause of maternal death worldwide, with major intriguing regional differences worldwide (1). They complicate approximately 2 to 8 per cent of all pregnancies (2,3,4) and comprise gestational hypertension (GH), preeclampsia (PE), superimposed PE and HELLP (Hemolysis, Elevated Liver enzymes and Low Platelets) syndrome, and in a varying percentage of cases they are related to intrauterine growth restriction. There is probably important heterogeneity in pathophysiology and clinical phenotype between subgroups and individual women, with maternal endothelial dysfunction as a central phenomenon, caused by an excessive maternal response to placental material. Furthermore a HDP identifies the woman at risk for cardiovascular disease later in life, probably due to shared risk factors and pathophysiology (2,4).

Hypertensive disorders of pregnancy can also have a major psychological impact on the woman and her family (5). As such, counseling on recurrence of a hypertensive disease in a future pregnancy is important. Consequently, many studies focused on the investigation of recurrence rates of HDP. Interpretation of these studies, however, is difficult because of many potential sources of bias, including patient selection and study methodologies. This causes every cohort to have a specific case mix of different clinical phenotypes and macro-ethnicities. Reported recurrence rates range from a few percent, up to 65% (6). Similarly, the performance of individualized risk prediction models has been disappointing, since hypertensive disorders of pregnancy have heterogeneous pathophysiology (7). In addition, suboptimal size and

generalizability of studies cause these prediction models to be unsatisfactory (7).

Recurrence rates of the individual syndromes of HDP in literature for preeclampsia can be found ranging up to 65% (6) and for HELLP syndrome: 2 – 3% (8,9).

Recurrence of GH has not been explored in many studies. Delivering a small for gestational age (SGA) child recurs in about 24 percent of pregnancies (10).

Individual Participant Data (IPD) meta-analysis is new to prognostic research (11). In contrast to conventional meta-analysis it uses the IPD of the original studies, thus enlarging the study population and increasing statistical power to detect subtle relationships. In contrast to aggregated data meta-analysis, it permits synthesis at an individual level, creating flexibility in choosing outcome and subgroups. Additionally, it allows redefinition of outcomes or predictors based on continuous variables and usage of information that did not reach publication in the original research.

Objective of the study:

The primary goal of this IPD meta-analysis study is to calculate the recurrence risk of hypertensive disorders of pregnancy. Secondly, we aim to show the recurrence of individual hypertensive syndromes.

MATERIALS AND METHODS

Sources

We performed a literature search in the electronic libraries PubMed (Medline) and Embase. Language restrictions or restrictions on publication date were not applied. The search covered all records until August 2012. The following terms were used: “preeclampsia” [MeSH] AND [early OR severe OR pre-term OR early onset OR 32 OR 34 OR 37] AND [history OR previous OR secondary OR subsequent OR recurrence]. Cross-references of the selected studies were checked to identify other studies of interest. All studies that described cohorts of women with a history of a hypertensive disorder that resulted in a delivery at any gestational age were eligible for inclusion. Inclusion was not restricted to any study design, apart from for case-control studies, where recurrence was a prerequisite. Studies that did not report recurrence of preeclampsia in the publication, but reasoned to have this information in the original data, were also considered eligible. If data between studies overlapped, only the larger of the two studies was included. Two independent reviewers (M.F.vO. and J.L.) screened the identified articles for eligibility based on title and abstract. Discrepancies were resolved by a third reviewer (W.G.).

Data collection

For each of the eligible articles, contact information of the first, second or last author was obtained through Medline, Embase or the internet. We approached the authors by email to inform them about the IPD meta-analysis project and to invite them to share their data. If authors were willing to participate, they were provided a more detailed study proposal, and asked to send their original database. Variables and

categories needed to be adequately labeled within the original database or in a separate data dictionary. Data of women who had a subsequent pregnancy after the hypertensive pregnancy were included. We focused on collecting demographic characteristics such as age and Body Mass Index (BMI), cardiovascular risk factors and the clinical syndrome of the index pregnancy. The quality of all included studies was evaluated with the Newcastle Ottawa Scale for cohort studies (12).

Definitions:

Hypertensive disorders of pregnancy were defined as gestational hypertension (GH), preeclampsia (PE), superimposed PE or HELLP syndrome. Non-hypertensive pregnancies with a small for gestational age child (SGA) were excluded. Chronic hypertension was not an exclusion criterion for the IPD, but was in some individual studies.

Preeclampsia was defined as hypertension (diastolic blood pressure ≥ 90 mmHg or systolic blood pressure ≥ 140 mmHg on two occasions, 4-5 hours apart) in combination with proteinuria (defined as a positive (0.3 g/l) proteinuria dipstick test, a protein/creatinine ratio of 30 mg/mmol or more in a random sample or an urine protein excretion of 300 mg or more per 24 hrs.) after 20 weeks of pregnancy (13).

Mild or severe PE were not separately defined, as only some of the studies made this distinction in their data. Women with hypertension beyond 20 weeks without proteinuria, or a significant rise in blood pressure if known chronic hypertension, were considered to have GH. Chronic hypertension was defined as the presence or history of preconceptional hypertension or detection of hypertension in the first half of pregnancy. De novo proteinuria or a sudden increase in proteinuria if already present, qualified women with chronic hypertension for superimposed preeclampsia

(13). HELLP syndrome was defined by hemolysis (elevated lactate dehydrogenase (LDH) levels (≥ 600 U/L), elevated liver enzymes by levels of aspartate transaminase (ASAT) or alanine transferase (ALAT) ≥ 70 U/L and low platelets $< 100,000/\text{mm}^3$ (14). HELLP syndrome in combination with hypertension was also classified as PE. SGA was defined as birth weight below the 10th percentile, according to the ACOG practice bulletin (15) and adjusted for gestational age based on a local reference population. The exact definitions for the hypertensive syndromes used in the included studies were not retrievable for a few studies, that were published as abstracts (20,22). Nevertheless we presume that the authors concur to the international accepted criteria, which were described above.

The primary outcome was the recurrence of any hypertensive disorder of pregnancy in the next subsequent pregnancy. Secondly, we aimed at showing the recurrence of individual hypertensive syndromes.

Statistical analysis

Most studies focused on detailed data of the subsequent pregnancy, whereas some studies only registered details of the index pregnancy. To utilize the data as best as possible, we combined data of the index- and subsequent pregnancy for BMI, smoking, medical history and chronic hypertension. If for example BMI is recorded at the time of the second pregnancy, but not at the index pregnancy, then we copied this BMI information.

Not all the dependent or independent variables have been registered in every database. Results are therefore accompanied with the number of cases in which the variable was registered (N). Proportions are presented as percentages of N, rather

than as percentages of the total population. For descriptive analysis we expressed continuous variables as mean with standard deviation (SD) or median with interquartile range (IQR), as appropriate. We decided not to use imputation, as the pattern of systematic missing values cannot be extrapolated between databases.

Differences in outcomes between the index and subsequent pregnancy were investigated using a random intercept random effects binomial regression model for dichotomous outcomes and using a random intercept random effects linear regression model for continuous outcomes. A random intercept was fitted per study to account for the fact that the baseline risk between studies may differ. Based on Akaike's Information Criterion (AIC) the random effects model was compared to a fixed effects model and the model with the lowest AIC was used as the final model for analysis for that outcome. Heterogeneity across studies was assessed using the I^2 measure and the values were interpreted as follows: 0% indicates no observed heterogeneity; 25%, 50%, and 75% indicate low, moderate, and high heterogeneity, respectively (16). P values less than 0.05 were considered to indicate statistical significance.

We have performed 3 post hoc sensitivity analyses: in one we performed separate analysis for studies with less or more than 200 inclusions, in one we excluded retrospective studies and in one we excluded studies with high or unclear risk of bias.

Statistical analysis was performed using SPSS 20.0 (SPSS Inc., Chicago, IL, USA) and R version 3.0.1 (The R Foundation for Statistical Computing).

RESULTS

Study selection:

Our search of Pubmed and Embase for eligible articles until August 2012 yielded 2819 non-duplicate hits. After screening title and abstract, a total of 94 articles were included. Of these articles, six had to be excluded due to overlapping databases, 39 due to failure to make contact, 14 due to unavailable data, two due to judgment of data as not useful by the authors and 11 authors decided not to share their data. Reasons for declining to share data included: no interest in participation, still in the publication process and a statement of 'no permission to send data outside the national borders'. Eventually, the IPD of 22 articles (17-39) were included in this IPD meta-analysis (Figure 1). The combined database comprised a total of 99,415 women.

An overview of the study characteristics of the included studies can be seen in Table 1. The majority of women was included from three registry-based studies that lack demographic and clinical detail (31,32,38). Two authors of included registry-based studies (21,38), were unable to provide the original data, but offered to perform a new data-extraction. They used the criteria from the initial study, creating an extended and more recent database. Eight prospective trials were included, investigating effects of aspirin (17,19,24,27,37), low-molecular-weight (LMW) heparin (25,27,37), vitamins (23,36) and close surveillance (37) on recurrence of hypertensive disease and maternal and fetal outcome. Quality characteristics of all included studies using the Newcastle Ottawa Scale (12) are shown in Figure 2.

To test for selection bias in the process of inclusion of IPD, a meta-analysis of recurrence rates in literature was performed. We used the originally included 88 articles (leaving out the overlapping datasets). Where available we used IPD (22 articles). Another 24 articles had to be excluded because applicable recurrence rates

were not published. Of these articles, only IPD that did not reach publication would have been useful, for example in specific subgroups. Regarding gestational hypertensive disorders: 64 articles, encompassing in total 152,213 women with a hypertensive disorder of pregnancy, reported on 27,558 recurrences, resulting in a recurrence rate of 18.1% (95% CI: 17.9 – 18.3). The range of recurrence rates in the cohorts was: 6 to 83% (29). Recurrence rates of individual syndromes of HDP were not reported very consistently over studies. References of the 64 articles in this meta-analysis are available as electronic supplementary material.

The baseline characteristics of the 99,415 women included in the IPD are shown in Table 2. The occurrence of major maternal complications (like pulmonary edema or maternal death) was recorded too sporadic to report on.

A hypertensive disorder of pregnancy (any type) recurred in 20,545 of 99,415 women (20.7% 95%CI: 20.4 – 20.9). The Odds Ratio between this recurrence risk and the recurrence risk calculated with the 64 included articles is: 1.18 (95%CI: 1.15-1.20, $p < .001$). Recurrence manifested as PE in 13,725 women ($N = 99,208$, 13.8%, 95%CI: 13.6 – 14.1) and GH in 6797 women ($N = 79,169$, 8.6% 95%CI: 8.4 – 8.8). HELLP complicated the HDP in 79 women ($N = 39,301$, 0.2%, 95%CI: 0.16 – 0.25). The delivery of a SGA child accompanied a hypertensive disorder in 1156 women ($N = 34,359$, 3.4%, 95%CI: 3.2 – 3.6). If we include non-hypertensive SGA as recurrence, SGA occurred in 4183 subsequent pregnancies ($N = 34,359$, 12%, 95%CI: 11.8 – 12.5). The different numbers of women in which the specific hypertensive syndrome was recorded (N) cause these percentages to not add up to the 20.7% overall recurrence. Premature delivery accompanying the recurrent hypertensive disorder occurred < 37 weeks in 3316 women ($N = 99,415$, 3.3% 95%CI:

3.2 – 3.5), < 34 weeks in 1224 women (N= 99,415, 1.2% 95%CI: 1.2 – 1.3) and < 28 weeks in 179 (N= 99,415, 0.18% 95%CI: 0.16 – 0.22). Figure 3 gives a more detailed overview of recurrence of the separate syndromes of hypertensive disorders and effects of prematurity on recurrence according to hypertensive syndrome and gestational age at onset at the index pregnancy.

Sensitivity analyses showed comparable recurrence rates when studies with unclear or high risk of bias were excluded from analysis and showed the same range of recurrence rates when discriminating studies by size and study design (data not shown).

Subgroup analysis

We compared recurrence amongst multiple and singleton pregnancies. Of the index pregnancies, 516 were multiple and 98.553 were singleton pregnancies. Recurrence occurred in 56 (10.9%) and 20.408 (20.7%), respectively (OR 0.53; 95% CI 0.40-0.70, p-value: <.001). Furthermore, we performed a subgroup analysis based on the presence of thrombophilia and the association between LMW-heparin use and recurrence. The combined database included 206 women with and 296 without thrombophilia. A statistically significant interaction was present between thrombophilia and LMW-heparin use (p-value: 0.005). Stratified analysis in those with thrombophilia showed that 26 of 56 women (46%) who used prophylactic LMW-heparin experienced recurrence of a hypertensive disorder of pregnancy, compared to 56 of 141 women (39.7%) who did not use LMW-heparin (OR 0.95; 95% CI 0.46-2.0, p-value: 0.89). In those without thrombophilia 8 of 52 (15.4%) that used

prophylactic LMW-heparin experienced recurrence, compared to 104 out of 244 (42.6%) that did not use LMW-heparin (OR 0.44; 95% CI 0.13-1.5, p-value: 0.20)

The clinical syndrome in the index and subsequent pregnancy

To assess differences between the clinical hypertensive syndrome in the index and subsequent pregnancy in women with recurrence, we compared the variables as shown in Table 3. Heterogeneity was high for intravenous anticonvulsive medication, caesarean section, and premature delivery <34 and <37 weeks' gestation, moderate for oral antihypertensive medication and premature delivery < 28 weeks' gestation, low for maximum diastolic blood pressure, small for gestational age < p10, and absent for the other outcomes. The maximum diastolic blood pressure was on average lower in the subsequent pregnancy, proteinuria > 300mg / 24hours occurred less often and oral antihypertensive or intravenous anticonvulsive medication was necessary less frequently. The subsequent pregnancy was complicated less often by a caesarean section, delivery of a small for gestational age child, premature delivery and perinatal mortality.

Similar to the sensitivity analyses for recurrence rate, the analyses showed comparable associations between the index and subsequent pregnancy after excluding studies with unclear or high risk of bias and showed the same range of associations when discriminating studies by size and study design (data not shown).

Chronic hypertension after pregnancy

Chronic hypertension occurring after pregnancy was reported in 5 studies (28,29,30,34,35). After exclusion of chronic hypertension before pregnancy, we included 581 women, who were normotensive before pregnancy, who experienced a

hypertensive disease of pregnancy and in whom chronic hypertension after pregnancy was registered. Of these, 236 women (41%) had recurrence of a hypertensive disorder in the next pregnancy. Women with recurrence developed chronic hypertension more often than women without recurrence (28.4 versus 9.6%, OR 4.2 95% CI: 2.6 – 6.7).

COMMENT:

This IPD meta-analysis explored the recurrence risk and predictors for recurrence of hypertensive disorders of pregnancy. The creation of this large combined database produced an opportunity to calculate an overall recurrence rate of HDP and to explore the role of individual risk factors. Some of the bias introduced in single cohort studies based on patient selection criteria and methodology could be reduced. The diversity of inclusion criteria of the included databases reflects the diverse world population.

Main findings

The recurrence rate of a hypertensive disorder of pregnancy (any type) is 20.7% (95%CI: 20.4 – 20.9). Concomitant HELLP syndrome or delivery of a SGA child increases recurrence of HDP. Also, decreasing gestational age at delivery in the index pregnancy, both the chance of having recurrence and the chance to deliver prematurely again increase. Multiple pregnancy at the index pregnancy, as a risk factor for the occurrence of HDP, is protective for recurrence. Use of LMW heparin was not protective for recurrence in our data, but the numbers are too small to draw conclusions. If the hypertensive disorder recurred, it was in general milder, regarding

most of the investigated variables. Women who were normotensive before pregnancy and who experienced recurrence of a hypertensive disorder, had a four times increased risk of developing chronic hypertension after pregnancy.

Strengths and weaknesses

Undertaking this study was not without challenges. Study selection was impaired and prolonged by difficulties in contacting authors of the original articles. Eligible studies date back to the eighties, causing data to be lost, authors to be retired and accurate contact information to be absent. We were able to include 22 of the 88 (25%) eligible articles. Unfortunately, several important studies were not included. The inclusion rate is adequate compared to other published IPD meta-analyses with inclusion rates of 12-24% (41,42), one study included 6 of 10 eligible studies, adding up to an inclusion rate of 60% (42).

The number of IPD in our study on the other hand, is much higher than in other IPDs, mostly containing several thousands of participants (40,41,42). The reason for this is our inclusion of nationwide registry based studies. Our combined database includes 99,415 women with a subsequent pregnancy, which is 69% of the 143,659 women that could be included in our regular meta-analysis of the 88 eligible studies. In comparison with the meta-analysis of these 88 studies, the recurrence rates differ statistically: 20.7 versus 18.1%, OR: 1.18 (95%CI: 1.15-1.20, $p < .001$). This suggests that some bias has occurred in the inclusion of data in our IPD. Thus, this IPD database may not be a completely representative cohort of the originally eligible studies, but it is close.

Another limitation of this study is the fact that we merged data that were obtained from very different study designs, settings and populations. This accounts

for the enormous range of recurrence rates between 6 and 83%. The upper bound of this range originates from a study with a very small and extreme high-risk population (29). It is unsurprising that heterogeneity across studies, assessed using the I^2 measure, was high for some of our results. The missing data that inevitably originate from merging databases challenged the statistical analysis. One of the methods that we used to overcome the problem of missing data is the merging of information on BMI, smoking, medical history and chronic hypertension between the two consecutive pregnancies. Although these characteristics can vary, we do think that on average, it gives a good indication of risk profile for the individual woman.

Furthermore, we included prophylactic trials in this IPD meta-analysis: 7 randomized controlled trials, which investigated the effect of aspirin (17,19,24) versus placebo, vitamins versus placebo (23,36), LMW-heparin plus aspirin versus aspirin alone (27) or LMW-heparin in thrombophilic women versus no treatment in non-thrombophilic women (37). The inclusion of these studies might be problematic when the treatments investigated in the different studies were effective in the reduction of hypertensive disorder recurrence. Two trials found a significant treatment effect. One trial found a significant reduction of recurrence in those treated with LMW-heparin plus aspirin compared with aspirin alone (27), but the numbers were very small (1 and 7 women experienced preeclampsia in the next pregnancy). The same applies for a trial reporting a significant effect of vitamin use compared to placebo (23). In addition, another vitamins trial shows conflicting results (36). Given the small sample sizes of the trials with a significant treatment effect and given these unclear relations to the recurrence of HDP, we do not expect that the inclusion of these studies hampers the generalizability of the results of this IPD meta-analysis in any way.

Variables that were infrequently registered, limited the analyses of the data. The percentages shown in Table 2 could therefore be difficult to interpret. HELLP syndrome was only recorded present in 512 women (0.5% of all included women) and SGA accompanied a HDP in 6448 cases (7%). This is unfortunate, as they are parts of the placental syndrome. The same applies for the inclusion of multiple pregnancy in the index pregnancy of 0.5%, being an exclusion criterion in some included studies. The presence of thrombophilia also requires an explanation. If registered, thrombophilia was present in 41%. The inclusion of studies with thrombophilia as one of the inclusion criteria resulted in this high percentage. We do not think this will act as bias, since it is only documented in 0.5% of the total cohort.

Another problem is reporting bias. Eclampsia was reported quite often (8% in 26.665 women in whom this complication was registered) in the index pregnancy. It is understandable that if eclampsia occurred, it is more likely to be registered than when it did not occur.

Sensitivity analysis in which we discriminated studies by study design and size showed some discordant results. This was mainly due to loss of power: the analyses of small studies and prospective studies separately comprised 972 and 1955 of the 915.415 inclusions respectively (1 and 2%). In contrast, the overall analysis was mainly dominated by three large retrospective studies (31,32,38). Also, the results were difficult to interpret as most prospective studies did not register clinical details of the index pregnancy. The sensitivity analysis that included only low risk of bias studies had results that were comparable to the overall results, mainly because only 3 small studies were excluded from analysis (26,27,29).

Counseling

Counseling couples about the chance of recurrence is important. This IPD meta-analysis did not include women, who refrained from a subsequent pregnancy. Three included databases contained information on the reason for not engaging in a subsequent pregnancy (30,34,35). Of 471 women, 140 (29%) refrained from a next pregnancy due to high perceived risk. This does not comply with the 21% recurrence risk and the fact that HDP recur in a milder form. The knowledge from this IPD meta-analysis can be used in counseling in the future.

In conclusion, interpretation of previous data of individual cohort studies was hampered by many sources of bias. IPD meta-analysis is a methodologically and logistically challenging approach. However, despite all the challenges and limitations stated above, the results are based on the largest database regarding recurrence of HDP so far. The opportunities of aggregated datasets are of paramount importance, because they allow the calculation of overall recurrence rates and also identify more accurately the role of individual risk factors on an individual level. The present IPD meta-analysis helped to create knowledge that can comprehensively be included in the counseling of couples after experiencing hypertensive disease of pregnancy.

REFERENCES:

1. Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. *Lancet* 2006;367:1066-74.
2. Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet* 2005;365:785-799.
3. Hauth JC, Ewell MG, Levine RL, Esterlitz JR, Sibai BM, Curet LB. Pregnancy outcomes in healthy nulliparous women who subsequently developed hypertension. *Obstet Gynecol* 2000; 95:24–28.
4. Steegers EA, von Dadelszen P, Duvekot JJ, Pijnenborg R. Pre-eclampsia. *Lancet* 2010;376:631-44.
5. Rep A, Ganzevoort W, Bonsel GJ, Wolf H, de Vries JI. Psychosocial impact of early-onset hypertensive disorders and related complications in pregnancy. *Am J Obstet Gynecol* 2007;197(2):158e1-6.
6. Sibai BM, Mercer B, Sarinoglu C. Severe preeclampsia in the second trimester: recurrence risk and long-term prognosis. *Am J Obstet Gynecol* 1991;165:1408–12.
7. Sep S, Smits L, Prins M, Peeters L. Prediction tests for recurrent hypertensive disease in pregnancy, a systematic review. *Hypertens Pregnancy* 2010;29(2):206-30.

8. Sibai BM, Ramadan MK, Chari RS, Friedman SA Pregnancies Complicated by HELLP Syndrome (Hemolysis, Elevated Liver Enzymes, and Low Platelets): Subsequent Pregnancy Outcome and Long-Term Prognosis. *Am J Obstet Gynecol* 1995;172:125-129.
9. van Pampus MG, Wolf H, Mayruhu G, Treffers PE, Bleker OP. Long-term follow-up in patients with a history of (H)ELLP syndrome. *Hypertens Pregnancy* 2001;20(1):15-23.
10. Ananth CV, Kaminsky L, Getahun D, Kirby RS, Vintzileos AM. Recurrence of fetal growth restriction in singleton and twin gestations. *Matern Fetal Neonatal Med* 2009;22:654–61.
11. Broeze KA, Opmeer BC, Bachmann LM, Broekmans FJ, Bossuyt PM, Coppus SF. Individual patient data meta-analysis of diagnostic and prognostic studies in obstetrics, gynaecology and reproductive medicine. *BMC Med Res Methodol* 2009;27:9-22.
12. Wells G, Shea B, O'connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. Ottawa Health Research Institute.
http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
13. ACOG practice bulletin. Diagnosis and management of preeclampsia and eclampsia. Number 33, January 2002. American College of Obstetricians and Gynecologists. *Int Gynaecol Obstet* 2002;77(1):67-75.

14. Sibai BM. The HELLP syndrome (hemolysis, elevated liver enzymes, and low platelets): much ado about nothing? *Am J Obstet Gynecol* 1990;162(2):311-6.
15. ACOG practice bulletin. Intrauterine growth restriction. Number 12, Januari 2000. American College of Obstetricians and Gynecologists. *Int Gynaecol Obstet* 2001;72(1):85-96.
16. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003; 327(7414):557-560.
17. Beroyz G, Casale R, Farreiros A, et al. CLASP: A randomised trial of low-dose aspirin for the prevention and treatment of pre-eclampsia among 9364 pregnant women. *Lancet* 1994;343:619-29.
18. Brown MA, Mackenzie C, Dunsmuir W, et al. Can we predict recurrence of pre-eclampsia or gestational hypertension? *BJOG* 2007;114:984-93.
19. Byaruhanga RN, Chipato T, Rusakaniko S. A randomized controlled trial of low-dose aspirin in women at risk from pre-eclampsia. *Int J Gynaecol Obstet* 1998;60:129-35.
20. Cameroni I, Crippa I, Roncaglia N, Locatelli A, Ornaghi S, Vergani P. Uterine artery Doppler in a risk population: What's its role in the prediction of small for gestational age fetuses? *Pregnancy Hypertens* 2011;1:260.

21. Campbell DM, MacGillivray I, Carr-Hill R. Pre-eclampsia in second pregnancy. *Br J Obstet Gynaecol* 1985;92:131-40.
22. Campbell D, Bhattacharya S, Lemon J. Recurrent preeclampsia - Once/twice? *Pregnancy Hypertens* 2010;1:S30.
23. Chappell LC, Seed PT, Briley AL, et al. Effect of antioxidants on the occurrence of pre-eclampsia in women at increased risk: A randomised trial. *Lancet* 1999;354:810-16.
24. Chiaffarino F, Parazzini F, Paladini D, et al. A small randomised trial of low-dose aspirin in women at high risk of pre-eclampsia. *Eur J Obstet Gynecol Reprod Biol* 2004;112:142-44.
25. Conserva V, Muggiasca M, Arrigoni L, Mantegazza V, Rossi E, Ferrazzi E. Recurrence and severity of abnormal pregnancy outcome in patients treated by low-molecular-weight heparin: A prospective pilot study. *J Matern Fetal Neonatal Med* 2012;25:1467-73.
26. Facchinetti F, Marozio L, Frusca T, et al. Maternal thrombophilia and the risk of recurrence of preeclampsia. *Am J Obstet Gynecol* 2009;200:46-5.
27. Ferrazzani S, D'Alessio MC, Fatigante G, G et al. Prophylaxis of recurrent preeclampsia: low-molecular-weight heparin plus low-dose aspirin versus low-dose aspirin alone. *Hypertens Pregnancy* 2006;25:115-27.

28. Figueiro-Filho EA, de Oliveira VM, Coelho LR, Breda I. Serum markers of inherited thrombophilia and antiphospholipid antibodies in pregnant women with previous history of severe pre-eclampsia. *Rev Bras Ginecol Obstet* 2012;34:40-46. Portuguese.
29. Gaugler-Senden IP, Berends AL, de Groot CJ, Steegers EAP. Severe, very early onset preeclampsia: subsequent pregnancies and future parental cardiovascular health. *Eur J Obstet Gynecol Reprod Biol* 2008;140:171-77.
30. Langenveld J, Buttinger A, van der Post J, Wolf H, Mol B, Ganzevoort W. Recurrence risk and prediction of a delivery under 34 weeks of gestation after a history of a severe hypertensive disorder. *BJOG* 2011;118(5):589-95.
31. Lykke JA, Paidas MJ, Langhoff-Roos J. Recurring complications in second pregnancy. *Obstet Gynecol* 2009;113:1217-24.
32. Mbah AK, Sharma PP, Alio AP, Fombo DW, Bruder K, Salihu HM. Previous cesarean section, gestational age at first delivery and subsequent risk of pre-eclampsia in obese mothers. *Arch Gynecol Obstet* 2012;284:1375-81.
33. Napolitano R, Rajakulasingam R, Memmo A, Bhide A, Thilaganathan B. Uterine artery Doppler screening for pre-eclampsia: comparison of the lower, mean and higher first-trimester pulsatility indices. *Ultrasound Obstet Gynecol* 2011;37:534-537.

34. van Oostwaard MF, Langenveld J, Bijloo R, et al. Prediction of recurrence of hypertensive disorders of pregnancy between 34 and 37 weeks of gestation: a retrospective cohort study. *BJOG* 2012;119(7):840-7.
35. van Oostwaard MF, Langenveld J, Schuit E, et al. Prediction of recurrence of hypertensive disorders of pregnancy in the term period, a retrospective cohort study. *Preg Hyper: An Int J Women's Card Health* 2014;4:194–202.
36. Poston L, Briley A, Seed P, Kelly F, Shennan A. Vitamin C and vitamin E in pregnant women at risk for pre-eclampsia (VIP trial): randomised placebo-controlled trial. *Lancet* 2006;367:1145-54.
37. Salim R, Czarnowicki T, Nachum Z, Shalev E. The impact of close surveillance on pregnancy outcome among women with a prior history of antepartum complications attributed to thrombosis: A cohort study. *Reprod Biol Endocrinol* 2008;6:55.
38. Trogstad L, Skrondal A, Stoltenberg C, Magnus P, Nesheim BI, Eskild A. Recurrence risk of preeclampsia in twin and singleton pregnancies. *Am J Med Genet A* 2004;126A(1):41-5.
39. Zhang J, Troendle JF, Levine RJ. Risks of hypertensive disorders in the second pregnancy. *Paediatr Perinat Epidemiol* 2001;15:226-31.

40. Broer SL, van Disseldorp J, Broeze KA, et al. Added value of ovarian reserve testing on patient characteristics in the prediction of ovarian response and ongoing pregnancy: an individual patient data approach. *Hum Reprod* 2013;1:26–36.
41. Broeze KA, Opmeer BC, Coppus SFPJ, et al. Chlamydia antibody testing and diagnosing tubal pathology in subfertile women: an individual patient data meta-analysis. *Hum Reprod Update* 2011;17(3):301–310.
42. Groeneveld E, Broeze KA, Lambers MJ, et al. Is aspirin effective in women undergoing in vitro fertilization (IVF)? Results from an individual patient data meta-analysis (IPD MA). *Hum Reprod Update* 2011;17(4):501–509.

Tables and figures:

Figure 1:
Inclusion of articles

Figure 2:
Risk of bias graph

Quality characteristics of all included studies, evaluated with the Newcastle Ottawa Scale for cohort studies. The item “selection of the on exposed cohort” was not relevant in this IPD and was therefore indicated as ‘low risk’ in all studies.

Figure 3A. Recurrence rates of the individual hypertensive syndromes of the 99,415 included patients according to the hypertensive syndrome at the index pregnancy

Figure 3B. Recurrence rates of the individual hypertensive syndromes of the 99,415 included patients according to the gestational age at onset (or diagnosis) of the hypertensive syndrome at the index pregnancy

Table 1:
Overview of study characteristics of the included studies

Table 2:
Baseline clinical characteristics and index pregnancy parameters of the 99,415 included patients

Table 3:
Differences in the clinical hypertensive syndrome between the index and subsequent pregnancy in recurrent disease

Electronic supplementary material:
References of the 60 articles included in the meta-analysis preceding the IPD meta-analyses

Table 1- Overview of study characteristics of the included studies

Study and Year	Country	Inclusion criteria	Exclusion criteria	Study design	Therapeutic trial	Total women original study	Total includable women* (recurrence rate)
Beroyz 1994 ¹⁷	United Kingdom	- Former PE - Former FGR - Chronic hypertension - Renal disease - Signs of PE or FGR in current pregnancy	- Bleeding disorders - Asthma - Allergy to aspirin	Randomized controlled trial	Aspirin vs placebo	9364	156 (59%)
Brown 2007 ¹⁸	Australia	- Former PE - Former GH - Chronic hypertension - Essential hypertension and superimposed PE	- None reported	Retrospective cohort study	-	1354	765 (32%)
Byaruhanga 1998 ¹⁹	South-Africa	- Former PE - Former GH - Chronic hypertension	- History of hypersensitivity to aspirin, peptic ulcer, bleeding disorders or chronic pulmonary disease - Use of NSAID's - Development of PE prior to trial entry	Randomized controlled trial	Aspirin vs placebo	250	213 (15%)
Cameroni 2011 ²⁰	Italy	- Former PE - Former FGR - Former placental abruption - Former stillbirth - Chronic hypertension	-	Retrospective cohort study		218	173 (10%)
Campbell 1985 and 2010 ^{21,22}	United Kingdom	- Total population of Aberdeen Maternity and Neonatal Data Bank	-	Retrospective cohort study	-	38130	7,725 Extended extraction** (26%)
Chappell 1999 ²³	United Kingdom	- Former PE < 37wks - Former HELLP or eclampsia at any GA - Abnormal uterine-artery doppler waveform in pregnancy	- Heparin or warfarin treatment - Fetal abnormalities - Multiple pregnancy	Randomized controlled trial	Antioxidants (Vit C and E) vs placebo	283	56 (50%)
Chiaffarino 2004 ²⁴	Italy	- Former severe PE or eclampsia - Former FGR - Former intrauterine foetal death	- Chronic disease other than hypertension, renal disease or diabetes - Allergy to aspirin - Fetal malformations - current twin pregnancy	Randomized controlled trial	Aspirin vs no treatment	40	15 (47%)
Conserva 2012 ²⁵	Italy	- Former PE - Former HELLP - Former GH - Former FGR - Former placental abruption - Former FGR and stillbirth	- acquired thrombophilia - clinical immune disease - treatment with LMWH in previous pregnancy - congenital fetal anomaly - non Caucasian ethnicity	Prophylactic trial	Enoxaparine	128	53 (11%)

Facchinetti 2009 ²⁶	Italy	<ul style="list-style-type: none"> -Former PE in singleton pregnancy -A complete evaluation for thrombophilia. -Current singleton pregnancy 	<ul style="list-style-type: none"> -History of thromboembolic diseases -Renal and/or cardiovascular disorder -Systemic lupus erythematosus -Diabetes -Any ethnic group other than white 	Prospective cohort study	-	172	172 (34%)
Ferrazzani 2006 ²⁷	Italy	<ul style="list-style-type: none"> -Previous severe preterm PE with associated FGR 	<ul style="list-style-type: none"> -Former HELLP syndrome 	Prophylactic trial	Aspirin vs aspirin and heparin	68	54 (15%)
Figueiro-Filho 2012 ²⁸	Brasil	<ul style="list-style-type: none"> -Previous severe PE with one of the following: -hospitalization < 32wks -imminent eclampsia -eclampsia -HELLP syndrome -systemic laboratory tests -preterm birth < 34 wks -admission of newborns in NICU -fetal loss -fetal growth restriction -oligohydramnia -abnormal uterine or umbilical artery doppler 	<ul style="list-style-type: none"> -Chronic hypertension -Systemic lupus erythematosus -Thrombophilia 	Prospective cohort study	-	113	67 (50%)
Gaugler-Senden 2008 ²⁹	The Netherlands	<ul style="list-style-type: none"> -Former severe PE < 24wks 	-	Retrospective cohort study	-	20	18 (83%)
Langenveld 2011 ³⁰	The Netherlands	<ul style="list-style-type: none"> -Former PE, GH or HELLP < 34wks in singleton pregnancy 	<ul style="list-style-type: none"> -Fetal abnormalities 	Retrospective cohort study	-	380	211 (55%)
Lykke 2009 ³¹	Denmark	<ul style="list-style-type: none"> -All women with two singleton deliveries in the National Patient Registry 	<ul style="list-style-type: none"> -cardiovascular diagnosis -diabetes -women who died or emigrated within 3 months of the second delivery 	Retrospective cohort study	-	536,419	26,939 (20%)
Mbah 2012 ³²	United States of America	<ul style="list-style-type: none"> -All women with two pregnancies registered in the Missouri maternally linked cohort database 1989 – 2005 	-	Retrospective cohort study	-	166712	23,390 (17%)
Napolitano 2011 ³³	United Kingdom	<ul style="list-style-type: none"> -all nulliparous women -parous women with former PE or FGR -concurrent maternal medical conditions ->7 years since last pregnancy 	<ul style="list-style-type: none"> -Miscarriage < 14 wks -Fetal chromosomal or structural abnormalities 	Prospective cohort study	-	6221	273 (26%)
van Oostwaard 2012 ³⁴	The Netherlands	<ul style="list-style-type: none"> -Former PE, GH, HELLP or FGR 34 - 37 wks 	<ul style="list-style-type: none"> -Fetal abnormalities 	Retrospective cohort study	-	425	189 (34%)

van Oostwaard 2014 ³⁵	The Netherlands	- Former PE, GH or HELLP > 37 wks	- Fetal abnormalities	Retrospective cohort study	-	638	312 (41%)
Poston 2006 ³⁶	United Kingdom	- Former PE < 37wks - Former HELLP or eclampsia at any GA - Other risk factors for hypertensive disease (Essential hypertension, Diabetes, renal disease, Antiphospholipid syndrome, abnormal uterine- artery doppler waveform in pregnancy or BMI>30)	- No informed consent - Warfarin treatment - Using vitamins before trial	Randomized controlled trial	Vitamins (Vit C and E) vs placebo	2404	556 (29%)
Salim 2008 ³⁷	Israel	- IUFD - SGA - severe PE or placental abruption - in any former pregnancy > 23wks	Former pregnancy with: - multiple gestation - major congenital or chromosomal anomalies - fetal infection / chorioamnionitis - hydrops - diabetes mellitus	Prospective cohort study	- Close surveillance - LMW heparin if thrombophilia - Aspirin added if antiphospholipid antibodies	97	19 (16%)
Trogstad 2004 ³⁸	Norway	- All women with two singleton deliveries in the Medical Birth Registry - And preeclampsia in the first pregnancy	- Multiple gestation (triplet or more) in the first pregnancy - Multiple gestation in the subsequent pregnancy (2 or more)	Retrospective cohort study	-	20,285	37,738 Extended extraction** (22%)
Zhang 2001 ³⁹	United States of America	- Women attending the prenatal care unit during inclusion period - And two consecutive pregnancies	-	Prospective cohort study	-	1641	321 (24%)

Total:

99,415

* Includable women: women with hypertensive disease (PE, GH, HELLP) in a former pregnancy, with a subsequent pregnancy, excluding control groups and cases without hypertensive complications in former pregnancies.

** New extended data extraction from the registry on behalf of this IPD, using the same inclusion criteria from the original cohort

Table 2. Baseline clinical characteristics and index pregnancy parameters of the 99,415 included patients.

	N*		% of total 99,415	References
Age at index pregnancy (years)	97,832	25 (SD: 5)		17 18 21 22 24 27 29 30 31 32 34 35 38 39
Smoking	27,304	5654 (21%)	6%	17 18 21 22 26 29 30 31 34 35 36 39
Ethnicity	- European	20,785 (81%)	21%	
	- Caribbean	113 (0.4%)	0.1%	
	- Asian	25,807 228 (0.9%)	0.2%	17 20 23 25 26 27 28 29 30 32
	- Sub-Saharan Africa	4545 (5%)	5%	33 34 35 36 39
	- Middle East	19 (0.1%)	0%	
Body mass index, BMI (kg/m ²)	32,544	25 (IQR: 22 - 29)		21 22 23 25 29 30 32 33 34 35 36 39
Chronic hypertension before pregnancy	26,879	2032 (8%)	2.0%	17 20 23 25 26 27 28 29 30 32 34 35 36 38 39
Thrombophilia	502	206 (41%)	0.2%	25 26 28 30 34 35 37
History of disease	- Diabetes Mellitus	90,749 1342 (1.5%)	1.3%	17 18 20 23 24 26
	- Coronary disease	51,387 167 (0.3%)	0.2%	27 28 29 30 32 33
	- Kidney disease	25,004 121 (0.5%)	0.1%	34 35 36 37 38 39
Pregnancy characteristics of index pregnancy				
Nulliparous	72,412	65,243 (90%)	66%	17 18 19 20 21 22 23 24 25 26 27 28 29 30 32 33 34 35 36 37 38 39
Multiple pregnancy	99,069	516 (0.5%)	0.5%	17 18 19 20 21 22 23 24 25 26 28 29 30 32 34 35 36 38 39
Gestational hypertension [†]	99,400	23,970 (24%)	24%	17 18 19 20 21 22 23 24 25 26 27 28 28 30 31 32 33 34 35 36 37 38 39 (all)
Preeclampsia [†]	99,202	75,172 (76%)	76%	17 18 19 20 21 22 23 24 25 26 27 28 28 30 31 32 33 34 35 36 37 38 39 (all)
Eclampsia [†]	26,665	2087 (8%)	2.1%	17 18 25 28 29 30 32 34 35 38
HELLP syndrome [†]	40,236	512 (1.3%)	0.5%	17 18 20 25 27 28 29 30 34 35 36 38
Placenta Abruption	51,803	1221 (2.4%)	1.2%	20 25 28 30 31 32 34 35 37 39
Maximum blood pressure (mmHg)	- Systolic	632 161 (SD: 21)	0.2%	
	- Diastolic	1028 103 (SD: 11)	0.1%	29 30 34 35 39
Use of medication	- oral antihypertensive	1446 738 (51%)	0.7%	
	- iv antihypertensive	687 141 (21%)	0.1%	18 29 30 34 35
	- iv anticonvulsive	1472 219 (15%)	0.2%	
Gestational age at delivery (weeks)	94,178	39 (SD: 20 days)		18 21 22 25 26 27 29 30 31 32 34 35 37 38 39

Birth weight (grams)		97,694	3185 (SD: 761)		18 21 22 25 26 27 29 30 31 32 34 35 37 38 39
Small for Gestational Age [†]		35,109	6448 (18%)	6.4%	17 18 20 21 22 23 24 25 27 28 29 30 32 33 34 35 37 38 39
Premature delivery	< 28 weeks	94,197	739 (0.8%)	0.7%	17 18 21 22 23 25
	< 34 weeks	94,353	5363 (5.7%)	5.4%	26 27 29 30 31 32
	< 37 weeks	94,965	14521 (15%)	15%	34 35 37 38 39
Caesarean Section		93,948	28,081 (30%)	28%	18 21 22 25 27 28 30 31 32 34 35 37 38
NICU admission neonate		5117	1157 (22%)	1.2%	18 38
Perinatal mortality		98,078	1608 (1.6%)	1.6%	17 18 20 21 22 24 25 27 29 30 31 32 34 35 37 38 39
Characteristics at the subsequent pregnancy					
Birth interval (months)		59,754	41 (SD: 25)		18 21 22 29 30 31 32 34 35 39
Change of partner		7344	668 (9%)	0.7%	20 21 22 23 35 39
Use of prophylaxis	- Aspirin	5663	737 (13%)	0.7%	17 19 20 21 22
	- LMW Heparin	1962	153 (8%)	0.2%	23 24 26 27 28
	- Both	1909	60 (3%)	0.1%	30 34 35 36 37

Continuous data are presented as means (SD); age, BMI and gestational age in median (IQR)

Proportions are shown as n (% of N)

* Number of women with available information

† Percentages sum up to more than 100% because of overlapping of disorders

Table 3. Differences in the clinical hypertensive syndrome between the index and subsequent pregnancy in recurrent disease


<i>Women with recurrence</i> <i>N= 20,545</i>	N*	Index Pregnancy	Subsequent pregnancy	P	I ² (%)	References	
Max systolic BP (mmHg)**	194	165 (19)	153 (17)	0.055	0	17 19 27 29 30 34 35 36	
Max diastolic BP (mmHg)**	331	107 (11)	100 (8.9)	<.001	25	29 30 34 35 39	
Proteinuria > 300mg/24h	119	82 (69%)	58 (49%)	0.011	0	18 29 27 30 34 35	
Thrombocytopenia < 100 *10 ⁹ /L	162	37 (23%)	18 (11%)	0.24	0	18 29 30 34 35	
Use of medication	- oral antihypertensive	504	295 (59%)	237 (47%)	0.002	71	17 18 19 23 25 28 29 30 34 35 36
	- iv antihypertensive	263	67 (26%)	27 (10%)	0.12	0	
	- iv anticonvulsive	508	82 (16%)	31 (6%)	<.001	72	
Hospital days***	154	5 (2-11)	3 (1-5)	0.17	0	34 35	
Caesarean section	18,488	6195 (34%)	6423 (35%)	<.001	99	17 18 19 20 21 22 25 26 28 29 30 31 32 34 35 37 38	
Small for gestational age < p10	6,542	996 (15%)	841 (13%)	<.001	45	17 18 20 21 22 23 25 27 28 29 30 32 33 34 35 37 38 39	
Perinatal mortality	20,111	466 (2.3%)	256 (1.3%)	<.001	0	17 18 20 21 22 24 25 27 29 30 31 32 33 34 35 37 38 39	
Premature delivery	< 28 weeks	18,638	267 (1.4%)	155 (0.8%)	<.001	54	18 21 22 25 26 27 29 30 31 32 34 35 37 38 39
	< 34 weeks	18,735	1868 (10%)	1106 (5.9%)	<.001	76	
	< 37 weeks	18,925	4312 (23%)	3125 (17%)	<.001	93	

* Number of women with available information for both of the pregnancies

** Data are presented as means (SD).

***Data are presented as median (IQR).

Significant differences are indicated in bold

Figure 1. Inclusion of articles






Figure 3A. Recurrence rates of the individual hypertensive syndromes of the 99,415 included patients according to the hypertensive syndrome at the index pregnancy



Data are shown as frequencies with percentages to the number of cases in which both variables were registered. Number of women with specific hypertensive disorder at the index pregnancy: Any type 99.241; PE 75.172; GH 23.970; HELLP 512; SGA 6448

Figure 3B. Recurrence rates of the individual hypertensive syndromes of the 99,415 included patients according to the gestational age at onset (or diagnosis) of the hypertensive syndrome at the index pregnancy



Data are shown as frequencies with percentages to the number of cases in which both variables were registered. Number of women with specific Gestational Age at the index pregnancy: Any GA 99.415; <37 wks 14.512; <34 wks 5354; <28 wks 730

Electronic supplementary material: References of the 60 articles included in the meta-analysis preceding the IPD meta-analyses.

Study	Recurrence
Aardenburg R, Spaanderman ME, van Eijndhoven HW, de Leeuw PW, Peeters LL. A Low Plasma Volume in Formerly Preeclamptic Women Predisposes to the Recurrence of Hypertensive Complications in the Next Pregnancy. <i>J Soc Gynecol Invest</i> 2006;13:598-603.	50.00
Ananth CV, Peltier MR, Chavez MR, Kirby RS, Getahun D, Vintzileos AM. Recurrence of ischemic placental disease. <i>Obstet Gynecol</i> 2007;110:128-33.	16.49
Beroz G, Casale R, Farreiros A, Palermo M, Margulies M, Voto L et al. CLASP: A randomised trial of low-dose aspirin for the prevention and treatment of pre-eclampsia among 9364 pregnant women. <i>Lancet</i> 1994;343:619-29.*	59.18
Brown MA, Mackenzie C, Dunsmuir W, Roberts L, Ikin K, Matthews J, Mangos G, Davis G. Can we predict recurrence of pre-eclampsia or gestational hypertension? <i>BJOG</i> 2007;114:984-93.*	31.90
Byaruhanga RN, Chipato T, Rusakaniko S. A randomized controlled trial of low-dose aspirin in women at risk from pre-eclampsia. <i>Int J Gynaecol Obstet</i> 1998;60:129-35.*	14.55
Cameroni I, Crippa I, Roncaglia N, Locatelli A, Ornaghi S, Vergani P. Uterine artery Doppler in a risk population: What's its role in the prediction of small for gestational age fetuses? <i>Pregnancy Hypertens</i> 2011;1:260.*	9.83
Campbell D, Bhattacharya S, Lemon J. Recurrent preeclampsia - Once/twice? <i>Pregnancy Hypertens</i> 2010;1:S30.*	25.61
Caritis S, Sibai B, Hauth J, Lindheimer MD, Klebanoff M, Thom E et al. Low-dose aspirin to prevent preeclampsia in women at high risk. <i>New Engl J Med</i> 1998;338:701-05.	6.00
Cathelain-Soland S, Coulon C, Subtil D, Houfflin-Debarge V, Deruelle P. Subsequent pregnancy outcome in women with a history of preeclampsia and/or HELLP syndrome. <i>Gynecol Obstet Fertil</i> 2010;38:166-72.	27.46
Chames MC, Haddad B, Barton JR, Livingston JC, Sibai BM, Suarez et al. Subsequent pregnancy outcome in women with a history of HELLP syndrome at (less-than or equal to) 28 weeks of gestation. <i>Am J Obstet Gynecol</i> 2003;188:1504-08.	54.84
Chappell LC, Seed PT, Briley AL, Kelly FJ, Lee R, Hunt BJ, Pamar K, Bewley SJ, Shennan AH, Steer PJ Poston L. Effect of antioxidants on the occurrence of pre-eclampsia in women at increased risk: A randomised trial. <i>Lancet</i> 1999;354:810-16.*	50.00
Chiaffarino F, Parazzini F, Paladini D, Acaia B, Ossola W, Marozio L, Facchinetti F, Del Giudice A. A small randomised trial of low-dose aspirin in women at high risk of pre-eclampsia. <i>Eur J Obstet Gynecol Reprod Biol</i> 2004;112:142-44.*	47.06
Conserva V, Muggiasca M, Arrigoni L, Mantegazza V, Rossi E, Ferrazzi E. Recurrence and severity of abnormal pregnancy outcome in patients treated by low-molecular-weight heparin: A prospective pilot study. <i>J Matern Fetal Neonatal Med</i> 2012;25:1467-73.*	11.32
de Vries JI, van Pampus MG, Hague WM, Bezemer PD, Joosten JH; For FRUIT investigators. Low-Molecular-Weight Heparin Added to Aspirin in the Prevention of Recurrent Early-Onset Preeclampsia in women with Inheritable Thrombophilia: the FRUIT-RCT. <i>J Thromb Haemost</i> 2012;10:64-72.	26.17
Dukler D, Porath A, Bashiri A, Erez O, Mazor M. Remote prognosis of primiparous women with preeclampsia. <i>Eur J Obstet Gynecol Reprod Biol</i> 2001;96:69-74.	25.79
Facchinetti F, Marozio L, Frusca T, Grandone E, - Venturini P - Tiscia GL, Zatti S, Benedetto C. Maternal thrombophilia and the risk of recurrence of preeclampsia. <i>Am J Obstet Gynecol</i> 2009;200:46-5.*	33.72
Ferrazzani S, D'Alessio MC, Fatigante G, Soreca G, De Carolis S, Paradisi G, Caruso A. Prophylaxis of recurrent preeclampsia: lowmolecularweight heparin plus lowdose aspirin versus lowdose aspirin alone. <i>Hypertens Pregnancy</i> 2006;25:115-27.*	14.81
Figueiro-Filho EA, de Oliveira VM, Coelho LR, Breda I. Serum markers of inherited thrombophilia and antiphospholipid antibodies in pregnant women with previous history of severe pre-eclampsia. <i>Rev Bras Ginecol Obstet</i> 2012;34:40-46.*	50.00
Gainger S, Saha SC, Dhaliwal L, Bagga R. Pregnancy outcome in subsequent pregnancies after eclampsia. <i>Pregnancy Hypertens</i> 2012;2:175.	37.74
Gaugler-Senden IP, Berends AL, de Groot CJ, Steegers EAP. Severe, very early onset preeclampsia: subsequent pregnancies and future parental cardiovascular health. <i>Eur J Obstet Gynecol Reprod Biol</i> 2008;140:171-77.*	83.33

- Gris JC, Chauleur C, Mares P, Quere I, Lefrant JY, Haddad B et al. Enoxaparin for the secondary prevention of placental vascular complications in women with severe pre-eclampsia: The pilot randomised controlled NOH-PE study. *Thromb Haemost* 2011;106:1053-1061. 11.17
- Gudnasson HM, Dubiel M, Gudmundsson S. Preeclampsia Abnormal uterine artery Doppler is related to recurrence of symptoms during the next pregnancy. *J Perinat Med* 2004;32:400-03. 30.22
- Habli M, Eftekhari N, Wiebracht E, Bombrys A, Khabbaz M, How H, Sibai B et al. Long-term maternal and subsequent pregnancy outcomes 5 years after hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome. *Am J Obstet Gynecol* 2009;201:385. 52.83
- Hargood JLFA, Brown MA. Pregnancy-induced hypertension: recurrence rate in second pregnancies. *Med J Aust* 1991;154:376-77. 47.14
- Hernandez-Diaz S, Toh S, Cnattingius S. Risk of pre-eclampsia in first and subsequent pregnancies: prospective cohort study. *BMJ (Clinical research ed.)* 2009;338:b2255. 9.14
- Hjartardottir S, Leifsson BG, Geirsson RT, Steinthorsdottir V. Recurrence of hypertensive disorder in second pregnancy. *Am J Obstet Gynecol* 2006;194:916-20. 39.51
- Hupuczi P, Rigo B, Sziller I, Szabo G, Szigeti Z, Papp Z. Follow-up analysis of pregnancies complicated by HELLP syndrome. *Fetal Diagn Ther* 2006;21:519-22. 59.26
- Kalk JJ, Huisjes AJ, de Groot CJ, van Beek E, van Pampus MG, Spaanderman ME et al. Recurrence rate of pre-eclampsia in women with thrombophilia influenced by low-molecular-weight heparin treatment? *Neth J Med* 2004;62:83-87. 41.38
- Kim J, Kim YH, Cho MK, Kim CH, Song TB. The usefulness of gestation-corrected hyperuricemia as a predictor of the development of preeclampsia on subsequent pregnancy. *Pregnancy Hypertens* 2012;2:336. 32.76
- Kreuwel JH, Scholten RR, Peeters LL, Spaanderman ME. Increased inter-pregnancy calciuria relates to recurrent preeclampsia. *Reprod Sci* 2011;18:360A. 22.92
- Kupfermanc M, Rimon E, Many A, Maslovitz S, Lessing JB, Gamzu R. Low molecular weight heparin versus no treatment in women with previous severe pregnancy complications and placental findings without thrombophilia. *Blood Coagul Fibrinolysis* 2011;22:123-26. 28.13
- Langenveld J, Buttinger A, van der Post J, Wolf H, Mol B, Ganzevoort W. Recurrence risk and prediction of a delivery under 34 weeks of gestation after a history of a severe hypertensive disorder. *BJOG* 2011 Apr;118(5):589-95.* 54.81
- Leeners B, Neumaier-Wagner PM, Kuse S, Mutze S, Rudnik-Schoneborn S, Zerres K et al. Recurrence risks of hypertensive diseases in pregnancy after HELLP syndrome. *J Perinat Med* 2011;39:673-78. 43.24
- Lojacono A, Valcamonica A, Tanzi P, Soregaroli M, Frusca T. Clinical follow-up and screening for autoimmune disorders in patients with previous severe early-onset preeclampsia. *Ital J Gynaecol Obstet* 1996;8:51-54. 25.45
- Lykke JA, Paidas MJ, Langhoff-Roos J. Recurring complications in second pregnancy. *Obstet Gynecol* 2009;113:1217-24.* 20.01
- Martinelli I, Ruggerenti P, Cetin I, Pardi G, Perna A, Vergani P et al. Heparin in pregnant women with previous placenta-mediated pregnancy complications: A prospective, randomized, multicenter, controlled clinical trial. *Blood* 2012;119:3269-75. 16.00
- Mbah AK, Sharma PP, Alio AP, Fombo DW, Bruder K, Salihu HM. Previous cesarean section, gestational age at first delivery and subsequent risk of pre-eclampsia in obese mothers. *Arch Gynecol Obstet* 2012;284:1375-81.* 16.75
- McDonald SDFA, Best CFAU, Lam K. The recurrence risk of severe de novo pre-eclampsia in singleton pregnancies: a population-based cohort. *BJOG* 2009;116:1578-84. 6.81
- Mello G, Parretti E, Fatini C, Riviello C, Gensini F, Marchionni M, Scarselli GF et al. Low-molecular-weight heparin lowers the recurrence rate of preeclampsia and restores the physiological vascular changes in angiotensin-converting enzyme DD women. *Hypertension* 2005;45:86-91. 25.13
- Mello G, Parretti E, Cioni R, Lagozio C, Mealli F, Pratesi M. Individual longitudinal patterns in biochemical and hematological markers for the early prediction of pre-eclampsia. *J Matern Fetal Neonatal Med* 2002;11:93-99. 17.50
- Mostello D, Kallogjeri D, Tungsiripat R, Leet T. Recurrence of preeclampsia: effects of gestational age at delivery of the first pregnancy, body mass index, paternity, and interval between births. *Am J Obstet Gynecol* 2008;199:55-57. 14.70
- Napolitano R, Rajakulasingam R, Memmo A, Bhide A, Thilaganathan B. Uterine artery Doppler screening for pre-eclampsia: comparison of the lower, mean and higher first-trimester pulsatility indices. *Ultrasound Obstet Gynecol* 2011; 37: 534-537.* 25.64
- Poston L, Briley A, Seed P, Kelly F, Shennan A. Vitamin C and vitamin E in pregnant women at risk for pre-

eclampsia (VIP trial): randomised placebo-controlled trial. *Lancet* 2006;367:1145-54.*

- Rijvers C, Marzano S, Winkens B, Bakker J, Peeters L. Asymmetric dimethylarginine (ADMA) in early pregnancy: No differences between formerly preeclamptics with and without recurrent disease. *Reprod Sci* 2010;17:237A-8A. 35.48
- Salim R, Czarnowicki T, Nachum Z, Shalev E. The impact of close surveillance on pregnancy outcome among women with a prior history of antepartum complications attributed to thrombosis: A cohort study. *Reprod Biol Endocrinol* 2008;6.* 15.79
- Scholten RR, Sep S, Peeters L, Hopman MTE, Lotgering FK, Spaanderman MEA. Prepregnancy low-plasma volume and predisposition to preeclampsia and fetal growth restriction. *Obstet Gynecol* 2011;117:1085-93. 19.10
- Sep S, Andrietti S, Smits L, Peeters L. Is Obesity really an independent risk factor for recurrent preeclampsia? *Reprod Sci* 2010;17:131A. 24.19
- Sep SJ, Smits LJ, Prins MH, Spaanderman ME, Peeters LL. Simple prepregnant prediction rule for recurrent early-onset hypertensive disease in pregnancy. *Reprod Sci* 2009;16:80-87. 9.69
- Sep SJS, Schreurs MPH, Bekkers SCAM, Kruse AJ, Smits LJ, Peeters LLH. Early-pregnancy changes in cardiac diastolic function in women with recurrent pre-eclampsia and in previously pre-eclamptic women without recurrent disease. *BJOG Int J Obstet Gynaecol* 2011;118:1112-19. 29.41
- Sibai BM, Koch MA, Freire S, Pinto e Silva JL, Rudge MV, Martins-Costa S et al. The impact of prior preeclampsia on the risk of superimposed preeclampsia and other adverse pregnancy outcomes in patients with chronic hypertension. *Am J Obstet Gynecol* 2011;204:345-46. 46.80
- Sibai BM, Sarinoglu C, Mercer BM. Eclampsia. VII. Pregnancy outcome after eclampsia and long-term prognosis. *Am J Obstet Gynecol* 1992;166:1757-63. 38.46
- Sibai BMFA, - el-Nazer AFAU - Gonzalez-Ruiz, Gonzalez-Ruiz A. Severe preeclampsia-eclampsia in young primigravid women: subsequent pregnancy outcome and remote prognosis. *Am J Obstet Gynecol* 1986;155:1011-16. 50.77
- Sibai BMFA, Mercer BFAU, Sarinoglu C. Severe preeclampsia in the second trimester: recurrence risk and long-term prognosis. *Am J Obstet Gynecol* 1991;165:1408-12. 17.31
- Spaanderman MEA, Aardenburg R, Ekhardt THA, Van Eyndhoven HWF, de Leeuw PW, Peeters LLH. Pre-pregnant prediction of recurrent preeclampsia in normotensive thrombophilic formerly preeclamptic women receiving prophylactic antithrombotic medication. *J Soc Gynecol Invest* 2005;12:112-17. 51.06
- Spinnato JA, - Freire S - Pinto e Silva JL, Cunha Rudge MV, Martins-Costa S, Koch M, Goco N, de Barros Santos C, Cecatti JG, Costa R, Ramos JG, Moss N, Sibai BM. Antioxidant therapy to prevent preeclampsia: a randomized controlled trial. *Obstet Gynecol* 2007;110:1311-18. 12.90
- Sullivan CA, Magann EF, Perry J, Roberts WE, Blake PG, Martin J. The recurrence risk of the syndrome of hemolysis, elevated liver enzymes, and low platelets (HELLP) in subsequent gestations. *Am J Obstet Gynecol* 1994;171:940-43. 48.45
- Trogstad L, Skrandal A, Stoltenberg C, Magnus P, Nesheim BI, Eskild A. Recurrence risk of preeclampsia in twin and singleton pregnancies. *Am J Med Genet A*. 2004 Apr 1;126A(1):41-5.* 21.53
- van Oostwaard MF, Langenveld J, Bijloo R, Wong KM, Scholten I, Loix S, Hukkelhoven CW, Vergouwe Y, Papatsonis DNM, Mol BW, Ganzevoort W. Prediction of recurrence of hypertensive disorders of pregnancy between 34 and 37 weeks of gestation: a retrospective cohort study. *BJOG* 2012 Jun;119(7):840-7.* 33.88
- Kuijk SMJ v, Nijdam ME, Janssen KJM, Sep SJS, Peeters LL, Delahaije DHJ et al. A model for preconceptional prediction of recurrent early-onset preeclampsia: Derivation and internal validation. *Reprod Sci* 2011;18:1154-59. 6.88
- van Oostwaard MF, - Langenveld J, Schuit E, Wigny K, Van Susante H, Beune I, Ramaekers R, Papatsonis DNM, Mol BWJ, Ganzevoort W. Prediction of recurrence of hypertensive disorders of pregnancy in the term period, a retrospective cohort study. *Preg Hyper: An Int J Women's Card Health* (2014), <http://dx.doi.org/10.1016/j.preghy.2014.04.001>. * 40.74
- van Pampus MG, Wolf H, Mayruhu G, Treffers PE, Bleker OP. Long-term follow-up in patients with a history of (H)ELLP syndrome. *Hypertens Pregnancy* 2001;20:15-23. 27.17
- van Rijn BB, Hoeks LB, Bots ML, Franx A, Bruinse HW. Outcomes of subsequent pregnancy after first pregnancy with earlyonset preeclampsia. *Am J Obstet Gynecol* 2006;195:723-28. 27.50
- Villar J, Purwar M, Merialdi M, Zavaleta N, Thi Nhu Ngoc N, Anthony J et al. World Health Organisation multicentre randomised trial of supplementation with vitamins C and e among pregnant women at high risk for pre-eclampsia in populations of low nutritional status from developing countries. *BJOG Int J Obstet Gynaecol* 2009;116:780-88. 62.68

Zhang J, Troendle JF, Levine RJ. Risks of hypertensive disorders in the second pregnancy. *Paediatr Perinat Epidemiol* 2001;15:226-31.*

24.30

Mean****18.10**

Recurrence was calculated from published data of patients with hypertensive disorders of pregnancy in the index pregnancy.

*Where available the individual patient data were used (22 articles).

**The mean was calculated from a recurrence of 27,558 in 152,213 patients with a hypertensive disorder of pregnancy

ACCEPTED MANUSCRIPT