

This is the peer reviewed version of the following article:

Visual versus auditory Simon effect: a behavioural and physiological investigation / D'Ascenzo, Stefania; Lugli, Luisa; Baroni, Giulia; Guidotti, Roberto; Rubichi, Sandro; Iani, Cristina; Nicoletti, Roberto. - In: THE QUARTERLY JOURNAL OF EXPERIMENTAL PSYCHOLOGY. - ISSN 1747-0218. - 71:4(2018), pp. 917-930. [10.1080/17470218.2017.1307429]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

02/05/2026 05:42

(Article begins on next page)

This is the final peer-reviewed accepted manuscript of:

D'Ascenzo S, Lugli L, Baroni G, et al. Visual versus auditory Simon effect: A behavioural and physiological investigation. Quarterly Journal of Experimental Psychology. 2018;71(4):917-930. doi:10.1080/17470218.2017.1307429

The final published version is available online at:

<https://doi.org/10.1080/17470218.2017.1307429>

Rights / License:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

This item was downloaded from IRIS Università di Bologna (<https://cris.unibo.it/>)

When citing, please refer to the published version.

1
2
3
4
5
6
7
8
9
10
11
12
13
14 **Visual versus auditory Simon effect: a behavioural and physiological investigation**
15
16
17

18 Stefania D'Ascenzo^{1,2,*}, Luisa Lugli², Giulia Baroni², Roberto Guidotti³, Sandro Rubichi¹, Cristina
19 Iani¹, Roberto Nicoletti²
20
21
22
23

24 ¹Department of Communication and Economics, University of Modena and Reggio Emilia, Italy
25

26 ²Department of Philosophy and Communication, University of Bologna, Italy
27

28 ³Department of Neuroscience, Imaging and Clinical Sciences, and Institute of Advanced
29 Biomedical Technologies, University G. D'Annunzio, Chieti, Italy
30
31
32
33
34

35 **Running head:** Visual vs. Auditory Simon effect
36
37
38
39
40
41

42 ***Corresponding author:**
43

44 Stefania D'Ascenzo

45 Department of Philosophy and Communication

46 University of Bologna, Italy

47 Phone number: +39 051 2092240

48 E-mail: stefania.dascenzo@unibo.it
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

The present study investigated whether the visual and auditory Simon effects could be accounted for by the same mechanism. In a single experiment we performed a detailed comparison of the visual and the auditory Simon effects arising in behavioural responses and in pupil dilation, a psychophysiological measure considered as a marker of the cognitive effort induced by conflict processing. To address our question, we performed sequential and distributional analyses on both reaction times and pupil dilation. Results confirmed that the mechanisms underlying the visual and auditory Simon effects are functionally equivalent in terms of the interaction between unconditional and conditional response processes. The two modalities, however, differ with respect to the strength of their activation and inhibition. Importantly, pupillary data mirrored the pattern observed in behavioural data for both tasks, adding physiological evidence to the current literature on the processing of visual and auditory information in a conflict task.

Keywords: Simon effect, visual and auditory modalities, sequential effects, pupil dilation, Reaction Time distributions.

Introduction

In daily life, our responses are guided by sensory stimuli presented in multiple modalities, and a crucial question within the field of cognitive neuroscience is how individuals process and respond to stimuli in the presence of irrelevant, and possibly conflicting, information in order to meet task goals, and how they continuously adjust performance to changing environmental demands. These abilities, known as cognitive control, have been studied by means of different tasks, such as the Eriksen flanker task (Eriksen & Eriksen, 1974), the Stroop color-naming task (Stroop, 1935) and the Simon task (Simon & Rudell, 1967), with the latter being extensively employed to explore the conflict encountered when different stimulus dimensions compete to activate a response (for reviews, Proctor & Vu, 2006; Rubichi, Vu, Nicoletti, & Proctor, 2006). In this task, participants are required to respond to a non-spatial stimulus dimension (e.g., colour) by pressing a spatially defined key (e.g., left or right). Even though stimulus location is task-irrelevant, responses are faster and more accurate when stimulus and response positions correspond (corresponding trials), compared to when they do not correspond (non-corresponding trials).

A large amount of evidence supports the notion that the Simon effect (i.e., the difference in terms of speed and accuracy between non-corresponding and corresponding trials) originates at the response-selection stage and results from the interaction between two parallel and independent processing routes connecting perception to action, that is an unconditional/direct route and a conditional/indirect route (e.g., De Jong, Liang, & Lauber, 1994; Kornblum, Hasbroucq, & Osman, 1990). In the unconditional route, the response is thought to be automatically activated by stimulus position through pre-existing stimulus-response (S-R) associations, which are independent from the instructions. Differently, in the conditional route, the required response is activated based on task-defined associations that connect a stimulus to a specific response. In corresponding trials, the two activated responses correspond and no conflict arises (Umiltà, Rubichi, & Nicoletti, 1999). The opposite holds for non-corresponding trials, when a conflict occurs because the two activated

1
2
3 responses differ. In these trials, the incorrect response needs to be aborted thus causing a slowing of
4
5 response time and an increased number of errors.
6

7 The classic Simon effect has been replicated across different paradigm variations. In fact, it
8
9 has been shown to emerge with visual (e.g., Wühr & Ansorge, 2005), auditory (e.g., Simon &
10 Small, 1969; Vu, Proctor, & Urcuioli, 2003; Buetti & Kerzel, 2008) and tactile (e.g., Salzer,
11 Aisenberg, Oron-Gilad & Henik, 2013) stimuli and with motor (using both upper and lower limbs;
12 Leuthold & Schröter, 2006), vocal (Wühr, 2006) and oculo-motor (e.g., Khalid & Ansorge, 2013;
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Lugli, Baroni, Nicoletti, & Umiltà, 2016) responses. As regards to the comparison between the
visual and auditory modalities, the observation that the auditory Simon effect is larger in magnitude
with respect to the visual one (see, e.g., Pick & Proctor, 1999; Proctor & Pick, 1998) has been taken
as an indication that the activation of spatially corresponding responses is stronger with auditory
stimuli than with visual ones (e.g., Vu et al., 2003). However, it should be noted that the auditory
Simon effect has been studied less than the visual Simon effect and the literature has not reached so
far a general agreement on the mechanisms underlying the auditory Simon effect. Consequently, the
debate on whether the two effects are generated by different mechanisms or simply differ in the
strength of response activation is still open.

 In the present study we performed a detailed comparison of the visual and the auditory
Simon effects by recording behavioural measures (reaction times, RTs, and error rate, ER) and
pupil dilation (PD), with the aim of investigating the mechanisms underlying both effects. We
specifically focused on those mechanisms responsible for shaping the RT distribution and for the
sequential modulations evident in the visual and the auditory versions of the task.

 As regards the RT distribution, in the current literature, different explanations have been
proposed. It is important to note that researches investigating the Simon effect typically focus on the
comparison between corresponding and non-corresponding trials at the level of mean of RT.
However, this analysis may mask relevant findings that can only be discovered when analyzing the
RT distribution of interference effects (cf. De Jong et al., 1994; Ridderinkhof, 2002; for reviews see

1
2
3 Dittrich, Kellen, & Stahl, 2014; Proctor, Miles, & Baroni, 2011). Indeed, this technique has
4
5 revealed that the Simon effect can show two different time courses: a decreasing (i.e., the
6
7 magnitude of the Simon effect decreases as RT increases) and an increasing or constant (i.e., the
8
9 magnitude of the Simon effect increases or remained unchanged as RT increases) time course. The
10
11 former is found when the unconditional response activation occurs soon after the stimulus onset and
12
13 then dissipates over time (e.g. see Proctor, Miles, & Baroni, 2011), while the latter is observed
14
15 when the irrelevant response needs more time to reach complete activation and thus to exert the
16
17 maximum influence on performance (e.g. Wascher, Schatz, Kuder, & Verleger, 2001; Wühr, 2006).
18
19

20
21 To explain these different effect functions, Wascher et al. (2001) proposed that the Simon
22
23 effect can be generated by two different and dissociable mechanisms: a visuomotor facilitation of
24
25 same-side responses and a cognitive interference of codes. The decreasing effect function, yielded
26
27 by the visuomotor process, would be generated by a 'natural' spatial-anatomical mapping, that is,
28
29 when the stimuli are visual and presented in the horizontal axis (Wascher et al., 2001, Experiment
30
31 1). On the contrary, the stable or increasing effect function, yielded by the cognitive process, would
32
33 be due to the lack of a 'natural' relation between a visual stimulus and anatomical effect, as, for
34
35 example, when the stimuli are presented in an auditory modality (Wascher et al., 2001, Experiment
36
37 2). Based on their results, Wascher et al. (2001) concluded that the visual Simon effects appeared to
38
39 be due to specific mechanisms of visuomotor information transmission, thus it is associated to the
40
41 automatic activation of the corresponding response through the unconditional route; while the
42
43 auditory Simon effect is attributed to cognitive interference arising within the conditional route
44
45 only, in which the relevant stimulus feature is translated into a response. Conversely, Leuthold and
46
47 Schröter (2006) proposed that the increasing Simon effect found with the auditory stimuli would
48
49 result from the interaction between the two processing routes, as occurs for the visual Simon effect.
50
51 Recently, Xiong and Proctor (2016) examined the auditory Simon effect as a function of tone
52
53 frequency and duration. They found a decreasing Simon effect only when auditory stimuli were
54
55
56
57
58
59
60

1
2
3 low-frequency tones (200 Hz and 500 Hz), with a short duration, suggesting that the automatic
4 activation account proposed by Wascher et al. (2001) applies also to auditory stimuli.
5
6

7 For what concerns correspondence sequence, to our knowledge, a direct comparison
8 between correspondence sequence effects in the visual and auditory Simon effects has not been
9 performed so far. We believe it might provide insights on the mechanisms responsible for the two
10 conditions. Specifically, it has been widely shown that the magnitude of the Simon effect depends
11 on correspondence sequence: the effect is larger following a corresponding trial while it disappears
12 or even reverses following a non-corresponding trial (e.g. Iani, Rubichi, Gherri, & Nicoletti, 2009;
13 Iani, Stella, & Rubichi, 2014a; Soetens, Maetens, & Zeischka, 2010; Stürmer, Leuthold, Soetens,
14 Schröter, & Sommer, 2002). According to some authors, whenever a conflict is detected, as occurs
15 in non-corresponding trials, the mechanism selectively suppresses the unconditional route in order
16 to avoid conflicts in subsequent trials (e.g., Stürmer et al., 2002; Iani et al., 2009). This reduces the
17 costs of subsequent non-corresponding trials and the benefits of subsequent corresponding trials;
18 hence the Simon effect is reduced or eliminated. However, since after corresponding trials no
19 conflict is detected, nothing is preventing the activation of the irrelevant location-based response
20 code. This enhances the costs of subsequent non-corresponding trials and the benefits of subsequent
21 corresponding trials. It should be noted, however, that some authors have proposed alternative
22 explanations of correspondence sequence considering them as reflecting S–R priming (Egner, 2007;
23 Mayr, Awh, & Laurey, 2003) or binding effects (e.g., Hommel, Proctor, & Vu, 2004; Notebaert,
24 Gevers, Verbruggen, & Liefoghe, 2006; Notebaert, Soetens, & Melis, 2001; Spapé, Band, &
25 Hommel, 2011) rather than conflict-driven adaptations in cognitive control. In the typical Simon
26 task, correspondence sequence is confounded when stimulus and response repetitions occur in
27 consecutive trials. Specifically, whereas sequences of two corresponding trials and sequences of
28 two non-corresponding trials are either complete repetitions of stimulus position and response or
29 complete changes of both stimulus position and response, mixed sequences are always partial
30 repetitions in which either stimulus position or response repeats. Since responses to both complete
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 repetitions and complete alternations are always faster than those to partial repetitions (Hommel,
4 2004; Pashler & Baylis, 1991), the advantage of correspondence-level repetition may be accounted
5 for by the repetition of specific stimulus and response features and, consequently, may be due to the
6 absence of unbinding costs (i.e. featuring integration account; Hommel, 2004). For this reason, in
7 the present study we used the same version of the Simon task that was previously shown to produce
8 strong conflict-driven adaptations even when stimulus and response repetitions were controlled for
9 (Iani et al., 2009; van Steenbergen & Band, 2013).

10
11 It is worth mentioning that RT distributions are affected by correspondence sequence. Ridderinkhof
12 (2002), indeed, used RT distributional analyses to further investigate the dynamics of the activation-
13 suppression processes in the visual Simon task. The author claimed that the active suppression
14 process would be engaged to control the automatic activation of the unconditional route. More
15 precisely, a decreasing time course of the Simon effect would be due to a relatively strong selective
16 suppression (i.e., Simon effect following non-corresponding trials), while a constant or increasing
17 function could be attributed to a weaker inhibition process (i.e., Simon effect following
18 corresponding trials). The pattern of sequential modulations found in the auditory task (see
19 Leuthold & Schroter, 2006) can be explained as well by the suppression account proposed by
20 Stürmer et al. (2002) who postulated that the unconditional route is suppressed following non-
21 corresponding stimuli, and the S-R processing is mainly mediated by the conditional route
22 (resulting in the complete absence of interference across the entire RT distribution).

23
24 Furthermore, it is noteworthy that we decided to measure PD because the current literature
25 provides consistent evidence that the task-evoked pupillary response indexes cognitive effort,
26 increasing its dilation as a function of task demands (i.e., more difficult is the task, larger is the PD;
27 e.g., Beatty & Kahneman, 1966; Kahneman & Beatty, 1967; Loewenfeld, 1993; for a review see
28 Beatty, 1982; Beatty & Lucero-Wagoner, 2000; Laeng, Sirois, & Gredebäck, 2012). In addition, in
29 recent studies on interference between stimuli and responses, a conflict-related PD has been
30 observed in the Stroop (see Laeng, Ørbo, Holmlund, & Miozzo, 2011), in the Eriksen flanker (van
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Bochove, van der Haegen, Notebaert, & Verguts, 2013; Wendt, Kiesel, Geringswald, Purmann, &
4
5 Fischer, 2014) and in the Simon tasks (van Steenbergen & Band, 2013; D'Ascenzo, Iani, Guidotti,
6
7 Laeng & Rubichi, 2016). In particular, van Steenbergen and Band (2013) used a visual Simon task
8
9 to investigate whether the PD could be considered as an indirect marker either of conflict- or
10
11 control-related processes. The authors analyzed sequential effects emerging in RTs and PD and
12
13 found that PD increased when the conflict was higher, that is for non-corresponding trials with
14
15 respect to corresponding ones. Consistent with the conflict monitoring theory (Botvinick, Braver,
16
17 Barch, Carter, & Cohen, 2001), they observed sequential effects on both RTs and PDs. Hence, their
18
19 results indicate that increased PD could be interpreted as a sensitive marker of conflict-related
20
21 processing. It has been pointed out, though, that the authors employed a visual Simon paradigm,
22
23 while it has still to be investigated whether the same processes are involved with a different sensory
24
25 modality. To our knowledge, in fact, neither the auditory Simon effect by means of PD has been
26
27 examined so far, nor, consequently a comparison between visual and auditory PD Simon effect has
28
29 been provided.
30
31
32
33

34 To this aim, in the present study we implemented an experiment in which a group of
35
36 participants performed a Simon task with visual stimuli, while another group performed a Simon
37
38 task with auditory stimuli. RTs, ER and PD in the two conditions were analysed as a function of
39
40 correspondence sequence. Furthermore, for RTs we analysed the time course of sequential
41
42 modulations (for a review see Dittrich et al., 2014) and subsequently, we investigated how the PD
43
44 changed depending on the RTs distribution.
45
46

47 As regards RTs, since it has been shown that the auditory modality is more automatically
48
49 alerting than the visual modality (e.g., Posner, Nissen, & Klein, 1976), we expected to find overall
50
51 faster responses in the auditory condition compared to the visual one. For the same reason, we also
52
53 hypothesized that auditory stimuli would produce a stronger activation of the corresponding
54
55 response through the unconditional route, this resulting in a larger Simon effect for auditory than for
56
57 visual stimuli (Simon, 1990; Pick & Proctor, 1999; Proctor & Pick, 1998; Vu et al., 2003). As
58
59
60

1
2
3 regards PDs, in line with the hypotheses on RTs, we expected to find an overall reduced PD
4
5 amplitude for the auditory condition compared to the visual one, indicating that auditory stimuli
6
7 require less effort to be processed. In addition, we expected to find, for both visual and auditory
8
9 stimuli, an increased PD in non-corresponding trials as compared to corresponding ones, replicating
10
11 the previous studies on the visual condition (see Laeng, Ørbo, Holmlund, & Miozzo 2011; van
12
13 Steenbergen & Band, 2013) and, interestingly, adding new evidence on the auditory condition. For
14
15 what concerns the Simon effect, as hypothesized for RTs, we expected a modality specific effect on
16
17 PD amplitude. Specifically, due to the stronger activation produced by corresponding auditory
18
19 stimuli compared to the visual ones, we expected smaller PD with the former compared to the latter
20
21 stimuli, resulting in a larger Simon effect for auditory than for visual stimuli.
22
23
24

25 As regards the mechanism responsible for shaping the RT distributions, according to the
26
27 above-mentioned literature, we expected to find a decreasing Simon effect as a function of RTs for
28
29 the visual condition, resulting from the visuomotor process (see Wascher et al., 2001, Experiment
30
31 1), and an increasing Simon effect as a function of RTs for the auditory condition, resulting from
32
33 the cognitive process (see Wascher et al., 2001, Experiment 2)¹ For what concerns the time course
34
35 of sequential modulations (e.g., Ridderinkhof, 2002; Stürmer et al., 2002), we hypothesized that for
36
37 the visual Simon condition, in line with Ridderinkhof (2002), a stable or increasing time course
38
39 would emerge following a corresponding trials, while a decreasing time course would emerge
40
41 following a non-corresponding trial; for the auditory Simon condition, in line with Stürmer et al.
42
43 (2002), an increasing time course would emerge following a corresponding trials, while a stable
44
45 time course would emerge following a non-corresponding trial. In addition, we analyzed PD
46
47 depending on RT distributions for each condition. Considering that a larger PD is an index of
48
49 greater effort, we expected to find, for both conditions, larger PDs in those bins for which a larger
50
51 Simon effect for RTs was evident.
52
53
54
55
56
57

58 ¹ To note Xiong and Proctor (2016) found a decreasing function for the RT Simon effect only when low frequency
59 tones were used. We did not expect to find the same pattern since we used tones of both low and high frequency.
60

1
2
3 As regards the mechanism responsible for producing sequential modulations, we expected
4 the magnitude of the Simon effect in RTs to be larger after a corresponding trial and smaller, absent
5 or reversed after a non-corresponding trial. We hypothesized that, if the mechanisms underlying the
6 visual and auditory Simon effects are comparable, the sequential modulation pattern should hold for
7 both visual (e.g., Stürmer et al., 2002) and auditory stimuli (e.g., Leuthold & Schröter, 2006). As
8 regards PD, we hypothesized that if the effects evident in the two modalities are due to the interplay
9 between unconditional and conditional processing, than a modulation of PD by correspondence
10 sequence should be evident in both conditions, with a larger Simon effect in PDs following
11 corresponding trials and a smaller effect following non-corresponding ones.
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **Methods**

26
27
28
29 **Participants.** Sixty-nine undergraduates from the University of Bologna participated as volunteers
30 to the experiment (41 female; two left handed; mean age: 20 years; standard deviation: 1.9 years).
31 All of them reported normal or corrected-to-normal vision and normal hearing. Thirty-two
32 participants took part in the visual condition and thirty-seven in the auditory condition. A different
33 number of participants was run in the two conditions because we wanted to have a sample size of
34 30² participants for each condition, after the application of the exclusion criteria, see Statistical
35 analysis section below. The local ethics committee approved the study and written informed consent
36 was obtained from all of the participants before participation.
37
38
39
40
41
42
43
44
45
46

47 **Apparatus and Stimuli.** Participants sat in front of a LCD monitor (1024x768) at a viewing
48 distance of 70 cm in a dimly lit room. Stimulus presentation and response collection were controlled
49 by the E-Prime® software system.
50
51
52
53
54
55
56

57 ² A power analysis using G*Power software (Faul et al., 2007) to calculate the sample size was conducted. It confirmed
58 that at least 36 subjects would be needed as a total sample in order to detect a statistically significant difference (if it
59 exists) between the two groups in the sequential modulations (α level $p=0.05$, power 95%, effect size = 0.25).
60

1
2
3 The stimuli for the visual condition were black capital letters “M” and “W” that were presented
4
5 1.07° to the left or right of a central fixation cross on a dark grey background (average luminosity of
6
7 each stimulus including the background was 127 units in the RGB system). The stimuli for the
8
9 auditory condition were “high” (1050 Hz) or “low” (650 Hz) tones with the loudness approximately
10
11 of 60 dB, presented through loudspeakers placed 15 cm to the left and to the right side of the
12
13 computer monitor. For both conditions, responses were made by pressing the “z” or the “m” keys
14
15 on the QWERTY keyboard with the left or the right index finger, respectively. The keyboard was
16
17 located centrally with respect to the body midline.
18
19

20 **Procedure.** For the visual (see van Steenbergen & Band, 2013) and the auditory Simon conditions,
21
22 participants were instructed to respond as quickly and as accurately as possible to the type of letter
23
24 and to the tone pitch, respectively, while ignoring their location. In the visual condition, half of the
25
26 participants responded to the "M" letter with the left hand and to the "W" letter with the right hand,
27
28 while the other half experienced the opposite mapping. In the auditory condition, half of the
29
30 participants responded to the "low" tone with the left hand and to the "high" tone with the right
31
32 hand, while the other half experienced the inverse mapping.
33
34
35

36 Both conditions consisted of 384 trials that were divided into six blocks of 64 trials each,
37
38 preceded by 8 practice trials. In each block, the trial sequence was controlled so that each trial could
39
40 be preceded with an equal probability by a corresponding (C) or a non-corresponding (NC) trial.
41
42 Hence, four different trial sequences were created (C–C, C–NC, NC–C, NC–NC, with italics
43
44 denoting current trial correspondence).
45
46

47 Each trial started with a white fixation cross that, after 900 ms, turned yellow for 200 ms (warning
48
49 cue) and then returned white for the remaining duration of the trial. The stimulus appeared for 100
50
51 ms together with the white fixation cross, that remained visible on the screen for the following 900
52
53 ms. Responses up to 1000 ms after stimulus presentation were recorded. A trial lasted 2100 ms.
54
55
56
57
58
59
60

1
2
3 **Pupillometry.** Participant's right eye pupil diameter was measured using the iViewX Remote Eye
4 Tracking Device (R.E.D.; SensoMotoric Instruments, SMI, Teltow, Germany). Data were recorded
5 at a sampling rate of 60 Hz using the iView X Software (SMI, Teltow, Germany).
6
7

8
9 The illumination of the testing room was kept constant during the whole session. A standard 5-point
10 eye tracker calibration routine was used at the beginning of the experiment. Participants were
11 instructed to keep their head as steady as possible.
12
13

14
15 A single measure (in mm) of pupil diameter was obtained for each sample. Pupil data were
16 pre-processed using a custom-made Python script to remove artefacts in the time series related to
17 eye blinks: data points with physiologically unlikely pupil sizes (smaller than 2 mm or larger than 8
18 mm,) together with the neighbouring data points (the preceding and following 80 ms) were
19 removed. Also, samples having more than 2.5 SDs from the mean pupil size within a trial were
20 removed from the time series. Trials with less than 50% of the data remaining after removal of
21 outliers were not included for further analysis. Resulting gaps in the data were replaced by linear
22 interpolation and the resulting time series were smoothed using a cubic spline. Baseline pupil
23 diameter was calculated as the average pupil size during the 200 ms (warning cue) preceding each
24 stimulus onset and was subtracted from the time series of the entire subsequent trial (0-1900 ms). If
25 the warning cue was discarded, the baseline was calculated using first 200 ms after stimulus onset.
26
27 We calculated the average of the trial's time duration as a function of the onset of the stimulus for
28 each condition, without considering trials excluded from behavioural analysis. Baseline-corrected
29 pupil diameter change in response to the stimulus is plotted in Figure 1, for the visual (A) and the
30 auditory Simon condition (B).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 (Figure 1 about here)
52
53
54
55

56 **Statistical Analyses.** Correct RTs, arcsine-transformed error rates (ERs) and mean PDs in the entire
57 epoch (0-1900 ms) for both conditions were submitted to a repeated-measures Analysis of Variance
58
59
60

1
2
3 (ANOVA) with Current trial correspondence (corresponding vs. non-corresponding) and Previous
4 trial correspondence (corresponding vs. non-corresponding) as within-participant factors and
5 Condition (visual vs. auditory) as between-participants factor³. When necessary, comparisons were
6 performed using Paired samples *t*-tests and by correcting the p-value for the number of comparisons
7 (Bonferroni correction).
8
9

10
11
12
13
14 To compute the time course of the sequential modulations in RTs we applied the
15 Vincentizing procedure (Ratcliff, 1979). For both the visual and the auditory conditions, individual
16 RTs were rank ordered as a function of Current and Previous trial correspondence and divided into
17 quintiles (bins). For each bin, we calculated the size of the Simon effect by subtracting the mean RT
18 for the corresponding condition from the mean RT for the non-corresponding condition. The
19 resulting values were submitted to a repeated-measure ANOVA with Bin (from 1 to 5) and Previous
20 trial correspondence as within-participant factors and Condition as between-participants factor. It
21 should be noted that, considering the way data were grouped, the main effect of Bin necessarily
22 turned out to be significant in all the analyses; therefore, it will not be discussed here or later on.
23
24
25
26
27
28
29
30
31
32

33
34 In addition, for both the visual and the auditory conditions, considering the quintiles (bins)
35 obtained by ranking RTs, we computed mean PD (baseline-corrected) in the entire epoch (0-1900
36 ms) as a function of each RT bin for both Current and Previous trial correspondence. The resulting
37 values were submitted to the same analysis reported above.
38
39
40
41
42

43 For the visual Simon condition, two participants were excluded from the analyses: one
44 because of the ER being higher than 20% of the total trials, and the other because 50% of his/her
45 total trials were discarded due to technical problems with the eye tracker recording. As a result, 30
46 out of 32 participants were included in the behavioural and pupillary analyses. For the auditory
47 Simon condition, seven participants were excluded from the analyses because of their high error
48
49
50
51
52
53
54
55
56

57
58 ³ We would like to point out that, to address our experimental questions, we decided to adopt a between-subject design to avoid practice effects from
59 the task performed as first to the task performed as second. However, we performed a follow-up experiment using a within-subject design (see
60 [Appendix for the details](#)).

1
2
3 rate (more than 20% of the total trials). As a result, 30 out of 37 participants were included in the
4
5 behavioural and pupillary analyses.
6

7 For both the visual and auditory Simon conditions, incorrect responses (7.2% and 6.4% of
8 the trials, for visual and auditory Simon condition, respectively) and latencies that were 2 standard
9 deviations above (3.7 % and 10.6 % of the trials, for visual and auditory Simon conditions,
10 respectively) or below (0.6% of the trials, for both visual and auditory Simon conditions) each
11 participant's mean were excluded from the analyses.
12
13
14
15
16
17

18 The exclusion criteria were decided a prior based on previous studies employing the same
19 task. For what concerns behavioural data see, for example, Iani, Milanese and Rubichi (2014b). For
20 what concerns pupillary data see, for example, Laeng et al. (2011, 2012) and Alnaes, Sneve,
21 Espeseth, Endstad, van de Pavert and Laeng (2014).
22
23
24
25
26
27

28 29 **Results**

30
31
32
33
34 **RTs.** The analysis showed a significant main effect of Current trial correspondence, $F(1, 58)$
35 = 106.00, $MSE = 49815$, $p < .001$, $\eta_p^2 = .65$, with faster responses in corresponding ($M = 349$; $SD =$
36 64) than in non-corresponding ($M = 378$; $SD = 68$) trials. The main effects of Previous trial
37 correspondence, $F(1, 58) = 1.09$, $MSE = 127.29$, and Condition, $F(1, 58) = 1.02$, $MSE = 16011.07$,
38 did not reach significance.
39
40
41
42
43
44

45 The interaction between Current trial correspondence and Condition was significant, $F(1,$
46 58) = 5.79, $MSE = 2721$, $p < .05$, $\eta_p^2 = .09$. T-tests showed that the difference between
47 corresponding ($M = 360$; $SD = 61$; $M = 338$; $SD = 64$; for visual and auditory conditions,
48 respectively) and non-corresponding trials ($M = 382$; $SD = 54$; $M = 373$; $SD = 80$; for visual and
49 auditory conditions, respectively) (i.e., the Simon effect) was of 22 ms in the visual modality ($t_{(29)} =$
50 6.368, $p_{\text{Bonferroni corrected}} < .001$) and of 35 ms in the auditory modality ($t_{(29)} = 8.090$, $p_{\text{Bonferroni corrected}}$
51
52
53
54
55
56
57
58
59
60

< .001). T-test showed that the difference between the two effect sizes was significant, ($t_{(58)} = -2.407, p < .05, d = -0.63$).

The interaction between Previous trial correspondence and Condition was also significant, $F(1, 58) = 15.67, MSE = 1836, p < .001, \eta_p^2 = .21$. T-tests showed that only in the visual condition responses were faster when the preceding trial was corresponding ($M = 368; SD = 54$) than when it was non-corresponding ($M = 375; SD = 53$), $t_{(29)} = -4.712, p_{\text{Bonferroni corrected}} < .001$. Importantly, the interaction between Current and Previous trial correspondence was significant, $F(1, 58) = 237, MSE = 50726, p < .001, \eta_p^2 = .80$. T-tests showed a significant Simon effect of 58 ms following a corresponding trial ($t_{(59)} = 15.669, p_{\text{Bonferroni corrected}} < .001$) and a null effect following a non-corresponding trial ($t_{(59)} = -0.072, p = .94$). Crucially, this two-way interaction was modulated by Condition, $F(1, 58) = 25.53, MSE = 5466, p < .001, \eta_p^2 = .30$ (see Figure 2A, visual and auditory conditions in the leftmost and rightmost panel, respectively). T-tests showed that the visual Simon effect was of 61 ms when the preceding trial was corresponding ($t_{(29)} = 14.888, p_{\text{Bonferroni corrected}} < .001$) and reversed to -16 ms when the preceding trial was non-corresponding ($t_{(29)} = -3.345, p_{\text{Bonferroni corrected}} = .008$), while the auditory Simon effect was of 55 ms when the preceding trial was corresponding ($t_{(29)} = 8.892, p_{\text{Bonferroni corrected}} < .001$) and decreased to 16 ms when the preceding trial was non-corresponding ($t_{(29)} = 4.674, p_{\text{Bonferroni corrected}} < .001$). T-tests showed that the Simon effect evident after a non-corresponding trial was significantly smaller in the visual ($M = -16, SD = 27.09$) than in the auditory ($M = 16, SD = 18.77; t_{(58)} = -5.411, p_{\text{Bonferroni corrected}} < .001, d = -1.42$) condition. No difference between conditions was evident after a corresponding trial (visual: $M = 61, SD = 22.33$; auditory: $M = 55, SD = 33.92; t_{(58)} = 0.758, p = .45, d = 0.20$).

ERs. The analysis showed significant main effects of Current trial correspondence, $F(1,58) = 67.28, MSE = 0.49, p < .001, \eta_p^2 = .53$, and Previous trial correspondence, $F(1, 58) = 9.81, MSE = 0.035, p < .005, \eta_p^2 = .14$, with higher error rates in non-corresponding ($M = 9.17\%; SD = 8.01\%$) than in corresponding trials ($M = 4.72\%; SD = 4.08\%$), and after corresponding ($M = 7.88\% SD = 8.04\%$) than after non-corresponding ($M = 6.01\%; SD = 4.92\%$) trials. The interaction between

Current and Previous trial correspondence was significant, $F(1, 58) = 103$, $MSE = 0.50$, $p < .001$, $\eta_p^2 = .64$. T-tests showed that the Simon effect was significant after a corresponding trial (9.10%; $t_{(59)} = 10.075$, $p_{\text{Bonferroni corrected}} < .001$), but did not reach significance after a non-corresponding trial (-0.21%, $t_{(59)} = -0.049$, $p = .96$). Importantly, the three-way interaction between Current and Previous trial correspondence and Condition was significant, $F(1, 58) = 39.35$, $MSE = 0.19$, $p < .001$, $\eta_p^2 = .40$ (see Figure 2B, visual and auditory conditions in the leftmost and rightmost panel, respectively). T-tests showed that the visual Simon effect was positive when the preceding trial was corresponding (12.5%; $t_{(29)} = 9.450$, $p_{\text{Bonferroni corrected}} < .001$) and negative when the preceding trial was non-corresponding (-2.3%; $t_{(29)} = -3.371$, $p_{\text{Bonferroni corrected}} = .012$). The auditory Simon effect was significant when the preceding trial was corresponding (5.7%; $t_{(29)} = 6.125$, $p_{\text{Bonferroni corrected}} < .001$) and failed to reach significance when the preceding trial was non-corresponding (1.9%; $t_{(29)} = 2.587$, $p_{\text{Bonferroni corrected}} = .09$). T-test showed that the Simon effect was comparable in the two conditions (visual: $M = 5.1\%$, $SD = 4$; auditory: $M = 3.8\%$, $SD = 5$; $t_{(58)} = 0.679$, $p = .92$, $d = 0.18$). However, considering sequential modulations, t-tests showed that after a corresponding trial the Simon effect in the visual condition ($M = 12.5\%$, $SD = 8.57$) was significantly larger than in the auditory condition ($M = 5.7\%$, $SD = 3.68$; $t_{(58)} = 3.906$, $p_{\text{Bonferroni corrected}} < .001$, $d = 1.03$). After a non-corresponding trial, the Simon effect in the visual condition ($M = -2.3\%$, $SD = 6.32$) was significantly smaller than in the auditory condition ($M = 1.9\%$, $SD = 4.87$; $t_{(58)} = -4.044$, $p_{\text{Bonferroni corrected}} < .001$, $d = -1.06$). No other main effects or interactions reached significance ($F_s < 1$).

PD. The analysis on mean PD (baseline-corrected) during the entire epoch (0-1900 ms) revealed a significant effect of Current trial correspondence, $F(1, 58) = 9.51$, $MSE = .00$, $p < .005$, $\eta_p^2 = .14$, with larger PDs in non-corresponding ($M = 0.079$ mm; $SD = 0.04$) than in corresponding ($M = 0.070$ mm; $SD = 0.04$) trials. Current trial correspondence did not interact with Condition, $F(1, 58) = 0.29$, $MSE = 0.00$, indicating that the difference between corresponding ($M = 0.072$; $SD = 0.03$; $M = 0.069$; $SD = 0.06$; for visual and auditory conditions, respectively) and non-corresponding trials ($M = 0.079$; $SD = 0.04$; $M = 0.079$; $SD = 0.06$; for visual and auditory

conditions, respectively) was comparable in the two conditions (visual: $M = 0.007$ mm; $SD = 0.018$; auditory: $M = 0.010$ mm, $SD = 0.023$).

Current and Previous trial correspondence interacted, $F(1,58) = 8.37$, $MSE = .01$, $p < .005$, $\eta_p^2 = .13$. T-tests showed that the Simon effect was significant following a corresponding trial (0.018 mm; $t_{(59)} = 4.133$, $p_{\text{Bonferroni corrected}} < .001$), whereas it did not reach significance following a non-corresponding trial (-0.001 mm; $t_{(59)} = -0.329$, $p = .743$). This pattern was evident for both conditions (see Figure 2C, visual and auditory conditions in the leftmost and rightmost panel, respectively), as indicated by the lack of a significant three-way interaction involving the Condition factor, $F(1, 58) = 1.58$, $MSE = 0.00$. No other main effects or interactions reached significance ($F_s < 1$).

(Figure 2 about here)

RT distributions. This analysis revealed a significant interaction between Current trial correspondence, Bin and Condition, $F(4, 232) = 60.144$, $MSE = 13164$, $p < .001$, $\eta_p^2 = .50$. T-test revealed that the visual Simon effect was significant in the first four bins ($t_{s(29)} > -13.10$, $p_s < .05$) but did not reach significance in the last bin ($t_{(29)} = 0.91$; $p = .37$). Differently, the auditory Simon effect was significant at all the bins ($t_{s(29)} > 5.59$, $p_s < .001$). Furthermore, Helmert contrasts showed that the size of the visual Simon effect decreased significantly from bin 1 to bin 5 (42, 35, 25, 10, and -5 ms, respectively), $F_s(1, 29) > 52.70$, $MSE_s > 20276.39$, $p_s < .001$, while the size of the auditory Simon effect increased significantly from bin 1 to bin 4 (18, 28, 36 and 47 ms, respectively), $F_s(1,29) > 40.18$, $p < .001$, and remained stable from bin 4 to bin 5 (47 and 50 ms, respectively), $F(1, 29) = .76$, $p = .39$ (see Figure 3A, visual and auditory conditions in the leftmost and rightmost panel, respectively).

Interestingly, the interaction between Previous and Current trial correspondence, Bin and Condition was significant, $F(4, 232) = 10.428$, $MSE = 1091$, $p < .001$, $\eta_p^2 = .15$. For the visual

Simon condition, t-test revealed that after a corresponding trial, the effect was significant across bins ($t_{s(29)} > -20.04$, $p_s < .001$), whereas after a non-corresponding trial, the effect was significant at bins 3, 4 and 5 ($t_{s(29)} > 3.68$, $p_s < .001$) and was not significant at bins 1 and 2 ($t_{s(29)} < -1.37$, $p_s > .180$). Helmert contrasts showed that the size of the Simon effect evident after a corresponding trial decreased significantly from bin 1 to bin 5 (78, 74, 67, 55 and 28 ms, respectively), $F_s(1, 29) > 61.17$, $p_s < .001$. Whereas the size of the Simon effect after a non-corresponding trial decreased significantly from bin 1 to bin 4 (7, -7, -20, -29 ms, respectively), $F_s(1, 29) > 33.39$, $MSE_s = 26301.74$, $p_s < .05$, but it did not differ significantly between bins 4 and 5 (-29 and -32 ms, respectively), $F(1, 29) = .495$, $MSE = 310.02$ (see Figure 3B, leftmost panel for the visual condition).

For the auditory Simon condition, t-tests revealed that the Simon effect was significant at all bins irrespective of previous trial correspondence (respectively, $t_{s(21)} > 5.91$, $p_s < .001$; $t_{s(21)} > 2.28$, $p_s < .05$). Furthermore, Helmert contrasts showed that the size of the Simon effect evident after a corresponding trial increased significantly from bin 1 to bin 4 (27, 41, 59 and 71 ms, respectively), $F_s(1, 29) > 62.59$, $MSE_s > 34415.17$, $p < .001$, but did not differ between bins 4 and 5 (71 and 71 ms, respectively), $F(1, 29) = .00$, $MSE = 0.00$). Conversely, the size of the Simon effect evident after a non-corresponding trial increased significantly from bin 1 to bin 2 (7, and 14 ms, respectively), $F(1, 29) = 5.57$, $MSE = 3560.68$, $p < .05$, but did not differ significantly in the other bins (14, 14, 19 and 22 ms, respectively), $F_s(1, 29) < 2.58$, $MSE_s < 1234.41$, (see Figure 3B, rightmost panel for the auditory condition).

PD depending on RTs distributions. The interaction between Current trial correspondence, Bin and Condition failed to reach significance $F(4, 55) = .797$, $MSE = .001$ (see Figure 4A, leftmost panel for the visual condition and rightmost panel for the auditory condition). Interestingly, the interaction between Previous and Current trial correspondence, Bin and Condition was significant $F(4, 55) = 3.351$, $MSE = .007$, $p < .01$, $\eta_p^2 = .22$. For the visual Simon condition, t-test revealed that after a corresponding trial, the effect was significant at bins 2, 3 and 4 ($t_{s(29)} > -3.008$, $p_s < .05$), and

1
 2
 3 it was not significant at bins 1 and 5 ($t_{s(29)} < -0.92$, $p_s > .78$), whereas after a non-corresponding
 4
 5 trial, the effect was close to significance at bin 5 ($t_{(29)} = 1.83$, $p = .078$) and it was not significant
 6
 7 from bin 1 to bin 4 ($t_{s(29)} < -0.788$, $p_s > .43$). Helmert contrasts showed that the size of the Simon
 8
 9 effect after a corresponding trial was stable across bins (-0.004, 0.022, 0.035, 0.022, 0.010 mm,
 10
 11 from bin 1 to bin 5 respectively), $F_s(1, 29) < 2.58$, $MSE_s < 0.02$. The size of the Simon effect after a
 12
 13 non-corresponding trial was also stable across bins (0.008, -0.001, -0.008, -0.007, -0.022 mm, from
 14
 15 bin 1 to bin 5 respectively), $F_s(1, 29) < 1.78$, $MSE_s < 0.01$ (see Figure 4B, leftmost panel for the
 16
 17 visual condition). For the auditory Simon condition, t-tests revealed that after a corresponding trial,
 18
 19 the effect was significant at bin 4 ($t_{(29)} > -4.379$, $p < .001$), and it was not significant at bins 1, 2, 3
 20
 21 and 5 ($t_{s(29)} < -0.28$, $p_s = .77$) whereas after a non-corresponding trial, the effect was significant at
 22
 23 bin 3 and 4 ($t_{(29)} = 2.089$, $p < .05$) and it was not significant at bin 1, 2 and 5 ($t_{s(29)} < -0.576$, $p_s =$
 24
 25 $.56$). Helmert contrasts showed that the size of the Simon effect evident after a corresponding trial
 26
 27 was stable across bins (0.003, 0.015, 0.005, 0.050, 0.009 mm, from bin 1 to bin 5 respectively),
 28
 29 $F_s(1, 29) < 3.36$, $MSE_s < 0.05$, whereas the size of the Simon effect after a non-corresponding trial
 30
 31 decreased significantly only from bin 3 to bin 4 (0.028, -0.023 mm, respectively), $F_s(1, 29) = 8.32$,
 32
 33 $MSE_s = 0.04$, $p < .01$, and it was stable at bins 1, 2 and 5 (0.006, -0.008, 0.007 mm, respectively),
 34
 35 $F_s(1, 29) < 8.32$, $MSE_s < 0.04$ (see Figure 4B, rightmost panel for the auditory condition).
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60

(Figure 3 about here)

General Discussion

The Simon effect has been widely investigated by employing visual stimuli, while the effect
 emerging with auditory stimuli remains, at least for some aspects, less studied. Specifically, while it
 is widely accepted that the visual Simon effect originates from the interplay between two processing
 routes leading to response activation, there is still an ongoing debate regarding the mechanisms

1
2
3 underlying the auditory Simon task (e.g., Wascher et al., 2001; Leuthold & Schröter, 2006; Xiong
4 & Proctor, 2016).

5
6
7 In the present study we performed a detailed comparison of the visual and the auditory
8 Simon effects recording behavioural measures (reaction times, RTs, and error rate, ER) and pupil
9 dilation (PD), which is considered as a physiological index of the cognitive effort induced by
10 conflict-related processing (i.e., van Steenbergen & Band, 2013), aiming to investigate the
11 mechanisms underlying both effects. We specifically focused on those mechanisms responsible for
12 shaping the RT distributions and for the sequential modulations evident in the visual and the
13 auditory versions of the task. Importantly, to our knowledge, this is the first study that examined
14 conflict processing by means of the pupillary response in the auditory Simon task. It should be
15 noted that to achieve our aim and to avoid practice effects from the task performed as first to the
16 task performed as second, a between-subjects design has been performed. However, in order to
17 investigate if practice effects could influence the subsequent task also in this specific paradigm, a
18 follow-up experiment using a within-subjects design was performed (see Appendix). Results of the
19 two experiments are in line, but the within-subjects design revealed an effect of the Order of
20 presentation of the task, supporting the notion that Simon task performance can be affected by a
21 similar task performed before (e.g., Iani et al., 2009; Marini, Iani, Nicoletti, & Rubichi, 2011;
22 Tagliabue, Zorzi, Umiltà, & Bassignani, 2000). Specifically, performing the auditory condition as
23 first speeded up performance as compared to when the visual condition was executed as first. This
24 result shows that performing the same task in a specific modality (i.e., auditory) can influence
25 performance of the same task in a different modality (i.e., visual modality; see for example Vu et
26 al., 2003). Consequently, we decided to focus our discussion on the results of the between-subjects
27 experiment.

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 Our behavioural results were in line with those of previous studies (e.g., Wühr & Ansorge,
55 2005; Leuthold & Schröter, 2006; Pick & Proctor, 1999): the Simon effect emerged for both visual
56 and auditory modalities. As hypothesized, despite we did not find an overall advantage of the
57
58
59
60

1
2
3 auditory condition compared to the visual one, the auditory Simon effect was significantly larger as
4
5 compared to the visual one (35 vs. 24 ms). Crucially, the analysis of sequential modulations showed
6
7 that after a corresponding trial the Simon effect for RTs did not differ across modalities (61 vs. 55
8
9 ms, respectively), while it was larger for the visual than for the auditory Simon condition on ERs
10
11 (12.5% vs. 5.7%, respectively). Differently, after a non-corresponding trial, the Simon effect in both
12
13 RTs and ERs was reversed in the visual modality and only reduced in the auditory modality (-16 ms
14
15 vs. 16 ms and -2.3% vs. 1.9 %, respectively). Overall, the sequential modulations replicated those of
16
17 previous studies using visual and auditory stimuli (e.g., Iani et al., 2009, 2012; Stürmer et al., 2002;
18
19 Leuthold & Schröter, 2006), hence supporting the idea that the conflict experienced in a trial
20
21 triggers adaptations aimed at eliminating the impact of spatial S–R correspondence on response
22
23 selection in the following trial. Specifically, these findings are consistent with the proposal by
24
25 Stürmer et al. (2002) that the unconditional processing route is under the control of an ancillary
26
27 monitoring mechanism (AMM), which detects response conflict and subsequently suppresses the
28
29 unconditional route (see also Ridderinkhof, 2002). It should, however, be noted that sequential
30
31 modulations have been observed by Hommel et al. (2004) in the absence of conflict adaptation
32
33 effects. The authors showed persistent sequential modulations in versions of the Simon task in
34
35 which the occurrence of response conflict on the previous trial was precluded and concluded that
36
37 congruency sequence effects in the Simon task do not require conflict-driven cognitive control
38
39 processes. Subsequently, Wühr (2005) showed that sequential modulations can also be obtained in
40
41 the absence of feature integration effects, concluding that, in a typical Simon task (as used in the
42
43 present study), sequential modulations might reflect additive effects of conflict adaptation and
44
45 feature integration (see also Wühr & Ansorge, 2005). Thus, it is not possible to exclude that our
46
47 results were influenced by repetition effect, since, with the present paradigm, it is not possible to
48
49 completely disentangle feature integration and conflict adaptation effects, and both may contribute
50
51 to the observed sequential modulations. Interestingly, the direct comparison between sequential
52
53 modulations in the visual and auditory versions of the Simon task provides some insights on the
54
55
56
57
58
59
60

1
2
3 mechanism responsible for these modulations in the two conditions. Indeed, if we consider
4
5 performance after a non-corresponding trial, we can notice that the visual Simon effect reversed
6
7 while the auditory Simon effect was simply reduced. This finding may suggest that in the two
8
9 conditions the same suppression mechanism produces different outcomes, probably because of the
10
11 different strength of the activation produced by stimuli in the two modalities. Specifically, the
12
13 activation of the corresponding response through the unconditional route produced by auditory
14
15 stimuli, that is thought to be stronger than the one produced by visual stimuli (see Vu et al., 2003),
16
17 could be more difficult to suppress, resulting in a reduced auditory Simon effect after non-
18
19 corresponding trials. Differently, since visual stimuli produce a weaker activation than auditory
20
21 stimuli, suppression may result in a reversed visual Simon effect after non-corresponding trials.
22
23 This inference is consistent with the notion that visual and auditory information processing differs
24
25 in many regards, as also manifested by relevant differences between visual and auditory spatial
26
27 attention (e.g., Neumann, van der Heijden, & Allport, 1986; Kanwisher & Wojciulik, 2000; Spence
28
29 & Driver, 1994; Wu, Weissman, Roberts, & Woldorff, 2007).

30
31
32 The difference reported in the behavioural results between the two conditions emerged partially also
33
34 for the pupillary data. Specifically, we did not find an overall reduced PD amplitude for the
35
36 auditory stimuli compared to visual stimuli. In addition, we replicated the previous finding in which
37
38 PD was larger in non-corresponding than in corresponding trials in the visual condition.
39
40 Interestingly, we found that the PD Simon effect was present also in the auditory condition but,
41
42 differently from RTs, no difference emerged between the magnitude of the visual and the auditory
43
44 PD Simon effects, suggesting that, probably, the two conditions required the same amount of
45
46 cognitive effort. Along with the results of previous studies (in which visual stimuli were used),
47
48 these results showed that pupil diameter might be used as an indirect marker of the cognitive effort
49
50 induced by conflict monitoring (e.g., Laeng et al., 2011; van Steenbergen & Band, 2013; Wendt et
51
52 al., 2014) independently of the stimulus modality. In addition, the results of the analysis of
53
54 sequential modulations showed that, in both conditions, a Simon effect emerged following a
55
56
57
58
59
60

1
2
3 corresponding trial, whereas a null effect was evident following a non-corresponding trial,
4
5 irrespective of the nature (visual or auditory) of the stimuli. Overall, this analysis revealed a pattern
6
7 of results that is in line with a previous study on the visual Simon effect (van Steenbergen & Band,
8
9 2013), and added new empirical evidences on the conflict-related processing of auditory stimuli.

11 The analysis of the RT distributions showed differences between the visual and auditory
12
13 conditions. More precisely, the visual Simon effect was present for fast RTs (bins 1 to 4) and
14
15 decreased with longer RTs (e.g., Ridderinkhof, 2002; Wascher et al., 2001), whereas the auditory
16
17 Simon effect was present at all bins and, compared to the visual Simon effect, increased with longer
18
19 RTs (e.g., Wascher et al., 2001) showing a longer-lasting effect. Furthermore, the RT distributions
20
21 were differently modulated by previous trial correspondence. Specifically, with visual stimuli, the
22
23 [visual](#) Simon effect following a corresponding trial was present at all bins and decreased with
24
25 increasing RTs, whereas, following a non-corresponding trial, it was absent at the fastest RTs and
26
27 reversed at longer RTs, showing a decreasing function. We interpreted these results in light of the
28
29 activation/inhibition account proposed by Ridderinkhof (2002). That is, the decreasing time course
30
31 evident after both corresponding and non-corresponding trials may have been due to a relatively
32
33 strong selective suppression of the unconditional route. Furthermore, it should be noted that after
34
35 non-corresponding trials the [visual](#) Simon effect reversed at the longer bins. Hence, it seems that the
36
37 inhibition process following non-corresponding trials was stronger compared to that after
38
39 corresponding trials (Ridderinkhof, 2002). However, the finding of a reverse [visual](#) Simon effect
40
41 following a non-corresponding trial could be explained by the fact that complete repetitions of non-
42
43 corresponding trials produce shorter RTs compared to partial repetitions. For auditory stimuli,
44
45 following a corresponding trial, the [auditory](#) Simon effect was present at all bins and increased with
46
47 increasing RTs, whereas, it increased in the first two bins to become stable with longer RTs
48
49 following a non-corresponding trial. This flat time course following non-corresponding trials is in
50
51 line with the account proposed by Stürmer et al. (2002), according to which after a non-
52
53
54
55
56
57
58
59
60

1
2
3 corresponding trial the direct route is suppressed and the response activation process is mediated by
4
5 the indirect route (see also Dittrich et al., 2014; Leuthold & Schröter, 2006).
6

7
8 It is important to note that such qualitatively different processes that have been inferred to exist for
9
10 the two conditions of the Simon task might be re-interpreted in the light of the Diffusion Model for
11
12 Conflict tasks (DMC) recently proposed by Ulrich, Schröter, Leuthold, and Birngruber (2015). The
13
14 model takes into account the idea of two simultaneous processes: one process (i.e., controlled
15
16 process) operates on task-relevant information while the other (i.e., automatic process) operates on
17
18 task-irrelevant information. Through a simulation model, the authors demonstrated that the shape of
19
20 the time course function depends on how the automatic activation spreads out in time. If this
21
22 automatic activation reaches its maximum relatively quickly, the time course will show a decreasing
23
24 trend. If, however, the maximum of the automatic activation is reached relatively late, the resulting
25
26 time course will be increasing. Ulrich and colleagues (2015) combined the concept of diffusion
27
28 processes with the idea of dual processing in conflict tasks to provide a novel account of the delta
29
30 functions. It should be noted that, if we consider the modulation of the RT distribution by previous
31
32 trial correspondence (sequential modulations), the model predicts the reduction of the Simon effect
33
34 when the amplitude of the automatic activation is diminished. However, it does not predict the
35
36 reversal of the Simon effect after a non-corresponding trial for slower responses, a result that we
37
38 found in our visual condition. For this reason, we believe that this model, although interesting,
39
40 needs to be further integrated.
41
42

43
44 In addition, the analysis of PD depending on the RT distributions showed that the **visual** Simon
45
46 effect in PD increased in the second bin to then decrease. The **auditory Simon** effect slightly
47
48 increased at the beginning and then decreased. It is possible that the difference in the time course
49
50 between the two stimulus modalities evident in RT did not emerge in PD because this index does
51
52 not reflect the different strength of activation characterizing the two modalities. Moreover,
53
54 considering the previous trial correspondence, in both conditions, the PD time course was stable
55
56 after both corresponding and non-corresponding trials; however it seems that it reflected the RT
57
58
59
60

1
2
3 distribution, with a larger PD for slower RTs. This result might confirm the role of PD as an index
4
5 of cognitive effort. However, it should be noted that PD, compared to RT, presents very small
6
7 variations and tends to be noisy, and, in this analysis, dividing the number of trial into five bins may
8
9 have decreased the statistical power to detect the effect.
10

11 To conclude, taken together the findings of the present study confirmed the widely accepted
12
13 account that the visual Simon effect can be explained in the framework of the dual-route model of
14
15 information processing. In addition, they seem to support the proposal that the visual and auditory
16
17 Simon effects rely on the same mechanism, with both involving automatic response activation
18
19 through the unconditional route (Leuthold & Schröter, 2006). Although the results of the RT
20
21 distributional analysis are in line with those of Wascher et al. (2001), with a decreasing effect
22
23 function emerging in the visual condition and an increasing effect function emerging in the auditory
24
25 condition, the results of our analysis of the sequential modulations of the RT distributions add more
26
27 evidence to the proposal of a control mechanism that, in both modalities, suppresses the
28
29 unconditional processing route following the detection of a conflict. In other words, our data seem
30
31 to indicate that, even though these modalities seem to differ with respect to the strength of the
32
33 activation and inhibition processes characterizing them, a common mechanism could underlie the
34
35 Simon effect arising in the two modalities (see also Buetti & Kerzel, 2008). Importantly, these
36
37 results were also corroborated by the pupillary measure that, in both conditions mirrored the pattern
38
39 observed in RTs and ERs.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Alnæs, D., Sneve, M. H., Espeseth, T., Endestad, T., van de Pavert, S. H. P., & Laeng, B. (2014). Pupil size signals mental effort deployed during multiple object tracking and predicts brain activity in the dorsal attention network and locus coeruleus. *Journal of Vision, 14*, 1–20. doi:10.1167/14.4.1
- Beatty, J. (1982). Task-evoked pupillary responses, processing load, and the structure of processing resources. *Psychological Bulletin, 91*, 276–292.
- Beatty, J., & Kahneman, D. (1966). Pupillary changes in two memory tasks. *Psychonomic Science, 5*, 371–372.
- Beatty, J., & Lucero-Wagoner, B. (2000). The pupillary system. In J.T. Cacioppo, L.G. Tassinary, & G. Berntson (Eds.), *Handbook of Psychophysiology* (pp. 142–162). Cambridge: Cambridge University Press.
- Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological Review, 108*, 624–652. doi:10.1037//0033-295X.108.3.624
- Buetti, S., & Kerzel, D. (2008). Time course of the Simon effect in pointing movements for horizontal, vertical, and acoustic stimuli: Evidence for a common mechanism. *Acta Psychologica, 129*, 420–428.
- D’Ascenzo, S., Iani, C., Guidotti, R., Laeng, B., & Rubichi, S. (2016). Practice-induced and sequential modulations in the Simon task: evidence from pupil dilation. *International Journal of Psychophysiology*. Epub 6 Agosto 2016. doi: 10.1016/j.ijpsycho.2016.08.002
- De Jong, R., Liang, C.-C., & Lauber, E. (1994). Conditional and unconditional automaticity: A dual-process model of effects of spatial stimulus–response correspondence. *Journal of Experimental Psychology: Human Perception & Performance, 20*, 731–750. doi:10.1037/0096-1523.20.4.731

- 1
2
3 Dittrich, K., Kellen, D., & Stahl, C. (2014). Analyzing distributional properties of interference
4 effects across modalities: chances and challenges. *Psychological Research*, 78, 387–399.
5 doi:10.1007/s00426-014-0551-y
6
7
8
9
10 Egner, T. (2007). Congruency sequence effects and cognitive control. *Cognitive Affective and*
11 *Behavioural Neuroscience*, 7, 380–390.
12
13
14 Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon identification of a target letter
15 in a non search task. *Perception & Psychophysics*, 16, 143–149. doi:10.3758/BF03203267
16
17
18 Hommel, B. (2004). Event files: Feature binding in and across perception and action. *Trends in*
19 *Cognitive Sciences*, 8, 494–500.
20
21
22 Hommel, B., Proctor, R. W., & Vu, K.-P. L. (2004). A feature-integration account of sequential
23 effects in the Simon task. *Psychological Research*, 68, 1–17.
24
25
26
27 Iani, C., Rubichi, R., Gherri, E., & Nicoletti, R. (2009). Co-occurrence of sequential and practice
28 effects in the Simon task: evidence for two independent mechanisms affecting response
29 selection. *Memory & Cognition*, 37, 358–367. doi:10.3758/MC.37.3.358
30
31
32
33 Iani, C., Stella, G., & Rubichi, S. (2014a). Response inhibition and adaptations to response conflict
34 in 6-to 8-year-old children: Evidence from the Simon effect. *Attention, Perception, &*
35 *Psychophysics*, 76, 1234–1241. doi: 10.3758/s13414-014-0656-9
36
37
38
39 Iani, C., Milanese N., & Rubichi, S. (2014b). The influence of prior practice and handedness on the
40 orthogonal Simon effect. *Frontiers in Psychology*, 5, 39. doi:10.3389/fpsyg.2014.00039
41
42
43
44 Kahneman, D., & Beatty, J. (1967). Pupillary responses in a pitch-discrimination task. *Perception*
45 *& Psychophysics*, 2, 101–105. doi:10.3758/BF03210302
46
47
48
49 Kanwisher, N., & Wojciulik, E. (2000). Visual attention: insights from brain imaging. *Nature*
50 *Reviews Neuroscience*, 1, 91–100.
51
52
53
54 Khalid, S., & Ansorge, U. (2013). The Simon effect of spatial words in eye movements:
55 Comparison of vertical and horizontal effects and of eye and finger responses. *Vision*
56 *Research*, 86, 6–14. doi:10.1016/j.visres.2013.04.001
57
58
59
60

- 1
2
3 Kornblum, S., Hasbroucq, T., & Osman, A. (1990). Dimensional overlap: Cognitive basis for
4 stimulus-response compatibility – A model and taxonomy. *Psychological Review*, *97*, 253–
5 270.
6
7
8
9
10 Laeng, B., Ørbo, M., Holmlund, T., & Miozzo, M. (2011). Pupillary Stroop effects. *Cognitive*
11 *Processing*, *12*, 13–21. doi:10.1007/s10339-010-0370-z
12
13
14 Laeng, B., Sirois, S., & Gredebäck, G. (2012). Pupillometry: a window to the preconscious?
15 *Perspectives on Psychological Science*, *7*, 18–27. doi:10.1177/1745691611427305
16
17
18
19 Leuthold, H., & Schröter, H. (2006). Electrophysiological evidence for response priming and
20 conflict regulation in the auditory Simon task. *Brain Research*, *1097*, 167–180.
21 doi:10.1016/j.brainres.2006.04.055
22
23
24
25
26 Loewenfeld, I. (1993). *The pupil: Anatomy, physiology, and clinical applications*. Detroit, MI:
27 Wayne State University Press.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- Lugli, L., Baroni, G., Nicoletti, R., & Umiltà, C.A. (2016). The Simon Effect with saccadic eye movements. *Experimental Psychology*, *63*(2), 107–116, doi: 10.1027/1618-3169/a000319.
- Marini, M., Iani, C., Nicoletti, R., & Rubichi, S. (2011). Between-task transfer of learning from spatial compatibility to a color Stroop task. *Experimental Psychology*, *58*(6), 473–479.
- Mayr, U., Awh, E., & Laurey, P. (2003). Conflict adaptation effects in the absence of executive control. *Nature Neuroscience*, *6*, 450–452.
- Neumann, O., van der Heijden, A. H. C., & Allport, D. A. (1986). Visual selective attention: introductory remarks. *Psychological Research*, *48*, 185–188. doi:10.1007/BF00309082
- Notebaert, W., Gevers, W., Verbruggen, F., & Liefoghe, B. (2006). Top-down and bottom-up sequential modulations of congruency effects. *Psychonomic Bulletin & Review*, *13*, 112–117.
- Notebaert, W., Soetens, E., & Melis, A. (2001). Sequential analysis of a Simon task—Evidence for an attention-shift account. *Psychological Research*, *65*, 170–184.

- 1
2
3 Pashler, H., & Baylis, G. C. (1991). Procedural learning: II. Intertrial repetition effects in speeded-
4 choice tasks. *Journal of Experimental Psychology: Learning, Memory, & Cognition*, *17*, 33–
5 48.
6
7
8
9
10 Pick, D. F., & Proctor, R. W. (1999). Age differences in the effects of irrelevant location
11 information. In M. Scerbo & M. W. Mouloua (Eds.), *Automation technology and human*
12 *performance* (pp. 258–261). Mahwah, NJ: Erlbaum.
13
14
15
16 Posner, M. I., Nissen, M. J., & Klein, R. M. (1976). Visual dominance: An information-processing
17 account of its origins and significance. *Psychological Review*, *83*, 157–171.
18 doi:10.1037/0033-295X.83.2.157
19
20
21
22
23 Proctor, R. W., & Pick, D. F. (1998). Lateralized warning tones produce typical irrelevant-location
24 effects on choice reactions. *Psychonomic Bulletin & Review*, *5*, 124–129.
25 doi:10.3758/BF03209467
26
27
28
29
30 Proctor, R. W., & Vu, K.-P. L. (2006). *Stimulus–response compatibility principle: Data, theory,*
31 *and application*. Boca Raton, FL: Taylor & Francis.
32
33
34 Proctor, R. W., Miles, J. D., & Baroni, G. (2011). Reaction time distribution analysis of spatial
35 correspondence effects. *Psychonomic Bulletin and Review*, *18*, 242–266. doi:10.3758/s13423-
36 011-0053-5
37
38
39
40
41 Ratcliff, R. (1979). Group reaction time distributions and an analysis of distribution statistics.
42 *Psychological Bulletin*, *86*, 446–461. doi:10.1037/0033-2909.86.3.446
43
44
45
46
47 Ridderinkhof, K. R. (2002). Activation and suppression in conflict tasks: Empirical clarification
48 through distributional analyses. In W. Prinz & B. Hommel (Eds.), *Attention and performance*
49 *XIX: Common mechanisms in perception and action* (pp. 494–519). Oxford: Oxford
50 University Press.
51
52
53
54
55 Rubichi, S., Vu, K., Nicoletti, R., & Proctor, R. (2006). Two-dimensional spatial coding.
56 *Psychonomic Bulletin & Review*, *13*, 201–216. doi:3758/BF03193832
57
58
59
60

- 1
2
3 Salzer, Y., Aisenberg, D., Oron-Gilad, T., & Henik, A. (2013). In Touch With the Simon Effect.
4
5 *Experimental Psychology, 61*, 165–179. doi:10.1027/1618-3169/a000236
6
- 7 Simon J. R., & Small, A. M. (1969). Processing auditory information: interference from an
8
9 irrelevant cue. *Journal of Applied Psychology, 53*, 433–435.
10
- 11 Simon, J. R. (1990). The effects of an irrelevant directional cue on human information processing.
12
13 In R.W. Proctor&T. G. Reeve (Eds.), *Stimulus–response compatibility: An integrated*
14
15 *perspective* (pp. 31–86). Amsterdam: North-Holland.
16
- 17 Simon, J. R., & Rudell, A. P. (1967). Auditory S–R compatibility: The effect of an irrelevant cue on
18
19 information processing. *Journal of Applied Psychology, 51*, 300–304. doi:10.1037/h0020586
20
21
- 22 Soetens, E., Maetens, K., & Zeischka, P. (2010). Practice-induced and sequential modulations of the
23
24 Simon effect. *Attention, Perception, & Psychophysics, 72*, 895–911.
25
26 doi:10.3758/APP.72.4.895
27
- 28 Spapé, M. M., Band, G. P. H., & Hommel, B. (2011). Compatibilitysequence effects in the Simon
29
30 task reflect episodic retrieval but not conflict adaptation: Evidence from LRP and N2.
31
32 *Biological Psychology, 88*, 116–123.
33
- 34 Spence, C. J., & Driver, J. (1994). Covert spatial orienting in audition. *Journal of Experimental*
35
36 *Psychology: Human Perception & Performance, 20*, 555–574.
37
38
- 39 Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Experimental Psychology, 18*,
40
41 643–662. doi:10.1037/h0054651
42
43
- 44 Stürmer, B., Leuthold, H., Soetens, E., Schröter, H., & Sommer, W. (2002). Control over location-
45
46 based response activation in the Simon task: Behavioral and electrophysiological evidence.
47
48 *Journal of Experimental Psychology: Human Perception and Performance, 28*, 1345–1363.
49
50 doi:10.1037/0096-1523.28.6.1345
51
52
- 53 Tagliabue, M., Zorzi, M., Umiltà, C., & Bassignani, C. (2000). The role of LTM links and STM
54
55 links in the Simon effect. *Journal of Experimental Psychology: Human Perception and*
56
57 *Performance, 26*, 648–670.
58
59
60

- 1
2
3 Ulrich, R., Schröter, H., Leuthold, H., & Birngruber, T. (2015). Automatic and controlled stimulus
4
5 processing in conflict tasks: Superimposed diffusion processes and delta functions. *Cognitive*
6
7 *Psychology*, 78, 148–174. doi: 10.1016/j.cogpsych.2015.02.005
8
- 9
10 Umiltà, C. A., Rubichi, S., & Nicoletti, R. (1999). Facilitation and interference components in the
11
12 Simon effect. *Archives Italiennes de Biologie*, 137, 139–149.
13
- 14 van Bochove, M., van der Haegen, L., Notebaert, W., & Verguts, T. (2013). Blinking predicts
15
16 enhanced cognitive control. *Cognitive, Affective, & Behavioral Neuroscience*, 13, 346–354.
17
18 doi:10.3758/s13415-012-0138-2
19
- 20 van Steenbergen, H., & Band, G. P. H. (2013). Pupil dilation in the Simon task as a marker of
21
22 conflict processing. *Frontiers in Human Neuroscience*, 7:215. doi:10.3389/fnhum.2013.00215
23
24
- 25 Vu, K.-P. L., Proctor, R. W., & Urcuioli, P. (2003). Transfer effects of incompatible location-
26
27 relevant mappings on a subsequent visual or auditory Simon task. *Memory & Cognition*, 31,
28
29 1146–1152. doi:10.3758/BF03196135
30
- 31 Wascher, E., Schatz, U., Kuder, T., & Verleger R. (2001). Validity and boundary conditions of
32
33 automatic response activation in the Simon task. *Journal of Experimental Psychology: Human*
34
35 *Perception and Performance*, 27, 731–751. doi:10.1037/0096-1523.27.3.731
36
37
- 38 Wendt, M., Kiesel, A., Geringswald, F., Purmann, S., & Fischer, R. (2014). Attentional adjustment
39
40 to conflict strength: evidence from the effects of manipulating flanker-target SOA on response
41
42 times and prestimulus pupil size. *Experimental Psychology*, 61, 55–67. doi:10.1027/1618-
43
44 3169/a000227
45
46
- 47 Wu, C. T., Weissman, D. H., Roberts, K. C., & Woldorff, M. G. (2007). The neural circuitry
48
49 underlying the executive control of auditory spatial attention. *Brain Research*, 1134, 187–198.
50
51 doi:10.1016/j.brainres.2006.11.088
52
53
- 54 Wühr, P. (2005). Evidence for gating of direct response activation in the Simon task. *Psychonomic*
55
56 *Bulletin & Review*, 12, 282–288.
57
58
59
60

1
2
3 Wühr, P. (2006). The Simon effect in vocal responses. *Acta Psychologica*, 121, 210–226.

4
5 doi:10.1016/j.actpsy.2004.12.003

6
7 Wühr, P., & Ansorge, U. (2005). Exploring trial-by-trial modulations of the Simon effect. *The*

8
9 *Quarterly Journal of Experimental Psychology*, 58A, 705–731.

10
11 doi:10.1080/02724980443000269

12
13 Xiong, A., & Proctor, R. W. (2016). Decreasing auditory Simon effects across reaction time

14
15 distributions. *Journal of Experimental Psychology: Human Perception and Performance*, 42,

16
17 23–38. doi: 10.1037/xhp0000117

Figure captions:

Figure 1 Baseline-corrected mean pupil dilation (PD) in the (A) visual and (B) auditory Simon conditions. Time 0 represents the onset of each stimulus. The vertical lines represent the point in time of each condition's mean reaction time. C, corresponding; NC, non-corresponding

Figure 2 Mean reaction time, RT (A), mean Error Rate, ER (B), and mean Pupil Dilation, PD (C), for Current trial correspondence as a function of Previous trial correspondence in the visual and auditory Simon conditions (leftmost and rightmost panel, respectively). Error bars indicate standard errors of the mean. Asterisks denote significant values (* $p < .05$; ** $p < .005$). C, corresponding; NC, non-corresponding

Figure 3 The Simon effect as a function of mean RT for each Bin considering (A) Current trial correspondence, and (B) Previous and Current trial correspondence, for the visual and auditory Simon conditions (leftmost and rightmost panel, respectively). Error bars indicate standard errors of the mean.

Figure 4 The Simon effect as a function of PD depending on RTs distributions for each Bin considering (A) Current trial correspondence, and (B) Previous and Current trial correspondence, for the visual and auditory Simon conditions (leftmost and rightmost panel, respectively). Error bars indicate standard errors of the mean.

APPENDIX

Visual vs. Auditory Simon effect: follow-up experiment using a within-subject design

In this experiment ($N = 24$), we balanced the order of presentation of the visual and the auditory Simon task across participants so that half of the participants performed the visual Simon task as first and the other half performed the auditory Simon task as first. Correct reaction times (RTs), arcsine-transformed error rates (ERs) and mean pupil dilation (PD) in the entire epoch (0-1900 ms) were submitted to a repeated-measures Analysis of Variance (ANOVA) with Condition (visual vs. auditory), Current trial correspondence (corresponding vs. non-corresponding) and Previous trial correspondence (corresponding vs. non-corresponding) as within-participant factors and Order of presentation (visual/auditory vs. auditory/visual) as between-participants factor. In addition, we performed a distributional analysis of RTs. For both the visual and the auditory conditions, individual RTs were rank ordered as a function of Current and Previous trial correspondence and divided into quintiles (bins). The resulting values were submitted to a repeated-measure ANOVA with Condition, Bin (from 1 to 5), Previous trial correspondence and Current trial correspondence as within-participant factors and Order of presentation as between-participants factor.

RTs. The analysis showed a significant main effect of Order of presentation, $F(1, 22) = 37.44$, $MSE = 1777.26$, $p < .001$, $\eta_p^2 = .63$, with faster responses for the auditory/visual order ($M = 382$; $SD = 59$) than for the visual/auditory order ($M = 338$; $SD = 53$). There were also main effects of Current trial correspondence, $F(1, 22) = 228.15$, $MSE = 29230$, $p < .001$, $\eta_p^2 = .91$, with faster responses in corresponding ($M = 348$; $SD = 58$) than in non-corresponding ($M = 373$; $SD = 60$) trials, and Previous trial correspondence, $F(1, 22) = 37.44$, $MSE = 1777.26$, $p < .001$, $\eta_p^2 = .63$, with faster responses after a corresponding ($M = 357$; $SD = 58$) than after a non-corresponding ($M = 363$; $SD = 58$) trial. The main effect of Condition did not reach significance, $F(1, 22) < 1$.

The interaction between Current trial correspondence and Order of presentation was significant, $F(1, 22) = 13.12$, $MSE = 1680.76$, $p < .005$, $\eta_p^2 = .37$. T-tests showed that the difference between corresponding ($M = 367$; $SD = 58$; $M = 329$; $SD = 52$; for visual/auditory and auditory/visual,

1
2
3 respectively) and non-corresponding ($M = 397$; $SD = 56$; $M = 348$; $SD = 53$; for visual/auditory and
4
5 auditory/visual, respectively) trials (i.e., the Simon effect) was of 30 ms when participants
6
7 performed the visual/auditory order ($t_{(11)} = 13.030$, $p_{\text{Bonferroni corrected}} < .001$), and of 19 ms when
8
9 participants performed the auditory/visual order ($t_{(11)} = 8.254$, $p_{\text{Bonferroni corrected}} < .001$). The
10
11 interaction between Previous trial correspondence and Order of presentation was also significant,
12
13 $F(1, 22) = 5.95$, $MSE = 282.45$, $p < .05$, $\eta_p^2 = .21$. T-tests showed that responses after a
14
15 corresponding trial were faster for the auditory/visual order ($M = 337$; $SD = 55$) than for the
16
17 visual/auditory order ($M = 378$; $SD = 63$, $t_{(23)} = 2.549$, $p_{\text{Bonferroni corrected}} < .05$), while responses after
18
19 a non-corresponding trial were faster for the auditory/visual order ($M = 340$; $SD = 52$) than for the
20
21 visual/auditory order ($M = 386$; $SD = 55$), $t_{(23)} = 2.835$, $p_{\text{Bonferroni corrected}} < .05$).
22
23

24
25 The difference between corresponding ($M = 353$; $SD = 52$; $M = 343$; $SD = 63$; for visual and
26
27 auditory conditions, respectively) and non-corresponding trials ($M = 382$; $SD = 47$; $M = 364$; $SD =$
28
29 70 ; for visual and auditory conditions, respectively) (i.e., the Simon effect) was of 28 ms in the
30
31 visual modality and of 21 ms in the auditory modality. These two effects did not differ as indicated
32
33 by the lack of a significant interaction between Current trial correspondence and Condition, $F(1, 22)$
34
35 $= 3.23$, $p = .08$. The interaction between Previous trial correspondence and Condition was
36
37 significant, $F(1, 22) = 8.54$, $MSE = 855.13$, $p < .01$, $\eta_p^2 = .28$. T-tests showed that only in the visual
38
39 condition responses were faster when the preceding trial was corresponding ($M = 362$; $SD = 56$)
40
41 than when it was non-corresponding ($M = 373$; $SD = 46$), $t_{(23)} = 4.976$, $p_{\text{Bonferroni corrected}} < .001$.
42
43 Importantly, the interaction between Current and Previous trial correspondence was significant, $F(1,$
44
45 $22) = 151.74$, $MSE = 24392$, $p < .001$, $\eta_p^2 = .87$. T-tests showed a significant Simon effect of 47 ms
46
47 following a corresponding trial ($t_{(23)} = 16.349$, $p_{\text{Bonferroni corrected}} < .001$) and a null effect of 2 ms
48
49 following a non-corresponding trial ($t_{(23)} = 0.851$, $p = .40$). Crucially, this two-way interaction was
50
51 modulated by Condition, $F(1, 22) = 131.30$, $MSE = 13358$, $p < .001$, $\eta_p^2 = .85$. T-tests showed that
52
53 the visual Simon effect was of 67 ms when the preceding trial was corresponding ($t_{(23)} = 16.261$, p
54
55 $\text{Bonferroni corrected} < .001$) and reversed to -11 ms when the preceding trial was non-corresponding ($t_{(23)}$
56
57 $\text{Bonferroni corrected} < .001$) and reversed to -11 ms when the preceding trial was non-corresponding ($t_{(23)}$
58
59 $\text{Bonferroni corrected} < .001$) and reversed to -11 ms when the preceding trial was non-corresponding ($t_{(23)}$
60

= -2.683, $p_{\text{Bonferroni corrected}} = .05$), while the auditory Simon effect was of 27 ms when the preceding trial was corresponding ($t_{(23)} = 7.879$, $p_{\text{Bonferroni corrected}} < .001$) and decreased to 15 ms when the preceding trial was non-corresponding ($t_{(23)} = 5.623$, $p_{\text{Bonferroni corrected}} < .001$). No other interaction was significant.

ERs. The analysis showed significant main effects of Current trial correspondence, $F(1,22) = 59.09$, $MSE = 0.30$, $p < .001$, $\eta_p^2 = .73$, and Condition, $F(1,22) = 19.07$, $MSE = 0.12$, $p < .001$, $\eta_p^2 = .46$ with higher error rates in non-corresponding ($M = 8.42\%$; $SD = 7.03$) compared to corresponding ($M = 4.72\%$; $SD = 4.20$) trials, and in the visual ($M = 8.09\%$; $SD = 7.36$) compared to the auditory ($M = 5.05\%$; $SD = 3.88\%$) condition.

The interaction between Condition and Order of presentation was significant, $F(1,22) = 5.80$, $MSE = 0.035$, $p < .05$, $\eta_p^2 = .21$, showing that for the visual/auditory order, errors did not differ between the visual ($M = 7.40\%$; $SD = 6.63$) and the auditory ($M = 5.86\%$; $SD = 4.63$) conditions ($t_{(11)} = 2.151$, $p_{\text{Bonferroni corrected}} = .11$), while for the auditory/visual order, participants made more errors in the visual ($M = 8.79\%$; $SD = 7.96$) than in the auditory ($M = 4.23\%$; $SD = 2.72$) condition ($t_{(11)} = 6.766$, $p_{\text{Bonferroni corrected}} < .001$).

The interaction between Current trial correspondence and Condition was significant, $F(1,22) = 8.45$, $MSE = 0.067$, $p < .01$, $\eta_p^2 = .28$. Participants made more errors in non-corresponding than in corresponding trials in both the visual ($M = 11.09\%$; $SD = 8.31$; $M = 5.10\%$; $SD = 4.60$, respectively; $t_{(23)} = 6.533$, $p_{\text{Bonferroni corrected}} < .001$) and auditory condition ($M = 5.75\%$; $SD = 3.91$, $M = 4.3\%$; $SD = 3.72$, respectively; $t_{(23)} = 2.802$, $p_{\text{Bonferroni corrected}} = .04$). In non-corresponding trials, participants made more errors in the visual than in the auditory condition ($t_{(23)} = 6.773$, $p_{\text{Bonferroni corrected}} < .001$), while no difference emerged for corresponding trials ($t_{(23)} = 0.896$, $p = .38$).

The interaction between Previous trial correspondence and Condition was also significant, $F(1,22) = 4.18$, $MSE = 0.014$, $p = .053$, $\eta_p^2 = .16$. In the visual condition, participants made more errors when the previous trial was corresponding ($M = 9.68\%$; $SD = 9.12$) than when it was non-corresponding ($M = 6.51\%$; $SD = 4.48$; $t_{(11)} = -4.996$, $p_{\text{Bonferroni corrected}} < .001$). In the auditory

condition, errors did not differ as a function of previous trial correspondence ($M = 5.06\%$; $SD = 3.78$; $M = 5.04\%$; $SD = 3.98$, respectively; $t_{(11)} = -0.251$, $p = .80$).

Importantly, the interaction between Current and Previous trial correspondence was significant, $F(1,22) = 104.38$, $MSE = 0.43$, $p < .001$, $\eta_p^2 = .83$. T-tests showed that the Simon effect was significant after a corresponding trial (8.3% ; $t_{(23)} = 11.386$, $p_{\text{Bonferroni corrected}} < .001$), but not after a non-corresponding trial (-0.91% , $t_{(23)} = -1.653$, $p = .11$). In addition, the three-way interaction between Current and Previous trial correspondence and Condition was significant, $F(1,22) = 37.68$, $MSE = 0.17$, $p < .001$, $\eta_p^2 = .63$. T-tests showed that the visual Simon effect was positive when the preceding trial was corresponding (13.8% ; $t_{(23)} = 10.909$, $p_{\text{Bonferroni corrected}} < .001$) but did not reach significance when the preceding trial was non-corresponding (-1.82% ; $t_{(23)} = -1.544$, $p = .14$). The auditory Simon effect was significant when the preceding trial was corresponding (2.82% ; $t_{(23)} = 4.748$, $p_{\text{Bonferroni corrected}} < .001$) and it was null when the preceding trial was non-corresponding (-0.0004% ; $t_{(23)} = 0.350$, $p = .73$).

PD. The analysis on mean PD (baseline-corrected) during the entire epoch (0-1900 ms) revealed a significant interaction between Current and Previous trial correspondence, $F(1,22) = 28.49$, $MSE = .008$, $p < .001$, $\eta_p^2 = .56$. T-tests showed that the Simon effect was significant following a corresponding trial (0.016 mm; $t_{(23)} = 3.38$, $p_{\text{Bonferroni corrected}} < .01$), whereas it failed to reach significance following a non-corresponding trial (-0.009 mm; $t_{(23)} = -0.023$, $p_{\text{Bonferroni corrected}} = .094$). This pattern was evident for both conditions, as indicated by the lack of a significant three-way interaction involving the Condition factor, $F(1, 22) = 1.68$, $MSE = 0.00$.

The interaction between Previous trial correspondence and Order of presentation was also significant, $F(1, 22) = 5.77$, $MSE = .003$, $p < .05$, $\eta_p^2 = .21$. T-tests showed that for the visual/auditory order pupil dilation was larger after a corresponding trial ($M = 0.076$ mm; $SD = 0.051$) than after non-corresponding trial ($M = 0.066$ mm; $SD = 0.057$, $t_{(11)} = -2.958$, $p_{\text{Bonferroni corrected}} = .05$). No other main effects or interactions reached significance ($F_s < 1$).

RT distributions. This analysis revealed a significant interaction between Condition, Bin and Current trial correspondence, $F(4, 19) = 25.819$, $MSE = 8851.97$, $p < .001$, $\eta_p^2 = .54$. T-tests revealed that the visual and the auditory Simon effect were significant at all bins ($t_{s(23)} > 2.586$, $p_s < .05$; $t_{s(23)} > 3.337$, $p_s < .001$, respectively). Furthermore, Helmert contrasts showed that the size of the visual Simon effect decreased significantly from bin 1 to bin 5 (39, 38, 31, 22 and 12 ms, respectively), $F_s(1, 23) > 16.245$, $MSEs > 4145.864$, $p_s < .001$, while the size of the auditory Simon effect increased significantly from bin 1 to bin 4 (11, 17, 20 and 27 ms respectively), $F_s(1,23) > 6.409$, $p < .05$, and remained stable from bin 4 to bin 5 (27 and 28 ms, respectively), $F(1, 23) = .12$, $p = .73$.

The interaction between Bin, Previous and Current trial correspondence was significant, $F(4, 19) = 5.938$, $MSE = 650.627$, $p < .001$, $\eta_p^2 = .21$. T-tests revealed that after a corresponding trial, the effect was significant across bins ($t_{s(23)} > 8.765$, $p_s < .001$), whereas after a non-corresponding trial, the effect was close to significance at bin 1 ($t_{(23)} = 1.978$, $p = .06$) and it was not significant in the other bins ($t_{s(23)} < 1.451$, $p_s < .16$). Helmert contrasts showed that the size of the Simon effect, evident after a corresponding trial, was stable from bin 1 to bin 4 (45, 50, 50 and 51 ms, respectively), $F_s(1, 23) > 0.848$, $p_s < .37$, and it decreased significantly from bin 4 to bin 5 (51 and 39 ms, respectively), $F(1, 23) = 21.686$, $p < .001$. Differently, the size of the Simon effect after a non-corresponding trial remained stable from bin 1 to bin 5 (5, 4, 1, -1 and 2 ms, respectively), $F_s(1, 23) > 0.208$, $p_s < .65$.

The interaction between Condition, Bin, Previous and Current trial correspondence did not reach significance, $F(4, 19) < 1$ $p = .94$. For the visual Simon condition, t-tests revealed that after a corresponding trial, the effect was significant across bins ($t_{s(23)} > 7.223$, $p_s < .001$), whereas after a non-corresponding trial, the effect was significant at bins 3, 4 and 5 ($t_s(23) > -2.223$, $p_s < .05$) but not at bins 1 and 2 ($t_s(23) < 0.485$, $p_s > .63$). Helmert contrasts showed that the size of the Simon effect, evident after a corresponding trial, increased significantly from bin 1 to bin 2 and it decreased significantly from bin 2 to bin 5 (76, 77, 73, 66 and 47 ms, respectively), $F_s(1, 23) >$

1
2
3 4.813, $p_s < .05$. Differently, the size of the Simon effect after a non-corresponding trial decreased
4 significantly from bin 1 to bin 4 (2, -2, -10, and -21 ms, respectively), $F_s(1, 23) > 13.170$, $MSEs =$
5 3034.056, $p_s < .001$, but it did not differ significantly between bins 4 and 5 (-21 and -23 ms,
6 respectively), $F(1, 23) = .264$, $MSE = 104.262$. For the auditory Simon condition, t-tests revealed
7 that after a corresponding trial, the Simon effect was significant at all bins ($t_s(23) > 4.16$, $p_s < .001$),
8 whereas after a non-corresponding trial, the effect was significant from bin 2 to bin 5 ($t_s(23) >$
9 2.794, $p_s < .05$) but not at bin 1 ($t_{(23)} = 1.792$, $p = .08$). Furthermore, Helmert contrasts showed that
10 the size of the Simon effect evident after a corresponding trial increased significantly from bin 1 to
11 bin 3 (15, 24 and 28 ms, respectively), $F_s(1, 23) > 3.938$, $MSEs > 1535.79$, $p < .059$, but did not
12 differ among the last three bins (28, 36 and 30 ms, respectively), $F_s(1, 23) > 1.405$, $MSEs >$
13 711.629. Conversely, the size of the Simon effect evident after a non-corresponding trial increased
14 significantly from bin 1 to bin 5 (8, 10, 13, 19 and 26 ms, respectively), $F(1, 23) > 4.091$, $MSE =$
15 1954.35, $p < .055$. Neither the main effect of Order of presentation nor the interactions involving
16 this factor reached significance ($F_s < 1$).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1 Baseline-corrected mean pupil dilation (PD) in the (A) visual and (B) auditory Simon conditions. Time 0 represents the onset of each stimulus. The vertical lines represent the point in time of each condition's mean reaction time. C, corresponding; NC, non-corresponding

78x36mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2 Mean reaction time, RT (A), mean Error Rate, ER (B), and mean Pupil Dilation, PD (C), for Current trial correspondence as a function of Previous trial correspondence in the visual and auditory Simon conditions (leftmost and rightmost panel, respectively). Error bars indicate standard errors of the mean. Asterisks denote significant values (*p < .05; ** p < .005). C, corresponding; NC, non-corresponding

129x104mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3 The Simon effect as a function of mean RT for each Bin considering (A) Current trial correspondence, and (B) Previous and Current trial correspondence, for the visual and auditory Simon conditions (leftmost and rightmost panel, respectively). Error bars indicate standard errors of the mean.

94x55mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4 The Simon effect as a function of PD depending on RTs distributions for each Bin considering (A) Current trial correspondence, and (B) Previous and Current trial correspondence, for the visual and auditory Simon conditions (leftmost and rightmost panel, respectively). Error bars indicate standard errors of the mean.

94x55mm (300 x 300 DPI)