

This is the peer reviewed version of the following article:

C4-plant foraging in Northern Italy: stable isotopes, Sr/Ca and Ba/Ca data of human osteological samples from Roccapelago (16th–18th century AD) / Lugli, Federico; Brunelli, Daniele; Cipriani, Anna; Bosi, Giovanna. - In: ARCHAEOLOGY. - ISSN 0003-813X. - 59:6(2017), pp. 1119-1134. [10.1111/arcm.12295]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

11/01/2026 15:46



C4-plant foraging in Northern Italy: stable isotopes, Sr/Ca and Ba/Ca data of human osteological samples from Roccapelago (16th–18th century AD)

Journal:	Archaeometry
Manuscript ID	ARCH-03-0027-2016.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	30-Aug-2016
Complete List of Authors:	Lugli, Federico; Università degli Studi di Modena e Reggio Emilia, Dipartimento di Scienze Chimiche e Geologiche Brunelli, Daniele; Università degli Studi di Modena e Reggio Emilia, Dipartimento di Scienze Chimiche e Geologiche Cipriani, Anna; Università degli Studi di Modena e Reggio Emilia, Dipartimento di Scienze Chimiche e Geologiche; Columbia University, Lamont-Doherty Earth Observatory Bosi, Giovanna; Università degli Studi di Modena e Reggio Emilia, Dipartimento di Scienze della Vita Traversari, Mirko; Università di Bologna, Dipartimento di Beni Culturali Gruppioni, Giorgio; Università di Bologna, Dipartimento di Beni Culturali
Keywords:	stable isotope analysis, strontium, barium, trace elements, diet, bone, Roccapelago, Italy, Early Modern, C4 plants
Abstract:	Human osteological samples (n = 23) taken from different anatomical parts of 11 individuals from the post-medieval (16th to 18th century AD) site of Roccapelago (Modena, Italy) were systematically analysed for $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and trace elements to investigate their diet. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ correlate and show a high variability between individuals, attesting for the dietary contribution of C4-plants. This is supported by pollen analysis of the burial site samples, which reveal the presence of maize. $\delta^{15}\text{N}$ correlates with Sr/Ca suggesting that the main protein source could have been milk and dairy. We therefore interpret the strong correlation between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ as the evidence of C4-plant foraging practice and exploitation of livestock for meat and milk combined with possible direct intake of C4-plants. The Roccapelago site represents an important case study to track post-medieval diet evolution and maize cultivation introduction in southern Europe as attested also by historical sources.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

**C₄-plant foraging in Northern Italy: stable isotopes, Sr/Ca and Ba/Ca data
of human osteological samples from Roccapelago (16th–18th century AD)**

Federico Lugli^{1,*}, Daniele Brunelli¹, Anna Cipriani^{1,2}, Giovanna Bosi³, Mirko Traversari⁴,
Giorgio Gruppioni⁴

¹ Department of Chemical and Geological Sciences, University of Modena and Reggio Emilia,
Modena, Italy.

² Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY, USA.

³ Department of Life Sciences, University of Modena and Reggio Emilia, Modena, Italy.

⁴ Department of Cultural Heritage, University of Bologna, Ravenna, Italy.

* Corresponding author

E-mail: federico.lugli@unimore.it

ABSTRACT

Human osteological samples ($n = 23$) taken from different anatomical parts of 11 individuals from the post-medieval (16th to 18th century AD) site of Roccapelago (Modena, Italy) were systematically analysed for $\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and trace elements to investigate their diet. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ correlate and show a high variability between individuals, attesting for the dietary contribution of C_4 -plants. This is supported by pollen analysis of the burial site samples, which reveal the presence of maize. $\delta^{15}\text{N}$ correlates with Sr/Ca suggesting that the main protein source could have been milk and dairy. We therefore interpret the strong correlation between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ as the evidence of C_4 -plant foraging practice and exploitation of livestock for meat and milk combined with possible direct intake of C_4 -plants. The Roccapelago site represents an important case study to track post-medieval diet evolution and maize cultivation introduction in southern Europe as attested also by historical sources.

KEYWORDS

stable isotope analysis; strontium; barium; trace elements; diet; bone; Roccapelago; Italy; Early Modern; C_4 plants

INTRODUCTION

Roccapelago was a secluded community (16th to 18th century AD) located in the Northern Apennines (Fig. 1) that practised animal husbandry and farming, but also exploited anthropic woods of chestnut as highlighted in historical documents. In 2011, excavation under the St. Paul's Conversion church in Roccapelago (Pievepelago district, Modena) brought to light a forgotten crypt used as a cemetery by the community between the 16th and the 18th centuries, where human bodies were massed in a pyramidal cumulus, and included infants, subadults and adults (Gruppioni et al. 2011). Some of these bodies are naturally (totally or partially) mummified due to the peculiar microclimate of the cryptal environment characterized by low humidity and intense aeration (Gruppioni et al. 2011), excluding external contamination and ensuring negligible diagenetic effects. In Northern Italy, the Roccapelago site is a unique case of natural mummification and a rare opportunity to reconstruct three centuries of the life, customs and traditions of a farmer community, which has lived during the arrival of maize in Europe (Rebourg et al. 2003). Several studies have followed this discovery including DNA (Cilli et al. 2015a), gut microbiota (Cilli et al. 2015b), clothing and artefacts (Schoenholzer Nichols 2016; Vernia 2016), palaeopathology (Traversari et al. 2016) and palaeoradiology (Petrella et al. 2016). However, little is known regarding the eating habits and detailed practices related to the diet of the inhabitants of Roccapelago (Pelù 2006; Traversari et al. 2016).

Stable isotope analyses of bone collagen are among the currently most widely used techniques applied to palaeodiet studies to unravel the type of diet and food sources of human and animals (e.g. DeNiro and Epstein 1981; DeNiro and Schoeniger 1983; Katzenberg et al. 1993; Ambrose et al. 1997). The isotope composition of bone collagen quantitatively reflects the isotope composition of the food ingested during the lifetime of the individual/animal (Richards et al. 2006). The most exploited isotopes for this purpose are $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$,

expressed as $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ per mil variation relative to two international standards (vPDB and AIR respectively).

$\delta^{13}\text{C}$ is used to distinguish C_3 from C_4 protein in the diet, because of the different carbon fixation of C_3 (such as wheat, rice, barley, rye and potato; $\delta^{13}\text{C}$ from -20 to -35‰) and C_4 -plants (such as maize, sorghum and millet; $\delta^{13}\text{C}$ from -9 to -14‰) during photosynthesis. Because carbon reservoirs (dissolved carbonate) in marine environments show a $\delta^{13}\text{C} \approx 0\text{‰}$ (atmospheric $\delta^{13}\text{C} \approx -7\text{‰}$), $\delta^{13}\text{C}$ values can also be used to discriminate marine vs. terrestrial resources in the diet, with the more negative values for terrestrial provenance (Katzenberg 2008). In terrestrial ecosystems, an offset of *c.a.* 5‰ in $\delta^{13}\text{C}$ is observed between the herbivore and the plants they consume; an offset of *c.a.* 1‰ occurs between a carnivore and its herbivore prey.

$\delta^{15}\text{N}$ values reflect the protein portion of the diet, given the lack of nitrogen in carbohydrates and lipids. Thus, $\delta^{15}\text{N}$ depends on the position of the individual in the trophic chain, with an enrichment of $+3\text{--}5\text{‰}$ for each step because of the preferential fixation of heavier molecules in living tissues. The length of aquatic trophic chains is the cause of the high $\delta^{15}\text{N}$ values in these environments, especially for the top predators (Hedges and Reynard 2007; Katzenberg 2008). Likewise, $\delta^{13}\text{C}$ shows a stepwise enrichment along the trophic chain of *c.a.* 1‰ , but this is generally too low to be detected.

Stable isotope variations of human collagen are generally explained as the result of two food sources mixed together modelled using simple linear mixing models with two end-members, chosen between the major food sources offered by the local environment. Thus, it is important to investigate the isotopic composition not only of human remains but also of the local fauna and flora. In more complicated contexts, where more than two food sources are part of the human diet it might be appropriate to use non-linear or bayesian mixing models (Phillips et al. 2014).

Trace elements have been used for decades to study the diet of individuals with mixed results

1 due to diagenetic effects on their abundances and to the fact that the behaviour of many trace
2 elements does not reflect the eating habits of the individual (Elias 1982; Pate and Hutton
3 1988; Pate et al. 1989; Tuross et al. 1989; Klepinger 1990; Pate 1994; Burton and Wright
4 1995; Ezzo 1994; Burton et al. 1999; Schutkowski et al. 1999; Balter et al. 2002; Hedges
5 2002; Mays 2003; Dolphin et al. 2005; Burton 2008; Arnay-De-La-Rosa et al. 2011; Austin et
6 al. 2013; L  sch et al. 2014). This is especially true for archaeological samples that sit in the
7 ground for years to thousands of years interacting with the chemicals dispersed in the soil and
8 with those transported by rain and ground water. In this regard, Roccapelago represents a
9 unique situation, where the dry and ventilated environmental conditions of the closed burial
10 site (crypt) led to the natural mummification of the individuals and kept them away from the
11 interaction with soil and water. In general, the exchange of Sr^{2+} (and other trace elements)
12 ions between soil and bone occurs in low-pH environment, where the hydroxyapatite (more
13 soluble at low-pH) interacts with soil solutions and underground waters. Hence, any
14 precipitation of secondary calcium phosphate may lead to the incorporation of trace elements
15 from both the soil and the bone tissue (Pate et al. 1989). Nevertheless, several works have
16 shown the reliability of Sr/Ca and Ba/Ca ratios as dietary proxies (e.g. Burton and Wright
17 1995; Burton et al. 1999; Balter et al. 2002). Strontium physiology and behaviour along the
18 food chain are well constrained (e.g. Burton and Wright 1995; Burton et al. 1999; Mays 2003;
19 Bentley 2006; L  sch et al. 2014). Divalent Sr^{2+} cations introduced in the body with food tend
20 to replace calcium (Ca^{2+}) in the hydroxyapatite of bones because of the close chemical and
21 physical behaviour of the two elements. Sr has no metabolic functions within the body and,
22 therefore, its concentration in bones is not regulated by homeostasis. However, Ca is
23 preferentially assimilated with respect to Sr during biological activities, a process known as
24 *biopurification* (Burton et al. 1999). This is because, the shift of Sr from the digestive system
25 to the bloodstream is less efficient than that of Ca, hence, only *ca.* 20% of the strontium
26 ingested is absorbed and fixed in bones. Therefore, the Sr/Ca ratio tends to decrease from low
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

to high trophic positions, reflecting this preferential assimilation of Ca. Furthermore, Burton and Wright (1995) demonstrated that, in a multi-component diet, Sr/Ca does not show a linear relationship with the abundance of consumed meat/plant. For this reason, they suggest that the Sr/Ca ratio better reflects the highest Ca source rather than the proportion of meat vs. plants in a multicomponent diet (Burton and Wright 1995; L \ddot{o} sch et al. 2014).

Barium behaves similarly to strontium (Burton 2008). However, because of its larger ionic radius Ba is less efficiently fixed in bones than Sr and, as a consequence, its biopurification degree is higher, resulting in lower Ba concentrations in bones compared to Sr (Burton 2008). Yet, because of their comparable metabolic behaviour, Ba can also be used as a marker of a Ca-source in a multi-component diet (Burton and Wright 1995). Since both indices (Ba/Ca and Sr/Ca) carry the same information about the diet, a strong correlation between the two is expected in a biological integer sample (Burton et al. 1999). On the contrary, when the seafood is the main dietary component, the Ba behaviour is quite different from Sr, causing a lack of correlation between the Ba/Ca and Sr/Ca ratios (Burton and Price 1991).

In this work, we analyzed the trace element composition (Sr/Ca and Ba/Ca), the $\delta^{13}\text{C}$ and the $\delta^{15}\text{N}$ of bones and teeth of 11 individuals from the Roccapelago site. The main aim of this study was to detect the food components and changes with time of their diet comparing collagen data from bone and dentine with the idea of possibly identifying the introduction of C_4 plants (likely maize) into the diet, given that this community lived during the arrival of maize from America to Europe and the spread of the same to Italy (Cazzola 1991; Rebourg et al. 2003). While bone collagen reflects the latest years of life because of constant bone remodelling, primary dentine, once formed, does not physiologically remodel and, therefore, records the childhood eating habits (Lee-Thorp 2008). Unfortunately, no animal remains were found in the Roccapelago site because of the strictly mortuary function of the context and it has not been possible to compare the trace element and isotopic ratios of human and local animals. However, the high variability of the diet found in studied individuals, gave us the

1 opportunity to infer at least two principal food sources and to reconstruct the probable local
2 foraging practices.
3
4
5
6
7
8
9

10 Archaeological and historical background of the Roccapelago site

11
12
13
14

15
16
17
18
19
20
21
22
23 Roccapelago is a village located near the small town of Pievepelago (Frignano area, Modena
24 Apennines) on the bank of the Scoltenna river, in the centre of the Pelago Valley (Fig. 1). The
25 most important and relevant building in Roccapelago is the Church of the Conversion of St.
26 Paul. The building was erected at the end of the 14th century, as a military fortification by
27 Obizzo di Montegarullo. Founded on the bare rock, the structure is located on a high peak
28 dominating the surrounding area (Gruppioni et al. 2011). In 1585, it lost all of its military
29 functions and became a proper Christian church. Between October 2009 and March 2011,
30 parallel to building restoration, the excavation of the site under the church by the
31 Archaeological Superintendence of Emilia Romagna brought a hidden funerary crypt to light.
32 This funeral context, dated through archaeological artefacts from the late 16th century to the
33 18th century, consists of bodies massed in a pyramidal cumulus, probably dropped through a
34 manhole from the church floor (Gruppioni et al. 2011). The presence of two small windows in
35 the crypt and the pyramidal arrangement of the bodies allowed drainage of decomposition
36 fluids and, consequently, a partial to total mummification of some individuals. The crypt bears
37 no traces of exogenous fluid flows suggesting no modification of the environment occurred
38 since the burial. The bodies belong to a wide age range of individuals (from infants to elders)
39 of both genders (Gruppioni et al. 2011), and from their clothing it has been inferred that they
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

were all common people from a rural community mostly dedicated to farming (Schoenholzer Nichols 2016). Therefore, the mummies of Roccapelago represent a significant discovery that allows studying the peasant practices and rural life in a post-medieval era (16th to 18th century AD).

MATERIALS AND METHODS

A total of 23 human samples (n = 7 tooth samples; n = 16 bone samples) were recovered from different anatomical areas of 11 individuals, chosen from the best preserved within the 281 (minimum number of individuals) of the site (Table 1). Given some restrictions in accessing the site, we could only sample a limited number of individuals and have chosen only among those individuals articulated in anatomical connection, so we can exclude an oversampling of the same individual. We stress also that these remains were retrieved from a crypt, with no interaction with soil and/or water.

Teeth were extracted with tongs from the maxilla or the mandible. Bone samples were separated from humerus and femur or tibia (when femur was absent) with a diamond wheel saw. Samples were split in two parts: one for trace element analyses and the other for stable isotope analyses. Both splits were cleaned mechanically with brushes. Bones samples for trace element analysis were cut with a diamond low-speed wheel saw in a 2 mm slice in order to expose the inner parts. Each slice was polished with alumina powder and abrasive papers at decreasing sizes: 800, 500, 5 and 1 µm, then rinsed with MilliQ[®] water and washed with 2 cycles of ultrasonic bath (15 min) with MilliQ[®] water. Samples were then dried for 48h at room temperature (protocol modified from Cucina et al. 2007).

Samples for stable isotope analysis were treated for collagen extraction. First, the samples were mechanically cleaned with a power dentist drill and crushed to obtain 150-200 mg of powder. Bone and dentin chunks were left in a 0.5 M HCl solution for 72h at 4°C (Longin 1971; Ambrose 1990; Bocherens et al. 1991; Iacumin et al. 2014). After 48h, HCl was replaced with new HCl. Samples were then washed with deionized water and left in a 0.125 M NaOH solution for 20h at room temperature. Residuals were washed again with deionized water, to reach neutrality, and put in a drying oven for 17h at 100°C. At last, the centrifuged supernatant was frozen and lyophilized. Stable isotope analysis of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were run and duplicated with a Carlo Erba EA1110 CHN analyzer, coupled to a Finnigan Delta S mass spectrometer. Bone and dentine samples were analyzed in order to obtain information about any possible variation in the diet of the studied individuals, from juvenile to adult age.

In situ Sr, Ba, Pb, Mg, Zn, Fe and Cu concentration of bone samples were measured at the Centro Interdipartimentale Grandi Strumenti (CIGS) of the University of Modena e Reggio Emilia using a 213 nm Nd:YAG laser ablation system (NewWave Research) coupled to a quadrupole ICP-MS (Thermo Fisher Scientific X-Series^{II}). NIST 1486 Bone Meal (powder) was prepared as a pellet under a manual press and used as reference material. Data were acquired for ^{88}Sr and ^{138}Ba and data reduction was obtained with the Plasma Lab software. Calcium concentrations, obtained at CIGS by an ESEM Quanta 200 in low vacuum, were used as internal standard for the correction of matrix-related effects in samples (Hanč et al. 2013). Cortical tissue in bone samples was investigated with 5 random ablation points per sample. Before analysis, samples were carefully pre-ablated to clean further the external surface, reducing any possible contamination. In order to ensure the accuracy of the method, a two-tailed *t*-test was performed on two certified elements of the standard 1486 Bone Meal: Zn and Sr. Statistical analysis showed no significant difference between expected and measured values ($p < 0.01$; mean of 3 analyses). The Ba concentration is not certified for NIST 1486, we therefore used data reported by Porte et al. (1997). The typical RSD error for laser ablation

trace element analyses is *c.a.* 10%.

Pollen analysis was performed in two ground samples collected from the crevices of a buried wall of the crypt contemporaneous with the latter mummies present in the site (18th century). The two soil assays where sampled, after manual removing of the exposed surface to avoid modern pollen contamination, from the most external part of the crevice, i.e. the closest to the room, and from the deeper part of the crevice between two stone blocks. Samples were treated according to the routine method in use at the Laboratory of Palynology and Palaeobotany of Modena (Florenzano et al. 2012). Cerealia pollen was identified under 1000x plain light microscope based on Beug (2004) and Faegri et al. (1989), with correction factor for glycerol jelly. About 500 pollen grains per samples were counted.

RESULTS

Stable isotopes

Data for bones and dentine for each individual are presented in Table 1 and Table 2.

$\delta^{13}\text{C}$ of bone samples ranges from -20.1 to -12.2‰ . $\delta^{15}\text{N}$ values are also highly variable, ranging from 5.5 to 9.5‰ (Table 1). For each individual, both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ from dentine show values quite similar to their respective bone samples (from -19.6 to -13.7‰ for $\delta^{13}\text{C}$ and from 5.5 to 8.9‰ for $\delta^{15}\text{N}$; Table 2). The quality of the data depends on the conservation state of collagen, that is evaluated based on the collagen yield and in modern bones is about

22 wt %. The mean of collagen yield for individuals from Roccapelago is 21.0 wt % \pm 2.9 (1 σ), very close to modern bones. Carbon and nitrogen can also be expressed as wt % of the combusted extracted collagen. Intact collagen is about 35 wt % for C and ranges between 11 wt % and 16 wt % for N. For the individuals here studied, C wt % ranges from 34 to 42 wt %, while N from 13 to 16 wt %. No C:N ratio falls outside the 2.9–3.6 range reported in the literature (Ambrose 1990; Van Klinken 1999). Routinely 1 σ -uncertainties of standards are \pm 0.2‰ for $\delta^{13}\text{C}$ and \pm 0.1‰ for $\delta^{15}\text{N}$.

Sr/Ca and Ba/Ca

Significant trace element ratios for diet studies of the Roccapelago individuals are presented in Table 1. However, for comparative purposes, we present in the Supplementary information other trace element data collected simultaneously during laser ablation ICP–MS analysis that, however, are less reliable and more problematic for palaeodiet reconstructions (Supplementary table 1).

The Sr/Ca [$\mu\text{g}/\text{mg}$] ratio in bone samples ranges from 0.45 to 0.79 and from 0.47 to 0.79 considering individual averages. The Ba/Ca [$\mu\text{g}/\text{mg}$] ratio ranges from 0.004 to 0.071 and from 0.004 to 0.065 considering individual averages (Table 1).

Pollen

About 40% of the recognized taxa are represented by pollens pertaining to entomophilous plants (e.g. roses, cornflowers, poppies among others). These plants are characterized by beautiful flowers, thus representing possible memories of offerings during burial rites. The pollen spectra also reflect the image of the neighbouring oak woods (with chestnut, deciduous *Quercus*, hazel), along with beech and conifers (including silver fir). In addition, our data show the presence of pastures (as suggested by high percentages of Cichorioideae and Poaceae wild grass group) and cereal fields (*Avena*/*Triticum* group, *Hordeum* group, *Secale cereale* and *Zea mays*). Cultivated fields might have been very close to the site, in accordance with geomorphological features observed on the western side of the site suitable for crop cultivations (Bosi et al. *in press*)

DISCUSSION

History and botany

The economic profile that characterized the population of Roccapelago was strongly influenced by the particular geographical situation, at least in the last 400 years. Located about 1200 m.a.s.l. between forests, Roccapelago has always used the sylvan environment for

1 subsistence, using the communication routes to move goods, animals and people (Pelù 2006).
2
3 Within the Frignano area, husbandry of cattle for milk and workforce and ovine (mainly
4 sheep) for milk and wool was a common practice; moreover, many pigs were bred for meat
5 and poultry for meat, eggs and feathers (Bellettini, 2012). One of the most important plant
6 species for the inhabitants of Roccapelago was the chestnut, extending to about 1000 m.a.s.l.,
7 on top of which were usually built shelters for shepherds and charcoal burners, including
8 species such as beech and oak. The cultivation of cereals (mainly barley and rye) was
9 moderately present with corn appearing in the historical cadastral records (also called *estimi*)
10 only during the 19th century (Cazzola 1997).
11
12 The archaeobotanical analysis of two ground samples from the crypt shows the presence of
13 maize pollen (Fig. 1). In Europe, maize is a relevant chronological marker (16th cent.);
14 moreover, it is one of the few outcrossing plants among the major cereals (Rebourg et al.
15 2003). In spite of the high pollen production of *Zea*, particularly significant in the oldest
16 varieties (estimated range pollen quantum per tassel; 14-50 million vs. 2-5 million in modern
17 hybrids), the atmospheric dispersion of maize pollen is quite narrow even in presence of
18 strong winds (Vogler et al. 2009). In fact, the large size of *Zea* pollen grains and their rapid
19 settling rate influence the airborne pollen dispersal. It has been demonstrated that at a distance
20 of 60 m from the source plant, maize pollen concentrations average about 1% of those at 1 m
21 and the quantity of pollen remaining airborne at 60 m is 5% of that one at 1 m (Raynor et al.
22 1972). The presence of maize in the two studied samples can be explained by the arrival of its
23 pollen through the two windows of the crypt (aerial transport), also favoured by the high
24 pollen production of the oldest landraces and suggesting cultivations close to the crypt itself.
25 We cannot exclude, however, that maize pollen could have arrived by unintentional anthropic
26 transport, perhaps through shrouds and clothes of the deceased.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Stable isotopes, Sr/Ca and Ba/Ca ratios

The high variability of the individual $\delta^{13}\text{C}$ values (-20.1 to -12.2‰ ; Fig. 2) suggests that at least two different sources have contributed to the Roccapelago inhabitant's diet: one with low $\delta^{13}\text{C}$ close to the range of C_3 plants (-23 to -32‰) and one with higher $\delta^{13}\text{C}$ in the range of C_4 plants (-10 to -16‰).

The variability of $\delta^{15}\text{N}$ (5.5 to 9.5‰) is smaller than $\delta^{13}\text{C}$ but reflects the combination of two different sources in the diet with different protein content (Fig. 2). The difference between the higher (9.5‰) and the lower (5.5‰) $\delta^{15}\text{N}$ is *c.a.* 4‰ , corresponding to an entire trophic step.

The high positive correlation between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ ($R^2 = 0.89$; $p < 0.01$) is usually used as a marker for marine fish consumption (e.g. Richards and Hedges 1999; Fornaciari 2008). However, within the Roccapelago community we exclude the consumption of aquatic resources. $\delta^{15}\text{N}$ is quite low when compared with the commonly high values typical of aquatic protein consumers (Richards and Hedges 1999; Hedges and Reynard 2007; Fornaciari 2008; Fuller et al. 2012) and none of the individuals cross the terrestrial–marine cut-off of 12.5‰ reported by Salamon et al. (2008) for an Italian medieval sample (Reitsema and Vercellotti 2012). Moreover, strong historical–archaeological evidences of fish consumption or presence are yet to be found. Fornaciari (2008) describes the diet of two Renaissance courts (Medici Gran Duke in Florence and Aragonese Princes in Naples) where individuals consumed fish and high protein foods (Fig. 3). Their $\delta^{15}\text{N}$ values are more positive (from *c.a.* 10 to 14‰) than those of Roccapelago, suggesting a much higher protein intake. $\delta^{13}\text{C}$ values are on average more negative (from *c.a.* -19 to -16.5‰) and this shift of $\delta^{13}\text{C}$ values from the classical C_3 environment range is interpreted as the effect of marine resources consumption

(Fornaciari 2008). The more negative $\delta^{15}\text{N}$ and more positive $\delta^{13}\text{C}$ values of Roccapelago inhabitants with respect to Medici and Aragonese lead us to a different interpretation of the correlation between the two variables. We suggest that the strong correlation in Roccapelago results from the intake of two protein sources from different trophic levels with different C_3 - C_4 proportion. In particular, the high $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ food source is identifiable with the greater C_4 protein portion. One of the possible explanations for this enrichment in both C_4 portion and protein intake could be the practice of C_4 -plant foraging and the exploitation of meat and dairy from C_4 -fed livestock (bovines or caprovines; Galli 1980) that leads to an increase of $\delta^{13}\text{C}$ in human collagen.

Few cereal plants with C_4 photosynthetic pathways are known in Emilia-Romagna during the 18th century: maize (*Zea mays*), sorghum (*Sorghum bicolor*) and millet (*Panicum miliaceum* and *Setaria italica*). Maize, in particular, is the most important from a historical point of view. Introduced in Emilia-Romagna and in the Po delta territories between the end of 16th century and the first half of 17th century, this cereal supplanted all the other local crops in northern Italy at the beginning of 17th century and had its vastest diffusion during the first half of the 18th century (Cazzola 1991). The recovery of *Zea* pollen in the layer contemporaneous to the mummies indicates that the individuals cannot be prior to the last decades of the 16th century AD. Historical sources testify for a spread of corn in the Este Dukedom since the 16th century. During the first half of the 17th century, maize became an important element of the rural economy (Cazzola 1991). In Italy and central Europe, both millet and maize were used as main food (Cortonesi 2002), especially in poor community during time of famine, and as animal forage. Parmentieri (1791) states that “cows eat maize forage greedily, and it makes them yield a lot of milk” (Barrière et al. 2006). Young (1792) says that farms can be categorized in “good” and “bad”, in relation to the presence or absence of maize in animal feeding (Barrière et al. 2006). Both historical observations are in support of the C_4 -plant feeding hypothesis.

From a geochemical point of view, we inferred the possible collagen stable isotope values of the Roccapelago livestock (Fig. 2) using our most enriched individuals (more positive $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of human collagen) as proxy of a nearly total animal protein diet. Taking into account a fractionation of 1‰ for the carbon and of 4‰ for the nitrogen, we obtain a livestock $\delta^{13}\text{C}$ of c.a. -13.2‰ and a $\delta^{15}\text{N}$ of c.a. 5.5‰ . We then extrapolate the possible livestock fodder composition in terms of C_3 and C_4 proportion, using a simple linear interpolation with two end-members (Tykot 2006). The $\delta^{13}\text{C}$ values of the two end-members are taken from Iacumin et al. (2014), with a $\delta^{13}\text{C}$ of -24.6‰ for the C_3 plants and a $\delta^{13}\text{C}$ of -10.4‰ for the C_4 plants. To correct for the diet-collagen offset, we consider an enrichment of $+5\text{‰}$ (Tykot 2006). Thus, our Roccapelago individuals fit in a model with a livestock fodder likely composed by 55% of C_3 plants and 45% of C_4 plants.

Individual depleted $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values reflect an almost total vegetarian diet, with very little intake of high-protein foods. Their primary vegetal food source can be reliably identified in local cultivated C_3 -cereals and forest products.

Stable isotopes of collagen primarily reflect the protein portion of the diet, for this reason we can not exclude the direct use of C_4 plants in the Roccapelago diet. This signal can be masked by the highest protein source that also consists of a high C_4 portion. In fact, high protein animal-derived food is the highest contributor to the collagen isotopic composition (Tykot 2006).

We do not observe any particular cluster in relation to sex or age, although the sample of our analyses is too small to provide a significant statistic. The differences between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of bone and dentine are in general very low. Taken together, all bone and dentine data do not show any specific trend in the protein intake or in the C_3 - C_4 proportion. This highlights an almost unchanged diet during the growth of the individual. Considering the values individually, however, we suspect a small variation in the eating habits for four individuals. Roc11 shows a $\Delta^{13}\text{C}_{\text{den-bone}}$ of -1.2‰ and a $\Delta^{15}\text{N}_{\text{den-bone}}$ of -0.6‰ , suggesting a lower intake of

protein during childhood. Roc10 shows an unchanged $\Delta^{13}\text{C}_{\text{den-bone}}$ (0.1‰) but a $\Delta^{15}\text{N}_{\text{den-bone}}$ of -0.9‰, probably related to a very small decrease in protein intake not detected by the $\delta^{13}\text{C}$ value or possibly to a little variation in the protein source mixing. Conversely, both Roc5 and Roc2 show an increase in $\delta^{15}\text{N}$ of *c.a.* 1‰ and a correspondent increase in $\delta^{13}\text{C}$ (*c.a.* 0.5‰), testifying a likely small increase in the intake of protein foods (Table 2).

Strontium and barium in bones derive from food ingestion and drinking water, as well as from possible diagenetic contamination, particularly in bone samples. However, the peculiarity of the archaeological context of Roccapelago, where bones were not directly buried in the ground and away from direct contact with ground and meteoric water and the natural mummification of the corpses with muscles still in place in some cases, let us assume that the Sr and Ba variabilities can be related to *in vivo* processes rather than post-burial diagenesis (cf. Pate and Hutton 1988; Pate et al. 1989). Moreover, the strong correlation ($R^2 = 0.83$; $p < 0.01$) between Sr/Ca and Ba/Ca (Fig. 4c), suggests that the samples are biologically integer, as demonstrated also in studies of Burton and Price (1991) and Burton et al (1999).

Sr/Ca individual values were compared with $\delta^{15}\text{N}$ to observe any possible correlation between the two variables. If we hypothetically assume an almost mono-component diet for the studied individuals, the correlation between the Sr/Ca ratio and the $\delta^{15}\text{N}$ can be interpreted as a mere trophic effect, recorded in both variables (Fig. 4a). However, in the case of a multi-component diet, the strong negative correlation ($R^2 = 0.61$; $p < 0.01$; 1 outlier removed) between individual Sr/Ca ratio and $\delta^{15}\text{N}$ may suggest that the protein intake and the major Ca-source were linked. Individuals with a lower Sr/Ca ratio, and a consequent highly biopurified-Ca diet, show also a higher protein diet (high $\delta^{15}\text{N}$; Fig. 4a). The diet of these individuals was probably composed by Ca-enriched or Sr-depleted foods, such as milk and dairy, in agreement with the livestock exploitation hypothesis. A similar correlation can be observed between Ba/Ca and $\delta^{15}\text{N}$ ($R^2 = 0.68$; $p < 0.01$; 1 outlier removed; Fig. 4b). Individuals presenting the highest Sr/Ca ratio probably used to eat low-Ca and/or high-Sr foods like cereals and legumes

(Schutkowski et al. 1999). One outlier value, removed from the previous regressions (solid black diamond in Fig. 4), is represented by a single individual with Sr/Ca ratio of 0.79, Ba/Ca ratio of 0.063 and $\delta^{15}\text{N}$ value of 9.0‰. This individual shows a higher Sr/Ca ratio than the other individuals with a similar $\delta^{15}\text{N}$. This may indicate that the main protein portion of the diet of this individual was represented by a Ca-depleted but high-protein food, like meat. However, no differences in clothes or ornaments was found in this individual to indicate that he was from a different rank in the society with a possibly different high-protein diet.

Given that Ba behaves quite differently from Sr in case of a seafood-based diet (Burton et al. 1999), the observed strong correlation between Ba/Ca and Sr/Ca ($R^2 = 0.83$; $p < 0.01$; Fig. 4c) confirms the lack of seafood in the diet of Roccapelago studied inhabitants as already showed by the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ data. The only individual that seems to not perfectly fit the regression is, again, the outlier mentioned before (Roc8); in fact, removing this individual from the fit, we observe an even stronger correlation ($R^2 = 0.92$; $p < 0.01$; 1 outlier removed). Such evidence corroborates the hypothesis of a different main protein-source for Roc8.

A closer look at the correlation of $\delta^{15}\text{N}$ with $\delta^{13}\text{C}$ (Fig. 2) shows the presence of sub-clusters of individuals. In particular, individuals Roc1, 2, 3, 4, 5, and 7 (group 1) plot together at low $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values; whereas individuals Roc6, 9 and 10 (group 2) and individuals Roc8 and 11 (group 3) cluster at higher $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$. Group 1 and group 3 define the two end-members of the correlation with group 2 lying in the middle between the two extremes. In the correlations of $\delta^{15}\text{N}$ with Sr/Ca and Ba/Ca and of Sr/Ca with Ba/Ca (Fig. 4), these clusters are fairly maintained. In fact, group 1 is still well defined in all plots while group 2 and 3 plot together at higher $\delta^{15}\text{N}$ and lower Sr/Ca and Ba/Ca values. Roc9 departs from group 2 and plots now in the middle of the correlations, while Roc8 becomes a total outlier following the group with the Ca-depleted diet (Group 1). These clustering of individuals can be interpreted as an ulterior proof of the diet-related significance of these ratios for the Roccapelago case study and, in general, an indication that the combined use of Sr, Ba and stable isotopes can

push the archaeological community further in understanding past eating habits.

CONCLUSIONS

Stable isotopes and trace element ratios can be a powerful tool to examine not only the diet of ancient population but also diet-related practices as the foraging of livestock. We interpret the isotopic trends shown in this case study (high correlation between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) as the mixing of at least two main food sources in the human diet, represented by C_3 -plants (lowest protein source) and C_3 - C_4 -fed livestock (highest protein source). In Roccapelago, C_4 -plants were probably used as forage for the livestock, but we cannot exclude a direct C_4 -plants intake, quite common in this kind of rural community. Sr/Ca and Ba/Ca ratios reveal that the main source of protein was likely represented by milk and dairy, in agreement with the exploitation of livestock, and no evidence of seafood.

ACKNOWLEDGMENTS

The authors would like to thank Giorgia Biviano, Alessio Zedde and Antonino Vazzana (University of Bologna, Ravenna Campus) for their precious help with bone sampling and

analysis; Emanuele Paganelli and Massimo Tonelli (University of Modena and Reggio Emilia – CIGS) for assistance with the analytical work; Antonietta Di Matteo and Paola Iacumin (University of Parma) for help with stable isotope analyses; Marta Bandini Mazzanti for help with the pollen analyses. Geochemical work supported by the Programma Giovani Ricercatori Rita Levi Montalcini to AC. Finally, we would like to thank two anonymous reviewers for their comments and suggestions.

REFERENCES

Ambrose, S. H., 1990, Preparation and Characterization of Bone and Tooth Collagen for Isotopic Analysis, *Journal of Archaeological Science*, **17**, 431–51.

Ambrose, S. H., Butler, B. M., Hanson, D. B., Hunter-Anderson, R. L., and Krueger, H. W., 1997, Stable Isotopic Analysis of Human Diet in the Marianas Archipelago, Western Pacific, *American Journal of Physical Anthropology*, **104**, 343–61.

Arnay-De-La-Rosa, M., González-Reimers, E., Yanes, Y., Romanek, C. S., Noakes, J. E., Galindo-Martín, L., 2011, Paleonutritional and paleodietary survey on prehistoric humans from Las Cañadas del Teide (Tenerife, Canary Islands) based on chemical and histological

analysis of bone, *Journal of Archaeological Science*, **38**, 884–895.

Austin, C., Smith, T. M., Bradman, A., Hinde, K., Joannes-Boyau, R., Bishop, D., Hare, D. J., Doble, P., Eskenazi, B., Arora, M., 2013, Barium distributions in teeth reveal early-life dietary transitions in primates, *Nature*, **498**, 216–219.

Balter, V., Bocherens, H., Person, A., Labourdette, N., Renard, M., Vandermeersch, B., 2002, Ecological and physiological variability of Sr/Ca and Ba/Ca in mammals of West European mid-Würmian food webs. *Palaeogeography Palaeoclimatology Palaeoecology*, **186**, 127–143.

Barrière, Y., Alber, D., Dolstra, O., Lapierre, C., Motto, M., Ordas, a., Van Waes, J., Vlaswinkel, L., Welcker, C., and Monod, J. P., 2006, Past and prospects of forage maize breeding in Europe. II. History, germplasm evolution and correlative agronomic changes, *Maydica*, **51**(3-4), 435–49.

Bellettini M. C., 2012. Il rapporto uomo-ambiente negli Statuti del Frignano (secoli XIV-XVI), In *Statuti del Frignano degli anni MCCCXXXVII-VIII. Vol. 1. Testo e studi*, (eds. A. Sorbelli, and F. Iacoli), 325–390, Iaccheri Editore, Pavullo, Modena.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Bentley, R. A., 2006, Strontium isotopes from the earth to the archaeological skeleton: A review, *Journal of Archaeological Method and Theory*, **13**(3), 135–87.

Beug, H. J., 2004, Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete, Verlag, München.

Bocherens, H., Fizet, M., Mariotti, A., Lange-Badré, B., Vandermeersch, B., Borel, J. P., Bellon, G. 1991, Isotopic biogeochemistry (^{13}C , ^{15}N) of fossil vertebrate collagen: implications for the study of fossil food web including Neandertal Man, *Journal of Human Evolution*, **20**, 481–92.

Bosi, G., Rinaldi, R., Torri, P., Bandini Mazzanti, M., *in press*, Informazioni botaniche dalla Cripta cimiteriale di Roccapelago, In *Le mummie di Roccapelago (XVI-XVIII sec.): vita e morte di una piccola comunità dell'Appennino Modenese. Archeologia e antropologia: una ricerca interdisciplinare*, Accademia dello Scoltenna, Pievepelago, Modena.

Burton, J. H., Price, T. D., 1991, Paleodietary applications of barium values in bone, In *Proceedings of the 27th International Symposium on Archaeometry Heidelberg, 1990*, (eds. E. Pernicka, and G. A. Wagner), 787–95, Berkhäuser-Verlag, Basel.

Burton, J. H., and Wright, L. E., 1995, Nonlinearity in the Relationship Between Bone Sr/Ca and Diet: Paleodietary Implications, *American Journal of Physical Anthropology*, **96**, 273–82.

Burton, J. H., Price, D. T., and Middleton, W. D., 1999, Correlation of Bone Ba/Ca and Sr/Ca due to Biological Purification of Calcium, *Journal of Archaeological Science*, **26**, 609–16.

Burton, J. H., 2008, Bone Chemistry And Trace Element Analysis, In *Biological Anthropology of the Human Skeleton*, second edition (eds. M. A. Katzenberg, and S. Saunders), 443–460, John Wiley & Sons, Hoboken, New Jersey.

Cazzola, F., 1991, L'introduzione del mais in Italia e la sua utilizzazione alimentare (sec. XVI-XVIII), In *La préparation alimentaire des céréales* (eds. S. Fournier, and F. Sigaut), 109–27, PACT, Centro Universitario Europeo per i Beni Culturali.

Cazzola, F., 1997, La ricchezza della terra. L'agricoltura emiliana fra tradizione e innovazione, In *Storia d'Italia. Le regioni dall'unità a oggi* (eds. R. Finzi), 53-123, Einaudi, Torino.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Cilli, E., De Fanti, S., Quagliariello, A., Sarno, S., Serventi, P., Traversari, M., Zedde, A., Luiselli, D., Gruppioni, G., 2015a, Genetic analysis of the population of Roccapelago - Modena (Italy) (16th - 18th c.), In *ArchaeoAnalytics. Chromatography and DNA analysis in archaeology*, 247–254, Esposende.

Cilli, E., De Filippo, C., Albanese, D., Lugli, F., Sordo, M., Viola, M.F., Traversari, M., Catalano, G., Serventi, P., De Fanti, S., Labate, D., Cipriani, A., Luiselli, D., Gruppioni, G., 2015b, Discovering past gut microbiomes through NGS analysis: the mummies of Roccapelago (MO), In *Abstract Book of the VI Congress of the Italian Society for Evolutionary Biology*, 58, EdiSES, Bologna.

Cortonesi, A., 2002, Agricoltura e tecniche nell'Italia medievale. I cereali, la vite, l'olivo, In *Uomini e campagne nell'Italia medievale* (ed. A. Cortonesi), 191-270, Laterza, Bari.

Cucina, A., Dudgeon, J., Neff, H., 2007, Methodological strategy for the analysis of human dental enamel by LA-ICP-MS, *Journal of Archaeological Sciences*, **34**, 1884–1888.

DeNiro, M., and Epstein, S., 1981, Influence of diet on the distribution of nitrogen isotopes in animals, *Geochimica et Cosmochimica Acta*, **45**(3), 341–51.

DeNiro, M. J., and Schoeniger, M. J., 1983, Stable Carbon and Nitrogen Isotope Ratios of Bone Collagen: Variations Within Individuals, Between Sexes, and Within Population Raised on Monotonous Diets, *Journal of Archaeological Science*, **10**, 199–203.

Dolphin, A. E., Goodman, A. H., Amarasiriwardena, D. D., 2005, Variation in elemental intensities among teeth and between pre- and postnatal regions of enamel, *American Journal of Physical Anthropology* **128**, 878–88.

Elias, R. W., Hirao, Y., Patterson, C. C., 1982, The circumvention of the natural biopurification of calcium along nutrient pathways by atmospheric inputs of industrial lead, *Geochimica et Cosmochimica Acta*, **46**, 2561- 2580.

Faegri, K., Kaland, P. E., Krzywinski, K., 1989, Textbook of Pollen Analysis, John Wiley & Sons, New York.

Fornaciari, G., 2008, Food and disease at the Renaissance courts of Naples and Florence: a paleonutritional study, *Appetite*, **51**(1), 10–4.

Fuller, B. T., Muldner, G., Van Neer, W., Ervynck, A., and Richards, M. P., 2012, Carbon and nitrogen stable isotope ratio analysis of freshwater, brackish and marine fish from Belgian archaeological sites (1st and 2nd millenium AD), *Journal Of Analytical Atomic Spectrometry*, **27**(5), 807.

Grupponi, G., Labate, D., Mercuri, L., Milani, V., Traversari, M., Vernia, B., 2011, Gli scavi della Chiesa di San Paolo di Roccapelago nell’Appennino Modenese: la cripta con i corpi mummificati naturalmente, *Pagani e Cristiani. Forme e attestazioni di religiosità del mondo antico in Emilia*, **10**, 219–45.

Hanć, A., Olszewska, A., and Barańkiewicz, D., 2013, Quantitative analysis of elements migration in human teeth with and without filling using LA-ICP-MS, *Microchemical Journal*, **110**, 61–9.

Hedges, R.E.M., 2002, Bone diagenesis: an overview of processes, *Archaeometry*, **44**, 319–328.

Hedges, R. E. M., and Reynard, L. M., 2007, Nitrogen isotopes and the trophic level of humans in archaeology, *Journal of Archaeological Science*, **34**(8), 1240–51.

Iacumin, P., Galli, E., Cavalli, F., and Cecere, L., 2014, C4-consumers in southern Europe: the case of Friuli V.G. (NE-Italy) during early and central Middle Ages., *American journal of physical anthropology*, **154**(4), 561–74.

Katzenberg, M. A., Saunders, S. R., and Fitzgerald, W. R., 1993, Age differences in stable carbon and nitrogen isotope ratios in a population of prehistoric maize horticulturists., *American journal of physical anthropology*, **90**(3), 267–81.

Katzenberg, M. A., 2008, Stable Isotope Analysis: A Tool For Studying Past Diet, Demography, And Life History, In *Biological Anthropology of The Human Skeleton*, second edition (eds. M. A. Katzenberg, and S. Saunders), 413–41, John Wiley & Sons, Hoboken, New Jersey.

Klepinger, L. L., 1990, Magnesium Ingestion and Bone Magnesium Concentration in

Paleodietary Reconstruction: Cautionary Evidence from an Animal Model, *Journal of Archaeological Science*, **17**, 513–7.

Lee-Thorp, J. A., 2008, On isotopes and old bones, *Archaeometry*, **50**(6), 925–50.

Longin R., 1971, New method of collagen extraction for radiocarbon dating, *Nature*, **230**, 241–42.

Lösch, S., Moghaddam, N., Grossschmidt, K., Risser, D. U., and Kanz, F., 2014, Stable Isotope and Trace Element Studies on Gladiators and Contemporary Romans from Ephesus (Turkey, 2nd and 3rd Ct. AD) - Implications for Differences in Diet., *PloS ONE*, **9**(10), e110489.

Mays, S., 2003, Bone strontium: calcium ratios and duration of breastfeeding in a Mediaeval skeletal population, *Journal of Archaeological Science*, **30**(6), 731–41.

Parmentier, A.A., 1791, Le maïs ou le blé de Turquie apprécié sous tous ses rapports, Imprimerie Impériale, Paris.

Pate, F.D., Hutton, J.T., 1988, The use of soil chemistry data to address post-mortem diagenesis in bone mineral, *Journal of Archaeological Science*, **15**, 729–739.

Pate, F.D., Hutton, J.T., Norrish, K., 1989, Ionic exchange between soil solution and bone: toward a predictive model, *Applied Geochemistry*, **4**, 303–316.

Pate, F.D., 1994, Bone chemistry and paleodiet, *Journal of Archaeological Method and Theory*, **1**, 161–209.

Pelù, P., 2006, Rapporti tra costa tirrenica e la bassa padana attraverso l'Appennino tosco-emiliano nel tardo Medioevo, In *Viabilità, traffici, Commercio, Mercati e fiere in Garfagnana dall'antichità all'unità d'Italia*, (ed. P. L. Raggi), 105–22, Aedes Muratoriana, Modena.

Petrella, E., Piciucchi, S., Feletti, F., Barone, D., Piraccini, A., Minghetti, C., Gruppioni, G., Poletti, V., Bertocco, M., Traversari, M., 2016, CT Scan of Thirteen Natural Mummies Dating Back to the XVI-XVIII Centuries: An Emerging Tool to Investigate Living Conditions and Diseases in History, *PLoS ONE*, **11**, e0154349.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Phillips, D. L., Inger, R., Bearhop, S., Jackson, A. L., Moore, J. W., Parnell, A. C., Semmens, B. X., and Ward, E. J., 2014, Best practices for use of stable isotope mixing models in food-web studies, *Canadian Journal of Zoology*, **92**, 823–35.

Porte, N., Mauerhofer, E., and Denschlag, H. O., 1997, Test of multielement analysis of bone samples using instrumental neutron activation analysis (INAA) and anti-Compton spectrometry, *Journal of Radioanalytical and Nuclear Chemistry*, **224**, 103–7.

Raynor, G. S., Ogden, E. C., Hayes, J. V., 1972, Dispersion and Deposition of Corn Pollen from Experimental Sources, *Agronomy Journal* **64**(4), 420–427.

Reitsema, L. J., and Vercellotti, G., 2012, Stable isotope evidence for sex- and status-based variations in diet and life history at medieval Trino Vercellese, Italy, *American Journal of Physical Anthropology*, **148**(4), 589–600.

Rebourg, C., Chastanet, M., Gouesnard, B., Welcker, C., Dubreuil, P., Charcosset, A., 2003, Maize introduction into Europe: the history reviewed in the light of molecular data,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Theoretical and Applied Genetics, **106**, 895–903.

Richards, M. P., and Hedges, R. E. M., 1999, Stable Isotope Evidence for Similarities in the Types of Marine Foods Used by Late Mesolithic Humans at Sites Along the Atlantic Coast of Europe, *Journal of Archaeological Science*, **26**(6), 717–22.

Richards, M. P., Fuller, B. T., and Molleson, T. I., 2006, Stable isotope palaeodietary study of humans and fauna from the multi-period (Iron Age, Viking and Late Medieval) site of Newark Bay, Orkney, *Journal of Archaeological Science*, **33**(1), 122–31.

Salamon, M., Coppa, A., McCormick, M., Rubini, M., Vargiu, R., and Tuross, N., 2008, The consilience of historical and isotopic approaches in reconstructing the medieval Mediterranean diet, *Journal of Archaeological Science*, **35**, 1667–72.

Schoenholzer Nichols, T., 2016, Confronto fra alcune camicie di Roccapelago e Monsampolo del Tronto, In *Roccapelago e le sue mummie: studio integrato della vita di una piccola comunità dell'Appennino tra XVI e XVIII secolo* (ed. F. Badiali), 259–268, Grafiche Sigem Srl, Modena.

Schutkowski, H., Herrmann, B., Wiedemann, F., Bocherens, H., and Grupe, G., 1999, Diet, Status and Decomposition at Weingarten: Trace Element and Isotope Analyses on Early Mediaeval Skeletal Material, *Journal of Archaeological Science*, **26**(6), 675–85.

Tykot, R. H., 2006, Isotope Analyses and the Histories of Maize, In *Histories of Maize* (eds. J. E. Staller, R. H. Tykot, and B. F. Benz), 131–42, Elsevier Academic Press.

Traversari, M., Feletti, F., Vazzana, A., Gruppioni, G., Frelat, M. A., 2016, Three cases of developmental dysplasia of the hip on partially mummified human remains (Roccapelago, Modena, 18th Century): a study of palaeopathological indicators through direct analysis and 3D virtual models, *BMSAP*, DOI: 10.1007/s13219-015-0140-7.

Tuross, N., Behrensmeyer, A.K., Eanes, E.D., 1989, Strontium increases and crystallinity changes in taphonomic and archaeological bone, *Journal of Archaeological Science*, **16**, 661–672.

Van Klinken, G. J., 1999, Bone Collagen Quality Indicators for Palaeodietary and Radiocarbon Measurement, *Journal of Archaeological Science*, **26**, 687–95.

Vernia, B., 2016, Le testimonianze devozionali: le medaglie rinvenute negli scavi di Roccapelago, In *Roccapelago e le sue mummie: studio integrato della vita di una piccola comunità dell'Appennino tra XVI e XVIII secolo* (ed. F. Badiali), 197–204, Grafiche Sigem Srl, Modena.

Vogler, A., Wettstein-Bättig, M., Aulinger-Leipner, I., Stamp, P., 2009, The airborne pollen flow of maize (*Zea mays* L.) in a multi-crop designed field plot, *Agricultural and Forest Meteorology*, **149**, 1776–80.

Young, A., 1792, *Travels in France*, Richardson, W. (ed.), London.

Figure 1. (a) Location of the village of Roccapelago (Modena) in Emilia-Romagna, Italy. (b) *Zea mays* pollen grain observed in the soil samples from Roccapelago crypt; scale: 10 μm – photos with Leica MC 120 HD at 1000X magnifications (by P. Torri). (c) Photo of the Conversion of St. Paul Church.


Figure 2. (a) Collagen stable isotopes of the Roccapelago individuals (black diamond) and fields of food end-members. C3 and C4 plant $\delta^{13}\text{C}$ values are from Iacumin et al. (2014) where $\delta^{15}\text{N}$ values are inferred from human data taking into account the trophic position. Plausible mixed C3-C4 fed livestock (bovines or caprovines) position is deduced from human and plant data. α (\pm standard error) is the intercept and β (\pm standard error) is the slope of the regression line. Circles outline clusters of individuals as explained in the text. (b) In the inset, the $\delta^{13}\text{C}$ value of a mammal with a pure C3-diet, a pure C4-diet and the modelled Roccapelago livestock mixed diet. Based on our data we extrapolate the $\delta^{13}\text{C}$ value of the Roccapelago livestock to be composed of a 45% C4 diet and 55% C3 diet.

Figure 3. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ comparison between Roccapelago, Medici and Aragonese (Fornaciari 2008) and Mainizza (Iacumin et al. 2014) diet. In dark gray the field of Roccapelago individuals. In light gray the field of the Mainizza settlement. Dashed field represents Medici and Aragonese families. In bold, two of the possible diet end-members for each site. Medici (16th–17th AD) and Aragonese (15th–17th AD) were italian noble families, Mainizza (10th–11th AD) is a rural settlement in Friuli V.G., NE Italy.

Figure 4. Correlations of $\delta^{15}\text{N}$ with Sr/Ca and Ba/Ca ratios and of Sr/Ca with Ba/Ca ratios in bones of Roccapelago individuals. Black diamond is the outlier Roc8 not considered in the regressions (see text for details). Labels represent the number of individuals (IDs from Table 1), wording “Roc” is omitted to improve the readability of the graph. Circles outline clusters


of individuals as explained in the text.

For Peer Review



(a) Location of the village of Roccapelago (Modena) in Emilia-Romagna, Italy. (b) Zea mays pollen grain observed in the soil samples from Roccapelago crypt; scale: 10 µm – photos with Leica MC 120 HD at 1000X magnifications (by P. Torri). (c) Photo of the Conversion of St. Paul Church.


Fig. 1
149x111mm (300 x 300 DPI)



(a) Collagen stable isotopes of the Roccipelago individuals (black diamond) and fields of food end-members. C3 and C4 plant $\delta^{13}\text{C}$ values are from Iacumin et al. (2014) where $\delta^{15}\text{N}$ values are inferred from human data taking into account the trophic position. Plausible mixed C3-C4 fed livestock (bovines or caprovines) position is deduced from human and plant data. α (\pm standard error) is the intercept and β (\pm standard error) is the slope of the regression line. Circles outline clusters of individuals as explained in the text. (b) In the inset, the $\delta^{13}\text{C}$ value of a mammal with a pure C3-diet, a pure C4-diet and the modelled Roccipelago livestock mixed diet. Based on our data we extrapolate the $\delta^{13}\text{C}$ value of the Roccipelago livestock to be composed of a 45% C4 diet and 55% C3 diet.


Fig. 2

135x116mm (300 x 300 DPI)



$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ comparison between Roccapelago, Medici and Aragonese (Fornaciari 2008) and Mainizza (Iacumin et al. 2014) diet. In dark gray the field of Roccapelago individuals. In light gray the field of the Mainizza settlement. Dashed field represents Medici and Aragonese families. In bold, two of the possible diet end-members for each site. Medici (16th–17th AD) and Aragonese (15th–17th AD) were italian noble families, Mainizza (10th–11th AD) is a rural settlement in Friuli V.G., NE Italy.

Fig.3
154x116mm (300 x 300 DPI)



Correlations of $\delta^{15}\text{N}$ with Sr/Ca and Ba/Ca ratios and of Sr/Ca with Ba/Ca ratios in bones of Roccapelago individuals. Black diamond is the outlier Roc8 not considered in the regressions (see text for details). Labels represent the number of individuals (IDs from Table 1), wording "Roc" is omitted to improve the readability of the graph. Circles outline clusters of individuals as explained in the text.

Fig. 4

153x121mm (300 x 300 DPI)

Table 1. Results of stable isotopes, Sr/Ca and Ba/Ca of human bone samples from Roccapelago.												
ID	Sex	Age	Body element	Bone sample data					Individual mean			
				$\delta^{13}\text{C}_{(\text{PDB}-1)}$	$\delta^{15}\text{N}_{(\text{AIR})}$	C:N	Sr/Ca ($\mu\text{g}/\text{mg}$)	Ba/Ca ($\mu\text{g}/\text{mg}$)	$\delta^{13}\text{C}_{(\text{PDB}-1)}$	$\delta^{15}\text{N}_{(\text{AIR})}$	Sr/Ca	Ba/Ca
Roc1	M	Adult	Femur	−19.2	6.2	2.9	0.71	0.070	−19.5	6.3	0.69	0.064
			Humerus	−19.8	6.5	3.0	0.68	0.058				
Roc2	M	Adult	Femur	−20.1	6.0	3.1	0.65	0.065	−19.9	6.1	0.63	0.065
			Humerus	−19.7	6.2	3.0	0.62	0.066				
Roc3	F	Adult	Humerus	−18.8	7.2	3.2	0.69	0.064	−18.8	7.2	0.69	0.064
Roc4	I	Adult	Femur	−19.9	5.6	2.9	0.65	0.062	−19.9	5.6	0.65	0.062
Roc5	I	Adult	Humerus	−19.7	5.5	3.0	0.69	0.060	−19.7	5.5	0.69	0.060
Roc6	F	Adult	Tibia	−16.4	8.4	3.2	0.47	0.012	−16.4	8.4	0.47	0.012
Roc7	M	Adult	Tibia	−19.3	6.6	3.0	0.58	0.043	−19.3	6.5	0.60	0.057
			Humerus	−19.3	6.4	2.9	0.63	0.071				
Roc8	F	Adult	Humerus	−12.2	9.0	3.1	0.79	0.063	−12.2	9.0	0.79	0.063
Roc9	F	Adult	Humerus	−16.4	8.1	2.9	0.55	0.040	−16.4	8.1	0.55	0.040
Roc10	M	Adult	Femur	−17.1	7.2	2.9	0.56	0.009	−17.0	7.5	0.44	0.007
			Humerus	−16.9	7.7	3.0	0.33	0.006				
Roc11	M	Adult	Femur	−12.6	9.7	3.1	0.45	0.004	−12.5	9.5	0.47	0.004
			Humerus	−12.5	9.3	3.1	0.48	0.004				

Table 2. Stable isotopes of human tooth samples from Roccapelago.

ID	Tooth	Dentine $\delta^{13}\text{C}_{(\text{PDB}-1)}$	Dentine $\delta^{15}\text{N}_{(\text{AIR})}$	C:N	$\Delta\text{C}_{\text{dentine-bone}}$	$\Delta\text{N}_{\text{dentine-bone}}$
Roc1	Canine	-19.2	6.8	3.1	0.3	0.5
Roc2	Canine	-19.5	7.2	3.0	0.5	1.1
Roc4	Premolar	-19.3	5.5	3.4	0.6	-0.1
Roc5	Premolar	-19.1	6.5	3.0	0.6	0.9
Roc7	Premolar	-19.6	6.8	3.0	-0.3	0.3
Roc10	Molar (1 st)	-16.8	6.6	3.1	0.1	-0.9
Roc11	Premolar	-13.7	8.9	2.9	-1.2	-0.6