

This is the peer reviewed version of the following article:

4Ts Score and EuroSCORE in cardiac surgery / Mattioli, Anna Vittoria; Manenti, Antonio; Farinetti, Alberto.
- In: JOURNAL OF THROMBOSIS AND THROMBOLYSIS. - ISSN 0929-5305. - 45:2(2018), pp. 291-292.
[10.1007/s11239-017-1604-z]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

08/01/2026 15:02

Dear Author,

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For fax submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style. Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL: [http://dx.doi.org/\[DOI\]](http://dx.doi.org/[DOI]).

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information go to: <http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

ArticleTitle	4Ts Score and EuroSCORE in cardiac surgery	
Article Sub-Title		
Article CopyRight	Springer Science+Business Media, LLC, part of Springer Nature (This will be the copyright line in the final PDF)	
Journal Name	Journal of Thrombosis and Thrombolysis	
Corresponding Author	Family Name	Mattioli
	Particle	
	Given Name	Anna Vittoria
	Suffix	
	Division	Department of Surgical, Medical and Dental, Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine
	Organization	University of Modena and Reggio Emilia
	Address	Via del pozzo, 71 41100, Modena, Italy
	Phone	0039/59/4224043
	Fax	
	Email	annavittoria.mattioli@unimore.it
	URL	
	ORCID	http://orcid.org/0000-0003-1487-9530
Author	Family Name	Manenti
	Particle	
	Given Name	Antonio
	Suffix	
	Division	Department of Surgical, Medical and Dental, Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine
	Organization	University of Modena and Reggio Emilia
	Address	Via del pozzo, 71 41100, Modena, Italy
	Phone	
	Fax	
	Email	
	URL	
	ORCID	
Author	Family Name	Farinetti
	Particle	
	Given Name	Alberto
	Suffix	
	Division	Department of Surgical, Medical and Dental, Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine
	Organization	University of Modena and Reggio Emilia
	Address	Via del pozzo, 71 41100, Modena, Italy
	Phone	
	Fax	
	Email	
	URL	
	ORCID	

Email
URL
ORCID

Schedule	Received
	Revised
	Accepted

Footnote Information	This is a Letter to article https://doi.org/10.1007/s11239-017-1592-z .
----------------------	---



LETTER TO THE EDITOR

4Ts Score and EuroSCORE in cardiac surgery

Anna Vittoria Mattioli¹ · Antonio Manenti¹ · Alberto Farinetti¹

© Springer Science+Business Media, LLC, part of Springer Nature 2017

Dear Sir,

We have read with great interest the paper “Initial and long term impact of a multi-disciplinary task force in the diagnosis and management of Heparin-Induced thrombocytopenia” by Lim et al. [1] and we found their conclusion of importance with a view to clinical prevention.

The authors underline how diagnosis and management of HIT has been changed after a retrospective audit on patients who underwent at least one laboratory testing for HIT.

The audit demonstrated a high frequency of PF4-ELISA testing (n = 54, 66%) in patients that had low 4T scores. Also, a very low percentage of PF4-ELISA-positive cases were confirmed to be SRA positive. If the 4T scoring criteria were initially used prior to PF4-ELISA testing, 54 (65.8%) PF4-ELISAs and 24 (55.8%) SRAs would not have been performed, resulting in a cost savings of \$11,850 [2].

On the basis of these data, Authors evaluated the initial and long-term impact and challenges of these institution-wide changes in the diagnosis and management of HIT in the inpatient setting at an academic medical center. They concluded that, despite their efforts, the number of PF4-ELISA testing was already higher than other academic medical centers of similar size [1]. In order to reduce test and to decrease costs they plan to create an electronic order set for 4T score for ELISA-PF4 testing.

With reference to the findings reported in the paper, we would like to make the following contribution to the discussion. We refer the results of our retrospective analysis in post-cardiac surgery patients about the incremental value

of 4Ts Score and EuroSCORE for prediction of heparin-induced thrombocytopenia.

The analysis was performed in a group of patients treated with unfractionated heparin undergoing cardiac surgery. Anti-PF4/heparin antibodies were tested in all patients using a commercial immunoassay (Asserachrom H PF4 ELISA kits) [3–5]. Of the 600 patients investigated, 131 (21.8%) were found to have anti-PF4/heparin antibodies in the post-operative period (5–7 days from surgery). All patients had the EuroSCORE [6] value and the 4Ts Score [7] evaluated before surgery. As expected the 4Ts Score identify as high probability patients that developed antibodies and we found 22 thrombotic events (16.79%). Within the group of patients that tested negative to anti-PF4/heparin antibodies, the incidence of thrombotic complications was lower (5%) (p > 0.01).

A 4T Score > 3 identified medium and high risk and predicted an increased risk of thrombosis in patients independently of the EuroSCORE value and add information for stratification. In these patients the ELISA test is suggested in order to identify the presence of antibodies. This group had a greater prevalence of antibodies and a greater prevalence of thrombotic complications at 30 days.

A low EuroSCORE value was associated with an exceedingly low hard-event rate (0.4% year) that increases significantly as a function of the 4T's score results. The 4Ts Score yielded incremental value for the prediction of hard events (X 56–88, p < 0.001) and significantly stratified patients. In patients with intermediate to high likelihood of complications as predicted by EuroSCORE the 4Ts Score significantly increased the prediction of events.

Our results suggested that cardiac surgery patients are at high risk for coagulation adverse effects, among which the generation of anti-PF4/heparin antibodies. This usually causes a late post-operative thrombocytopenia, and, when persisting, thrombotic complications at different levels [8–10].

The evaluation of the risk of patients had to include the 4Ts Score and the EuroSCORE in order to identify patients more likely to developed anti-PF4/heparin antibodies and

This is a Letter to article <https://doi.org/10.1007/s11239-017-1592-z>.

✉ Anna Vittoria Mattioli
 annavittoria.mattioli@unimore.it

¹ Department of Surgical, Medical and Dental, Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Via del pozzo, 71 41100 Modena, Italy

thrombotic complications. These patients need a more careful follow-up and the ELISA test for anti-PF4/heparin antibodies could help in the management of post-operative anticoagulant therapy. On the other side patients with low risk could avoid PF4/ELISA test. We support the hypothesis by Lim et al. that a HIT task force can reduce the number of inappropriate HIT testing with significant cost savings.

Compliance with ethical standards

Conflict of interest Anna Vittoria Mattioli, Antonio Manenti and Alberto Farinetti declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

- Lim MY, Foster J, Greenberg CS (2017) Initial and long term impact of a multi-disciplinary task force in the diagnosis and management of heparin-induced thrombocytopenia. *J Thromb Thrombolysis*. <https://doi.org/10.1007/s11239-017-1592-z>
- Pierce W, Mazur J, Greenberg C, Mueller J, Foster J, Lazarchick J (2013) Evaluation of heparin-induced thrombocytopenia (HIT) laboratory testing and the 4Ts scoring system in the intensive care unit. *Ann Clin Lab Sci* 43:429–435
- Mattioli AV, Bonetti L, Sternieri S, Mattioli G (2000) Heparin-induced thrombocytopenia in patients treated with unfractionated heparin: prevalence of thrombosis in a 1 year follow-up. *Ital Heart J* 1:39–42
- Warkentin TE, Heddle NM (2003) Laboratory diagnosis of immune heparin-induced thrombocytopenia. *Curr Hematol Rep* 2:148–157
- Mattioli AV, Bonetti L, Carletti U, Ambrosio G, Mattioli G (2009) Thrombotic events in patients with antiplatelet factor 4/heparin antibodies. *Heart* 95(16):1350–1354. <https://doi.org/10.1136/hrt.2008.160549>
- Nashef SAM, Roques F, Michel P, Gauducheau E, Lemeshow S, Salamon R, for the EuroSCORE study group (1999) European system for cardiac operative risk evaluation (EuroSCORE). *Eur J Cardio-thorac Surg* 16:9–13
- Lee Y, Weeks P, Gass J, Sieg A (2017) Evaluation of 4T's scoring system in the identification of heparin-induced thrombocytopenia in patients with mechanical circulatory support. *Thromb Res* 160:66–68. <https://doi.org/10.1016/j.thromres.2017.10.017>
- Mattioli AV, Bonetti L, Zennaro M, Ambrosio G, Mattioli G (2009) Heparin/PF4 antibodies formation after heparin treatment: temporal aspects and long-term follow-up. *Am Heart J* 157(3):589–595. <https://doi.org/10.1016/j.ahj.2008.11.007>
- Greinacher A, Selleng K, Warkentin TE (2017) Autoimmune heparin-induced thrombocytopenia. *J Thromb Haemost* 15:2099–2114
- Selleng S, Selleng K (2016) Heparin-induced thrombocytopenia in cardiac surgery and critically ill patients. *Thromb Haemost* 116(5):843–851