

This is a pre print version of the following article:

Impact of bacterial spot outbreaks on the phytosanitary quality of tomato and pepper seeds / Giovanardi, Davide; Biondi, Enrico; Ignjatov, Maja; Jevtic, Radivoje; Stefani, Emilio. - In: PLANT PATHOLOGY. - ISSN 1365-3059. - 67:5(2018), pp. 1168-1176. [10.1111/ppa.12839]

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

10/01/2026 20:07

# 1    **Impact of bacterial spot outbreaks on the phytosanitary quality of tomato and pepper seeds**

2

3    D. Giovanardi <sup>1</sup>, E. Biondi <sup>2</sup>, M. Ignjatov <sup>3</sup>, R. Jevtić <sup>3</sup>, E. Stefani <sup>1</sup>.

4

5    <sup>1</sup>Department of Life Sciences, University of Modena and Reggio Emilia, via Amendola 2, 42122  
6    Reggio Emilia, Italy. <sup>2</sup>Department of Agricultural Sciences, University of Bologna, viale Fanin 44,  
7    40127 Bologna, Italy. <sup>3</sup>Institute of Field and Vegetable Crops, Maksima Gorkog 30, 21000 Novi  
8    Sad, Serbia.

9

10    E-mail address: [davide.giovanardi@unimore.it](mailto:davide.giovanardi@unimore.it)

11

## 12    **ABSTRACT**

13    The impact of disease outbreaks on the phytosanitary quality of seeds was investigated for the  
14    following pathosystems: tomato-*Xanthomonas vesicatoria* and pepper-*Xanthomonas euvesicatoria*.  
15    This study, which was performed in Italy and in Serbia, aimed to evaluate the season-to-season  
16    transmission of phytopathogenic regulated bacteria associated with phytosanitary risks posed by  
17    seeds produced in areas where bacterial infections are possible. For each pathosystem, field plots  
18    were experimentally inoculated to simulate an initial infection rate of 1%, 5% and 15%. The area  
19    under the disease progress curve (AUDPC) was calculated for each field plot, the produced seeds  
20    were analysed to determine the contamination level and rate, and the plant-to-seed transmission was  
21    evaluated by a seedling grow-out (SGO) assay. To investigate transmission under field conditions, a  
22    second-year experiment was performed, wherein seeds collected from the first year were used to  
23    establish new field plots. During the first growing season, AUDPC values were positively  
24    correlated with the percentages of initial infection for each pathosystem. Seed contamination levels  
25    ranged from 34 to 100 CFU g<sup>-1</sup>, and the contamination rate ranged from 1.50% up to 3.17% for  
26    *Xanthomonas euvesicatoria*, whereas processing and fresh market tomato seeds produced both in

Italy and Serbia were not infected by *Xanthomonas vesicatoria*. During SGO assays and the second cropping year, no symptoms were observed in either tomato or pepper plants. Therefore, the calculated pepper seed contamination rate for *Xanthomonas euvesicatoria* appeared to be less than the threshold necessary to initiate a disease outbreak. Finally, all seeds obtained during the second cropping year were uninfected.

**Keywords:** Seed-borne bacteria, bacterial leaf spot, pathogen transmission, phytosanitary seed quality.

## INTRODUCTION

Infected seeds are the most important pathway for the introduction and spread of several plant pathogenic bacteria that may affect both fruit quality and plant viability, thus causing economic losses worldwide (Gitaitis & Walcott, 2007). Among other bacterial diseases, tomato can be affected by bacterial spot caused by *Xanthomonas vesicatoria* (e.g., Doidge, 1920; Vauterin *et al.* 1995) (Xv) and *X. euvesicatoria* (Jones *et al.* 2004) (Xe). The latter is particularly aggressive against pepper (Ignjatov *et al.* 2010). These bacteria are spread primarily through contaminated/infected seeds as the primary source of *inoculum* (Dutta *et al.* 2014) and are listed as quarantine organisms by the European Plant Protection Organization (EPPO). These pathogens can survive in seeds for extended periods (Bashan *et al.* 1982a); such survival ability allows them long-distance dissemination and facilitates their introduction into pathogen-free areas through seed trade. Seed testing and certification (EPPO, 2013) and seed production in pathogen-free areas with no conducive environmental conditions are the most important management strategies for the preventive control of these pathogens. Sanitation of potentially contaminated seeds is only partially achieved using common disinfection methods, e.g., heat treatment, fermentation of fruit pulp as in the case of tomato (Chambers & Merriman, 1975; Dhanvantari, 1989), or chlorine/acid treatment for pepper seeds (Dempsey & Walker, 1973). So far, there is no method available that can ensure

the complete eradication of pathogens from naturally infected seeds without dramatically reducing seed germination (Dhanvantari, 1989). Despite the use of phytosanitary certification and quarantine procedures in domestic and international seed trade, which can considerably reduce disease incidence, severe epidemics are occasionally reported (Gitaitis & Walcott, 2007).

Transplant production studies on the transmissibility of *Clavibacter michiganensis* subsp. *michiganensis* (Smith) (Davis *et al.* 1984; Strider, 1969), the causal agent of tomato bacterial canker, revealed that a single infected tomato seed among 10,000 is sufficient to initiate an epidemic under favourable conditions (Chang *et al.* 1991). Similar studies are currently not available for Xv or for Xe. Very few studies have been conducted under field conditions to evaluate the seed contamination threshold necessary for pathogen transmission from seed to plants. Chang *et al.* (1991) demonstrated a systemic infection in tomato transplants grown from infected seeds containing approx.  $10^4$  CFU/g<sup>-1</sup> of *C. michiganensis* subsp. *michiganensis*. Conversely, knowledge regarding the Xv and Xe seed contamination thresholds needed for disease expression on tomato and pepper plants under field conditions is lacking. Bacterial spot is a polycyclic disease. Secondary *inocula* might be abundantly produced during the cropping period and may play a key role in the short-distance spread of both bacterial species. In this phase, endophytic bacterial cells are released either by guttation droplets or through infected stomata and spread via splashing water and wind-driven rain. Xv and Xe seed contamination/infection can occur via two pathways: penetration through fruit lenticels (Bashan *et al.* 1982a) and/or floral structures (Dutta *et al.* 2016). For both causal agents of bacterial spot of tomato and pepper plants (Xv and Xe), seed contamination mainly occurs on the seed surface from infected pulp rather than as internal seed infections (EPPO, 2013). However, neither the transmission of such bacteria from diseased plants to seeds under field conditions nor the correlation between disease quantity, and the contamination level of bacteria on/in seeds have been explored to date. This extensive study was performed in confined experimental fields approved by local phytosanitary authorities and located in Northern Italy (Emilia Romagna Region) and Serbia (Vojvodina Province), two regions where tomato and pepper

79 represent important crops and where bacterial diseases may cause significant economic losses. The  
80 objectives of this study were (i) to evaluate and quantify disease outbreaks under field conditions,  
81 (ii) to assess disease progression and the correlation between disease quantity and the phytosanitary  
82 quality of seeds, and (iii) to investigate the relationship between the assessed seed contamination  
83 level and the risks of possible disease outbreaks in the following year's crop. These data will be  
84 useful for seed companies for the production of seeds with an acceptable phytosanitary quality  
85 when disease symptoms are expressed during a vegetative season. The outcome of this study may  
86 also aid in the identification of some additional aspects of Xv and Xe plant-to-seed and seed-to-  
87 plant transmission in tomato and pepper, respectively, thus shedding light on the epidemiology of  
88 these diseases.

89

## 90 **MATERIALS AND METHODS**

### 91 **Bacterial strains and plant material**

92 The virulent strains IPV-BO 2684 and KFB29 of Xv and MI-A-6 of Xe, belonging to the bacterial  
93 collections of the Department of Agricultural Sciences (University of Bologna, Italy) and the  
94 Institute of Field and Vegetable Crops (Novi Sad, Serbia), respectively, were used for this study. To  
95 confirm their pathogenicity prior to experiments, Xv and Xe were inoculated and re-isolated from  
96 symptomatic fruits and leaves of tomato and pepper, respectively. All strains were routinely grown  
97 on glucose-yeast extract-calcium carbonate medium (GYCA) (Dye, 1962) at  $27.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$  for 48-  
98 72 h.

99 Tomato and pepper cultivars were selected for their high susceptibility to bacterial diseases and  
100 adaptation in their respective countries. In Italy, processing tomato cv. VF10 was used, whereas in  
101 Serbia, fresh market tomato cv. Jabučar and bell pepper cv. Amphora were used. Certified tomato  
102 and pepper seeds were kindly provided by commercial sources: processing tomato seeds were  
103 provided by ISI Sementi s.p.a. (Fidenza, Italy), and fresh market tomato and bell pepper seed were  
104 provided by NS-SEME (Novi Sad, Serbia).

105

## 106 **Field experiments**

107 The respective plant health authorities were notified of the field experiments, which were conducted  
108 in confined experimental fields under the supervision of the local Phytosanitary Service (Cadriano  
109 Experimental Station, Bologna, Italy; IFVCS Experimental Station, Novi Sad, Serbia).

110 Field experiments were performed in Italy on processing tomato-Xv and in Serbia on fresh market  
111 tomato-Xv and pepper-Xe pathosystems. The experiments started in 2013 and were conducted  
112 during two cropping seasons.

113 In the first cropping year, for each pathosystem, three plots of 96 plants each (12 plants in each of 8  
114 rows) were set. Tomato and pepper seedlings were planted according to the common agricultural  
115 procedures followed by farmers for commercial purposes. For processing tomato, each plot  
116 consisted of 8 rows of 12 plants spaced 0.3 m apart with 0.7 m between rows. For fresh market  
117 tomato and bell pepper, each plot consisted of 8 rows of 12 plants spaced 0.5 m apart with 1.1 m  
118 between rows. Rows of maize were grown between plots to avoid cross contamination. One, five  
119 and fourteen plants of each plot in both Italy and Serbia were arbitrarily selected, labelled and  
120 experimentally inoculated to obtain approximately 1%, 5% and 15% of infected plants equitably  
121 distributed among the plots (see supplementary material S1). The plants were experimentally  
122 inoculated five weeks after transplanting. At this point, the phenological growth stage of the tomato  
123 and pepper plants was the beginning of flowering and intensive leaf growth and bud shooting,  
124 respectively. Late in the afternoon, the labelled plants were spray-inoculated until run-off with a  
125 water suspension containing each pathogen at a concentration of approximately  $10^8$  CFU ml<sup>-1</sup>.  
126 Specifically, Xv and Xe strains were grown on GYC agar for 48 h at 28 °C, and the concentration  
127 of bacterial cells was determined using a spectrophotometer (Spectronic 20; Bausch and Lomb,  
128 Rochester, NY) (optical density at 600 nm = 0.3;  $\approx 1 \times 10^8$  CFU ml<sup>-1</sup>). After spraying, each  
129 inoculated plant was sealed in a polyethylene (PE) bag to facilitate water congestion and pathogen  
130 penetration mainly through stomata and hydathodes into the host. The following day, early in the

131 morning, the PE bags were removed. A successful inoculation was demonstrated by symptom  
132 development, sampling of symptomatic leaves and analysis to confirm infection by the respective  
133 pathogens. Specifically, to confirm pathogen identity between the *inoculum* source and re-isolated  
134 strains, rep-PCR using the primer BOX A1R (5'-CTACGGCAAGGCGACGCTGACG-3') was  
135 performed according to Versalovic *et al.* 1994. The plants were cultivated according to the local  
136 best agricultural practices. In Italy (processing tomatoes), two harvests were manually performed 7  
137 days apart; in Serbia, fresh market tomatoes and bell peppers were weekly harvested for a period of  
138 2 months (12 harvests in total).

139 In the second cropping season, the tomato and pepper seeds, generated during the previous year  
140 from each experimentally infected field plot, were sown in blotters. The seedlings (at the 3<sup>rd</sup>-4<sup>th</sup> leaf  
141 stage) were then transplanted in new experimental fields, which were designed as in the previous  
142 year; seedlings generated from commercially certified seeds were also transplanted in an additional  
143 plot of 96 plants as a negative control. Rows of maize were grown between plots to avoid cross  
144 contamination. Again, the plants were cultivated according to the local best agricultural practices.  
145 Fruits were harvested, and seeds were extracted according to the procedures described below. Agro-  
146 climatic parameters were monitored throughout both growing seasons.

147

#### 148 **Phytopathometric evaluations**

149 During the first cropping year, phytopathometric assessments in experimental fields were weekly  
150 performed, starting from the first appearance of symptoms until harvest. During these assessments,  
151 the increase in disease incidence and severity was recorded. Therefore, diseased plants were  
152 counted (incidence), and each one was assigned to a disease severity class. Disease severity in  
153 tomato and pepper affected by *Xanthomonas* spp. strains was evaluated according to five  
154 phytopathometric classes, ranging from 0 to 4 (0 = no symptoms; 1 = 1-10 spots on up to 3 leaves; 2  
155 = 11-30 spots on 4 to 10 leaves; 3 = more than 30 spots and some confluent necrosis on 5 to 20  
156 leaves; 4 = confluent necrosis on more than 20 leaves or branch desiccation) (Giovanardi *et al.*

2015). Disease scores were calculated as  $\Sigma$  of Q, where Q represents the combination of the disease severity and the incidence (severity  $\times$  incidence) at each assessment. The area under the disease progression curve (AUDPC) (Van der Plank, 1963) was then calculated according to Madden *et al.* (2007) from the first phytopathometric evaluation to the last assessment before harvest. Moreover, the AUDPCs were statistically evaluated by ANOVA (p 0.05) using SPSS 15.0 software for Windows®.

#### Seed extraction

Tomato and pepper seeds were produced according to common commercial procedures (Opeña *et al.* 2001). For both tomato cultivars, seeds were extracted according to the fermentation technique as follows: harvested tomatoes were left in a dark store at  $23.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$  for 2 weeks for post-harvest full maturation before seed extraction. Then, seeds were manually extracted from the pulp, and the mixture of seeds and placental tissue was maintained at  $25.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$  for 24 h. The seeds were then separated from the fermentation mixture, thoroughly washed under running tap water and dried overnight in the dark at  $25.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$  on sterilized paper trays. For bell pepper, seeds were manually collected after pericarp removal, thoroughly washed under running tap water and subsequently dried for 48 h in the dark at  $25.0 \pm 1\text{ }^{\circ}\text{C}$ . The seeds produced were weighed and counted for each plot. Finally, all dried seeds were stored in paper boxes and kept in the dark at  $4.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$ .

#### Seed analysis and contamination rate of seed plots

For each seed lot generated from the experimental plots, seed analysis was performed by dilution plating and molecular assays according to the EPPO (2013) standard diagnostic protocol for *Xanthomonas* spp. causing bacterial spot of tomato and pepper. Seed samples ( $n = 10,000$ ) from each field plot were soaked in a Stomacher bag with a ratio of 4 ml of sterilised PBS-T ( $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$  19.57 g l<sup>-1</sup>,  $\text{KH}_2\text{PO}_4$  1.65 g l<sup>-1</sup>, Tween 20 0.5 g l<sup>-1</sup>; pH 7.4) per g of seeds for 14 h at  $4.0\text{ }^{\circ}\text{C} \pm$



183 1 °C (EPPO, 2013). The seeds, contained in Stomacher bags, were then crushed with a hammer for  
184 2 min, and the maceration fluid was centrifuged at 10,000 x g for 20 min at 4.0 °C ± 1 °C. The  
185 resulting pellet was then resuspended in 1 ml of sterilized PBS-T. Each seed extract was divided  
186 into two aliquots of 900 and 100 µl. The 900-µl aliquots were used for DNA extraction and  
187 purification with the DNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to the  
188 manufacturer's instructions.

189 Two µl of purified DNA was then subjected to conventional PCR assay using the specific primer  
190 sets for the detection of Xv (Bs-XvF: 5'-CCA TGT GCC GTT GAA ATA CTT G -3'; Bs-XvR: 5'-  
191 ACA AGA GAT GTT GCT ATG ATT TGC-3') and Xe (Bs-XeF: 5'-CAT GAA GAA CTC GGC  
192 GTA TCG-3'; Bs-XeR: 5'-GTC GGA CAT AGT GGA CAC ATA C- 3') (Koenraadt *et al.* 2009)  
193 according to EPPO (2013). Expected amplicon sizes were 138 bp and 173 bp for the Xv-specific  
194 and Xe-specific primer sets, respectively. For each sample, the 100-µl aliquot was ten-fold diluted  
195 and streaked onto GYCA medium. The tomato and pepper seed extracts were also streaked onto  
196 mTMB (McGuire *et al.* 1986), a specific semi-selective medium for Xv/Xe. After incubating for  
197 72-96 h at 28 °C, putative Xv and Xe colonies were selected for purification on GYC agar and  
198 further identification of axenic colonies with specific primer sets, as previously described. To  
199 confirm pathogen identity between the inoculum source and re-isolated strains, rep-PCR with the  
200 BOX A1R primer set was performed (Versalovic *et al.* 1994). Seed samples were assayed in  
201 triplicate for each contaminated plot.

202 The contamination level (CFU) of each seed sample ( $n = 10,000$ ) was preliminary calculated as the  
203 mean number of colonies of the four ten-fold dilutions (*i.e.*, 1:1; 1:10; 1:100 and 1:1000).  
204 Considering the weight per thousand of grain (WTG), each seed contamination level was expressed  
205 as CFU g<sup>-1</sup>. The three seed sample ( $n = 10,000$ ) replicates of each plot were used to calculate the  
206 mean contamination level expressed as CFU g<sup>-1</sup>.

207 Thereafter, to assess the contamination rates of seeds from each infected field plot (*i.e.*, 1, 5 and  
208 15%), 50 replicates of 100 seeds from each plot were analysed by conventional PCR. From each

209 plot, the contamination rates of seeds were calculated according to the formula  $p = 1 - (Y/N)^{1/n}$ , where  
210  $N$  is the number of analysed replicates,  $n$  the number of seeds in a replicate and  $Y$  the number of  
211 healthy groups (Maury *et al.* 1985; Darrasse *et al.* 2007). Each 100-seed sample was soaked in a  
212 Stomacher bag in 3 ml of sterilized PBS-T for 14 h at  $4.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$  (EPPO, 2013). The seed  
213 samples were then processed and analysed by conventional PCR with specific primer sets for the  
214 detection of Xv and Xe, as described above.

215

#### 216 **Seed germination rate assay**

217 Seeds generated from both first and second cropping season in Italy and Serbia were tested to  
218 determine their germination rate; *in vitro* germination was carried out according to International  
219 Rules for Seed Testing standards (ISTA, 2009). One hundred seeds for each pathosystem were  
220 placed on Whatman No. 5 filter paper (Sigma-Aldrich, Saint Louis, Missouri) in Petri dishes and  
221 dampened with 5 ml of sterilized distilled water (SDW); the plates were then placed at  $25.0\text{ }^{\circ}\text{C} \pm 1$   
222  $^{\circ}\text{C}$  in the dark. Germinated seedlings were counted every day for 14 days. The *in vitro* germination  
223 assays were repeated four times for each seed sample, and the results were collected and statistically  
224 evaluated by ANOVA (p 0.05) using SPSS 15.0 software for Windows®.

225

#### 226 **Seedling grow-out (SGO) assay**

227 To test the seed transmission of Xv and Xe, a SGO assay was performed on blotters by sowing seed  
228 samples ( $n = 1,000$ ) into pots containing peat. The seedlings were then kept in a climatic chamber  
229 for 4 weeks at  $28.0\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$ , with a relative humidity (RH) of up to 90%. After 28 days, the  
230 seedlings were inspected for typical Xv or Xe symptoms. After 4 weeks, all seedlings were  
231 collected; then, segments of approx. 2 cm were cut from each stem and pooled in Stomacher bags  
232 with 30 ml of sterilized PBS-T. The samples were then crushed by hammering and incubated at  
233 room temperature for 30 min. The stem macerates were centrifuged at  $10,000 \times g$  for 20 min at  $4.0$   
234  $^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$  and the resulting pellets were resuspended in 1 ml of PBS-T. Each sample was analysed

235 by dilution plating and molecular assays, as previously described. Seed samples from each  
236 contaminated plot were tested in triplicate.

237

## 238 **RESULTS**

### 239 **Field experiments**

240 In the first cropping season in Italy, the first symptoms were detected on processing tomato leaves 2  
241 weeks after the inoculation with Xv by surveying the experimental fields. These symptoms were  
242 observed as necrotic spots and marginal necrotic areas from which the Xv strain IPV-BO 2684 was  
243 re-isolated. Strain identity was confirmed by rep-PCR assay using the BOX A1R primer set. At  
244 harvesting time, 14 weeks after transplanting, the final disease incidence calculated for bacterial  
245 spot was approx. 12%, 60% and 85% in the plots with an initial infection percentage of 1%, 5% and  
246 15%, respectively. Disease severity, calculated according to a disease index scale and assessed for  
247 each plant, progressively increased from one observation to the next; therefore, the calculated  
248 disease quantity (Q) constantly increased until the last assessment. The increase in the disease  
249 progression curve for Xv in infected plots directly correlated with the percentage of initial infection  
250 (Figure 1). The AUDPC value of processing tomato plants inoculated with Xv in the field plot with  
251 a 1% initial infection rate was 249, which was approx. six- and ten-fold lower than the field plots  
252 with an initial infection rate of 5% and 15% (AUDPC, 1512 and 2654, respectively) (Table 1). Xv  
253 infections produced spots on a limited number of fruits.

254 During the second cropping season, no bacterial spot symptoms were recorded in any of the plots  
255 where processing tomato plants were obtained from seeds produced during the previous year in  
256 diseased plots.

257 In Serbia, during the first cropping season, leaf spot symptoms on fresh market tomato and bell  
258 pepper appeared 2 weeks after the experimental inoculation and increased until the last assessment,  
259 which was 18 weeks after transplanting. Symptoms developed on both leaves and fruits. In addition,  
260 both the Xv strain KFB29 and the Xe strain MI-A-6 were re-isolated from symptomatic plant

261 tissues, and their identity was confirmed by rep-PCR assay using the BOX A1R primer set. Disease  
262 quantity (Q) increased in all infected fresh market tomato plots (Figure 2) until 100% of the plants  
263 were diseased (data not shown). Similar results were observed for bell pepper: disease quantity (Q)  
264 increased in all plots (Figure 3) until it reached 100% (data not shown). The AUDPC value of Xv-  
265 infected tomato plants in the field with a 1% initial infection rate was 8589, approx. two times  
266 lower than that of the fields with 5% (AUDPC = 15074) and 15% (AUDPC = 18788) initial  
267 infection rates.

268 For the bell pepper-Xe pathosystem, even though the calculated AUDPC of the field plot with an  
269 initial infection rate of 1% was 5743, approx. 1.5 times lower than that of the field plot with a 5%  
270 initial infection rate (AUDPC = 8522) and almost 3 times lower than that calculated in the plot with  
271 a 15% initial infection rate (AUDPC = 13632), AUDPC values did not show statistical differences  
272 among the different plots (Table 1).

273 During the second cropping season, no symptoms were recorded in any of the fresh market tomato  
274 and bell pepper plots generated with seeds produced in the previous year from diseased plots.

275 The environmental conditions of the two cropping seasons are reported in the supplementary  
276 material (Figures S2 and S3). In Italy, during both cropping seasons, the minimum, maximum and  
277 mean temperatures were average for that period, whereas the RH was below average, with values  
278 <70% from May 20<sup>th</sup> to August 31<sup>st</sup> in 2013 and from July 1<sup>st</sup> to August 31<sup>st</sup> in the second cropping  
279 season. In Serbia, during both cropping seasons, the minimum, maximum and mean temperatures  
280 were average for that period, whereas the RH was below average, with values <80% from July 15<sup>th</sup>  
281 to August 26<sup>th</sup> in 2013 and from July 1<sup>st</sup> to August 19<sup>th</sup> in the second cropping season.

282

### 283 **Harvest and seed production**

284 In the first cropping season in Italy, *ca.* 65,000 seeds per plot (*ca.* 170 g per plot) were extracted  
285 from processing tomato fruits of Xv-infected fields. In the second cropping season, *ca.* 44,000 seeds

per plot (*ca.* 114 g per plot) were extracted from the fruits of Xv-infected processing tomato fields. From the negative control field plot, *ca.* 37,000 seeds (*ca.* 95 g) were extracted. In the first cropping season in Serbia, *ca.* 60,000 seeds per plot (*ca.* 156 g per plot) of fresh market tomato and *ca.* 40,000 seeds per plot (*ca.* 280 g per plot) of bell pepper were produced. In the second cropping season, *ca.* 50,000 seeds per plot (*ca.* 130 g per plot) of fresh market tomato and *ca.* 74,000 seeds per plot (*ca.* 233 g per plot) of bell pepper were extracted. From the negative control field plots, *ca.* 57,000 (*ca.* 149 g) of fresh market tomato seeds and *ca.* 39,000 (*ca.* 419 g) of bell pepper seeds were extracted. The WTG was 2.61 g, 2.60 g and 6.96 g for processing tomato, fresh market tomato and bell pepper seeds, respectively.

296

#### 297 **Seed analysis and contamination rate of seed lots**

Tomato and pepper seed lots produced during both cropping seasons were tested according to the EPPO (2013) standard diagnostic protocol for *Xanthomonas* spp. Specifically, for each seed lot, three seed samples ( $n = 10,000$ ) were analysed by direct isolation and molecular assays, and the results are presented in Table 2. For seeds produced during the first cropping season, Xv was not detected by either PCR or direct isolation on GYCA and mTMB media from tomato seeds produced in Italy and in Serbia. For pepper, Xe was consistently detected by PCR and recovered on mTMB and GYCA media from all seed lots produced in the different field plots. The seed contamination level was assessed as 34 (SD = 13), 37 (SD = 22) and 100 (SD = 48) CFU g<sup>-1</sup> in the 1%, 5% and 15% infected plots, respectively. For seeds produced during the second cropping season, direct isolation and molecular analyses did not result in the detection of Xv and Xe in any plot in Italy or in Serbia.

Additionally, to determine the contamination rates of each seed lot, 50 replicates of 100 seeds were analysed. The contamination rate of seed lots produced during the first cropping season in Italy and Serbia was determined according to the formula  $p = 1 - (Y/N)^{1/n}$  (Maury *et al.* 1985). No seed

contamination (0%) was observed for processing or fresh market tomato produced in Xv-infected plots. For pepper seeds, the seed contamination rate was 1.50%, 3.17% and 3.17% (Table 2) in the plots infected with Xe at 1%, 5% and 15%, respectively. The seed contamination rate of all seed lots produced during the second cropping season in Italy and Serbia tested negative for Xv and Xe.

#### **Seed germination rate assay**

All seed lots produced during both cropping seasons in Italy and Serbia were subjected to quality testing by assessing their germination rate and their ability to produce marketable seedlings. The germination rate of tomato and pepper seeds produced during the first cropping season was *ca.* 95-98%, with no significant differences ( $P \geq 0.05$ ) among seed lots produced in plots with different disease quantities. Similar results were obtained from seed lots produced during the second cropping year. No significant differences ( $P \geq 0.05$ ) in the germination rate (ranging from 90 to 98%) were observed among the different seed lots produced and the uninfected plots (negative controls).

#### **SGO assay**

The assessment of disease incidence during the SGO assays did not reveal any disease symptom on seedlings obtained from seeds produced in both the first and second cropping years in Italy and Serbia. Additionally, microbiological and molecular analyses performed on each seedling macerates and their DNA extracts were all negative for the presence of Xv and Xe.

### **DISCUSSION**

Xv and Xe are phytopathogenic bacteria of great concern to seed companies and farmers. Xv has been recorded in several important seed-producing countries (*e.g.*, Brazil, Mexico, USA, India, Thailand), whereas Xe exhibited a far more restricted distribution (EFSA Panel on Plant Health, 2014). Both Xv and Xe are seed-borne pathogens, and infested seeds serve as a main source of

338 primary inoculum in transplant and fruit production systems (Jones *et al.* 1993; Leite *et al.* 1995;  
339 Dutta *et al.* 2014).

340 Seed companies are devoted to producing tomato and pepper seeds under pathogen exclusion  
341 conditions, either in areas where the pathogens have never been reported, or under strict hygienic  
342 practices (Gitaitis & Walcott, 2007). Nonetheless, disease symptoms may appear in seed production  
343 fields due to the use of seeds that are contaminated at a level under the pathogen detection threshold  
344 or because the production area is not sufficiently surveyed. It remains unknown whether a  
345 consistent positive correlation exists between seed contamination rate and disease outbreak and  
346 whether a similar correlation exists between disease quantity, as measured in fields devoted to seed  
347 production, and the phytosanitary quality of the resulting seeds.

348 In our study devoted to simulate bacterial spot outbreaks occurring from external sources of  
349 *inoculum*, the results obtained during the first cropping season revealed a positive correlation  
350 between the incidence of initial infection, disease progression and disease quantity/score (Q) for  
351 both Xv and Xe, as confirmed by the calculated AUDPC values. The disease scores measured on  
352 tomato were remarkably different between the Italian and Serbian fields; these differences might be  
353 explained by the length of the growing cycle, which is considerably longer for fresh market tomato  
354 (7 weeks longer) than that for processing tomato, and by the presence of more favourable  
355 environmental conditions for disease outbreak in Serbia than in Italy (*e.g.*, higher RH).

356 Similar disease progression occurred in bell pepper, since monitoring and harvesting of peppers  
357 continued for an additional 8 weeks after processing tomato harvest. In addition, even though  
358 AUDPC values from 1% infected plots appeared lower than those from 5% and 15% infected plots,  
359 the differences were not significant. This could be due to the length of the growing cycle, but also to  
360 the more effective spread ability of Xe in bell pepper in comparison to that of Xv in fresh market  
361 tomato at the same environmental conditions.

362 In Italy, Xv infections produced spots on a limited number of fruits: this might be explained by the  
363 low daily mean RH (< 60%), which was consistently measured at anthesis and fruit development

364 and ripening (July-August 2013). Conversely, during the same physiological growth stages in  
365 Serbia, a daily mean RH > 80% was consistently recorded. Therefore, the RH conditions, which  
366 were more favourable for bacterial infection than those recorded in Italy, might explain the high  
367 disease incidence and severity expression on both fresh market tomato and bell pepper. Given that  
368 no outside sources of *inoculum* were present during the field experiments, bacterial disease  
369 development in the plots was the exclusive result of infection from the artificial *inoculum* sources,  
370 as previously discussed by Kocks & Zadoks (1996) for other xanthomonads. Indeed, the appearance  
371 of typical *foci* in the field plots around the experimental *inoculum* sources (experimentally  
372 inoculated plants were marked) ultimately supports this conclusion (Zadoks & Van den Bosch,  
373 1994). Finally, the pathogens were re-isolated from infected plants, and their identities were  
374 confirmed by rep-PCR as the experimental *inoculum* sources (*i.e.*, strains IPV-BO 2684 and KFB29  
375 of Xv and MI-A-6 of Xe).

376 Despite the positive correlation observed between the incidence of initial infection, disease  
377 progression and AUDPC values for both Xv and Xe, the analyses of seeds produced during the first  
378 cropping season did not exhibit any apparent correlation between disease quantity over time  
379 (AUDPC), as measured in the production fields, and contamination rates of produced seeds for  
380 either the tomato or pepper pathosystems. In the case of tomato, none of the seed analyses detected  
381 Xv, though 58% and 63% of the total produced seeds in Italy and in Serbia were tested,  
382 respectively, and the disease observed in the field plots was remarkably severe and present on all  
383 aerial parts as leaves, fruits, petioles and stems. Conversely, seed contamination rates were between  
384 1.50% and 3.17% for pepper seed lots produced in diseased plots. These pepper seed lots tested  
385 positive for Xe detection by conventional PCR and by isolation on semi-selective media. These  
386 results confirm a seed contamination by Xe; however, the infection is characterized by relatively  
387 low rates and non-uniform distribution of Xe among the seed lots. Pathogen re-isolation indicated  
388 that the seed population density for Xe was *ca.* 34 to 100 CFU g<sup>-1</sup>.



389 The seed extraction protocol based on the fermentation of tomato pulp (Opeña *et al.* 2001) is  
390 thought to play a major role in the viability and detectability of infecting bacteria. This method is  
391 able to decrease the viability of bacterial populations contaminating the seeds, leading to a  
392 consistent reduction in the density of viable bacterial cells on tomato seed surfaces (Chambers &  
393 Merriman, 1975). Bacteria in the viable but non-culturable (VBNC) state, which fail to grow on the  
394 routine bacteriological media, have been described (Oliver, 2005). These bacteria are in a state of  
395 very low metabolic activity and do not divide. This feature occurs in response to stress (*e.g.*, due to  
396 adverse nutrient, temperature, osmotic, oxygen, and light conditions) (Stokell & Steck, 2012). In  
397 our case, we analysed seeds immediately after their production, and they were stored according to  
398 the best possible procedures (*i.e.*, dry seeds stored at 4 °C in the dark). Therefore, we presume that  
399 no stress or other adverse conditions may have caused the development of a VBNC status in both  
400 Xv and Xe. An additional confirmation of this conclusion was obtained by molecular assays. These  
401 assays could detect the DNA present in VBNC, but the results were negative for the presence of the  
402 pathogen. Basically, when the pathogen was re-isolated on semi-selective medium, PCR  
403 consistently detected its presence in DNA extracts.

404 Throughout the second cropping season, no bacterial spot symptoms were observed in pepper or  
405 tomato plants. Moreover, in seeds extracted in the second cropping season, the pathogens were not  
406 detected by any of three biologically different tests: microbiological, molecular and SGO assays.  
407 Therefore, the lack of symptom development in all field plots indicates that seeds were not infected  
408 or contaminated by Xv or that the pathogen transmission was negligible due to a low bacterial load  
409 of Xe on pepper seeds, which was not sufficient to cause disease development in the next cropping  
410 season. Since Bashan *et al.* (1982b) observed that Xv can survive in tomato seeds for a long time,  
411 up to 8 years, we may presume that the viability of the pathogen in seeds used in the following year  
412 was not dissimilar to the contamination level assessed after seed production.

413 In the past, *inoculum* transmission thresholds in seeds have been studied in few pathosystems  
414 addressed to identify the correlation between seed infection rates and disease outbreaks, as in the

415 case of *Pseudomonas savastanoi* pv. *phaseolicola* in bean seeds (Taylor *et al.* 1979) or  
416 *Xanthomonas campestris* pv. *campestris* in crucifer seeds (Shaad *et al.* 1980). In these studies, the  
417 *inoculum* thresholds were set either arbitrarily (*e.g.* using experimental seed inoculation) or simply  
418 with field observations. In the case of tomato and pepper seeds, the correlation between *inoculum*  
419 thresholds of Xv and Xe and disease outbreaks has not been previously evaluated under field  
420 conditions. Additionally, to the best of our knowledge, no studies in the literature have elucidated  
421 the quantitative/qualitative correlation between disease quantity observed in tomato/pepper fields,  
422 the concentration of *inoculum* in the seeds produced in those fields and the bacterial spot outbreak  
423 risks posed by these seeds on the next crop.

424 This is the first report on the evaluation of a seed contamination threshold for bacterial spot in  
425 pepper. The results from this study showed that a seed contamination level higher than 100 CFU g<sup>-1</sup>  
426 is needed for a disease outbreak. Due to the highly polycyclic nature of the disease, it is important  
427 to emphasise that the threshold level may be variable, considering that pepper growing areas have  
428 quite different climatic conditions and/or different agronomic practices (*e.g.*, higher seeding rates).

429 Conversely, for the Xv-tomato pathosystem, our data showed a total lack of *inoculum* transmission,  
430 confirmed by the absence of living cells on semi-selective media used for re-isolation. Xv, during  
431 SGO assays and under field conditions, was not transmitted both to seedlings and to seeds in either  
432 Italy or Serbia. The results obtained from molecular and microbiological assays on seed extracts of  
433 processing tomatoes also suggest the important role of extraction protocols in seed sanitation (*i.e.*, a  
434 fermentation step). The fermenting process could be considered an appropriate seed surface  
435 disinfection step that does not affect seed quality, as confirmed by *in vitro* germination results. The  
436 results obtained from SGO assays, in particular those related to the Xe-pepper pathosystem, for  
437 which plant-to-seed transmission was successfully achieved, suggest that the bacterial loads in  
438 contaminated seeds were not sufficient to develop symptoms, even using optimal controlled  
439 conditions for both host and pathogen. As observed in the field, the pathogen population on seeds of  
440 the two strains used were confirmed to be insufficient for disease outbreak.

441 This study provides new information on the seed transmission of bacterial spot as well as a deeper  
442 knowledge of epidemics in the field. Our data will facilitate a better understanding of the  
443 epidemiology of *Xanthomonas* spp. seed-borne bacteria.

444

#### 445 **ACKNOWLEDGEMENTS**

446 This research was financed in the framework of the EU Project “Development of seed testing  
447 methods for pests and pathogens of plant health concern (TESTA)” ([www.seedtesta.eu](http://www.seedtesta.eu)). The  
448 authors wish to thank Assosementi, Italian Member of the European Seed Association (ESA) and  
449 the International Seed Federation (ISF) for assistance and for providing seeds for the experiments  
450 and facilities for workshops and dissemination of results.

451 **LITERATURE CITED**

- 452 1. Bashan, Y., Diab, S., Okon, Y. 1982a. Survival of *Xanthomonas campestris* pv. *vesicatoria* in  
453 pepper seeds and roots, in symptomless and dry leaves, in on-host plants and in the soil. *Plant and*  
454 *Soil* **68**: 161–170.
- 455 2. Bashan, Y., Okon, Y., Henis, Y. 1982b. Long-term survival of *Pseudomonas syringe* pv. *tomato*  
456 and *Xanthomonas campestris* pv. *vesicatoria* in tomato and pepper seeds. *Phytopathology* **72**: 1143–  
457 1144.
- 458 3. Chambers, S.C., Merriman, P.R. 1975. Perennation and control of *Pseudomonas tomato* in  
459 Victoria. *Australian Journal of Agricultural Research* **26**: 657–663.
- 460 4. Chang, R., Ries, S., Pataky, J. 1991. Dissemination of *Clavibacter michiganensis* subsp.  
461 *michiganensis* by practices used to produce tomato transplants. *Phytopathology* **81**: 1276–1281.
- 462 5. Darrasse, A., Bureau, C., Samson, R., Morris, C., Jacques, M.-A. 2007. Contamination of bean  
463 seeds by *Xanthomonas axonopodis* pv. *phaseoli* associated with low bacterial densities in the  
464 phyllosphere under field and greenhouse conditions. *European Journal Plant Pathology* **119**: 203–  
465 215.
- 466 6. Davis, M. J., Gillaspie, A. G., Jr Vidaver, A. K., Harris, R. W. (1984). *Clavibacter*: a new genus  
467 containing some phytopathogenic coryneform bacteria, including *Clavibacter xyli* subsp. *xyli* sp.  
468 nov, subsp. nov. and *Clavibacter xyli* subsp. *cynodontis* subsp. nov., pathogens that cause ratoon  
469 stunting disease of sugarcane and bermudagrass stunting disease. *International Journal of*  
470 *Systematic Bacteriology* **34**: 107–117.
- 471 7. Dempsey, A.W., Walker, J.T. 1973. Efficacy of calcium and sodium hypochlorite for seed  
472 treatment of pepper. *Horticultural Science* **8**: 328–329.
- 473 8. Dhanvantari, B.N. 1989. Effect of seed extraction methods and seed treatments on control of  
474 tomato bacterial canker. *Canadian Journal of Plant Pathology* **11**: 400–408.

- 475 9. Dutta, B., Gitaitis, R., Sanders, H., Booth, C., Smith, S., Langston, D.B. 2014. Role of blossom  
476 colonization in pepper seed infestation by *Xanthomonas euvesicatoria*. *Phytopathology* **104**: 232-  
477 239.
- 478 10. Dutta, B., Gitaitis, R.D., Smith, S., Van Keith Searcy, J., Langston, D. 2016. Pathogen  
479 distribution, incubation period and seedling transmission resulting from secondary contamination of  
480 pepper seeds with *Xanthomonas euvesicatoria*. *Seed Science & Technology* **44(1)**: 1-10.  
481 [doi:10.15258/sst.2016.44.1.11](https://doi.org/10.15258/sst.2016.44.1.11)
- 482 11. Dye, D.W. 1962. The inadequacy of the usual determinative tests for the identification of  
483 *Xanthomonas* spp. *New Zealand Journal of Science* **5**: 393-416.
- 484 12. EFSA PLH Panel (EFSA Panel on Plant Health), 2014. Scientific Opinion on the pest  
485 categorisation of *Xanthomonas campestris* pv. *vesicatoria* (Doidge) Dye. *EFSA Journal* **12(6)**:  
486 3720, 26 pp. doi:10.2903/j.efsa.2014.3720..
- 487 13. EPPO, 2013a. PM 7/110 (1) *Xanthomonas* spp. (*Xanthomonas euvesicatoria*, *Xanthomonas*  
488 *gardneri*, *Xanthomonas perforans*, *Xanthomonas vesicatoria*) causing bacterial spot of tomato and  
489 sweet pepper. *EPPO Bulletin* **43**: 7–20. doi:10.1111/epp.12018.
- 490 14. Giovanardi, D., Biondi, E., Perez, S., Ignjatov, M., Gasic, K., Ferrari, M., Jevtic R., Stefani E.  
491 2015. Seed transmission of *Xanthomonas vesicatoria* and *Clavibacter michiganensis* subsp.  
492 *michiganensis* in tomato and *Xanthomonas euvesicatoria* in pepper and implementation of seed  
493 disinfection methods. In: D. Marčić, M. Glavendekić, P. Nicot (Eds.). *Proceedings of the VII*  
494 *Congress on Plant Protection. Plant Protection Society of Serbia, IOBC-EPRS, IOBC-WPRS,*  
495 *Belgrade, Serbia*: 53-58.
- 496 15. Gitaitis, R., Walcott, R.R. 2007. The epidemiology and management of seedborne bacterial  
497 diseases. *Annual Review of Phytopathology* **45**: 371-397.
- 498 16. Ignjatov, M., Gasić, K., Ivanović, M., Šević, M., Obradović, A., Milošević, M. 2010.  
499 Characterisation of *Xanthomonas euvesicatoria* strains pathogens of pepper in Serbia. *Pesticidi i*  
500 *Fitomedicina* **25**: 139–149.

501 17. ISTA (International Rules for Seed Testing). 2009. International Seed Testing Association,  
502 Zurich.

503 18. Jones, J. B., Jones, J. P., and Woltz, S. S. 1993. The effect of selected copper bactericides on  
504 population dynamics of *Xanthomonas campestris* pv. *vesicatoria* on tomato leaflets. Proceedings of  
505 Florida State Horticultural Society, 106: 160-163.

506 19. Jones, J.B., Lacy, G.H., Bouzar, H., Stall, R.E, Schaad, N.W. 2004. Reclassification of the  
507 *xanthomonads* associated with bacterial spot disease of tomato and pepper. *Systematic and Applied*  
508 *Microbiology* **27**: 755–762.

509 20. Kocks, C.G., Zadoks, J.C. 1996. Cabbage refuse piles as sources of *inoculum* for black rot  
510 epidemics. *Plant Disease* **80**: 789–792.

511 21. Koenraad, H., van Betteray, B., Germain, R., Hiddink, G., Jones, J.B., Oosterhof, J.,  
512 Rijlaarsdam, A., Roorda, P., Woudt, B. 2009. Development of specific primers for the molecular  
513 detection of bacterial spot of pepper and tomato. Proceedings of the 2<sup>nd</sup> International Symposium on  
514 Tomato Diseases. (Eds. Saygili, H., Sahin, F. and Aysan, Y.). *Acta Horticulturae* **808**: 99–102

515 22. Leite, R. P., Jones, J. B., Somodi, G. C., Minsavage, G. V., and Stall, R. E. 1995. Detection of  
516 *Xanthomonas campestris* pv. *vesicatoria* associated with pepper and tomato seed by DNA  
517 amplification. *Plant Disease* **79**: 917-922.

518 23. Madden, L.V., Hughes, G., Van den Bosch, F. 2007. The study of plant disease epidemics.  
519 APS, St Paul, MN.

520 24. McGuire, R.G., Jones, J.B., Sasser, M. 1986. Tween media for semiselective isolation of  
521 *Xanthomonas campestris* pv. *vesicatoria* from soil and plant material. *Plant Disease* **70**: 887–891.

522 25. . Oliver, J. D. (2005). The viable but nonculturable state in bacteria. *Journal of Microbiology*,  
523 **43**: 93–100.

524 26. Opeña, R.T., Chen, J.T., Kalb, T., Hanson, P. 2001. Seed Production of Open-Pollinated  
525 Tomato Lines. Several International Cooperator's Guides. Asian Vegetable Research and  
526 Development Center, pub # 01-528.

- 527 27. Schaad, N.W., Sitterley, W.R., Mumaydan, H. 1980. Relation of incidence of seedborne  
528 *Xanthomonas campestris* to black rot of crucifers. *Plant Disease* **64**: 91-92.
- 529 28. Smith, E.F. 1910. A new tomato disease of economic importance. *Science* **803**: 794–796.
- 530 29. Stokell, J.R., and Steck, T.R. 2012. Viable but nonculturable bacteria. In: eLS. John Wiley &  
531 Sons Ltd, Chichester. doi: 10.1002/9780470015902.a0000407.pub2
- 532 30. Strider, D. L. (1969). Bacterial canker of tomato caused by *Corynebacterium michiganense*: a  
533 literature review and bibliography. Technical bulletin 193, North Carolina Agricultural  
534 Experimental Station.
- 535 31. Taylor, J.D., Dudley, C.L., Presly, L. 1979. Studies of halo-blight seed infection and disease  
536 transmission in dwarf beans. *Annals of Applied Biology* **93**: 267-277.
- 537 32. Van der Plank, J. E. 1963. Plant Diseases: Epidemics and Control. Academic Press, New York.
- 538 33. Vauterin, L., Hoste, B., Kersters, K., Swings, J. 1995. Reclassification of *Xanthomonas*.  
539 *International Journal of Systematic Bacteriology* **30**: 472–489.
- 540 34. Versalovic, J., Koeuth, T., Lupski, J. R. 1991. Distribution of repetitive DNA sequences in  
541 eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acid Research* **19**: 6823–  
542 6831.
- 543 35. Zadoks, J. C., Van den Bosch, F. 1994. On the spread of plant disease: A theory on foci.  
544 *Annual Review of Phytopathology* **32**: 503-521.

545

## 546   **Figures**

547

548   Figure 1. Disease progression curves of bacterial leaf spot caused by *Xanthomonas vesicatoria* on  
549   processing tomato plants (cv. VF 10) from the beginning of July to the 1<sup>st</sup> of August during the first  
550   year of experiments in Italy. Bars represent standard deviations at each phytopathometric  
551   assessment. The percentage values in the legend represent the initial percentage of inoculated plants  
552   in each plot of the experimental field.

553

554   Figure 2. Disease progression curves of bacterial leaf spot caused by *Xanthomonas vesicatoria* on  
555   fresh market tomato plants (cv. Jabučar) from the beginning of July to the second week of  
556   September during the first year of experiments in Serbia. Bars represent standard deviations at each  
557   phytopathometric assessment. The percentage values in the legend represent the initial percentage  
558   of inoculated plants in each plot of the experimental field.

559

560   Figure 3. Disease progression curves of bacterial leaf spot caused by *Xanthomonas euvesicatoria* on  
561   bell pepper plants (cv. Amphora) from the beginning of July to the end of September during the first  
562   year of experiments in Serbia. Bars represent standard deviations at each phytopathometric  
563   assessment. The percentage values in the legend represent the initial percentage of inoculated plants  
564   in each plot of the experimental field.

565

566



567 **Tables**

568

569 Table 1. AUDPC values calculated for the different pathosystems in Italy and Serbia during the first  
570 cropping season. Different letters indicate different statistical classes (Duncan's test, p 0.05).

571

572 Table 2. Contamination levels and rates of *Xanthomonas vesicatoria* (Xv) and *Xanthomonas*  
573 *euvesicatoria* (Xe) in tomato and pepper seeds, respectively, produced in Italy and Serbia during the  
574 first cropping season from field plots experimentally infected at different initial contamination  
575 incidences (1%, 5% and 15%). The bacterial colonies (CFU) are related to 1.0 g<sup>-1</sup> of seeds. To  
576 determine the seed contamination rate of each seed lot, 50 replicates of 100 seeds were tested. Then,  
577 their contamination rate (p) was calculated according to the formula  $p = 1 - (Y/N)^{1/n}$ , where N is the  
578 number of analysed replicates, n is the number of seeds in a replicate and Y is the number of non-  
579 contaminated sub-samples (Maury *et al.* 1985).

580

581

582

583

584

585

586

587

588

589

590

591

592

593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615

## **SUPPLEMENTARY MATERIAL**

Figure S1. Design of experimental field plots used during the first cropping season in Italy and Serbia. This design was used to evaluate and quantify disease outbreaks under field conditions and assess disease progression from different initial infection percentage of approx. 1%, 5% and 15% by *Xanthomonas vesicatoria* in processing and table tomato and *Xanthomonas euvesicatoria* in bell pepper. The position of each experimentally inoculated plant is highlighted by a cross.

Figure S2. Meteorological data reported in Italy (A) and Serbia (B) during the first cropping season in 2013 from the 1<sup>st</sup> of April to the last harvest.

Figure S3. Meteorological data reported in Italy (A) and Serbia (B) during the second cropping season in 2015 from the 1<sup>st</sup> of April to last harvest.