

This is the peer reviewed version of the following article:

Non-blood sources of cell-free DNA for cancer molecular profiling in clinical pathology and oncology /
Ponti, G.; Manfredini, M.; Tomasi, A.. - In: CRITICAL REVIEWS IN ONCOLOGY HEMATOLOGY. - ISSN 1040-
8428. - 141:(2019), pp. 36-42. [10.1016/j.critrevonc.2019.06.005]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

18/12/2025 18:57

Accepted Manuscript


Title: Non-blood sources of cell-free DNA for cancer molecular profiling in clinical pathology and oncology

Author: Giovanni Ponti

PII: S1040-8428(18)30346-9
DOI: <https://doi.org/10.1016/j.critrevonc.2019.06.005>
Reference: ONCH 2767

To appear in: *Critical Reviews in Oncology/Hematology*

Received date: 24 July 2018
Revised date: 19 September 2018
Accepted date: 4 June 2019



Please cite this article as: Ponti G, Non-blood sources of cell-free DNA for cancer molecular profiling in clinical pathology and oncology, *Critical Reviews in Oncology / Hematology* (2019), <https://doi.org/10.1016/j.critrevonc.2019.06.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Non-blood sources of cell-free DNA for cancer molecular profiling in clinical pathology and oncology

Corresponding author

Giovanni Ponti

Department of Surgical, Medical, Dental & Morphological Sciences with Interest Transplant, Oncological & Regenerative Medicine, Clinical Pathology Unit, University of Modena & Reggio Emilia, Via del Pozzo n 71, Modena, 41100, Italy.

giovanni.ponti@unimore.it

Highlights

- Liquid biopsy can quantify and qualify cell-free and tumour-derived DNA fragments in the bloodstream.
- CfDNA quantification and mutation analysis can be applied as novel oncologic biomarkers.
- Non-blood sources of cfDNA include cerebrospinal fluids, urine, sputum, saliva, pleural effusion, stool and seminal fluid.
- Seminal plasma cfDNA quantification can be used as prostate cancer (PCa) biomarker.
- cfDNA analysis from non-blood biological liquids will become routine clinical practice for cancer patient diagnosis and management.

Table of contents

1. Introduction	5
2. Biology of Cell free DNA and cell tumoral DNA	8
3. Sources of cfDNA in oncology	11
3.1 cfDNA in saliva and sputum	11
3.2 cfDNA in urine	11
3.3 cfDNA in cerebrospinal fluid	13
3.4 cfDNA in stool	13
3.5 CtDNA in pleural fluids	14
3.6 CtDNA in ascites	15
3.7 cfDNA in seminal plasma	15
4. cfDNA and cancer patient management	18
4.1 Measuring disease burden	18
4.2 Patient stratification and prediction of therapeutic response	19
5. Future directions: Potential non-invasive sources of DNA in clinical pathology and oncology: Oral mucosa and skin	20
6. Conclusion	21
References	22

Abstract

Liquid biopsy can quantify and qualify cell-free (cfDNA) and tumour-derived (ctDNA) DNA fragments in the bloodstream. CfDNA quantification and mutation analysis can be applied to diagnosis, follow-up and therapeutic management as novel oncologic biomarkers. However, some tumor-types release a low amount of DNA into the bloodstream, hampering diagnosis through standard liquid biopsy procedures. Several tumors, as such as brain, kidney, prostate, and thyroid cancer, are in direct contact with other body fluids and may be alternative sources for cfDNA and ctDNA. Non-blood sources of cfDNA/ctDNA useful as novel oncologic biomarkers include cerebrospinal fluids, urine, sputum, saliva, pleural effusion, stool and seminal fluid. Seminal plasma cfDNA, which can be analyzed with cost-effective procedures, may provide powerful information capable to revolutionize prostate cancer (PCa) patient diagnosis and management. In the near future, cfDNA analysis from non-blood biological liquids will become routine clinical practice for cancer patient diagnosis and management.

Keywords: Cell-free DNA, non blood cfDNA, Liquid biopsy, biomarkers, cancer diagnosis, cancer management.

1. Introduction

Over recent decades there has been a rapid expansion of knowledge in the field of molecular biology and biochemistry, leading to the development of precision medicine and tailored therapies.(Lu and Liang 2016) The understanding of oncogenic pathways and neoplastic genetic signatures shed the conceptual basis for the development of liquid biopsy procedures and oncological-targeted therapies. Today, immunohistochemical and biochemical tumor

characterization are part of the routine process to define patient diagnosis and prognosis.(G. Ponti et al. 2016; Hofman et al. 2018; G. Ponti, Manfredini, Pastorino, Maccaferri, et al. 2018)

The identification of new diagnostic and prognostic markers from non-tumoral biological samples is an interesting field in continuous evolution. Liquid biopsy procedures have been developed to identify oncologic biomarkers in several body fluids such as blood, plasma, cerebrospinal liquid, seminal fluid, saliva and urine.(Ulrich and Paweletz 2018; Stewart et al. 2018; Burgener et al. 2017) The concept that body fluids may reveal the presence of several systemic diseases dates back to ancient Greek and Indian history. The development of humoral theory was attributed to Hippocrates (ca. 460–370 BCE), and its concepts were the base of Western medicine from antiquity through to the 19th century. “Humoral” derives from the word “humor,” which in ancient greek means “fluid”, and health was defined as the proper humoral balance for that individual. The most extensively studied body fluids are blood and urine, but many other liquids, such as cerebrospinal fluid, saliva and seminal fluids have been analyzed in order to determine their diagnostic and prognostic potential.

An example of a routinely applied liquid biopsy for patient management is the dosage of prostate specific antigen (PSA) in the bloodstream. However, the PSA protein is present both in healthy and in neoplastic prostate cells and therefore has a low sensitivity and specificity for prostate cancer (PCa) diagnosis. In contrast, tumor-derived cell-free DNA (ctDNA) based techniques are more specific and can be applied to many body fluids in order to identify novel oncologic biomarkers for diagnosis, prognosis and monitoring of tumor therapeutic response.

The presence of DNA within the non-cellular fraction of peripheral blood, termed cell-free DNA (cfDNA), was initially identified more than 50 years ago.(Ulrich and Paweletz 2018; Stewart et al. 2018; Burgener et al. 2017) Decades passed before cfDNA levels were found to be elevated within the serum of cancer patients, with at least a portion of the cfDNA being tumor-derived (ctDNA). The advantage of ctDNA analysis from liquids other than blood is the collection of

targeted information from a body fluid directly in contact with the tumor source (eg. cerebrospinal fluid for central nervous system neoplasm). Furthermore, contrary to what occurs with ctDNA in the bloodstream, ctDNA in other bodily fluids are not diluted and higher concentrations of ctDNA are derived from active secretion and/or necrotic phenomena.

ctDNA harbors cancer-specific genetic and epigenetic alterations that allow its detection and quantification using a variety of emerging techniques. The promise of convenient non-invasive access to the complex and dynamic molecular features of cancer through peripheral blood has galvanized translational researchers around this topic with compelling routes to clinical implementation, particularly in the post -treatment surveillance setting. (Ulrich and Paweletz 2018; Stewart et al. 2018; Burgener et al. 2017) Although analysis methods must contend with small quantities of ctDNA present in most patients and the relative over-abundance of background cfDNA derived from normal tissues, recent technical innovations have led to dramatic improvements in the sensitivity of ctDNA detection. The collection of biologic samples to isolate cfDNA is often repeatable, allowing for real-time and dynamic monitoring of molecular changes.(Zeng et al. 2018) As a result, ever more studies are investigating the clinical utility of ctDNA for applications in treatment response assessment,(Dawson et al. 2013) identification of emerging resistance mechanisms,(Mok et al. 2015; Tabernero et al. 2015) and minimal residual disease detection.(Guo et al. 2016) In addition, ctDNA is released from multiple tumor regions, and may thereby represent the whole molecular picture of a patient's malignancy, potentially solving the problem of intra-tumor heterogeneity,(L De Mattos-Arruda et al. 2014; Leticia De Mattos-Arruda et al. 2015; Jamal-Hanjani et al. 2016) which may lead to false-negative results and suboptimal therapy selection, and characterization of clonal heterogeneity and selection. (Ulrich and Paweletz 2018; Stewart et al. 2018; Burgener et al. 2017).

In this review we explore the application of non-blood derived ctDNA and cfDNA in the discovery of novel oncological biomarkers, describing the existing evidence and focusing on the

recent development of seminal plasma analysis.

ACCEPTED MANUSCRIPT

2. Biology of Cell free DNA and cell tumoral DNA

Cell-free DNA (cfDNA) was first identified in human plasma in 1948 and is believed to be released from cells throughout the body into the blood stream.(MANDEL and METAIS 1948) Most cfDNA in the blood originates from hematopoietic cells(Allam et al. 2014; Stroun et al. 1989) however many other organs and tissues contribute to the total amount of blood cfDNA.(Stroun et al. 2001)

The origin of cfDNA has not fully been explored, but there are many mechanisms by which DNA gets into the circulation. Blood cfDNA fragments of healthy individuals are primarily of lymphoid and myeloid origin.(Anker et al. 1999) Once outside of the cell, cfDNA is steadily degraded by nucleases, possibly with the help of macrophages(Chused, Steinberg, and Talal 1972) and excreted into urine via the renal system.(Thierry et al. 2016; Botezatu et al. 2000) Multiple studies have demonstrated that the majority of cfDNAs are short molecules around 166-167bp, although longer fragments also exist.(Jiang and Lo 2016; Snyder et al. 2016) While necrosis results in longer fragments (>10.000 bp), apoptosis is associated with DNA fragments of about 180bp, or multiples of this length, which appear as a ladder electrophoresis pattern. cfDNA is likely associated with nucleosomes as the length of 167 bp corresponds approximately to the length of DNA wrapped around a histone.(Snyder et al. 2016) In addition to cell death, neutrophils can mediate the immune response by releasing neutrophil extracellular traps (NETs) that can trap and kill various pathogens.(Kaplan and Radic 2012) These are extracellular network structures composed of both nuclear and mitochondrial DNA fibers, which are covered by various proteins such as histones and proteases.(Kaplan and Radic 2012) Another way of releasing DNA into the circulation is active release of newly synthesized DNA via vesicles and lipoproteonucleotide complexes.(Fuchs et al. 2012) Based on the appearance in the circulation, cfDNA molecules can be divided into three basic categories: free DNA fragments, vesicle-bound DNA and DNA-macromolecular complexes.

In cancer, a portion of cfDNA originates from tumor cells, referred to as circulating-tumor DNA (ctDNA), and can harbor specific mutations corresponding to the patient's tumor. ctDNA profiling has recently become an area of increasing clinical relevance in oncology, in particular due to advances in the sensitivity of sequencing technologies, allowing a reliable identification of cancer mutations. Several studies showed high concordance between individual mutations found in ctDNA samples and tumor tissue.(Bettegowda et al. 2014) Cancer patients have much higher cfDNA concentrations (0 to >1,000 ng/mL) than healthy individuals (0-100 ng/mL) and the total amount varies among patients with comparable cancer type and stage.(Bettegowda et al. 2014) Direct correlations between ctDNA level and tumor burden, stage, vascularity and therapy response have been described for several cancers.

cfDNA can be found in plasma as well as other body fluids such as urine, cerebral spinal fluid (CSF), pleural fluid, and saliva, among others.(Stewart and Tsui 2018) Depending on tumor type, different rates of cfDNA/ctDNA have been reported in blood samples. CtDNA was detected in over 75% of patients with advanced pancreatic, ovarian, colorectal, bladder, gastroesophageal, breast, melanoma, hepatocellular, and head and neck cancers. However, levels below 50% were found in primary brain tumors, in kidney, prostate, and thyroid cancers,(Bettegowda et al. 2014) and may be due to the influence other factors, such as blood-brain barrier restriction mechanisms, inflammatory or autoimmune processes that increase non-tumor cfDNA fraction, effectively diluting the ctDNA amount. The identification of ctDNA from biologic fluids other than blood, enables prompt diagnosis and a reliable genetic identification of cancer types, characterized by shedding low concentrations of ctDNA into the blood stream. These tumors are often in direct contact with other body fluids, such as urine, saliva, cerebrospinal fluid, pleural effusion or seminal liquid, releasing a considerable amount of ctDNA and creating a privileged source for non-invasive ctDNA quantification and characterization.

3. Sources of cfDNA in oncology

3.1 cfDNA in saliva and sputum

Saliva provides good-quality genomic DNA, which is comparable to blood as a template for genotyping.(Bahlo et al. 2010; Hu et al. 2012; Siravegna et al. 2017) Salivary DNA has been used for the detection of germline mutations in various screening studies, such as for breast and brain cancers.(Dean et al. 2015; Adel Fahmideh et al. 2015) Numerous studies have shown that sputum tumor DNA could be a promising tool for early detection of lung cancer. Recent studies have shown proof-of-principle for ctDNA as a biomarker in head and neck squamous cell carcinoma (HNSCC), with both blood and saliva serving as a diagnostic medium. In a cohort of 93 HNSCC patients, plasma or saliva samples were collected to identify somatic mutations (TP53, PIK3CA, CDKN2A, HRAS, NRAS) and HPV (HPV-16, -18).(Y. Wang, Springer, Mulvey, et al. 2015) Tumor DNA was detected in 76% of saliva samples (n=93) and 87% of plasma samples (n=47) respectively, and TP53 was the most common mutant detected (86%). Patients with oral cavity cancers were all shown to have tumor-specific DNA in their saliva samples (100%, n=46). Saliva was shown to be a more sensitive predictor than plasma for early stage disease, with 100% detection in saliva versus 70% in plasma.(Y. Wang, Springer, Mulvey, et al. 2015)

3.2 cfDNA in urine

Several reports have analyzed the value of urinary cfDNA (UCF-DNA) for the diagnosis of bladder cancer (BC).(Todenhöfer et al. 2018) At the present time, the standard diagnostic methods for BC are cystoscopy and urine cytology. However, urine cytology has poor sensitivity (except for high grade tumors)(Bier et al. 2018) and, although flexible cystoscopy has been recently introduced, the procedure is both invasive and uncomfortable. Therefore, there is growing interest in new non-invasive diagnostic tools that have better sensitivity and specificity for BC. To overcome these

difficulties, several urine-based biomarkers, such as bladder tumor antigen, nuclear matrix protein 22 (NMP22), and fluorescence *in situ* hybridization (FISH), have been studied but none of them have been able to demonstrate superiority to cystoscopy or cytology.(Breen et al. 2015)

The main source of UCF-DNA is thought to be apoptotic and necrotic cancer cells.(Bryzgunova and Laktionov, n.d.) UCF-DNA might be gathered as a result of renal cfDNA transport from the blood or direct contact from urinary tracts. Most of the UCF-DNA of urinary tract cancers patients is not derived from blood, but is rather derived directly from tumor cells (ctDNA). Kim TW et al. demonstrated that the non-invasive quantification of IQ motif containing GTPase activating protein 3 (IQGAP3) urinary nucleic acids (NA) could be a suitable tool for distinguishing between BC patients and patients with non-cancer-associated hematuria.(Kim et al. 2018) In other reports, UCF-DNA profiling using PCR-based detection or high throughput sequencing technology of tumor-associated genes, has been indicated as a technique to yield promising results in bladder and PCa. Recently Christensen *et al.* applied a digital droplet PCR (ddPCR) approach for the detection of common mutations in UCF-DNA, suggesting that the levels of mutant ctDNA in the urine of non-muscle-invasive bladder cancer (NMIBC) patients were positively correlated with tumor stage, grade and size, and that a high initial level of mutant urinary ctDNA was predictive of future disease progression.(Christensen et al. 2018) In a cystectomy patient group, high mutant UCF-DNA was able to predict future disease recurrence, the association being more pronounced with ctDNA.

PCa represents one of the most common tumors in European men and one of the leading causes of male cancer associated deaths.(Alberts, Schoots, and Roobol 2015) At present, the only noninvasive approach currently used for the diagnosis of PCa is the determination of PSA (prostate-specific antigen) in blood, which has been shown to reduce PCa mortality. However, the use of PSA has recently been questioned because of its low accuracy, especially in terms of specificity.(Alberts, Schoots, and Roobol 2015; Ilic et al. 2018) Previous studies identified UCF-DNA integrity as a

potentially good marker for the early diagnosis of noninvasive PCa, with an overall diagnostic accuracy of about 80% in small patient series.(Casadio et al. 2013; Salvi et al. 2015) UCF-DNA is easily quantifiable in PCa patients and could prove to be an important source of biomarkers, such as gene mutations or epigenetic modifications, that may accurately assist in distinguishing PCa from other benign diseases of the urogenital tract.

3.3 cfDNA in cerebrospinal fluid

Tumor-derived DNA typically constitutes a small fraction of all cfDNA in plasma, while the proportion of such DNA in cerebrospinal fluid (CSF) is much higher due to the lower background of normal DNA.(Leticia De Mattos-Arruda et al. 2015; Y. Wang, Springer, Zhang, et al. 2015) Owing to the blood–brain barrier, CSF cfDNA is unable to circulate fully within the blood system, resulting in a limited amount of cfDNA from central nervous system (CNS) being released into blood plasma. Thus, blood plasma is not the best biologic liquid for the detection and characterization of intracranial lesions.(Pentsova et al. 2016; Pan et al. 2015; Li et al. 2018; Yang et al. 2014; Alix-Panabières and Pantel 2016) CSF circulates throughout the CNS and may provide potential information on intracranial lesions. Few studies indicated that CSF could be an important method of liquid biopsy in patients with CNS cancers, even though lumbar puncture is an invasive procedure that should be performed only if necessary. Moreover, selected gene profiles in CSF could be consistent with their primary tumors.(Pentsova et al. 2016; Pan et al. 2015; Li et al. 2018; Yang et al. 2014; Alix-Panabières and Pantel 2016) Li YS et al. demonstrated that CSF liquid biopsy has potential clinical applications for diagnosis and characterization of leptomeningeal metastasis (LM) of non-small cell lung cancer (NSCLC) with mutated Epidermal Growth Factor Receptor (EGFR).(Li et al. 2018) Moreover, CSF cfDNA has a potential role in revealing dynamic changes of the tumor burden of LM throughout treatment, and further studies should be conducted to explore its roles in prediction and prognosis. (Li et al. 2018)

3.4 cfDNA in stool

Colorectal cancer (CRC), a common cause of cancer-related mortality worldwide, is preventable with effective screening and removal of precursor lesions.(Berger and Ahlquist 2012) Yet, screening efforts have been hampered by low participation rates and by performance limitations of the screening tools themselves. Stool DNA testing has emerged as a biologically rational and user-friendly strategy for the non-invasive detection of both CRC and critical precursor lesions.(Berger and Ahlquist 2012; Ahlquist et al. 2008; Redwood et al. 2016) Unlike most conventional screening tools, stool DNA testing detects proximal and distal colorectal neoplasms with higher sensitivity compared to fecal immunochemical testing for hemoglobin (FIT), for the detection of screening-relevant colorectal neoplasia (SRN). Several key technical advances have led to increasingly accurate approaches for stool DNA testing, including the use of a DNA preservative buffer with stool collection, efficient target capture and amplification methods, broadly informative multi-marker panels, and automated assay components.(Imperiale et al. 2004; Imperiale et al. 2014; Ahlquist et al. 2008) Based on previous studies, advanced multi-marker stool DNA tests including methylated markers, mutation markers and an assessment of faecal haemoglobin, have been shown to detect CRC at sensitivities of 85% and higher and adenomas >1 cm at 60% and higher in a case-control environment.(Berger and Ahlquist 2012) In a large population cross-sectional study, Imperiale et al. demonstrated that the sensitivity of the DNA test for the detection of both CRC (92.3%) and advanced precancerous lesions (42.4%) exceeded that of FIT by an absolute difference of nearly 20 percentage points.(Imperiale et al. 2014)

3.5 CtDNA in pleural fluids

Pleural effusion is a common complication of lung cancer. The collection of pleural effusion fluid or of bronchial washing samples with physiological saline solutions is currently used in

diagnosing cancers of the respiratory system.(Kimura et al. 2006; Soh et al. 2006) The detection of *EGFR* mutations in cytological samples of pleural effusion fluid is feasible, although often difficult owing to the limited number of cancer cells that are usually available for analysis. Kimura et al. demonstrated that pleural effusion cfDNA can be used to detect *EGFR* mutations and that the *EGFR* mutation status may be a useful predictor of the gefitinib response in patients with non-small cell lung cancer (NSCLC).(Kimura et al. 2006) In another study, the feasibility of identifying *EGFR* mutations in tumor derived DNA collected through bronchial washings, termed cytology cell free DNA (ccfDNA), was examined.(Kawahara et al. 2015) The results demonstrated the high sensitivity and specificity (88% and 100%, respectively) of this approach compared with the analysis of DNA from tumor tissue, suggesting that activating *EGFR* mutations can be accurately detected in ccfDNA. Thus, ccfDNA might be a valuable alternative to cytological samples, although larger investigations are needed to validate this diagnostic approach. A limited number of studies have investigated the diagnostic, prognostic, or predictive value of miRNAs in pleural effusion fluid from patients with NSCLC.(Han et al. 2013; T. Wang et al. 2012) In one study, the authors found that a signature comprising five miRNAs in the effusion samples was predictive of the overall survival of patients with NSCLC and malignant pleural effusion.

3.6 CtDNA in ascites

Detection and characterization of cfDNA in ascites has only been analyzed in one preliminary study.(Husain et al. 2017) Husain H. et al. isolated cfDNA from ascites and demonstrated the presence of tumorigenic copy number variations (CNVs) in cancer-associated genes in a small series of 6 metastatic cancer patients, providing a rationale for the study of ascites as a source of ctDNA for the comprehensive analysis of relevant targets.(Husain et al. 2017) In addition, in their small patient series, the ascitic cfDNA of one patient was characterized by a 15-fold amplification of the *EGFR* gene in ascites cfDNA analysis, which was not present in two separate lung biopsies of the tumor. Even though the significance of detecting discordant alterations in primary tissue

versus blood versus ascites is unclear, the authors hypothesized that ascites may provide important genomic information regarding an individual's cancer that may complement and expand data obtained from tissue biopsies.(Husain et al. 2017)

3.7 cfDNA in seminal plasma

The quantification of serum and plasma cfDNA in PCa patients has been found to be significantly higher with respect to age-matched healthy controls.(G. Ponti, Maccaferri, Manfredini, Kaleci, et al. 2018) It has been proven that increased levels of cfDNA are released and can be isolated from serum and plasma in PCa patients. Further, the level of cfDNA in PCa patients was found to be significantly higher than in benign prostatic hyperplasia (BPH) patients.(Wyatt et al. 2017; Feng et al. 2013)

It has recently been demonstrated for the first time that human seminal fluid can be a valuable source of cfDNA for the identification of novel oncological biomarkers. Seminal plasma cfDNA from PCa patients is significantly more concentrated than age-matched healthy individuals, but tumor stage has been found to be independent of other parameters, such as age at diagnosis. (Giovanni Ponti, Maccaferri, Micali, Manfredini, et al. 2018; G Ponti, Maccaferri, Manfredini, Cotugno, et al. 2018; Giovanni Ponti, Maccaferri, Mandrioli, et al. 2018) Fluorometric and electrophoretic assessments allow a reliable quantification and qualification of seminal plasma cfDNA, that could be routinely adopted for PCa screening programs. In the same study, seminal cfDNA of PCa patients yielded higher values than those of age-matched healthy volunteers, 1721.27 ng/ μ l ng/ μ l and 75.4 ng/ μ l, respectively.(G. Ponti, Maccaferri, Mandrioli, et al. 2018) This allows a reliable characterization and differentiation between a cohort of patients affected by PCa and the age-matched control group. The aforementioned average seminal cfDNA is notably higher than average blood cfDNA concentrations of PCa patients, being approximately 100 times higher

compared to mean blood cfDNA values reported in literature (1.8–35 ng/μl).(Costa et al. 2017) In another study, seminal cfDNA was compared to BPH patients, revealing a significant difference in the concentration levels of cfDNA in PCa and BPH patient cohorts. A possible cut-off level of 450 ng/ul seminal cfDNA was proposed as able to discriminate between the two distinct groups. In addition, the electrophoresis of seminal plasma cfDNA enabled the discrimination between PCa patients and BPH or age-matched healthy individuals, because of a distinct electrophoretic pattern in PCa patients.

Seminal cfDNA levels are a potential clinical biomarker in early PCa diagnosis, referring important information regarding tumor characterization, patient prognosis and management, for the determination of therapeutic strategies and subsequent follow-up.

4. cfDNA and cancer patient management

There are several important applications in which liquid biopsy might confer an advantage to patients with cancer: the potential use of cfDNA as a biomarker include applications in the diagnostic, prognostic and predictive settings. Therefore, CfDNA quantification and analysis should be included in a management strategy that considers the patient's clinical status, the clinical relevance of test results, and local feasibility of the different testing methods. Liquid biopsy can be considered at the time of initial diagnosis in all patients who need tumor molecular profiling, but it is particularly recommended when tumor tissue is scarce, unavailable, or a significant delay is expected in obtaining tumor tissue.(Stewart and Tsui 2018) CfDNA originating from biologic samples such as blood or other biological body fluids can be analyzed through quantitative and qualitative analysis, guiding cancer patient management, as demonstrated in many cancer types. For instance, in NSCLC treatment-naïve patients, *EGFR* and *ALK* rearrangement assessment using ctDNA is currently a recommended procedures for cancer management. (Rolfo et al. 2018; Marmarelis et al. 2017)

Several biologic fluids are under investigation for the diagnostic, prognostic and therapeutic value of cfDNA quantification and analysis. UCF-DNA and scfDNA are the most striking examples of non-blood biologic fluids, containing large amounts of cfDNA, whose quantification has a great prognostic and therapeutic value.

Cancer is a dynamic disease and ctDNA released by tumor cells strictly reflects the heterogeneity of both primary cancer and metastases.(Stewart et al. 2018; Stewart and Tsui 2018) In this scenario, to improve the standard of patient care, the institution of a Molecular Tumor Board (MTB) has been proposed in several hospitals to discuss the best treatment option for the patient, considering molecular testing results, including those from liquid biopsies. Some initial experiences have been reported.(Rolfo et al. 2018; Harada et al. 2017) Rolfo et al. created an MTB that retrospectively evaluated 141 patients, recommending a treatment in 78 (55%) of patients.(Rolfo et

al. 2018) The group of Harada et al. also used MTB for the selection of cancer patients who should be advised to perform genetic testing; resulting in the approval of 132/191 cases for NGS analysis.(Harada et al. 2017)

4.1 Measuring disease burden

The clinical utility of liquid biopsy has been explored in different clinical phases for several tumors, from early disease detection to the identification of prognostic factors in early stages and to the molecular characterization of metastatic disease and eventual relapse. The presence of ctDNA itself is indicative of disease and the amount of ctDNA can also be an indicator of the amount of disease. As previously discussed, the amount of ctDNA is correlated with tumor stage, and many groups have observed that higher levels of ctDNA have been associated with worse survival outcomes in patients(Lecomte et al. 2002; Gray et al. 2015; Gautschi et al. 2007; Stewart and Tsui 2018), and can be used as a measure of disease burden, along with imaging studies. Overall, ctDNA has been shown to be a better predictor of prognosis than other tumour markers and that ctDNA concentration increase correlates with poorer clinical and radiological outcomes.

In CRC patients and in PCa patients, those with higher ctDNA levels for selected genes had worse outcomes with respect to those with lower ctDNA levels.(Romanel et al. 2015) While this observation is probably due to the correlation between ctDNA levels and tumor burden, the ability of cancer cells to shed DNA in circulation may also reflect disease aggressiveness.

4.2 Patient stratification and prediction of therapeutic response

Identifying molecular biomarkers in early tumor patients is needed in order to develop more personalized follow-up and treatment schedules. Detection of gene mutations in liquid biopsy for early detection of recurrence requires highly sensitive techniques. Molecular profiling tests prior to treatment provides the possibility of stratifying patients based on prognosis for the administration of adjuvant therapy (Romanel et al. 2015; Scherer et al. 2016) or for the selection of specific targeted therapies. By measuring several mutations, either in blood or in other body fluids, the changes in their ratios can provide some insight into the tumor's evolution and continued heterogeneity during treatment. (Misale et al. 2012; Chabon et al. 2016) This can also be extended to identify the appearance of resistance mechanisms. Serial studies of CRC found positive selection of blood *KRAS* mutations during anti-EGFR therapy and a decline in their representation after withdrawal. (Misale et al. 2012) Similar results have been recorded in NSCLC patients treated with EGFR inhibitors, where resistance mutations were identified in ctDNA prior to clinical progression. (Sorensen et al. 2014) A 54-gene panel detected ctDNA in 58% of patients with multiple types of cancer, with 68% of those having an actionable mutation by an FDA-approved drug. (Schwaederle et al. 2016) Interestingly, DNA derived from NSCLC tumors can be detected with high sensitivity in urine and plasma, enabling diagnostic detection and monitoring of therapeutic responses from non-blood body fluids. (Reckamp et al. 2016)

ctDNA has also been shown to be relevant to predict recurrence after resection of locally advanced rectal cancer or liver metastases from CRC. (Parkinson et al. 2016) Overman et al. showed that post-operative detection of ctDNA was significantly correlated with relapse-free survival (RFS). Tie et al. showed that ctDNA analysis appears to be strongly predictive of recurrence among patients with both lower (pathological complete response) and higher risk (node positive) disease. (Parkinson et al. 2016) In this context, ctDNA could be exploited to closely monitor patients before and after surgery to identify high-risk patients for disease recurrence.

5. Future directions: Potential non-invasive sources of DNA in clinical pathology and oncology: Oral mucosa and skin

Guthrie/FTA card-based blood spots, buccal scrapes, and finger nail clippings are DNA-containing specimens that are uniquely accessible and thus attractive as alternative tissue sources.(Klassen et al. 2012) As an alternative source to blood or buccal swabs, or both, there have been many trials to discover a non-invasive method to collect DNA reference samples from various specimens of the human body, such as nails, hair and skin scales.(Hogervorst et al. 2014; Blumenberg 2012; Bond 2007; Ghatak, Muthukumaran, and Nachimuthu 2013; Albujja et al. 2018) The value of DNA profiling from skin or adnexal samples has been previously analyzed and reviewed for forensic purposes only and crime scene investigations, however the application of these DNA analysis methods to skin oncology may allow direct genomic tumor profiling from tumor derived skin scales, nail clippings of nail tumors or hair sampled from the tumoral skin surface.

Previous trials have attempted to overcome the disadvantages of traditional samples (eg. blood samples) with the advantages of new, less invasive sampling materials to ensure the largest possible inclusion of subjects or volunteers in the studies. In particular, some attempts have been made to collect reference samples from skin surface cells (Zamir et al. 2004; Kopka et al. 2011) supporting this methodology as easy and simple in an accessible area. Such a method could be extremely useful among certain cultures (e.g., in the Middle East), where the use of traditional samples or buccal swabs can be considered invasive or is otherwise socially unacceptable. Skin lifts are a promising source of non-blood tumoral DNA. In previously mentioned studies, all the genetic profiles generated from skin lifts and swabs were consistent with their corresponding buccal swab and blood samples, indicating that both body regions and recovery methods yield accurate profiles (Zamir et al. 2004; Kopka et al. 2011). The accuracy of these newer techniques is necessary for future evaluations of the diagnostic value of the tumoral DNA recovered from alternative non-blood sources.

6. Conclusion

Applications of liquid biopsies in oncology have emerged and developed at an incredible rate over the past 5 years. Many studies have shown that cfDNA quantification is able to improve cancer diagnosis and patient management, however, several tumor-types, as such as kidney, prostate, and thyroid cancer, shed a low amount of cfDNA into the bloodstream hampering the diagnosis through standard liquid biopsy procedures. Research on cfDNA quantification from other body fluids is a promising field for the identification of new biomarkers that will allow prompt diagnosis and a better management of cancer patients. The current authors recently demonstrated, for the first time, that human seminal fluid can be a valuable source of cfDNA for the identification of novel oncological biomarkers and seminal plasma cfDNA from PCa patients is significantly more concentrated than age-matched healthy individuals and patients affected by BPH.(G. Ponti, Maccaferri, Manfredini, Cotugno, et al. 2018)

In the near future, the quantification and analysis of cfDNA will probably become part of routine diagnostic and clinical management of cancer patients. In order to accomplish this goal, it will be important to standardize ctDNA quantification methods and allow ctDNA detection for rare molecular alterations in order to anticipate drug resistance.

Conflict of interest

The authors have no conflict of interest

References

- Adel Fahmideh, Maral, Catharina Lavebratt, Joachim Schüz, Martin Rösli, Tore Tynes, Michael A Grotzer, Christoffer Johansen, et al. 2015. "CCDC26, CDKN2BAS, RTEL1 and TERT Polymorphisms in Pediatric Brain Tumor Susceptibility." *Carcinogenesis* 36 (8): 876–82. doi:10.1093/carcin/bgv074.
- Ahlquist, David A, Daniel J Sargent, Charles L Loprinzi, Theodore R Levin, Douglas K Rex, Dennis J Ahnen, Kandice Knigge, et al. 2008. "Stool DNA and Occult Blood Testing for Screen Detection of Colorectal Neoplasia." *Annals of Internal Medicine* 149 (7): 441–50, W81. <http://www.ncbi.nlm.nih.gov/pubmed/18838724>.
- Alberts, Arnout R, Ivo G Schoots, and Monique J Roobol. 2015. "Prostate-Specific Antigen-Based Prostate Cancer Screening: Past and Future." *International Journal of Urology : Official Journal of the Japanese Urological Association* 22 (6): 524–32. doi:10.1111/iju.12750.
- Albujja, Mohammed H, Abdul Aziz Bin Dukhyil, Abdul Rauf Chaudhary, Ahmed Ch Kassab, Ahmed M Refaat, Saranya Ramesh Babu, Mohammad K Okla, and Sachil Kumar. 2018. "Evaluation of Skin Surface as an Alternative Source of Reference DNA Samples: A Pilot Study." *Journal of Forensic Sciences* 63 (1): 227–33. doi:10.1111/1556-4029.13468.
- Alix-Panabières, Catherine, and Klaus Pantel. 2016. "Clinical Applications of Circulating Tumor Cells and Circulating Tumor DNA as Liquid Biopsy." *Cancer Discovery* 6 (5): 479–91. doi:10.1158/2159-8290.CD-15-1483.
- Allam, Ramanjaneyulu, Santhosh V R Kumar, Murthy N Darisipudi, and Hans-Joachim Anders. 2014. "Extracellular Histones in Tissue Injury and Inflammation." *Journal of Molecular*

Medicine (Berlin, Germany) 92 (5): 465–72. doi:10.1007/s00109-014-1148-z.

Anker, P, H Mulcahy, X Q Chen, and M Stroun. 1999. “Detection of Circulating Tumour DNA in the Blood (Plasma/Serum) of Cancer Patients.” *Cancer Metastasis Reviews* 18 (1): 65–73. <http://www.ncbi.nlm.nih.gov/pubmed/10505546>.

Bahlo, Melanie, Jim Stankovich, Patrick Danoy, Peter F Hickey, Bruce V Taylor, Sharon R Browning, Australian and New Zealand Multiple Sclerosis Genetics Consortium (ANZgene), Matthew A Brown, and Justin P Rubio. 2010. “Saliva-Derived DNA Performs Well in Large-Scale, High-Density Single-Nucleotide Polymorphism Microarray Studies.” *Cancer Epidemiology, Biomarkers & Prevention : A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology* 19 (3): 794–98. doi:10.1158/1055-9965.EPI-09-0812.

Berger, Barry M, and David A Ahlquist. 2012. “Stool DNA Screening for Colorectal Neoplasia: Biological and Technical Basis for High Detection Rates.” *Pathology* 44 (2): 80–88. doi:10.1097/PAT.0b013e3283502fdf.

Bettegowda, Chetan, Mark Sausen, Rebecca J. Leary, Isaac Kinde, Yuxuan Wang, Nishant Agrawal, Bjarne R. Bartlett, et al. 2014. “Detection of Circulating Tumor DNA in Early- and Late-Stage Human Malignancies.” *Science Translational Medicine* 6 (224): 224ra24. doi:10.1126/scitranslmed.3007094.

Bier, Simone, Jörg Hennenlotter, Michael Esser, Sarah Mohrhardt, Steffen Rausch, Christian Schwentner, Moritz Maas, et al. 2018. “Performance of Urinary Markers for Detection of Upper Tract Urothelial Carcinoma: Is Upper Tract Urine More Accurate than Urine from the Bladder?” *Disease Markers* 2018: 1–5. doi:10.1155/2018/5823870.

Blumenberg, Miroslav. 2012. “SKINOMICS: Transcriptional Profiling in Dermatology and Skin Biology.” *Current Genomics* 13 (5): 363–68. doi:10.2174/138920212801619241.

- Bond, John W. 2007. "Value of DNA Evidence in Detecting Crime." *Journal of Forensic Sciences* 52 (1): 128–36. doi:10.1111/j.1556-4029.2006.00323.x.
- Botezatu, I, O Serdyuk, G Potapova, V Shelepov, R Alechina, Y Molyaka, V Ananév, et al. 2000. "Genetic Analysis of DNA Excreted in Urine: A New Approach for Detecting Specific Genomic DNA Sequences from Cells Dying in an Organism." *Clinical Chemistry* 46 (8 Pt 1): 1078–84. <http://www.ncbi.nlm.nih.gov/pubmed/10926886>.
- Brawley, Otis W. 2018. "Prostate Cancer Screening: And the Pendulum Swings." *Cancer*, May. doi:10.1002/cncr.31380.
- Breen, Vivienne, Nikola Kasabov, Ashish M Kamat, Elsie Jacobson, James M Suttie, Paul J O'Sullivan, Laimonis Kavalieris, and David G Darling. 2015. "A Holistic Comparative Analysis of Diagnostic Tests for Urothelial Carcinoma: A Study of Cxbladder Detect, UroVysion® FISH, NMP22® and Cytology Based on Imputation of Multiple Datasets." *BMC Medical Research Methodology* 15 (May): 45. doi:10.1186/s12874-015-0036-8.
- Bryzgunova, O E, and P P Laktionov. n.d. "Extracellular Nucleic Acids in Urine: Sources, Structure, Diagnostic Potential." *Acta Naturae* 7 (3): 48–54. <http://www.ncbi.nlm.nih.gov/pubmed/26483959>.
- Burgener, Justin M, Ariana Rostami, Daniel D De Carvalho, and Scott V Bratman. 2017. "Cell-Free DNA as a Post-Treatment Surveillance Strategy: Current Status." *Seminars in Oncology* 44 (5): 330–46. doi:10.1053/j.seminoncol.2018.01.009.
- Casadio, Valentina, Daniele Calistri, Samanta Salvi, Roberta Gunelli, Elisa Carretta, Dino Amadori, Rosella Silvestrini, and Wainer Zoli. 2013. "Urine Cell-Free DNA Integrity as a Marker for Early Prostate Cancer Diagnosis: A Pilot Study." *BioMed Research International* 2013: 270457. doi:10.1155/2013/270457.

- Chabon, Jacob J, Andrew D Simmons, Alexander F Lovejoy, Mohammad S Esfahani, Aaron M Newman, Henry J Haringsma, David M Kurtz, et al. 2016. "Corrigendum: Circulating Tumour DNA Profiling Reveals Heterogeneity of EGFR Inhibitor Resistance Mechanisms in Lung Cancer Patients." *Nature Communications* 7 (November): 13513. doi:10.1038/ncomms13513.
- Christensen, Emil, Iver Nordentoft, Søren Vang, Karin Birkenkamp-Demtröder, Jørgen Bjerggaard Jensen, Mads Agerbæk, Jakob Skou Pedersen, and Lars Dyrskjød. 2018. "Optimized Targeted Sequencing of Cell-Free Plasma DNA from Bladder Cancer Patients." *Scientific Reports* 8 (1): 1917. doi:10.1038/s41598-018-20282-8.
- Chused, T M, A D Steinberg, and N Talal. 1972. "The Clearance and Localization of Nucleic Acids by New Zealand and Normal Mice." *Clinical and Experimental Immunology* 12 (4): 465–76. <http://www.ncbi.nlm.nih.gov/pubmed/4650369>.
- Costa, F, F Barbisan, C E Assmann, N K F Araújo, A R de Oliveira, J P Signori, F Rogalski, B Bonadiman, M S Fernandes, and I B M da Cruz. 2017. "Seminal Cell-Free DNA Levels Measured by PicoGreen Fluorochrome Are Associated with Sperm Fertility Criteria." *Zygote (Cambridge, England)* 25 (2): 111–19. doi:10.1017/S0967199416000307.
- Dawson, Sarah-Jane, Dana W Y Tsui, Muhammed Murtaza, Heather Biggs, Oscar M Rueda, Suet-Feung Chin, Mark J Dunning, et al. 2013. "Analysis of Circulating Tumor DNA to Monitor Metastatic Breast Cancer." *The New England Journal of Medicine* 368 (13): 1199–1209. doi:10.1056/NEJMoa1213261.
- De Mattos-Arruda, L, B Weigelt, J Cortes, H H Won, C K Y Ng, P Nuciforo, F-C Bidard, et al. 2014. "Capturing Intra-Tumor Genetic Heterogeneity by de Novo Mutation Profiling of Circulating Cell-Free Tumor DNA: A Proof-of-Principle." *Annals of Oncology : Official Journal of the European Society for Medical Oncology* 25 (9): 1729–35. doi:10.1093/annonc/mdu239.

- De Mattos-Arruda, Leticia, Regina Mayor, Charlotte K Y Ng, Britta Weigelt, Francisco Martínez-Ricarte, Davis Torrejon, Mafalda Oliveira, et al. 2015. "Cerebrospinal Fluid-Derived Circulating Tumour DNA Better Represents the Genomic Alterations of Brain Tumours than Plasma." *Nature Communications* 6 (November): 8839. doi:10.1038/ncomms9839.
- Dean, Michael, Joseph Boland, Meredith Yeager, Kate M Im, Lisa Garland, Maria Rodriguez-Herrera, Mylen Perez, et al. 2015. "Addressing Health Disparities in Hispanic Breast Cancer: Accurate and Inexpensive Sequencing of BRCA1 and BRCA2." *GigaScience* 4: 50. doi:10.1186/s13742-015-0088-z.
- Feng, Jiang, Feng Gang, Xiao Li, Tang Jin, Huang Houbao, Cao Yu, and Li Guorong. 2013. "Plasma Cell-Free DNA and Its DNA Integrity as Biomarker to Distinguish Prostate Cancer from Benign Prostatic Hyperplasia in Patients with Increased Serum Prostate-Specific Antigen." *International Urology and Nephrology* 45 (4): 1023–28. doi:10.1007/s11255-013-0491-2.
- Fuchs, Tobias A, Johanna A Kremer Hovinga, Daphne Schatzberg, Denisa D Wagner, and Bernhard Lämmle. 2012. "Circulating DNA and Myeloperoxidase Indicate Disease Activity in Patients with Thrombotic Microangiopathies." *Blood* 120 (6): 1157–64. doi:10.1182/blood-2012-02-412197.
- Gautschi, O, B Huegli, A Ziegler, M Gugger, J Heighway, D Ratschiller, P C Mack, et al. 2007. "Origin and Prognostic Value of Circulating KRAS Mutations in Lung Cancer Patients." *Cancer Letters* 254 (2): 265–73. doi:10.1016/j.canlet.2007.03.008.
- Ghatak, Souvik, Rajendra Bose Muthukumaran, and Senthil Kumar Nachimuthu. 2013. "A Simple Method of Genomic DNA Extraction from Human Samples for PCR-RFLP Analysis." *Journal of Biomolecular Techniques : JBT* 24 (4): 224–31. doi:10.7171/jbt.13-2404-001.
- Gray, Elin S, Helen Rizos, Anna L Reid, Suzanah C Boyd, Michelle R Pereira, Johnny Lo, Varsha

- Tembe, et al. 2015. “Circulating Tumor DNA to Monitor Treatment Response and Detect Acquired Resistance in Patients with Metastatic Melanoma.” *Oncotarget* 6 (39): 42008–18. doi:10.18632/oncotarget.5788.
- Guo, Nannan, Feng Lou, Yongfu Ma, Jie Li, Bo Yang, Wei Chen, Hua Ye, et al. 2016. “Circulating Tumor DNA Detection in Lung Cancer Patients before and after Surgery.” *Scientific Reports* 6: 33519. doi:10.1038/srep33519.
- Han, Hye-Suk, Jieun Yun, Sung-nam Lim, Joung-Ho Han, Ki Hyeong Lee, Seung Taik Kim, Min-Ho Kang, et al. 2013. “Downregulation of Cell-Free MiR-198 as a Diagnostic Biomarker for Lung Adenocarcinoma-Associated Malignant Pleural Effusion.” *International Journal of Cancer* 133 (3): 645–52. doi:10.1002/ijc.28054.
- Harada, Shuko, Rebecca Arend, Qian Dai, Jessica A Levesque, Thomas S Winokur, Rongjun Guo, Martin J Heslin, et al. 2017. “Implementation and Utilization of the Molecular Tumor Board to Guide Precision Medicine.” *Oncotarget* 8 (34): 57845–54. doi:10.18632/oncotarget.18471.
- Helgstrand, John T, Martin A Røder, Nina Klemann, Birgitte G Toft, Daphne Y Lichtensztajn, James D Brooks, Klaus Brasso, Ben Vainer, and Peter Iversen. 2018. “Trends in Incidence and 5-Year Mortality in Men with Newly Diagnosed, Metastatic Prostate Cancer-A Population-Based Analysis of 2 National Cohorts.” *Cancer*, May. doi:10.1002/cncr.31384.
- Hofman, Véronique, Sandra Lassalle, Coraline Bence, Elodie Long-Mira, Sacha Nahon-Estève, Simon Heeke, Virginie Lespinet-Fabre, Catherine Butori, Marius Ilié, and Paul Hofman. 2018. “Any Place for Immunohistochemistry within the Predictive Biomarkers of Treatment in Lung Cancer Patients?” *Cancers* 10 (3). doi:10.3390/cancers10030070.
- Hogervorst, Janneke G F, Roger W L Godschalk, Piet A van den Brandt, Matty P Weijenberg, Bas A J Verhage, Leonie Jonkers, Joy Goessens, et al. 2014. “DNA from Nails for Genetic Analyses in Large-Scale Epidemiologic Studies.” *Cancer Epidemiology, Biomarkers &*

Prevention : A Publication of the American Association for Cancer Research, Cosponsored by the American Society of Preventive Oncology 23 (12): 2703–12. doi:10.1158/1055-9965.EPI-14-0552.

Hovelson, Daniel H, Chia-Jen Liu, Yugang Wang, Qing Kang, James Henderson, Amy Gursky, Scott Brockman, et al. 2017. “Rapid, Ultra Low Coverage Copy Number Profiling of Cell-Free DNA as a Precision Oncology Screening Strategy.” *Oncotarget* 8 (52): 89848–66. doi:10.18632/oncotarget.21163.

Hu, Yueshan, Erik A Ehli, Kelly Nelson, Krista Bohlen, Christophina Lynch, Patty Huizenga, Julie Kittlelsrud, Timothy J Soundy, and Gareth E Davies. 2012. “Genotyping Performance between Saliva and Blood-Derived Genomic DNAs on the DMET Array: A Comparison.” *PloS One* 7 (3): e33968. doi:10.1371/journal.pone.0033968.

Husain, Hatim, David Nykin, Nam Bui, Daniel Quan, German Gomez, Brian Woodward, Sumathi Venkatapathy, et al. 2017. “Cell-Free DNA from Ascites and Pleural Effusions: Molecular Insights into Genomic Aberrations and Disease Biology.” *Molecular Cancer Therapeutics* 16 (5): 948–55. doi:10.1158/1535-7163.MCT-16-0436.

Ilic, Dragan, Mia Djulbegovic, Jae Hung Jung, Eu Chang Hwang, Qi Zhou, Anne Cleves, Thomas Agoritsas, and Philipp Dahm. 2018. “Prostate Cancer Screening with Prostate-Specific Antigen (PSA) Test: A Systematic Review and Meta-Analysis.” *BMJ*, September, k3519. doi:10.1136/bmj.k3519.

Imperiale, Thomas F, David F Ransohoff, Steven H Itzkowitz, Theodore R Levin, Philip Lavin, Graham P Lidgard, David A Ahlquist, and Barry M Berger. 2014. “Multitarget Stool DNA Testing for Colorectal-Cancer Screening.” *The New England Journal of Medicine* 370 (14): 1287–97. doi:10.1056/NEJMoa1311194.

Imperiale, Thomas F, David F Ransohoff, Steven H Itzkowitz, Barry A Turnbull, Michael E Ross,

and Colorectal Cancer Study Group. 2004. “Fecal DNA versus Fecal Occult Blood for Colorectal-Cancer Screening in an Average-Risk Population.” *The New England Journal of Medicine* 351 (26): 2704–14. doi:10.1056/NEJMoa033403.

Jamal-Hanjani, M, G A Wilson, S Horswell, R Mitter, O Sakarya, T Constantin, R Salari, et al. 2016. “Detection of Ubiquitous and Heterogeneous Mutations in Cell-Free DNA from Patients with Early-Stage Non-Small-Cell Lung Cancer.” *Annals of Oncology : Official Journal of the European Society for Medical Oncology* 27 (5): 862–67. doi:10.1093/annonc/mdw037.

Jiang, Peiyong, and Y M Dennis Lo. 2016. “The Long and Short of Circulating Cell-Free DNA and the Ins and Outs of Molecular Diagnostics.” *Trends in Genetics : TIG* 32 (6): 360–71. doi:10.1016/j.tig.2016.03.009.

Kaplan, Mariana J, and Marko Radic. 2012. “Neutrophil Extracellular Traps: Double-Edged Swords of Innate Immunity.” *Journal of Immunology (Baltimore, Md. : 1950)* 189 (6): 2689–95. doi:10.4049/jimmunol.1201719.

Kawahara, Akihiko, Chihiro Fukumitsu, Tomoki Taira, Hideyuki Abe, Yorihiro Takase, Kazuya Murata, Tomohiko Yamaguchi, et al. 2015. “Epidermal Growth Factor Receptor Mutation Status in Cell-Free DNA Supernatant of Bronchial Washings and Brushings.” *Cancer Cytopathology* 123 (10): 620–28. doi:10.1002/cncy.21583.

Kim, Won Tae, Ye Hwan Kim, Pildu Jeong, Sung-Pil Seo, Ho-Won Kang, Yong-June Kim, Seok Joong Yun, et al. 2018. “Urinary Cell-Free Nucleic Acid IQGAP3: A New Non-Invasive Diagnostic Marker for Bladder Cancer.” *Oncotarget* 9 (18): 14354–65. doi:10.18632/oncotarget.24436.

Kimura, H, Y Fujiwara, T Sone, H Kunitoh, T Tamura, K Kasahara, and K Nishio. 2006. “EGFR Mutation Status in Tumour-Derived DNA from Pleural Effusion Fluid Is a Practical Basis for Predicting the Response to Gefitinib.” *British Journal of Cancer* 95 (10): 1390–95.

doi:10.1038/sj.bjc.6603428.

Klassen, Tara L, Eva-Lotta von Rüden, Janice Drabek, Jeffrey L Noebels, and Alica M Goldman.

2012. "Comparative Analytical Utility of DNA Derived from Alternative Human Specimens for Molecular Autopsy and Diagnostics." *The Journal of Molecular Diagnostics : JMD* 14 (5): 451–57. doi:10.1016/j.jmoldx.2012.04.005.

Kopka, Julieta, Monika Leder, Stella M Jaureguiberry, Gottfried Brem, and Gabriel O Boselli.

2011. "New Optimized DNA Extraction Protocol for Fingerprints Deposited on a Special Self-Adhesive Security Seal and Other Latent Samples Used for Human Identification." *Journal of Forensic Sciences* 56 (5): 1235–40. doi:10.1111/j.1556-4029.2011.01853.x.

Lecomte, Thierry, Anne Berger, Franck Zinzindohoué, Stéphanie Micard, Bruno Landi, Hélène

Blons, Philippe Beaune, Paul-Henri Cugnenc, and Pierre Laurent-Puig. 2002. "Detection of Free-Circulating Tumor-Associated DNA in Plasma of Colorectal Cancer Patients and Its Association with Prognosis." *International Journal of Cancer* 100 (5): 542–48. doi:10.1002/ijc.10526.

Li, Y S, B Y Jiang, J J Yang, X C Zhang, Z Zhang, J Y Ye, W Z Zhong, et al. 2018. "Unique

Genetic Profiles from Cerebrospinal Fluid Cell-Free DNA in Leptomeningeal Metastases of EGFR-Mutant Non-Small-Cell Lung Cancer: A New Medium of Liquid Biopsy." *Annals of Oncology : Official Journal of the European Society for Medical Oncology* 29 (4): 945–52. doi:10.1093/annonc/mdy009.

Lu, Jun-Liang, and Zhi-Yong Liang. 2016. "Circulating Free DNA in the Era of Precision

Oncology: Pre- and Post-Analytical Concerns." *Chronic Diseases and Translational Medicine* 2 (4): 223–30. doi:10.1016/j.cdtm.2016.12.001.

MANDEL, P, and P METAIS. 1948. "[Not Available]." *Comptes Rendus Des Seances de La*

Societe de Biologie et de Ses Filiales 142 (3–4): 241–43.

<http://www.ncbi.nlm.nih.gov/pubmed/18875018>.

Marmarelis, Melina, Jeffrey C Thompson, Charu Aggarwal, Tracey L Evans, Erica Carpenter, Roger B Cohen, Corey J Langer, and Joshua Bauml. 2017. "Emerging Uses of Circulating Tumor DNA in Advanced Stage Non-Small Cell Lung Cancer." *Annals of Translational Medicine* 5 (18): 380. doi:10.21037/atm.2017.07.29.

Mehra, Niven, David Dolling, Semini Sumanasuriya, Rossitza Christova, Lorna Pope, Suzanne Carreira, George Seed, et al. 2018. "Plasma Cell-Free DNA Concentration and Outcomes from Taxane Therapy in Metastatic Castration-Resistant Prostate Cancer from Two Phase III Trials (FIRSTANA and PROSELICA)." *European Urology*, February. doi:10.1016/j.eururo.2018.02.013.

Misale, Sandra, Rona Yaeger, Sebastijan Hobor, Elisa Scala, Manickam Janakiraman, David Liska, Emanuele Valtorta, et al. 2012. "Emergence of KRAS Mutations and Acquired Resistance to Anti-EGFR Therapy in Colorectal Cancer." *Nature* 486 (7404): 532–36. doi:10.1038/nature11156.

Mok, Tony, Yi-Long Wu, Jin Soo Lee, Chong-Jen Yu, Virote Sriuranpong, Jennifer Sandoval-Tan, Guia Ladrera, et al. 2015. "Detection and Dynamic Changes of EGFR Mutations from Circulating Tumor DNA as a Predictor of Survival Outcomes in NSCLC Patients Treated with First-Line Intercalated Erlotinib and Chemotherapy." *Clinical Cancer Research : An Official Journal of the American Association for Cancer Research* 21 (14): 3196–3203. doi:10.1158/1078-0432.CCR-14-2594.

Pan, Wenying, Wei Gu, Seema Nagpal, Melanie Hayden Gephart, and Stephen R Quake. 2015. "Brain Tumor Mutations Detected in Cerebral Spinal Fluid." *Clinical Chemistry* 61 (3): 514–22. doi:10.1373/clinchem.2014.235457.

Parkinson, Christine A, Davina Gale, Anna M Piskorz, Heather Biggs, Charlotte Hodgkin, Helen

Addley, Sue Freeman, et al. 2016. "Exploratory Analysis of TP53 Mutations in Circulating Tumour DNA as Biomarkers of Treatment Response for Patients with Relapsed High-Grade Serous Ovarian Carcinoma: A Retrospective Study." *PLoS Medicine* 13 (12): e1002198. doi:10.1371/journal.pmed.1002198.

Pentsova, Elena I, Ronak H Shah, Jiabin Tang, Adrienne Boire, Daoqi You, Samuel Briggs, Antonio Omuro, et al. 2016. "Evaluating Cancer of the Central Nervous System Through Next-Generation Sequencing of Cerebrospinal Fluid." *Journal of Clinical Oncology : Official Journal of the American Society of Clinical Oncology* 34 (20): 2404–15. doi:10.1200/JCO.2016.66.6487.

Ponti, G., M. Maccaferri, M. Mandrioli, M. Manfredini, S. Micali, M. Cotugno, G. Bianchi, et al. 2018. "Seminal Cell-Free DNA Assessment as a Novel Prostate Cancer Biomarker." *Pathology and Oncology Research*. doi:10.1007/s12253-018-0416-6.

Ponti, G., M. Maccaferri, M. Manfredini, M. Cotugno, G. Pellacani, A. Conti, S. Micali, M. Mandrioli, and A. Tomasi. 2018. "Seminal Cell-Free DNA Molecular Profile as a Novel Diagnostic and Prognostic Prostate Cancer Biomarkers." *Medical Hypotheses* 114. doi:10.1016/j.mehy.2018.02.034.

Ponti, G., M. Maccaferri, M. Manfredini, S. Kaleci, M. Mandrioli, G. Pellacani, T. Ozben, et al. 2018. "The Value of Fluorimetry (Qubit) and Spectrophotometry (NanoDrop) in the Quantification of Cell-Free DNA (CfDNA) in Malignant Melanoma and Prostate Cancer Patients." *Clinica Chimica Acta* 479. doi:10.1016/j.cca.2018.01.007.

Ponti, G., M. Manfredini, L. Pastorino, M. Maccaferri, A. Tomasi, and G. Pellacani. 2018. "PTCH1 Germline Mutations and the Basaloid Follicular Hamartoma Values in the Tumor Spectrum of Basal Cell Carcinoma Syndrome (NBCCS)." *Anticancer Research* 38 (1). doi:10.21873/anticancer.12246.

- Ponti, G., M. Manfredini, G. Pellacani, and A. Tomasi. 2016. "Role of Microsatellite Instability, Immunohistochemistry and Mismatch Repair Germline Aberrations in Immunosuppressed Transplant Patients: A Phenocopy Dilemma in Muir-Torre Syndrome." *Clinical Chemistry and Laboratory Medicine* 54 (11). doi:10.1515/cclm-2015-1210.
- Ponti, G, M Maccaferri, M Manfredini, M Cotugno, G Pellacani, A Conti, S Micali, M Mandrioli, and A Tomasi. 2018. "Seminal Cell-Free DNA Molecular Profile as a Novel Diagnostic and Prognostic Prostate Cancer Biomarkers." *Medical Hypotheses* 114 (May): 69. doi:10.1016/j.mehy.2018.02.034.
- Ponti, Giovanni, Monia Maccaferri, Mauro Mandrioli, Marco Manfredini, Salvatore Micali, Michele Cotugno, Giampaolo Bianchi, et al. 2018. "Seminal Cell-Free DNA Assessment as a Novel Prostate Cancer Biomarker." *Pathology Oncology Research : POR*, May. doi:10.1007/s12253-018-0416-6.
- Ponti, Giovanni, Monia Maccaferri, Salvatore Micali, Marco Manfredini, Riccardo Milandri, Giampaolo Bianchi, Giovanni Pellacani, et al. 2018. "Seminal Cell Free DNA Concentration Levels Discriminate Between Prostate Cancer and Benign Prostatic Hyperplasia." *Anticancer Research* 38 (9): 5121–25. doi:10.21873/anticancer.12833.
- Pös, Ondrej, Orsolya Biró, Tomas Szemes, and Bálint Nagy. 2018. "Circulating Cell-Free Nucleic Acids: Characteristics and Applications." *European Journal of Human Genetics : EJHG*, April. doi:10.1038/s41431-018-0132-4.
- Reckamp, Karen L, Vladislava O Melnikova, Chris Karlovich, Lecia V Sequist, D Ross Camidge, Heather Wakelee, Maurice Perol, et al. 2016. "A Highly Sensitive and Quantitative Test Platform for Detection of NSCLC EGFR Mutations in Urine and Plasma." *Journal of Thoracic Oncology : Official Publication of the International Association for the Study of Lung Cancer* 11 (10): 1690–1700. doi:10.1016/j.jtho.2016.05.035.

- Redwood, Diana G, Elvin D Asay, Ian D Blake, Pamela E Sacco, Claudia M Christensen, Frank D Sacco, James J Tiesinga, et al. 2016. "Stool DNA Testing for Screening Detection of Colorectal Neoplasia in Alaska Native People." *Mayo Clinic Proceedings* 91 (1): 61–70. doi:10.1016/j.mayocp.2015.10.008.
- Rolfo, Christian, Philip C Mack, Giorgio V Scagliotti, Paul Baas, Fabrice Barlesi, Trever G Bivona, Roy S Herbst, et al. 2018. "Liquid Biopsy for Advanced Non-Small Cell Lung Cancer (NSCLC): A Statement Paper from the IASLC." *Journal of Thoracic Oncology : Official Publication of the International Association for the Study of Lung Cancer* 13 (9): 1248–68. doi:10.1016/j.jtho.2018.05.030.
- Romanel, Alessandro, Delila Gasi Tandefelt, Vincenza Conteduca, Anuradha Jayaram, Nicola Casiraghi, Daniel Wetterskog, Samanta Salvi, et al. 2015. "Plasma AR and Abiraterone-Resistant Prostate Cancer." *Science Translational Medicine* 7 (312): 312re10. doi:10.1126/scitranslmed.aac9511.
- Salvi, Samanta, Giorgia Gurioli, Filippo Martignano, Flavia Foca, Roberta Gunelli, Giacomo Cicchetti, Ugo De Giorgi, Wainer Zoli, Daniele Calistri, and Valentina Casadio. 2015. "Urine Cell-Free DNA Integrity Analysis for Early Detection of Prostate Cancer Patients." *Disease Markers* 2015: 574120. doi:10.1155/2015/574120.
- Scherer, Florian, David M Kurtz, Aaron M Newman, Henning Stehr, Alexander F M Craig, Mohammad Shahrokh Esfahani, Alexander F Lovejoy, et al. 2016. "Distinct Biological Subtypes and Patterns of Genome Evolution in Lymphoma Revealed by Circulating Tumor DNA." *Science Translational Medicine* 8 (364): 364ra155. doi:10.1126/scitranslmed.aai8545.
- Schwaederle, Maria, Hatim Husain, Paul T Fanta, David E Piccioni, Santosh Kesari, Richard B Schwab, Kimberly C Banks, et al. 2016. "Detection Rate of Actionable Mutations in Diverse Cancers Using a Biopsy-Free (Blood) Circulating Tumor Cell DNA Assay." *Oncotarget* 7 (9):

9707–17. doi:10.18632/oncotarget.7110.

Siravegna, Giulia, Silvia Marsoni, Salvatore Siena, and Alberto Bardelli. 2017. “Integrating Liquid Biopsies into the Management of Cancer.” *Nature Reviews. Clinical Oncology* 14 (9): 531–48. doi:10.1038/nrclinonc.2017.14.

Snyder, Matthew W, Martin Kircher, Andrew J Hill, Riza M Daza, and Jay Shendure. 2016. “Cell-Free DNA Comprises an In Vivo Nucleosome Footprint That Informs Its Tissues-Of-Origin.” *Cell* 164 (1–2): 57–68. doi:10.1016/j.cell.2015.11.050.

Soh, Junichi, Shinichi Toyooka, Keisuke Aoe, Hiroaki Asano, Syuji Ichihara, Hideki Katayama, Akio Hiraki, et al. 2006. “Usefulness of EGFR Mutation Screening in Pleural Fluid to Predict the Clinical Outcome of Gefitinib Treated Patients with Lung Cancer.” *International Journal of Cancer* 119 (10): 2353–58. doi:10.1002/ijc.22190.

Sorensen, Boe S, Lin Wu, Wen Wei, Julie Tsai, Britta Weber, Ebba Nexø, and Peter Meldgaard. 2014. “Monitoring of Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor-Sensitizing and Resistance Mutations in the Plasma DNA of Patients with Advanced Non-Small Cell Lung Cancer during Treatment with Erlotinib.” *Cancer* 120 (24): 3896–3901. doi:10.1002/cncr.28964.

Stewart, Caitlin M, Prachi D Kothari, Florent Mouliere, Richard Mair, Saira Somnay, Ryma Benayed, Ahmet Zehir, et al. 2018. “The Value of Cell-Free DNA for Molecular Pathology.” *The Journal of Pathology* 244 (5): 616–27. doi:10.1002/path.5048.

Stewart, Caitlin M, and Dana W Y Tsui. 2018. “Circulating Cell-Free DNA for Non-Invasive Cancer Management.” *Cancer Genetics*, March. doi:10.1016/j.cancergen.2018.02.005.

Stroun, M, P Anker, P Maurice, J Lyautey, C Lederrey, and M Beljanski. 1989. “Neoplastic Characteristics of the DNA Found in the Plasma of Cancer Patients.” *Oncology* 46 (5): 318–

22. doi:10.1159/000226740.

Stroun, M, J Lyautey, C Lederrey, A Olson-Sand, and P Anker. 2001. "About the Possible Origin and Mechanism of Circulating DNA Apoptosis and Active DNA Release." *Clinica Chimica Acta; International Journal of Clinical Chemistry* 313 (1–2): 139–42.
<http://www.ncbi.nlm.nih.gov/pubmed/11694251>.

Tabernero, Josep, Heinz-Josef Lenz, Salvatore Siena, Alberto Sobrero, Alfredo Falcone, Marc Ychou, Yves Humblet, et al. 2015. "Analysis of Circulating DNA and Protein Biomarkers to Predict the Clinical Activity of Regorafenib and Assess Prognosis in Patients with Metastatic Colorectal Cancer: A Retrospective, Exploratory Analysis of the CORRECT Trial." *The Lancet. Oncology* 16 (8): 937–48. doi:10.1016/S1470-2045(15)00138-2.

Thierry, A R, S El Messaoudi, P B Gahan, P Anker, and M Stroun. 2016. "Origins, Structures, and Functions of Circulating DNA in Oncology." *Cancer Metastasis Reviews* 35 (3): 347–76.
doi:10.1007/s10555-016-9629-x.

Todenhöfer, Tilman, Werner J Struss, Roland Seiler, Alexander William Wyatt, and Peter C Black. 2018. "Liquid Biopsy-Analysis of Circulating Tumor DNA (CtDNA) in Bladder Cancer." *Bladder Cancer (Amsterdam, Netherlands)* 4 (1): 19–29. doi:10.3233/BLC-170140.

Ulrich, Bryan C, and Cloud P Paweletz. 2018. "Cell-Free DNA in Oncology: Gearing up for Clinic." *Annals of Laboratory Medicine* 38 (1): 1–8. doi:10.3343/alm.2018.38.1.1.

Wan, Jonathan C M, Charles Massie, Javier Garcia-Corbacho, Florent Mouliere, James D Brenton, Carlos Caldas, Simon Pacey, Richard Baird, and Nitzan Rosenfeld. 2017. "Liquid Biopsies Come of Age: Towards Implementation of Circulating Tumour DNA." *Nature Reviews. Cancer* 17 (4): 223–38. doi:10.1038/nrc.2017.7.

Wang, Tingting, Mingming Lv, Sunan Shen, Sheng Zhou, Ping Wang, Yueqiu Chen, Baorui Liu,

Like Yu, and Yayi Hou. 2012. “Cell-Free MicroRNA Expression Profiles in Malignant Effusion Associated with Patient Survival in Non-Small Cell Lung Cancer.” *PloS One* 7 (8): e43268. doi:10.1371/journal.pone.0043268.

Wang, Yuxuan, Simeon Springer, Carolyn L Mulvey, Natalie Silliman, Joy Schaefer, Mark Sausen, Nathan James, et al. 2015. “Detection of Somatic Mutations and HPV in the Saliva and Plasma of Patients with Head and Neck Squamous Cell Carcinomas.” *Science Translational Medicine* 7 (293): 293ra104. doi:10.1126/scitranslmed.aaa8507.

Wang, Yuxuan, Simeon Springer, Ming Zhang, K Wyatt McMahon, Isaac Kinde, Lisa Dobbyn, Janine Ptak, et al. 2015. “Detection of Tumor-Derived DNA in Cerebrospinal Fluid of Patients with Primary Tumors of the Brain and Spinal Cord.” *Proceedings of the National Academy of Sciences of the United States of America* 112 (31): 9704–9. doi:10.1073/pnas.1511694112.

Wyatt, Alexander W, Matti Annala, Rahul Aggarwal, Kevin Beja, Felix Feng, Jack Youngren, Adam Foye, et al. 2017. “Concordance of Circulating Tumor DNA and Matched Metastatic Tissue Biopsy in Prostate Cancer.” *Journal of the National Cancer Institute* 109 (12). doi:10.1093/jnci/djx118.

Yang, Haihong, Linbo Cai, Yalei Zhang, Hongyu Tan, Qiuhua Deng, Meiling Zhao, and Xin Xu. 2014. “Sensitive Detection of EGFR Mutations in Cerebrospinal Fluid from Lung Adenocarcinoma Patients with Brain Metastases.” *The Journal of Molecular Diagnostics : JMD* 16 (5): 558–63. doi:10.1016/j.jmoldx.2014.04.008.

Zamir, Ashira, Carla Oz, Ehud Wolf, Asya Vinokurov, and Baruch Glattstein. 2004. “A Possible Source of Reference DNA from Archived Treated Adhesive Lifters.” *Journal of Forensic Sciences* 49 (1): 68–70. <http://www.ncbi.nlm.nih.gov/pubmed/14979346>.

Zeng, Hu, Bo He, Chengqi Yi, and Jinying Peng. 2018. “Liquid Biopsies: DNA Methylation Analyses in Circulating Cell-Free DNA.” *Journal of Genetics and Genomics = Yi Chuan Xue*

Bao, March. doi:10.1016/j.jgg.2018.02.007.

ACCEPTED MANUSCRIPT