

This is the peer reviewed version of the following article:

Effects of soiling and weathering on the albedo of building envelope materials: Lessons learned from natural exposure in two European cities and tuning of a laboratory simulation practice / Paolini, R.; Terraneo, G.; Ferrari, C.; Sleiman, M.; Muscio, A.; Metrangolo, P.; Poli, T.; Destailats, H.; Zinzi, M.; Levinson, R.. - In: SOLAR ENERGY MATERIALS AND SOLAR CELLS. - ISSN 0927-0248. - 205:(2020), pp. 1-12. [10.1016/j.solmat.2019.110264]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

10/01/2025 18:32

The 13th European Nutrition Conference, FENS 2019, was held at the Dublin Convention Centre, 15–18 October 2019

Insights into the association of potassium intake with blood pressure: results of a dose-response meta-analysis of randomized controlled trials

Tommaso Filippini¹, Duarte Torres^{2,3}, Carla Lopes^{2,4}, Catarina Carvalho^{2,3}, Pedro Moreira^{2,3}, Androniki Naska⁵, Maria-Iosifina Kasdagli⁵, Marcella Malavolti¹, Nicola Orsini⁶ and Marco Vinceti^{1,7}

¹Environmental, Genetic and Nutritional Epidemiology Research Center (CREAGEN), Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy,

²EPIUnit - Institute of Public Health, University of Porto, Porto, Portugal,

³Faculty of Food and Nutrition Sciences, University of Porto, Porto, Portugal,

⁴Department of Public Health and Forensic Sciences, and Medical Education, Unit of Epidemiology, Faculty of Medicine, University of Porto, Porto, Portugal,

⁵Department of Hygiene, Epidemiology and Medical Statistics, School of Medicine, National and Kapodistrian University of Athens, Athens, Greece,

⁶Department of Public Health Sciences, Karolinska Institute, Stockholm, Sweden and

⁷Department of Epidemiology, Boston University School of Public Health, Boston, USA

Abstract

Introduction: Observational studies provide evidence for an association between potassium intake and BP levels. However, uncertainties still exist about the size and the shape of this relation. Conversely, experimental studies have not been used to estimate dose-response curves, since standard methods can only be applied in trials including at least three exposure groups.

Materials and Methods: We carried out a systematic review of the evidence concerning the effect of potassium supplementation on blood pressure in epidemiologic experimental studies. Following a PubMed search up to June 20, 2019, we included randomized controlled trials (RCTs) encompassing potassium supplementation as the only intervention for at least four weeks. We used a restricted cubic spline model and the 'one-stage' approach to perform a dose-response meta-analysis, a newly-developed statistical procedure which allows inclusion of studies with as few as two categories of exposure (Stat Methods Med Res. 2019;28:1579–1596). Finally, we repeated the analyses stratifying for hypertensive status and use of anti-hypertensive medication.

Results: Overall, we included 33 studies carried out in adult population, with potassium supplementation ranging from 30 to 140 mmol/day. RCTs' duration ranged from 4 up to 26 weeks. Most of the studies have a cross-over design (N = 24), include hypertensive individuals (N = 27) and subjects not under anti-hypertensive medication (N = 27). Overall, an increase of 40, 80 and 120 mmol/day of potassium resulted in reductions of SBP by -5.64 (95% CI - 8.78, -2.50), -4.62 (-6.41, -2.84) and -2.54 mmHg (95% CI -5.14, + 0.06), respectively. Higher potassium intakes also resulted in reduced DBP levels by -3.57 (95% CI -5.55, -1.59), -3.07 (95% CI -5.07, -1.08), and -1.92 mmHg (95% CI -5.65, 1.81). The effect of increasing potassium intake on BP was larger among hypertensives than normotensives, and among pharmacologically untreated hypertensives compared to their treated counterparts. Subgroup analyses according to study design (parallel vs. crossover) yielded similar results.

Discussion: With the application of advanced dose-response modeling on RCT results, we found a U-shaped relation between potassium intake and blood pressure. A low to moderate increase in potassium intake resulted in a progressive reduction in blood pressure, which was reversed at higher levels of potassium supplementation. The effect was stronger among untreated hypertensives.

Supported by grant GP-EFSA-AFSCO-2017-01 GA09 of the European Food Safety Authority - EFSA. The text reflects only the authors' view; and EFSA is not responsible for any use that may be made of the information it contains.

Conflict of Interest

There is no conflict of interest