

This is a pre print version of the following article:

Obesity risk during collective quarantine for the COVID-19 epidemic / Mattioli, Anna Vittoria; Pinti, Marcello; Farinetti, Alberto; Nasi, Milena. - In: OBESITY MEDICINE. - ISSN 2451-8476. - 20:(2020), pp. 100263-100264. [10.1016/j.obmed.2020.100263]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

10/01/2026 21:35

Journal Pre-proof

Obesity risk during collective quarantine for the COVID-19 epidemic

Anna Vittoria Mattioli, Marcello Pinti, Alberto Farinetti, Milena Nasi

PII: S2451-8476(20)30083-X

DOI: <https://doi.org/10.1016/j.obmed.2020.100263>

Reference: OBMED 100263

To appear in: *Obesity Medicine*

Received Date: 3 June 2020


Revised Date: 5 June 2020

Accepted Date: 5 June 2020

Please cite this article as: Mattioli, A.V., Pinti, M., Farinetti, A., Nasi, M., Obesity risk during collective quarantine for the COVID-19 epidemic, *Obesity Medicine*, <https://doi.org/10.1016/j.obmed.2020.100263>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd.



Obesity risk during collective quarantine for the COVID-19 epidemic.

Anna Vittoria Mattioli^a, Marcello Pinti ^b, Alberto Farinetti ^c, Milena Nasi^a

Affiliation:

- a) Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Italy
- b) Department of life science, University of Modena and Reggio Emilia, Italy
- c) Department of Medical and Surgical Sciences for Children and Adults, University of Modena and Reggio Emilia, Modena, Italy.

Running title: quarantine and obesity

Address for correspondence:

Prof Anna Vittoria Mattioli,
Surgical, Medical and Dental Department of Morphological Sciences related to Transplant,
Oncology and Regenerative Medicine
University of Modena and Reggio Emilia,
Via del pozzo, 71 41124 Modena (Italy)
Ph +390594222841 Fax +390594222841
E-mail: annavittoria.mattioli@unimore.it

Author contribution. AVM, MN, MP,AF conceived of the idea at the basis of the article, AVM, MN developed the different part of the manuscript, AVM, MN, MP,AF performed the final supervision. All authors contributed to and approved the final manuscript.

Mail address, e mail and Orcid of authors:

Anna Vittoria Mattioli:

Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Via del pozzo, 71 41124 Modena (Italy)

E mail: annavittoria.mattioli@unimore.it

Orcid: 0000-0003-1487-9530

Marcello Pinti:

Department of Life science, University of Modena and Reggio Emilia, Via Campi, 287 41124 Modena (Italy)

E mail: marcello.pinti@unimore.it

Orcid: 0000-0001-9118-1262

Alberto Farinetti

Department of Medical and Surgical Sciences for Children and Adults, University of Modena and Reggio Emilia, Via del pozzo, 71 41124 Modena (Italy)

E mail: alberto.farinetti@unimore.it

Orcid: 0000-0003-2133-3595

Milena Nasi

Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Via del pozzo, 71 41124 Modena (Italy)

E mail: milena.nasi@unimore.it

Orcid: 0000-0003-3079-8001

Word count**Main text:** 996 words (max 1000)**Tables:** 1**References:** max 20**Abstract**

In March 2020, when COVID-19 epidemics involved several countries, the WHO defined the infection as a pandemic. Government adopted measures to prevent the diffusion of infection; i.e. quarantine and isolation. One of the consequences of quarantine-induced stress is a change in lifestyle and eating habits leading to obesity. The present commentary briefly analyzes the effects of quarantine on obesity.

Key-words: quarantine; COVID-19; obesity; stress; lifestyle; gender

In March 2020, when COVID-19 epidemics involved several countries, the WHO defined the infection as a pandemic. (Li 2020) Government adopted measures to prevent the diffusion of infection; i.e. quarantine and isolation. (WHO, 2020a)

“Quarantine” refers to the separation of persons (or communities) who have been exposed to an infectious disease. “Isolation” refers to the separation of persons who are infected. (Mattioli and Ballerini Puviani, 2020a; WHO, 2020a)

The present commentary briefly analyzes the effects of quarantine on obesity.

Little is known about the effects of quarantine on obesity, although everyone agrees that this corresponds to a period of severe stress and stress can be linked to an increased risk of obesity. (Mattioli and Ballerini Puviani, 2020a; Abbas et al, 2020) One of the consequences of quarantine stress is a change in lifestyle and eating habits. (Mattioli et al, 2020b) Laitinen and coworkers (2002) reported that stress-related eating (defined as trying to make oneself feel better by eating or drinking in a stressful situation) was significantly associated with obesity, mainly in women. There may be a gender-specific response to stress in which women are more likely to use food to deal with stress, whereas men are more likely to use other oral behaviors such as alcohol consumption or smoking as strategies to cope with stress. (Conway et al, 1981)

Modern women have multiple roles that are both time- and energy-consuming. These lead to an increase in psychosocial stressors (e.g. anxiety, depression and marital stress), which are known to increase the overall cardiovascular risk. Furthermore, women rarely follow a healthy lifestyle based on an adequate diet, regular physical activity and weight management because they are both costly and time consuming. Adherence to healthy lifestyle is strongly influenced by socio-economic level, social role, and education (Sciomer et al, 2019; Mattioli et al, 2017)

During quarantine, due to the anxiety caused by the hypothesis of a future food shortage, people buy large quantities of packaged and long-lasting food rather than fresh food.

(Brooks et al, 2020) Foods with a long shelf life have a high content of salt, sugar or trans fats which increases the risk of obesity. In addition some people developed the desire to consume a specific kind of food, a behavior defined as “food craving”. Food craving is a *“multidimensional experience as it includes cognitive (e.g., thinking about food), emotional (e.g., desire to eat or changes in mood), behavioral (e.g., seeking and consuming food), and physiological (e.g., salivation) aspect”* (Rodríguez-Martín and Meule, 2015) The desire for carbohydrates increases the production of serotonin which has a positive effect on mood, and this effect is proportional to the glycemic index of food. (Rodríguez-Martín and Meule, 2015; Mattioli et al 2020c) During the quarantine, patients suffering from obesity experienced immense stress that made them more vulnerable to a sedentary lifestyle and unhealthy nutrition, thus predisposing them to further weight gain. Furthermore, the next economic downturn will also lead to greater consumption of unhealthy foods as it is cheaper. This will lead to a further increase in the prevalence of obesity, especially in the weaker sections of society. (Mattioli et al, 2020b; Mediouni, 2020; Torres and Nowson, 2007) Growing evidence has reported an important role of obesity in Covid-19's prognosis. (Torres and Nowson, 2007; Abbas et al, 2020)

What are the consequences of quarantine-related stress on the immune system and on health? Dantzer et al. (2018) summarized the relationship between resilience and immunity, where resilience is the Janus face of vulnerability factors in response to stressing conditions. While, short-term stress enhances immune protection, the humoral and cellular immune responses are suppressed by long-term stress.

The reduction of the immune function is associated with a reduce protection against infection and with a exacerbation of pathological immune responses. (Dantzer et al, 2018) Thus, aged people or people with other chronic pathologies such as cancer, cardiovascular, neurodegenerative, or metabolic disease, are more susceptible to immune alterations induced by outbreak, with relevant clinical side effects. It is well known that obesity is

associated to a chronic, sterile, low-grade inflammation which contributes to the activation of atherosclerotic plaques, making patients affected by obesity more vulnerable to cardiovascular events. (Abbas et al, 2020; Nasi et al, 2019) (figure 1) Inflammatory markers were seen to increase by short-term as well as long-term stress. In both cases, an increase of cortisol and glucocorticoid hormones have been observed, however, acute stress is associated to a decrease of proinflammatory cytokines whereas chronic stress is associated to an increase of these biomarkers. (Cossarizza et al, 2020; Dantzer et al, 2018; Marsland et al, 2017)

The innate and adaptive immune response of the host against SARS-Cov-2 is crucial to control the infection. The uncontrolled inflammatory response causes tissue damages at local and systemic level, and it is widely accepted that the so-called “cytokine storm” is associated with the severity and the outcome of the COVID-19. (Cossarizza et al 2020)

During quarantine, due to Governments limitation of outdoor exercise and social activities subjects reduced the physical activity. Regular physical activity reduces inflammation, and contribute to maintain normal weight and to reduce visceral fat accumulation. (Mattioli et al, 2020d, Nasi et al, 2019) Limited physical activity can be associated with several metabolic effects that would increase the cardiovascular risk. (Rahmati-Ahmadabad and Hosseini, 2020)

To prevent damages induced by physical inactivity, WHO (WHO 2020b) has published a guide for self-quarantine people without any symptoms or diagnosis of acute respiratory illness, containing practical advice on how to stay active and reduce sedentary behavior at home.

Staying active and maintaining a home-physical exercise routine will be important for mental and physical health. WHO suggest following on-line exercise classes, and using video- or app-guided aerobics training at home. (WHO 2020b)

The Internet and apps could also be helpful in weight management by monitoring diet and daily income.

Conclusions

We must be prepared for an increase in obesity following the pandemic, and this increase in obesity will be associated with an increase of the cardiovascular risk burden. It is mandatory to promote a healthy diet and physical activity at home. After the quarantine we must re-evaluate the metabolic and biometric parameters in obese subjects and we must also stratify the cardiovascular risk

Acknowledgements

We would like to thank Mrs Janet Ann Carter for reviewing the paper.

We thank everyone who is working to resolve the SARS - CoV - 2 pandemic.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest.

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

References

- Abbas, A.M.; Fathy, S.K.; Fawzy, A.T.; Salem, A.S.; Shawky, M.S. 2020. The mutual effects of COVID-19 and obesity. *Obes Med*
<https://doi.org/10.1016/j.obmed.2020.100250>
- Brooks, S.K., Webster, R.K., Smith, L.E., Woodland, L., Wessely, S., Greenberg, N., Rubin G.J. 2020 The psychological impact of quarantine and how to reduce it: rapid review of the evidence *Lancet* Mar 14;395(10227):912-920. doi: 10.1016/S0140-6736(20)30460-8. Epub 2020 Feb 26.
- Conway, T.L., Vickers, R.R., Ward, H.W., Rahe, R.H. 1981 Occupational stress and variation in cigarette, coffee, alcohol consumption. *J Health Soc Behav* 1981;22:155– 65.
- Cossarizza, A., De Biasi, S., Guaraldi, G., Girardis, M., Mussini, C., Modena Covid-19 Working Group. 2020 SARS-CoV-2, the Virus that Causes COVID-19: Cytometry and the New Challenge for Global Health. *Cytometry*; 97(4):340-343. doi: 10.1002/cyto.a.2400
- Dantzer, R., Cohen, S., Russo, S.J., Dinan, T.G. 2018 Resilience and immunity. *Brain Behav Immun.* 74:28-42. doi:10.1016/j.bbi.2018.08.010
- Laitinen, J., Ek, E., Sovio, U. 2002. Stress-related eating and drinking behavior and body mass index and predictors of this behavior. *Prev Med.* 34:29 –39.
- Marsland, A.L., C. Walsh, K. Lockwood, N.A. John-Henderson. 2017 The effects of acute psychological stress on circulating and stimulated inflammatory markers: A systematic review and meta-analysis. *Brain Behav Immun.* 64(208-219. doi: 10.1016/j.bbi.2017.01.011
- Mattioli, A.V., Coppi, F., Migaldi, M., Scicchitano, P., Ciccone, M.M., Farinetti, A. 2017 Relationship between Mediterranean diet and asymptomatic peripheral arterial disease in a population of pre-menopausal women. *Nutr Metab Cardiovasc Dis.* 27 (11) 985-990. doi: 10.1016/j.numecd.2017.09.011

Mattioli, A.V., Ballerini Puviani, M. 2020a. Lifestyle at time of COVID-19, how could quarantine affect cardiovascular risk.. Am J Lifestyle Med. Apr 17;14(3):240-242. doi: 10.1177/1559827620918808

Mattioli, A.V., Sciomer, S., Cocchi, C., Maffei, S., Gallina, S. 2020b. "Quarantine during COVID-19 outbreak: changes in Diet and physical activity increase the risk of cardiovascular disease" Nutr Metab Cardiovasc Dis doi.org/10.1016/j.numecd.2020.05.020

Mattioli, A.V., Nasi, M., Cocchi, C., Farinetti, A. 2020c. COVID 19 outbreak: impact of the quarantine-induced stress on cardiovascular disease risk burden. Future cardiology Apr 30;10.2217/fca-2020-0055. doi: 10.2217/fca-2020-0055. Online ahead of print.

Mattioli, A.V., Ballerini Puviani, M., Nasi, M., Farinetti, A. 2020d. COVID-19 pandemic: the effects of quarantine on cardiovascular risk. Eur J Clin Nutr. May 5;1-4. doi: 10.1038/s41430-020-0646-z.

Mediouni, M., Kaczor-Urbanowicz, K.E., Madiouni, R. 2020 Future Epidemic: DEPREOBESITY. Obesity Medicine. September;19:100240

Nasi, M., Patrizi, G., Pizzi, C., Landolfo, M., Boriani, G., Dei Cas, A. et al. 2019. The role of physical activity in individuals with cardiovascular risk factors: An opinion paper from Italian Society of Cardiology-Emilia Romagna-Marche and SIC-Sport. J Cardiovasc Med. 20:631-639. doi: 10.2459/JCM.0000000000000855

Rahmati-Ahmadabad, S., Hosseini, F., 2020. Exercise against SARS-CoV-2 (COVID-19): does workout intensity matter? (A mini review of some indirect evidence related to obesity). Obes Med <https://doi.org/10.1016/j.obmed.2020.100245>

Rodríguez-Martín, B.C., Meule, A. 2015. Food craving: new contributions on its assessment, moderators, and consequences. Front Psychol.6:21. Published 2015 Jan 22. doi:10.3389/fpsyg.2015.00021

Sciomer, S., Moscucci, F., Maffei, S., Gallina, S., Mattioli A.V. 2019. Cardiovascular risk factors prevention in women: the life style paradox and stereotypes to defeat. Eur J Prev Cardiol. 26(6):609-610. doi: 10.1177/2047487318810560

Torres, S.J., Nowson, C.A. 2007. Relationship between stress, eating behavior, and obesity. Nutrition. 23(11- 12):887-894. <https://doi.org/10.1016/j.nut.2007.08.008>

WHO 2020a. WHO announces COVID-19 outbreak a pandemic.

<http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>


WHO 2020b. WHO guideline “Stay physically active during self-quarantine”

<http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-coronavirus-2019-ncov-technical-guidance/stay-physically-active-during-self-quarantine>

Figure 1

Stress alter glucose metabolism, promote insulin resistance and influence multiple appetite-related hormones and hypothalamic neuropeptides

Journal Pre-proof



Highlights

- Quarantine and isolation are effective measures to reduce diffusion of infection and to prevent pandemic. However these conditions can induce depression, anxiety, anger, and stress.
- Quarantine forces people to stay at home with a subsequent reduction of outside exercise and physical activity
- Anxiety and stress are associated with unhealthy lifestyle involving both nutrition and physical activity leading to obesity

Please wait...

If this message is not eventually replaced by the proper contents of the document, your PDF viewer may not be able to display this type of document.

You can upgrade to the latest version of Adobe Reader for Windows®, Mac, or Linux® by visiting http://www.adobe.com/go/reader_download.

For more assistance with Adobe Reader visit <http://www.adobe.com/go/acrreader>.

Windows is either a registered trademark or a trademark of Microsoft Corporation in the United States and/or other countries. Mac is a trademark of Apple Inc., registered in the United States and other countries. Linux is the registered trademark of Linus Torvalds in the U.S. and other countries.