



# Extreme-tolerance mechanisms in meiofaunal organisms: a case study with tardigrades, rotifers and nematodes

Lorena Rebecchi · Chiara Boschetti · Diane R. Nelson

Received: 1 August 2019 / Revised: 20 November 2019 / Accepted: 27 November 2019 / Published online: 16 December 2019  
© Springer Nature Switzerland AG 2019

**Abstract** To persist in extreme environments, some meiofaunal taxa have adopted outstanding resistance strategies. Recent years have seen increased enthusiasm for understanding extreme-resistance mechanisms evolved by tardigrades, nematodes and rotifers, such as the capability to tolerate complete desiccation and freezing by entering a state of reversible suspension of metabolism called anhydrobiosis and cryobiosis, respectively. In contrast, the less common phenomenon of diapause, which includes encystment and cyclomorphosis, is defined by a suspension of growth and development with a reduction in metabolic activity induced by stressful

environmental conditions. Because of their unique resistance, tardigrades and rotifers have been proposed as model organisms in the fields of exobiology and space research. They are also increasingly considered in medical research with the hope that their resistance mechanisms could be used to improve the tolerance of human cells to extreme stress. This review will analyse the dormancy strategies in tardigrades, rotifers and nematodes with emphasis on mechanisms of extreme stress tolerance to identify convergent and unique strategies occurring in these distinct groups. We also examine the ecological and evolutionary consequences of extreme tolerance by summarizing recent advances in this field.

---

Guest editors: Nabil Majdi, Jenny M. Schmid-Araya & Walter Traunspurger / Patterns and Processes of Meiofauna in Freshwater Ecosystems

---

L. Rebecchi  
Department of Life Sciences, Modena and Reggio Emilia University, Modena, Italy

L. Rebecchi (✉)  
Department of Life Sciences, University of Modena and Reggio Emilia, Via G. Campi 213/D, 41125 Modena, Italy  
e-mail: lorena.rebecchi@unimore.it

C. Boschetti  
School of Biological and Marine Sciences, University of Plymouth, Plymouth, UK

D. R. Nelson  
Department of Biological Sciences, East Tennessee State University, Johnson City, TN, USA

**Keywords** Anhydrobiosis · Cryptobiosis · Desiccation · Diapause · Dormancy · Encystment

## Introduction

Tardigrades, rotifers and nematodes are considered permanent and essential members of freshwater and terrestrial meiofaunal communities that can undergo dormancy during their life stages (Rundle et al., 2002; Hengherr & Schill, 2018; Schill & Hengherr, 2018; Bertolani et al., 2019).

Tardigrades, commonly called “water bears”, are micrometazoans categorized into two main classes

(Eutardigrada and Heterotardigrada) with 1298 species described from marine, freshwater and terrestrial habitats (Degma et al., 2019). The highest number of species belongs to the class eutardigrades and to the family Echiniscidae within the heterotardigrades and has been described from terrestrial habitats, where they are inactive unless surrounded by a film of water. The smallest numbers are true limnic species, but several species are limnoterrestrial and can colonize both terrestrial and freshwater habitats (Nelson et al., 2015). Rotifera, also called “wheel animals”, is a phylum of microscopic metazoans, comprising about 2000 species (Segers, 2007) traditionally divided in three main classes: (1) Bdelloidea live in freshwater and terrestrial ephemeral aquatic environments and only reproduce by apomictic parthenogenesis; (2) Monogononta live in freshwater and marine environments and reproduce by cyclical parthenogenesis and (3) Seisonida, with only a few exclusively marine species (Ricci, 1987; Wallace & Snell, 1991; Melone et al., 1998; Mark Welch & Meselson, 2000; Ricci & Melone, 2000; Segers, 2007). A fourth class, the exclusively parasitic Acanthocephala, has recently been added, although its exact relationship with the other taxa is still debated (e.g. Sørensen et al., 2005; Sielaff et al., 2016). The majority of nematodes, also called “roundworms”, are small free-living animals inhabiting the thin layer of water surrounding soil particles and in aquatic sediments, although some taxa have become endoparasitic and can reach metres in length (Lee, 2002). Some taxa have evolved the ability to resist desiccation during various stages of their life cycles (Womersley, 1987; Ricci & Pagani, 1997; Shannon et al., 2005; Erkut et al., 2011).

Tardigrades, rotifers and nematodes are meiofaunal aquatic animals common in lakes, rivers, streams and ponds, but paradoxically they are able to colonize and persist in desiccation-prone environments, such as freshwater (e.g. temporary ponds, Antarctic lakes, cryoconite holes) and terrestrial (e.g. mosses and lichens) habitats where liquid water is not always available (Rundle et al., 2002; Nelson et al., 2015, 2018). In these habitats, water loss can occur via evaporation or freezing, with diel, seasonal, annual or longer fluctuations in the duration of the wet phase. Since tardigrades, rotifers and nematodes are incapable of active migration to more suitable habitats, occupancy of these unpredictable habitats requires organisms to be versatile, tolerant or to possess

specific and exceptional resistance and adaptive strategies (Fontaneto, 2019). Accordingly, life in these environments is adapted to a dual existence, flourishing when the habitat contains liquid water, and dormant when liquid water is not available and dormant states are linked to a temporary suspension of active life with reduction or interruption of metabolism and/or arrested development. Dormancy includes any form of resting stage, regardless of the cues required for induction or termination (Hand, 1991; Cáceres, 1997). Tardigrades, rotifers and nematodes exhibit both forms of dormancy: quiescence (cryptobiosis) and diapause (encystment, cyclomorphosis and resting eggs) (e.g. Crowe & Madin, 1975; Ricci, 1987; Guidetti et al., 2011a).

Among the various forms of dormancy, cryptobiosis (“hidden life”, Keilin, 1959) is under exogenous control, being directly induced and maintained by adverse environmental conditions, and it is immediately reversed by the removal of the external stimuli. It originated independently several times in the history of life, as it is present in diverse groups of bacteria, metazoans, fungi and plants (Clegg, 2001). Cryptobiosis includes different strategies such as anhydrobiosis, cryobiosis, anoxybiosis and osmobiosis directly induced by desiccation, sub-zero temperatures, low oxygen pressure and osmotic extremes, respectively (Keilin, 1959; Wright et al., 1992). Cryptobiosis allows tardigrades, rotifers and nematodes to survive periods of desiccation, whereas few freshwater and marine species are known to have this adaptive strategy (Ricci & Pagani, 1997; Ricci, 1998; Eyres et al., 2015; Guidetti et al., 2011a, b; Clausen et al., 2014; Møbjerg et al., 2011). Conversely, encystment and the production of resting eggs are a state of diapause controlled by both exogenous and endogenous stimuli and is more common in freshwater and marine species. Although tardigrades, rotifers and nematodes exhibit both forms of dormancy, there are differences among taxa. Tardigrades, as well as insects, can undergo both diapause (encystment and cyclomorphosis) (Guidetti & Møbjerg, 2019) and the production of resting eggs (Hansen & Katholm, 2002; Altiero et al., 2010). In comparison, in rotifers the two main types of dormancy are restricted to two separate taxa. The class Bdelloidea can resist adverse environmental conditions via quiescence and directly respond to environmental stimuli at any life stage, from eggs to adults, although with age-dependent degrees of

resistance (Ricci, 1987, 1998; Örstan, 1995, 1998), while the other main class, the Monogononta, only engage in diapause via the production of resting eggs, which tend to stop at a specific and common developmental stage and are generally very resistant to various environmental stresses, including desiccation (e.g. Balompapueng et al., 1997; Cáceres, 1997; Schröder, 2005; García-Roger et al., 2006, Boschetti et al., 2010, Ziv et al., 2017). Within nematodes, dormancy is more scattered across taxa. For example some genera or species can survive desiccation (e.g. Wharton, 1996; Tyson et al., 2012), while other species only have limited resistance at specific life stages (e.g. Erkut et al., 2011; Erkut & Kurzchalia, 2015).

This review analyses the dormancy strategies in tardigrades, rotifers and nematodes with emphasis on mechanisms of stress tolerance in order to identify convergent strategies occurring in these animal taxa. The review also considers the ecological and evolutionary consequences of extreme tolerance by summarizing recent advances in this field.


### Diapause: encystment

In terrestrial and freshwater tardigrades, encystment is an adaptive strategy that involves profound morphological changes that occur during the molting process, resulting in the dormant organism lying within retained cuticular exuvia. During this state, the organism also presents a very low or undetectable metabolism, even if the cyst is not desiccated (Ziv et al., 2017), highlighting possible physiological similarities between diapause and quiescence. Although encystment is rare in moss-dwelling tardigrades, it has been confirmed in grassland and leaf litter habitats but is more common in freshwater sediments (Guidetti et al., 2006). Encystment has been verified in limnic eutardigrades and a few heterotardigrade and eutardigrade limnoterrestrial species, however the phenomenon may be widespread but relatively unstudied (Bertolani et al., 2019; Guidetti & Møbjerg, 2019). In addition, the marine intertidal heterotardigrade *Echiniscoides sigismundi* Plate, 1888, a cryptic species complex, produces two or three new cuticles during cyst formation (Clausen et al., 2014).

In response to the gradual onset of adverse environmental conditions (e.g. temperature, oxygen tension, pH), encystment in tardigrades begins with the ejection of the sclerified parts of the buccal-pharyngeal apparatus (“simplex stage”). Instead of undergoing normal ecdysis, however, one to three new cuticles are serially produced in addition to the retained external (old) cuticle (Fig. 1). The animal’s size is reduced by longitudinal contraction, body movements cease completely, metabolism is significantly reduced and the mouth and cloaca are closed. Modified claws and buccal-pharyngeal apparatus are synthesized, but non-functional. At this stage, the cyst resembles an onion or a Russian doll (“Matryoshka”) (Guidetti et al., 2006), often with one cuticle becoming hardened and pigmented. Encystment ends as environmental conditions improve, and the tardigrade gradually resynthesizes a normal cuticle, claws and feeding apparatus and leaves the cyst. Unknown endogenous stimuli may also play a role in the process.

### Ecology of cysts

Limnic eutardigrades that frequently encyst belong to the genera *Dactylobiotus*, *Pseudobiotus*, *Isohypsibius*, *Hypsibius*, *Thulinus* and *Bertolanius* (See Table 9.1 in Guidetti & Møbjerg, 2019 for a list of encysting



**Fig. 1** **a** In toto cyst of the freshwater eutardigrade *Hypsibius* sp. (phase contrast). **b** In toto cyst of the freshwater eutardigrade *Dactylobiotus parthenogeneticus* (phase contrast). Bar = 10  $\mu$ m

tardigrade species reported in the literature). Detailed steps in encystment in *Dactylobiotus* and moss-dwelling/freshwater *Bertolanius* were provided by Guidetti et al. (2006). *Dactylobiotus* has only one type of cyst, which is dark-reddish brown (Szymańska, 1995; Guidetti et al. 2006, 2008), whereas *Bertolanius*, which has both limnic and moss-dwelling species, forms two types of cysts (“white/type 1” in cold periods and “red/type 2” in warm periods) that Westh and Kristensen (1992) correlated with seasonal environmental changes in Greenland. Cyst formation in *Bertolanius* is cyclic and it is a part of cyclomorphosis, defined as cyclic and reversible morphological modifications within a single species (Kristensen, 1982; Rebecchi & Bertolani, 1994; Hansen & Katholm, 2002). The production of extra cuticles isolates and protects the animals from environmental factors. Since the cysts remain viable for several months, encystment enhances tardigrade survival of freezing in winter and desiccation in summer (since limnic tardigrades often disappear in summer). Although encystment is best studied in limnic species, which do not undergo anhydrobiosis (but a few species can withstand cryobiosis), most of the species that produce cysts can also enter anhydrobiosis (Guidetti et al., 2011a). Since diapause (encystment) and cryptobiosis are dormancy states that can be present in a single species, their evolution was not mutually exclusive. Although we are beginning to understand the molecular mediators involved in cryptobiosis, the molecular mechanisms involved in encystment (see Rozema et al., 2019) are unknown.

### Extreme-resistance strategy: anhydrobiosis

The most widespread and best-known form of extreme-stress resistance evolved by tardigrades, rotifers and nematodes is the capability to tolerate complete desiccation (drying to  $< 0.1 \text{ g H}_2\text{O g}^{-1}$  dry mass) by entering in a state of reversible suspension of metabolism called anhydrobiosis (“life without water”) without the loss of viability. At the end of dehydration process, tardigrades have lost 97% of their body water (Westh & Ramløv, 1991; Horikawa et al., 2008), and similar values have been shown for the anhydrobiotic nematodes *Ditylenchus dipsaci* (Kuhn, 1857), *Aphelenchus avenae* Bastian, 1865 and *Panagrolaimus superbus* (Fuchs, 1930) (Crowe &

Madin, 1975; Wharton, 1996; Banton & Tunnacliffe, 2012).

Anhydrobiosis indicates a fundamental concept about the nature of living systems since an anhydrobiotic organism lacks all dynamic features of living organisms due to the absence of detectable metabolism. In that sense it is not alive, but it is not dead because rehydration produces a living organism and a kind of resuscitation routinely occurs (Clegg, 2001; Tunnacliffe & Lapinski, 2003). Consequently, anhydrobiotic organisms have two distinct living physiological states: active and anhydrobiotic.

Despite its clear adaptive potentiality, anhydrobiosis can be found only in a restricted number of metazoans whose sizes generally do not exceed 1 mm, with the exception of a few taxa that can reach 5–7 mm in length, such as the larvae of the African midge *Polypedylum vanderplanki* Hinton 1951 (Watanabe et al., 2004). These apparent morphological and ecological characteristics could be linked to limiting factors required for tolerating physical and physiological constraints imposed by complete dehydration (Alpert, 2005). In animals, desiccation tolerance occurs either in the whole animal at any stage of their life cycle, from the egg to the adult stage (tardigrades, bdelloid rotifers and nematodes), in which case the animals are defined as holo-anhydrobiotic (Jönsson, 2005; Rebecchi et al., 2007), or at a specific life stage, usually egg/embryo/larval stage (shrimps, the midge *P. vanderplanki*, monogonont rotifers, some nematodes).

As described above, anhydrobiosis allows tardigrades, rotifers and nematodes to colonize and persist in various otherwise unavailable environments. A high number of species colonize habitats subjected to periodic desiccation (e.g. lichens, mosses and ephemeral lakes and ponds) that are prohibitive for most other animals. In these habitats, they perform all activities of routine life only when there is at least a small layer of water around the body of the animals. For example, mosses and lichens provide habitats featuring a myriad of small pockets of water; as their surroundings lose water through evaporation, animals lose water with them. Consequently, their life cycle consists of active periods for growth and reproduction, interrupted by periods of metabolic inactivity (Jönsson, 2005; Glime, 2017). When rehydrated by dew, rain or melting snow, they can return to their active state in a few minutes to a few hours. Therefore, during

their life, holo-anhydrobiotic animals can enter anhydrobiosis several times (e.g. Ricci, 1987; Womersley, 1987). An experimental study evidenced that the moss-dwelling eutardigrade *Richtersius coronifer* (Richters, 1903) may survive up to 6 repeated desiccations, with a declining survival rate with an increasing number of desiccation events (Czernekova and Jönsson, 2016). Interestingly, repeated desiccation seems also to improve the long-term survival of rotifer populations. Populations that are regularly subjected to desiccation grow faster than permanently hydrated corresponding cohorts, suggesting that diapause is not only a strategy to survive harsh environmental conditions, but it also has ecological advantages to the organisms that managed to evolve this strategy (Ricci et al., 2007; Sommer et al., 2019).

The time for recovery to active life after a period of anhydrobiosis is directly related to the environmental condition during the desiccation phase (e.g. humidity rate during the desiccation process) in which higher stressors lead to longer recovery time, and to the time spent in anhydrobiosis (Rebecchi et al., 2009a). The recovery time is probably function of the metabolic activities linked to the repair of damages caused by desiccation and/or to the restoration of metabolic pathways (see Mattimore & Battista, 1996).

Among anhydrobiotic tardigrade and rotifers studied, desiccation tolerance varied from zero to high tolerance (e.g. Ricci, 1987; Wright, 1989a; Bertolani et al., 2004; Rebecchi et al., 2006). These gradients are correlated with the abiotic factors (e.g. humidity) of the substrate inhabited since species living in constantly wet or submerged mosses usually show lower anhydrobiotic performance than those living in mosses growing on trees and rocks (Guidetti et al., 2011b; Eyres et al., 2015). In addition, anhydrobiotic capability is similar among species belonging to distant evolutionary lines, but they can be very different among closely related species. However, species with similar ecological requirements share a close similarity in anhydrobiotic performances (Wright, 1991, 2001; Ricci, 1998, 2001; Fontaneto et al., 2004; Ricci & Caprioli, 2005; Fontaneto & Ambrosini, 2010; Guidetti et al., 2011b; Eyres et al., 2015).

Therefore in both rotifers and tardigrades, we hypothesize that anhydrobiosis is more likely linked to local adaptations to habitats than to phylogenetic relationships suggesting that anhydrobiotic

capabilities have been evolved once and secondarily lost in some lineages.

Some species of nematodes within the genus *Panagrolaimus* Fuchs, 1930 can survive immediate desiccation (e.g. Ricci and Pagani, 1997) and are referred to as fast-desiccation strategists, while others (e.g. *A. avenae*) require a period of slow-drying (pre-conditioning) and are referred to as slow desiccation strategists (e.g. Womersley, 1987; Shannon et al., 2005). Similar patterns were detected in tardigrade and rotifer species when experimentally desiccated under laboratory conditions (e.g. Ricci, 1987, 2001; Wright 1989a; Eyres et al., 2015; Hashimoto et al., 2016; Boothby et al., 2017). Full anhydrobiotic nematodes can undergo desiccation at any stage of their life cycles, but recent studies have suggested that some species, traditionally considered intolerant to desiccation, can actually survive desiccation at least in some stages of their life cycle [e.g. the dauer larvae of the model species *Caenorhabditis elegans* (Maupas, 1900)] (Erkut et al., 2011). As in tardigrades and rotifers, the anhydrobiotic abilities of different taxa of nematodes do not appear to be related to their phylogeny, suggesting that the evolutionary processes have affected the loss or maintenance of this remarkable ability. Although traditionally less studied, the recent characterization of some of the molecular strategies of diapause in nematodes, and especially in the well-known and well-characterized model organism *C. elegans*, allows a better understanding of how diapause is induced, maintained and what its effects are, as well as common mechanisms to different organisms (e.g. Fielenbach & Antebi, 2008; Hand et al., 2016).

In the desiccated state, holo-anhydrobiotic animals are biostable for decades (e.g. tardigrades 20 years; Guidetti & Jönsson, 2002; Bertolani et al., 2004; Rebecchi et al., 2006; Jørgensen et al., 2007) even though recently the consistent long-term survival of at least some taxa under desiccation has been debated (Jönsson & Bertolani, 2001; Fontaneto et al., 2012a). For example, a comparative study of the survival rate of different taxa and the statistical model developed from it suggested that recovery of bdelloid rotifers, tardigrades and nematodes found in lichens within collections in museums decreases to almost zero after desiccation periods of up to 10 years; this is significantly longer than the life span of single individuals in the active state, but is not as long as anecdotally

suggested by other studies, and not as long as in other taxa like resting eggs of monogonont rotifers (Cáceres, 1997; Fontaneto et al., 2012a). These data confirm that these organisms do survive long periods of desiccation but that the rate and general conditions of desiccation, as well as the substrate and the storage conditions during diapause, influence survival in a significant way (e.g. Ricci & Caprioli, 2001; Fontaneto et al., 2012a).

Other than its effect on longevity, anhydrobiosis can have an impact on ageing in meiofauna as illustrated by the “Sleeping Beauty” and “Picture of Dorian Grey” models derived from experimental data on a few species of holo-anhydrobiotic organisms (for a review, see Kaczmarek et al., 2019). The first model predicts that anhydrobiotic organisms do not age during anhydrobiosis in at least some tardigrade and bdelloid rotifer species (Ricci & Covino, 2005; Hengherr et al., 2008a, b). The latter model predicts that anhydrobiotic organisms age, at least in the initial stages of the anhydrobiosis process, as in some species of nematodes (Ricci & Pagani, 1997). Nevertheless, a comprehensive comparative analysis that considers all taxa and strategies is still lacking.

Numerous studies have focused on molecular changes during ageing in tardigrades, rotifers and nematodes, especially from the molecular approach, and the potential “rejuvenation” of stressed animals, but the full picture is very complex and still poorly understood. Early studies highlighted general changes in protein patterns with age (Carmona et al., 1989), and recent advances have started uncovering specific changes in regulatory molecules (e.g. Snell et al., 2014), protein modifications like carbonylation (Krisko & Radman, 2019) and improved physiological characteristics like fecundity (Ricci & Covino, 2005; Ricci & Perletti, 2006). Based on these and other studies, rotifers can be added to the list of useful model organisms which can be used to study ageing (Snell et al., 2015), although the exact links between molecular changes and ageing are still not fully characterized. Even more obscure at the moment are the precise links between the ability of some types of dormancy to stop or reverse ageing. For example, both desiccation and starvation seem to stop or reverse ageing in bdelloid rotifers, allowing dormant bdelloids to “wake up” with similar or higher fitness than animals in the pre-stressed condition (Ricci & Covino, 2005; Ricci & Perletti, 2006; Sommer et al., 2019).

Some recent advances suggest that some of the mechanisms and molecules involved in the organism’s protection during desiccation, like antioxidants or LEA proteins, can also prevent at least some aspects correlated with ageing (e.g. Kaneko et al. 2005; Snare et al., 2013). Ageing is generally better characterized in nematodes, although the majority of studies are limited to model species like *C. elegans* (e.g. Schafitzel & Hertweck, 2006; Hughes et al., 2007; Mack et al., 2018) and therefore lack the more direct link between ageing and dormancy in stress-resistant animals from natural habitats. Interestingly, where data are available, they suggest that the rejuvenation effect of desiccation is not present in at least some anhydrobiotic nematodes of the genus *Panagrolaimus* (Ricci and Pagani, 1997), making the understating of the relationship between desiccation resistance and ageing even more fascinating and interesting.

Even though dehydration can have a major effect on survival, ageing and longevity, the anhydrobiotic process per se can induce molecular damages that accumulate with time, reducing the viability of desiccated animals (França et al., 2007; Tyson et al., 2007; Neumann et al., 2009; Rebecchi et al., 2009a; Marotta et al., 2010; Hespels et al., 2014). The amount of these damages is directly impacted by high temperature, high humidity level and high oxygen partial pressure. In tardigrades, the time required to recover active life after a period of desiccation is affected by these abiotic conditions and can be related to the metabolic activities necessary to repair molecular damages and to catabolise damaged molecules (Rebecchi et al., 2009a; Guidetti et al., 2011a). Different strategies and molecules seem to be involved in the reduction and/or repair of molecular damage (see below).

Even more striking, in the dry state, anhydrobiotic organisms show extraordinary resistance to physical and chemical extremes (very low sub-zero temperature, high pressure, radiation, extreme pH, toxic chemicals, lack of geomagnetic field) that may far exceed the tolerance ranges of active organisms (Wharton et al., 2003; Jönsson et al., 2005, 2013; Watanabe et al., 2006; Rebecchi et al., 2007; Gladyshev and Meselson, 2008; Rebecchi et al., 2009b; Altiero et al., 2011; Guidetti et al., 2011a; Krisko et al., 2012; Rebecchi, 2013; Hashimoto et al., 2016; Erdmann et al., 2017; Jönsson & Wojcik, 2017; Giovannini et al., 2018).


In tardigrades, a strong correlation between the capability to withstand desiccation and the capability to withstand sub-zero temperatures ( $-20^{\circ}\text{C}$ ,  $-80^{\circ}\text{C}$ ) was detected, and species that were not able to enter anhydrobiosis showed low or no capability to withstand sub-zero temperatures (Guidetti et al., 2011a, b). This direct relationship could be related to the fact that during both desiccation and freezing stresses, tardigrades are under the same selective pressure induced by a wide variation in body fluid osmolality and in cell volume (Sømme, 1996; Guidetti et al., 2011b). Nevertheless, the freeze resistance of anhydrobiotic tardigrades should be distinguished from cryobiosis, which is the ability of active hydrated animals in contact with water to freeze and survive after thawing (Guidetti et al., 2011b).

The aggregate of all these characteristics, especially radiation tolerance, has led to the characterization of tardigrades as the “toughest animals on the Earth” (Copley, 1999) and to make them an emerging model for space biology (Jönsson, 2007; Horikawa et al., 2008; Erdmann and Kaczmarek, 2017), more recently joined by bdelloid rotifers. Tardigrades and rotifers have been exposed to space stressors in Low Earth Orbit several times, on board of the International Space Station and FOTON (Ricci & Boschetti, 2003; Ricci et al., 2005; Leandro et al., 2007; Jönsson et al., 2008; Selch et al., 2008; Rebecchi et al., 2009b, 2011; Persson et al., 2011; Guidetti et al., 2012; Vukich et al., 2012).


### Morphological, physiological and molecular adaptations enabling anhydrobiosis

The evolution of a series of behavioural, morphological, physiological and molecular/biochemical adaptations provided anhydrobiotic organisms with mechanisms to withstand the deleterious effects caused by the drastic loss of water. The majority of holo-anhydrobiotic organisms cannot survive a desiccation rate that is too rapid (as shown in a few hours in laboratory experiments, even though the rate is species-dependent (Wright, 1989a, b, c; Wright et al., 1992; Jönsson & Järemo, 2003; Banton & Tunnacliffe, 2012; Boothby et al., 2017), so they have evolved different strategies to slow down the rate of water evaporation.

To reduce the rate, the tardigrade shrivels into a barrel-shaped structure (“tun”), about one-third of its original size, by contracting the body anterior-posteriorly and withdrawing the legs and head (Figs. 2 and 3). Tun formation produces a new spatial organization of some internal organs (such as the pharyngeal bulb), and epidermal cells, storage cells, ovarian cells and digestive system cells undergo shrinkage, containing electron dense cytoplasm (Czerneková et al., 2016). Lipids and polysaccharides dominate in the reserve material of the storage cells, whereas the amount of protein is small (Czerneková et al., 2016). The tun minimizes the permeability and evaporative surface of the organism by removing the high permeability areas of the cuticle from direct contact with the air, resulting in a slow rate of desiccation (Wright, 1988a, b, 1989a, b, c, 2001). Differences in the reduction of cuticle permeability detected among tardigrade species are related to the level of desiccation tolerance of each species and to the morphology of the cuticle in eutardigrades and heterotardigrades (Wright, 1989a, b). The permeability slump of the cuticle permits animals to lose water slowly, allowing animals to produce bioprotectants. Somewhat similarly to tardigrades, bdelloid rotifers contract their




**Fig. 2** **a** In toto specimen of the limnoterrestrial eutardigrade *Acutuncus antarcticus* (in vivo and Nomarski contrast). **b** In toto female of the lichen-dwelling eutardigrade *Ramazzottius* cf. *oberhaeuseri*; the ovary containing three oocytes (asterisk). (in vivo and Nomarski contrast). **c** Tun (desiccated animal) of the eutardigrade *Ramazzottius* cf. *oberhaeuseri* (in vivo). Arrow: buccal-pharyngeal apparatus; arrowhead: midgut; cross: gonad. Bar = 10  $\mu\text{m}$



**Fig. 3** Scanning electron micrographs of the moss-dwelling heterotardigrade *Echiniscus* sp. **a** Dorsal view of an in toto and hydrated specimen. **b** Dorsal view of an in toto desiccated specimen (tun). **c** Ventral view of an in toto desiccated specimen (tun). Bar = 10  $\mu\text{m}$

body into a compact shape by withdrawing their cephalic and caudal extremities into the trunk, facilitated by muscle contractions and by a coordinated



**Fig. 4** Scanning electron micrographs of the rotifer *Adineta tuberculosa*. **a** In toto and hydrated specimen. **b** In toto desiccated specimen (tun). Bar = 100  $\mu\text{m}$ . (Courtesy of Giulio Melone and Diego Fontaneto)

morphological arrangement of internal structures (Ricci, 2001; Ricci et al., 2003; Marotta et al., 2010; Fig. 4). A decrease in permeability (the permeability slump) during the early stages of desiccation was detected in the anhydrobiotic plant-parasitic nematode *D. dipsaci* during which the surface of the animal body was coated with an extracuticular layer of lipid (triglyceride) that produced a slow rate of water loss necessary for its survival (Wharton et al., 2008). Nematodes tend to coil their body (Crowe, 1971) and certain nematodes are also reported to congregate into masses of “nematode wool”, with better survival of specimens in the centre of the mass (Ellenby, 1968). The aggregation effect has also been experimentally produced in tardigrades (Ivarsson & Jönsson, 2004), but not yet verified in nature.

As water evaporates and dry conditions set in, holo-anhydrobiotic organisms start generating a variety of protective agents, collectively termed bioprotectants, which they accumulate in and around the cells of their body. It was initially thought that non-reducing disaccharides, like trehalose, were solely responsible for preventing damage (e.g. Crowe et al., 1984, 1992), but more recent studies point to a complex picture of molecular adaptations. These bioprotectant molecules include: sugars, mostly disaccharides such as trehalose; a unique repertoire of proteins generally lacking persistent tertiary structure classified as



intrinsically disordered proteins (IDPs) or proteins with intrinsically disordered regions (IDRs) and represented by Late Embryogenesis Abundant proteins (LEA), Heat Shock proteins (HSPs), cytoplasmic abundant heat soluble (CAHS) proteins, secretory abundant heat soluble (SAHS) proteins and mitochondrial abundant heat soluble (MAHS) proteins; antioxidants, and molecules involved in protection from or repair of DNA damage (e.g. Lapinski & Tunnacliffe, 2003; Schill et al., 2004; Jönsson & Schill, 2007; Pouchkina-Stantcheva et al., 2007; Förster et al., 2009, 2011; Schokraie et al., 2010; Boschetti et al., 2011; Altiero et al., 2012; Yamaguchi et al., 2012; Boschetti et al., 2013; Rebecchi 2013; Wang et al., 2014; Tanaka et al., 2015; Hashimoto et al., 2016; Boothby et al., 2017; Schill & Hengherr, 2018). A recent study showed that one of the most stress-tolerant tardigrade species (*Ramazzottius varieornatus* Bertolani and Kinchin, 1993) has a tardigrade-unique DNA-associating protein, termed Dsup, which is able to suppress the incidence of DNA breaks caused by radiation (Hashimoto et al., 2016). Accumulation of these xeroprotectants is generally slow and gradual, taking place in parallel with the drying process, although a few taxa, like the nematode *P. superbus*, seem to express a full repertoire of protective molecules and have therefore been defined as fast-desiccation strategist (Shannon et al., 2005; Banton & Tunnacliffe, 2012). Even though many organic compounds have been identified in tardigrades, rotifers and nematodes, the biochemical and molecular mechanisms involved in complete desiccation tolerance are currently little known and constitute an intriguing challenge for biologists. For instance, it is well known that in some species of tardigrades the synthesis of the disaccharide trehalose counteracts the loss of water, as well as other environmental extremes (Westh & Ramløv, 1991; Hengherr et al., 2008b; Jönsson & Persson, 2010; Welnicz et al., 2011; Cesari et al., 2012; Schill & Hengherr, 2018). In any case, the absolute trehalose levels detected in tardigrades are much lower than those reported for other anhydrobiotic organisms. This sugar has a double role in desiccation-tolerant organisms. As the trehalose replaces water, it protects biomolecules and the integrity of membranes during dehydration, and participates in the formation of a glassy matrix that reduces the rates of chemical reactions and inhibits free radical production (Crowe et al., 1984; Teramoto

et al., 2008). The CAHS, SAHS and MAHS proteins have been detected so far only in eutardigrades (Tanaka et al., 2015; Boothby et al., 2017). However, the distribution of the encoding genes of these proteins is scattered among tardigrades, suggesting species- or at least taxon-specific adaptations (Yoshida et al., 2017; Kamilari et al., 2019). The CAHS and SAHS proteins probably form a molecular shield inside and outside cells, respectively, whereas MAHS proteins are defined as potent specific mitochondrial protectants (Boothby et al., 2017). The heterologous expression of some CAHS proteins in both prokaryotic and eukaryotic cells allows an increase in their tolerance to desiccation, and purified CAHS proteins protect desiccation-sensitive proteins in vitro (Boothby et al., 2017). The very low or null metabolic activity of anhydrobionts limits the production/accumulation of products of metabolism such as Reactive Oxygen Species (ROS). Even though the origin of ROS in anhydrobiosis is not yet well known, their production can occur both during the dehydration, in a permanent desiccated state as well as during rehydration, so an efficient antioxidant mechanism is necessary (França et al., 2007; Cornette et al., 2010; Rebecchi, 2013). For example, in desiccated specimens of the eutardigrade *Paramacrobiotus spatialis* Guidetti et al., 2019, glutathione peroxidase was the most abundant antioxidant enzyme in hydrated animals, followed by the enzyme superoxide dismutase and glutathione content (Rizzo et al., 2010). With regard to the repair of DNA damages, desiccation enhanced the expression of DNA-repair proteins in tardigrades (Wang et al., 2014; Kamilari et al., 2019).

Rotifers possess similar strategies, but there are also marked differences. Neither trehalose nor the metazoan genes for its synthesis have been found in bdelloid rotifers (Lapinski & Tunnacliffe, 2003), although trehalose has been found in monogonont rotifers (Caprioli et al., 2004), suggesting that non-reducing disaccharides are not necessary for successful recovery from desiccation. Instead, other molecules are now thought to be essential to protect molecules, cells and tissues and to repair any damage caused by anhydrobiosis. It is becoming clear that no single class of protectant/repair molecules is sufficient, but successful desiccation depends on the coordinated action of all of them. These molecules, which are usually upregulated upon desiccation, include different types of LEA proteins which perform

different and often still uncharacterized functions, other (non-LEA) protein families, often at least partially unstructured (IDPs and proteins containing IDRs) and with still uncharacterized functions, other types of hydrophilins or chaperones, different types of antioxidants, molecules involved in DNA repair as well as, probably, other still unknown molecules and mechanisms (e.g. Browne et al., 2002; Browne et al., 2004; Goyal et al., 2005; Pouchkina-Stantcheva et al., 2007; Denekamp et al., 2010, 2011; Boschetti et al., 2011, 2012; Hanson et al., 2013; Hespeels et al., 2014).

All these molecules, some of which are taxon-specific while others are common to all analysed taxa (e.g. Denekamp et al., 2010; Mali et al., 2010; Boschetti et al., 2011, 2012; Hanson et al., 2013; Hashimoto et al., 2016; Boothby et al., 2017; Hashimoto and Kunieda, 2017; Kamilari et al., 2019), seem to be necessary for an integrated and effective response to anhydrobiosis.

Interestingly, the majority of the previously mentioned studies were based on the analyses of one or a relatively small subset of genes, but recent technological advances have allowed analyses of whole genomes and transcriptomes and are uncovering an even more fascinating story, suggesting that some animals, and bdelloid rotifers in particular, have been acquiring genes, which code for protective/repair molecules, from organisms that are not direct ancestors and can even belong to different taxa, in a process known as Horizontal Gene Transfer (HGT, also called lateral gene transfer, LGT). Horizontal gene transfer was previously known only in bacterial and archaeal organisms and was thought to be absent in eukaryotic organisms, but recent studies suggest that it is more widespread than previously thought (e.g. Dunning Hotopp, 2011; Boto, 2014; Drezen et al., 2017) and that these “foreign” genes can indeed contribute to the resistance to desiccation of some organisms, especially bdelloid rotifers. Initial suggestions that bdelloid rotifers possess a very high percentage of genes acquired via HGT (Gladyshev et al., 2008; Boschetti et al., 2012) and “domesticated” (Barbosa et al., 2006) have now been confirmed and expanded (Flot et al., 2013; Eyres et al., 2015; Hespeels et al., 2015; Nowell et al., 2018). This unusual characteristic is made even more interesting by the recent understanding of the role that these foreign genes play in stress resistance: many foreign genes are over-expressed during

desiccation or rehydration and might therefore at least partially responsible for their successful anhydrobiotic capabilities (Boschetti et al., 2011, 2012; Eyres et al., 2015), although the precise link between desiccation resistance and levels of HGT is still unclear (e.g. Eyres et al., 2015; Nowell et al., 2018). Indeed, recent studies have found genes involved in trehalose synthesis in bdelloids, but they seem to have been originated by HGT (Hespeels et al., 2015), while other foreign genes can add various biochemical capabilities, some of which might improve the desiccation abilities of bdelloids (Boschetti et al., 2012; Szydlowski et al., 2015). This unusual high level of foreign genes seems to be a characteristic only of bdelloid rotifers: other taxa have been analysed and, although a few foreign genes are present (e.g. Boothby et al., 2015; Bemm et al., 2016; Koutsovoulos et al., 2016; Nowell et al., 2018), they are not so abundant, and the details of their contribution to successful desiccation is still being characterized (e.g. Yoshida et al., 2017; Nowell et al., 2018; Kamilari et al., 2019).

### Ecological and evolutionary consequences of extreme tolerance of meiofaunal organisms

The evolution of anhydrobiosis is the result of trade-offs between the selective advantages of this adaptive strategy, the energetic costs and the physical and physiological constraints related to the process (Jönsson, 2005; Guidetti et al., 2011a). Energy is probably necessary to produce and accumulate bio-protectants during the initial phase of anhydrobiosis and to catabolize them during the exit phase (rehydration). There are few data on anhydrobiotic energetic costs, but a substantial energetic cost of anhydrobiosis was shown in the tardigrade *Richtersius coronifer* (Jönsson and Rebecchi, 2002) and in some species of nematodes (Madin and Crowe, 1975; Demeure et al., 1978). Little is known about rotifers, but the presence of lipid droplets (Wurdak et al., 1978) and the differential expression of some genes potentially involved in lipid metabolism or protection (Denekamp et al., 2009) in monogonont resting eggs and in desiccated bdelloids (Marotta et al., 2010) suggest that costs are present in these taxa as well.

The anhydrobiotic process requires energy that is withdrawn from other physiological functions such as growth and reproduction. This should have strong

effects on the life histories of holo-anhydrobiotic organisms. Even though there is no direct evidence for a trade-off between anhydrobiosis and fitness, the few ecological studies on this topic are consistent with the hypothesis that fitness of desiccation-tolerant organisms is lower (Jönsson, 2005; Alpert, 2006; Guidetti et al., 2007). In tardigrades, both positive and negative relationships between body size (as an indication of age) and desiccation performance have been demonstrated at the intraspecific and interspecific levels. In species living in the same moss and with high anhydrobiotic performance, desiccation survival increases in *R. coronifer* with an increase of the body size, whereas it decreases in *Ramazzottius oberhaeuseri* (Doyère, 1840) (Jönsson et al., 2001; Jönsson and Rebecchi, 2002). These contrasting models could be due to genetic differences and/or contingent factors, such as nutritional state, level of molecular protectants and some life history traits, including age, reproductive stage and phenotypic plasticity. Lastly, differences in anhydrobiotic performances among geographically isolated populations of eutardigrades have been reported, but the literature data are conflicting (Horikawa & Higashi, 2004; Jönsson et al., 2001) probably due to the presence of cryptic species and differences in ecological conditions of the microhabitats.

Interestingly, as previously mentioned, bdelloid rotifers seem to be different, i.e. desiccation improves individual and population fitness (Ricci et al., 2007; Sommer et al., 2019), but with still unknown mechanisms.

In addition, low fitness associated with a long-lifespan could slow down rates of evolution in comparison to organisms with similar lifespans but without the capability to perform anhydrobiosis. Furthermore, ancestral genetic traits may reappear after a long time in anhydrobiosis, jumping generations and contributing to the longer existence of unchanged traits (Kaczmarek et al., 2019). Anhydrobiosis represents an “escape in time” from habitat conditions hostile to active life, opposed to an “escape in space” performed by organisms with an ability to migrate away from unfavourable conditions (Jönsson, 2005). In addition, it limits selection and creates a “seedbank” for maintaining haplotypes in time and space (environment) (Guidetti et al., 2011a). These advantages are reinforced by the ageing models (“Sleeping Beauty” and “Picture of Dorian Grey”)

allowing organisms to withstand adverse conditions for a long time and the capability to restore active life and reproduction when environmental conditions become suitable (Kaczmarek et al., 2019). Such scenarios are in line with the hypothesis that anhydrobiotic organisms almost avoid environmental selection since they are active only under favourable environmental conditions (Pilato, 1979). Therefore, anhydrobiotic periods could have an impact on generation time, which in turn influences the potential rate of evolution. This could be the cause of the surprising morphological uniformity at the species, genus and family level of terrestrial anhydrobiotic tardigrades in contrast to marine species that, in practice, are not able to enter anhydrobiosis (Kaczmarek et al., 2019). Interestingly, molecular analyses have suggested that bdelloid and monogonont rotifers might have different diversification and mutation rates, although it is still unclear if this is due to the different dormancy patterns (quiescence vs dormancy, respectively) or the different reproductive strategies (obligately vs cyclical parthenogenesis, respectively) of these taxa or other, still unknown, factors, and if they are indeed common (Barraclough et al., 2007; Swanstrom et al., 2011; Fontaneto et al., 2012b).

Further selective advantages of anhydrobiosis can be cited. Anhydrobiosis allows the reduction of predators, competitors and parasites since stochastic habitats are colonized only by a reduced number of species (Guidetti et al., 2011a; Wilson & Sherman, 2013). Since holo-anhydrobiotic organisms are aquatic animals, desiccation tolerance allows them to colonize and persist in terrestrial habitats other than in stochastic and extreme habitats. Moreover, the capability to withstand extreme conditions by entering anhydrobiosis increases the number of possible “refugia” that can be utilized by the species during long harsh environmental conditions, with a decrease in the rate of extinction and the loss of diversity (Guidetti et al., 2011a). Anhydrobiosis increases passive dispersal capability since dormant anhydrobiotic animals and eggs can act as propagules, be transported over long distances and cross physical barriers for months without losing viability, which active animals cannot do, and establish new populations in new territories (Guidetti et al., 2011a; Mogle et al., 2018; Fontaneto, 2019). This capability is aided by the fact that mostly holo-anhydrobionts reproduce via telytokous parthenogenesis, a reproductive strategy favourably

adapted to colonize new and isolated habitats with a single individual (Bertolani, 2001; Ricci & Fontaneto, 2009; Fontaneto, 2019). This could influence the biogeographical pattern of holo-anhydrobionts that supports the hypothesis that “everything is everywhere”. This hypothesis was confirmed for many bdelloid rotifers (Fontaneto et al., 2008), monogonont rotifers (e.g. Gómez et al. 2002; Mills et al., 2017) and for few tardigrade species (Kaczmarek et al., 2019), although some caution should be exercised, as other variables, including sampling effort or hidden diversity, might influence results and should therefore be carefully considered (Fontaneto et al., 2007, 2008, 2009; Mills et al., 2017). Nevertheless, despite that tardigrades are able to disperse by wind as are other terrestrial anhydrobionts (Nkem et al., 2006; Rivas et al., 2019), most tardigrade species have a narrow species range, with a large number of endemic species (e.g. Guidetti et al., 2019). In contrast, biogeographic patterns were detected in several anhydrobiotic taxa of nematodes (Faurby and Barber, 2015; Zullini, 2018). In any case, the paucity of faunistic data, the presence of cryptic species and the high level of confounding factors make the distribution patterns more complex with the exigency to collect further experimental and faunistic data. Some of these effects on the life history, like the ability to “escape in time”, ageing, “refugia”, generation time, selection and avoidance of predators, are also valid for other dormant stages, for example resting eggs and cysts, even when they are not desiccated, highlighting common ecological effects of dormancy, irrespective of the physiological adaptations of each taxon or response to stress.

These conditions, together with the capability of desiccation-tolerant organisms to repopulate habitats when liquid water returns, affect community dynamics and produce substantial modifications in the structure of biological communities, even leading to modifications in the functional integrity of the ecosystems (Irons et al., 1993; Walsh et al., 2014).

### **Implications/application of extreme tolerance of meiofaunal organisms**

A better understanding of the life strategies of anhydrobiotic animals both at the ontogenetic and phylogenetic levels can provide answers to many

fundamental questions as well as useful practical outcomes in many branches of applied sciences. Understanding desiccation tolerance in anhydrobiotic organisms will enable us to induce or engineer tolerance in sensitive species and to produce subsequent long-term stabilization and preservation of biological material in a dry state. This is a topic of considerable practical importance both in medical and commercial fields since drying is widely used in the food and pharmaceutical industries as a long-term preservation technique (Saragusty & Loi, 2019).

Based on knowledge accumulated from anhydrobiotic organisms, much of the research on stabilizing cellular membranes and proteins has centred on trehalose, which preserves cell membranes, and proteins, which can allow fluids to solidify without forming crystals through glass transition or vitrification, forming a large number of hydrogen bonds with membranes and proteins, and by replacing water molecules during the drying process (Hengherr et al., 2009), although many other protein families, as well as molecules, perform many functions, some of which still uncharacterized (e.g. Tompa, 2002; Tunnacliffe & Wise, 2007; Tunnacliffe et al., 2010). Some anhydrobiotic organisms naturally possess the molecular mechanisms to produce these sugars and load and unload them to and from the cells of the body or the intracellular spaces. Since the first report that biomolecules, membranes and organisms can be stabilized in a dry state, due to the presence of trehalose, an array of possible applications for trehalose have been reported, ranging from the stabilization of vaccines, lysosomes, platelets, spermatozoa and oocytes to the hypothermic storage of human organs (Crowe et al., 2005; Schill et al., 2009; Saragusty & Loi, 2019). The determination of the properties of trehalose and the debate whether trehalose alone is sufficient to preserve biomolecules (e.g. García de Castro & Tunnacliffe, 2000; Ratnakumar & Tunnacliffe, 2006; Pouchkina-Stantcheva et al., 2007; Tapia et al., 2015, Chau et al., 2016) have stimulated the continuation of basic research to discover the secret of life without water. Recently, Boothby et al. (2017) indicated that the heterologous expression of some CAHS proteins in both prokaryotic and eukaryotic cells is sufficient to increase desiccation tolerance in these sensitive systems, and purified CAHS proteins protect desiccation-sensitive proteins *in vitro*. Moreover, Hashimoto et al. (2016) found that

tardigrade DNA-associating protein (Dsup) suppresses X-ray-induced DNA damage by 40% and improves radiotolerance of human cultured cells. The story of the already known bioprotectants tells us that the tolerant ability of anhydrobiotic animals could be transferred to more sensitive organisms at least partly by transferring the corresponding genes. Recent rapid progress of molecular analyses should accelerate the elucidation of the mechanisms at the basis of extreme stresses, including complete desiccation stress, providing novel clues that open new avenues to confer stress resistance to intolerant species, including humans.

## Conclusions

Various extreme-tolerance mechanisms have evolved in meiofauna, enabling micrometazoans like tardigrades, rotifers and nematodes, to reduce or interrupt metabolism and survive stressful environments. In response to the gradual onset of adverse environmental conditions (e.g. water availability, temperature, oxygen tension, pH), these organisms undergo complex molecular, physiological, morphological and behavioural changes, which can share common characteristics but also present some differences. For example, tardigrades undergo encystment, an adaptive strategy that involves profound morphological changes that occur during the molting process, resulting in the dormant organism lying within retained cuticular exuvia. On the other side, cryptobiosis happens at any stage of the life cycle of the organisms and includes different strategies such as anhydrobiosis, cryobiosis, anoxybiosis and osmobiosis directly induced by desiccation, sub-zero temperatures, low oxygen pressure and osmotic extremes, respectively. The most widespread and best-known form of these is anhydrobiosis, the capability evolved by tardigrades, rotifers and nematodes to tolerate complete desiccation by entering in a state of reversible suspension of metabolism without the loss of viability.

When dormant, these taxa show extraordinary resistance to physical and chemical extremes that may far exceed the tolerance ranges of active organisms, therefore the two dormancy strategies, quiescence and diapause, allows tardigrades, rotifers and nematodes to colonize and persist in various otherwise unavailable environments. Interestingly, while

dormant, some taxa do not age, although the specific effects of dormancy on ageing varies with the taxa and is poorly understood but this ability make tardigrades, rotifers and nematodes very useful model organisms that can be used to study the ageing process. Furthermore, the evolution of anhydrobiosis resulted in selective advantages but also in energetic costs with effects on growth, reproduction, life history and fitness, in turn affecting the rate of evolution, but more studies are needed to fully understand the ecological and evolutionary implications of these resistance strategies on these taxa.

Furthermore, novel findings have also contributed to expand other aspects of these taxa, with potential exciting applications in other fields: the evolution of a series of behavioural, morphological, physiological and molecular/biochemical adaptations provided anhydrobiotic organisms with different unusual mechanisms to withstand desiccation. To prevent cell damage during dehydration, bioprotectant molecules that accumulate in and around the cells of their body are generated; the identification of these molecules and their mechanisms are the focus of much current research, including the role of horizontal gene transfer. It is becoming clear that no single class of protectant/repair molecules is sufficient, but successful desiccation depends on the coordinated action of all of them. These meiofauna taxa are becoming popular model organisms in the fields of exobiology and medical research, with the hope that they might also help to improve the tolerance of human cells to extreme stress in the future.

**Acknowledgements** We are grateful to Prof. Giulio Melone (University of Milan, Italy) and Dr. Diego Fontaneto (National Research Council of Italy, Italy) for pictures on rotifers, and Dr. Edoardo Massa (University of Modena and Reggio Emilia, Italy) for the editing of all pictures. Finally, we thank the reviewers for their helpful suggestions which improved the quality of the manuscript.

## References

- Alpert, P., 2005. The limits and frontiers of desiccation-tolerant life. *Integrative and Comparative Biology* 45: 685–695.
- Alpert, P., 2006. Constraints of tolerance: why are desiccation-tolerant organisms so small or so rare? *Journal of Experimental Biology* 209: 1575–1584.

- Altiero, T., R. Bertolani & L. Rebecchi, 2010. Hatching phenology and resting eggs in *Paramacrobiotus richtersi* (Eutardigrada, Macrobiotidae). *Journal of Zoology*, London 280: 290–296.
- Altiero, T., R. Guidetti, V. Caselli, M. Cesari & L. Rebecchi, 2011. Ultraviolet radiation tolerance in hydrated and desiccated eutardigrades. *Journal of Zoological Systematics and Evolutionary Research* 49(1): 104–110.
- Altiero, T., R. Guidetti, D. Boschini & L. Rebecchi, 2012. Heat-shock proteins expression in encysted and anhydrobiotic eutardigrades. *Journal of Limnology* 71: 211–215.
- Balompapueng, M. D., A. Hagiwara, Y. Nozaki & K. Hirayama, 1997. Preservation of resting eggs of the euryhaline rotifer *Brachionus plicatilis* O. F. Muller by canning. *Hydrobiologia* 358: 163–166.
- Banton, M. C. & A. Tunnacliffe, 2012. MAPK phosphorylation is implicated in the adaptation to desiccation stress in nematodes. *The Journal of Experimental Biology* 215: 4288–4298.
- Barbosa, E. G. G., A. Crisp, S. E. Broadbent, M. Carrillo, C. Boschetti & A. Tunnacliffe, 2006. A functional difference between native and horizontally acquired genes in bdelloid rotifers. *Gene* 590: 186–191.
- Barraclough, T. G., D. Fontaneto, C. Ricci & E. A. Herniou, 2007. Evidence for inefficient selection against deleterious mutations in Cytochrome Oxidase I of asexual bdelloid rotifers. *Molecular Biology and Evolution* 24(9): 1952–1962.
- Bemm, F., C. L. Weiss, J. Schultz & F. Forster, 2016. Genome of a tardigrade: horizontal gene transfer or bacterial contamination? *Proceedings of the National Academy of Sciences of the United States of America* 113: E3054–E3056.
- Bertolani, R., 2001. Evolution of the reproductive mechanisms in tardigrades. A review. *Zoologischer Anzeiger* 240: 247–252.
- Bertolani, R., R. Guidetti, K. I. Jönsson, T. Altiero, D. Boschini & L. Rebecchi, 2004. Experiences with dormancy in tardigrades. *Journal of Limnology* 63: 16–25.
- Bertolani, R., R. Guidetti, T. Altiero, D. R. Nelson & L. Rebecchi, 2019. Chapter 3. Dormancy in freshwater tardigrades. In Alekseev, V. & B. Pinel-Alloul (eds), *Dormancy in Aquatic Organisms. Theory, Human Use and Modeling.*, Monographiae Biologicae Springer, Cham: 92.
- Boothby, T. C., J. R. Tenlen, F. W. Smith, J. R. Wang, K. A. Patanella, E. O. Nishimura, S. C. Tintori, Q. Li, C. D. Jones, M. Yandell, D. N. Messina, J. Glasscock & B. Goldstein, 2015. Evidence for extensive horizontal gene transfer from the draft genome of a tardigrade. *Proceedings of the National Academy of Sciences of the United States of America* 112(52): 15976–15981.
- Boothby, T. C., H. Tapia, A. L. Brozena, S. Piszkiwicz, A. E. Smith, I. Giovannini, L. Rebecchi, G. J. Pielak, D. Koshland & B. Goldstein, 2017. Tardigrades use intrinsically disordered proteins to survive desiccation. *Molecular Cell* 65: 975–984.
- Boschetti, C., F. Leasi & C. Ricci, 2010. Developmental stages in diapausing eggs: an investigation across monogonont rotifer species. *Hydrobiologia* 662: 149–155.
- Boschetti, C., N. Pouchkina-Stantcheva, P. Hoffmann & A. Tunnacliffe, 2011. Foreign genes and novel hydrophilic protein genes participate in the desiccation response of the bdelloid rotifer *Adineta ricciae*. *Journal of Experimental Biology* 214(1): 59–68.
- Boschetti, C., A. Carr, A. Crisp, I. Eyres, Y. Wang-Koh, E. Lubzens, T. G. Barraclough, G. Micklem & A. Tunnacliffe, 2012. Biochemical diversification through foreign gene expression in bdelloid rotifers. *PLoS Genetics* 8(11): e1003035.
- Boschetti, C., A. Crisp, G. Micklem, M. J. Wise & A. Tunnacliffe, 2013. Anhydrobiosis: the curious case of the bdelloid rotifer. *Cryobiology and Cryotechnology* 59(1): 29–34.
- Boto, L., 2014. Horizontal gene transfer in the acquisition of novel traits by metazoans. *Proceedings of the Biological Society of London B* 281: 20132450.
- Browne, J. A., A. Tunnacliffe & A. Burnell, 2002. Plant desiccation gene found in a nematode. *Nature* 416: 38.
- Browne, J. A., K. M. Dolan, T. Tyson, K. Goyal, A. Tunnacliffe & A. M. Burnell, 2004. Dehydration-specific induction of apheliphilic protein genes in the anhydrobiotic nematode *Aphelenchus avenae*. *Eukaryotic Cell* 3(4): 966–975.
- Cáceres, C. E., 1997. Dormancy in invertebrates. *Invertebrate Biology* 116: 371–383.
- Caprioli, M., A. K. Katholm, G. Melone, H. Ramlov, C. Ricci & N. Santo, 2004. Trehalose in desiccated rotifers: a comparison between a bdelloid and a monogonont species. *Comparative Biochemistry and Physiology, Part A* 139: 527–532.
- Carmona, M. J., M. Serra & M. R. Miracle, 1989. Protein patterns in rotifers: the timing of aging. *Hydrobiologia* 186(187): 325–330.
- Cesari, M., T. Altiero & L. Rebecchi, 2012. Identification of the trehalose-6-phosphate synthase (tps) gene in desiccation tolerant and intolerant tardigrades. *Italian Journal of Zoology* 79: 530–540.
- Chau, D. Y. S., A. R. Dennis, H. Lin & A. Tunnacliffe, 2016. Determination of water content in dehydrated mammalian cells using terahertz pulse imaging: a feasibility study. *Current Pharmaceutical Biotechnology* 17(2): 200–207.
- Clausen, L. K. B., K. N. Andersen, T. L. Hygum, A. Jørgensen & N. Møbjerg, 2014. First record of cysts in the tidal tardigrade *Echiniscoides sigismundi*. *Helgoland Marine Research* 68(4): 531–537.
- Clegg, J. S., 2001. Cryptobiosis – a peculiar state of biological organization. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology* 128(4): 613–624.
- Copley, J., 1999. Indestructible. *New Scientist* 164: 45–46.
- Cornette, R., M. Kanamori, M. Watanabe, Y. Nakahara, O. Gusev, K. Mitsumasu, K. Kadono-Okuda, M. Shimomura, K. Kikawada & T. Okuda, 2010. Identification of anhydrobiosis-related genes from an expressed sequence tag database cryptobiotic midge *Polypedium vanderplanki* (Diptera: Chironomidae). *Journal of Biological Chemistry* 285: 35889–35899.
- Crowe, J. H., 1971. Anhydrobiosis: an unsolved problem. *American Naturalist* 105: 563–573.
- Crowe, J. H. & K. A. C. Madin, 1975. Anhydrobiosis in nematodes: evaporative water loss and survival. *Journal of Experimental Zoology* 193: 323–333.

- Crowe, J. H., L. M. Crowe & D. Chapman, 1984. Preservation of membranes in anhydrobiotic organisms: the role of trehalose. *Science* 223(4637): 701–703.
- Crowe, J. H., F. A. Hoekstra & L. M. Crowe, 1992. Anhydrobiosis. *Annual Review of Physiology* 54: 579–599.
- Crowe, J. H., L. M. Crowe, W. F. Wolkers, A. E. Oliver, X. Ma, J. H. Auh, M. Tang, S. Zhu, J. Norris & F. Tablin, 2005. Stabilization of dry mammalian cells: lessons from nature. *Integrative and Comparative Biology* 45(5): 810–820.
- Czernekova, M. & K. I. Jönsson, 2016. Experimentally induced repeated anhydrobiosis in the eutardigrade *Richtersius coronifer*. *PLoS ONE* 11(11): e0164062.
- Czerneková, M., K. I. Jönsson, L. Chajec, S. Student & I. Poprawa, 2016. The structure of the desiccated *Richtersius coronifer* (Richters, 1903). *Protoplasma* 254: 1367–1377.
- Degma, P., R. Bertolani & R. Guidetti, 2019. Actual checklist of Tardigrada species (2009–2019, 36th Edition: 01-09-2019). [https://doi.org/10.25431/11380\\_1178608](https://doi.org/10.25431/11380_1178608). Accessed 11 Sept 2019.
- Demeure, Y., G. Reversat, S. D. Van Gundy & D. W. Freckman, 1978. The relationship between nematode reserves and their survival to desiccation. *Nematropica* 8: 7–8.
- Denekamp, N. Y., M. A. S. Thorne, M. S. Clark, M. Kube, R. Reinhardt & E. Lubzens, 2009. Discovering genes associated with dormancy in the monogonont rotifer *Brachionus plicatilis*. *BMC Genomics* 10: 108.
- Denekamp, N. Y., R. Reinhardt, M. Kube & E. Lubzens, 2010. Late embryogenesis abundant (LEA) proteins in nondesiccated, encysted, and diapausing embryos of rotifers. *Biology of Reproduction* 82: 714–724.
- Denekamp, N. Y., R. Reinhardt, M. W. Albrecht, M. Drungowski, M. Kube & E. Lubzens, 2011. The expression pattern – of dormancy associated genes in multiple life-history stages in the rotifer *Brachionus plicatilis*. *Hydrobiologia* 662(1): 51–63.
- Drezen, J.-M., J. Gauthier, T. Josse, A. Bézier, E. Herniou & E. Huguet, 2017. Foreign DNA acquisition by invertebrate genomes. *Journal of Invertebrate Pathology* 147: 157–168.
- Dunning Hotopp, J. C., 2011. Horizontal gene transfer between bacteria and animals. *Trends in Genetics* 27: 157–163.
- Ellenby, C., 1968. Desiccation survival in the plant parasitic nematodes, *Heterodera rostochiensis* Wollenweber and *Ditylenchus dipsaci* (Kuhn) Filipjev. *Proceedings of the Royal Society of London, B Biological Sciences* 169: 203–213.
- Erdmann, W. & Ł. Kaczmarek, 2017. Tardigrades in space research – past and future. *Origins of Life and Evolution of Biospheres* 47: 545–553.
- Erdmann, W., B. Idzikowski, W. Kowalski, B. Szymański, J. Z. Kosicki & Ł. Kaczmarek, 2017. Can the tardigrade *Hypsibius dujardini* survive in the absence of the geomagnetic field? *PLoS ONE* 12(9): e0183380.
- Erkut, C. & T. V. Kurzchalia, 2015. The *C. elegans* dauer larva as a paradigm to study metabolic suppression and desiccation tolerance. *Planta* 242(2): 389–396.
- Erkut, C., S. Penkov, H. Khesbak, D. Vorkel, J. M. Verbavatz, K. Fahmy & T. V. Kurzchalia, 2011. Trehalose renders the dauer larva of *Caenorhabditis elegans* resistant to extreme desiccation. *Current Biology* 21: 1331–1336.
- Eyres, I., C. Boschetti, A. Crisp, T. P. Smith, D. Fontaneto, A. Tunnacliffe & T. G. Barraclough, 2015. Horizontal gene transfer in bdelloid rotifers is ancient, ongoing and more frequent in species from desiccating habitats. *BMC Biology* 13(1): 90.
- Faurby, S. & P. H. Barber, 2015. Extreme population subdivision despite high colonization ability: contrasting regional patterns in intertidal tardigrades from the west coast of North America. *Journal of Biogeography* 42: 1006–1017.
- Fielenbach, N. & N. Antebi, 2008. *C. elegans* dauer formation and the molecular basis of plasticity. *Genes & Development* 22: 2149–2165.
- Flot, J. F., B. Hespeels, X. Li, B. Noel, I. Arkhipova, E. G. J. Danchin, A. Hejnol, B. Henrissat, R. Koszul, J.-M. Aury, J.-F. Flot & B. Hespeels, 2013. Genomic evidence for ameiotic evolution in the bdelloid rotifer *Adineta vaga*. *Nature* 500: 453–457.
- Fontaneto, D., 2019. Long-distance passive dispersal in microscopic aquatic animals. *Movement Ecology* 7: 10.
- Fontaneto, D. & R. Ambrosini, 2010. Spatial niche partitioning in epibiont rotifers on the waterlouse *Asellus aquaticus*. *Limnology and Oceanography* 55: 1327–1337.
- Fontaneto, D., G. Melone & A. Cardini, 2004. Shape diversity in the trophy of different species of Rotaria (Rotifera, Bdelloidea): a geometric morphometric study. *Italian Journal of Zoology* 71: 63–72.
- Fontaneto, D., E. A. Herniou, C. Boschetti, M. Caprioli, G. Melone, C. Ricci & T. G. Barraclough, 2007. Independently evolving species in asexual bdelloid rotifers. *PLoS Biology* 5(4): e87.
- Fontaneto, D., T. G. Barraclough, K. Chen, C. Ricci & E. A. Herniou, 2008. Molecular evidence for broad-scale distributions in bdelloid rotifers: everything is not everywhere but most things are very widespread. *Molecular Ecology* 17: 3136–3146.
- Fontaneto, D., M. Kaya, E. A. Herniou & T. G. Barraclough, 2009. Extreme levels of hidden diversity in microscopic animals (Rotifera) revealed by DNA taxonomy. *Molecular Phylogenetics and Evolution* 53: 182–189.
- Fontaneto, D., N. Bunnefeld & M. Westberg, 2012a. Long-term survival of microscopic animals under desiccation is not so long. *Astrobiology* 12(9): 863–869.
- Fontaneto, D., C. Q. Tang, U. Obertegger, F. Leasi & T. G. Barraclough, 2012b. Different diversification rates between sexual and asexual organisms. *Evolutionary Biology* 39: 262–270.
- Förster, F., C. Liang, A. Shkumatov, D. Beisser, J. C. Engelman, M. Schnölzer, M. Frohme, T. Müller, R. O. Schill & T. Dandekar, 2009. Tardigrade workbench: comparing stress-related proteins, sequence-similar and functional protein clusters as well as RNA elements in tardigrades. *BMC Genomics* 10(1): 469.
- Förster, F., D. Beisser, M. Frohme, R. O. Schill & T. Dandekar, 2011. Bioinformatics identifies tardigrade molecular adaptations including the DNA-j family and first steps towards dynamical modelling. *Journal of Zoological Systematics and Evolutionary Research* 49: 120–126.
- França, M. B., A. D. Panek & E. C. A. Eleutherio, 2007. Oxidative stress and its effects during dehydration. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* 146(4): 621–631.

- García de Castro, A. & A. Tunnacliffe, 2000. Intracellular trehalose improves osmotolerance but not desiccation tolerance in mammalian cells. *FEBS Letters* 487: 199–202.
- García-Roger, E. M., M. J. Carmona & M. Serra, 2006. Patterns in rotifer diapausing egg banks: density and viability. *Journal of Experimental Marine Biology and Ecology* 336: 198–210.
- Giovannini, I. T., R. Guidetti Altiero & L. Rebecchi, 2018. Will the Antarctic tardigrade *Acutuncus antarcticus* be able to withstand environmental stresses related to global climate change? *Journal of Experimental Biology* 221: 1–11.
- Gladyshev, E. & M. Meselson, 2008. Extreme resistance of bdelloid rotifers to ionizing radiation. *Proceedings of the National Academy of Sciences of the United States of America* 105(13): 5139–5144.
- Gladyshev, E. A., M. Meselson & I. R. Arkipova, 2008. Massive horizontal gene transfer in bdelloid rotifers. *Science* 320: 1210–1213.
- Glime, J. M., 2017. Ch 4-6. Adaptive strategies: life cycles. In Glime, J. M. (ed.) *Bryophyte Ecology*. Volume 1. Physiological 4-6-1 Ecology. Last updated 6 March 2017. <http://digitalcommons.mtu.edu/bryophyte-ecology/>.
- Gómez, A., M. Serra, G. R. Carvalho & D. H. Lunt, 2002. Speciation in ancient cryptic species complexes: evidence from the molecular phylogeny of *Brachionus plicatilis* (Rotifera). *Evolution* 56(7): 1431–1444.
- Goyal, K., L. J. Walton & A. Tunnacliffe, 2005. LEA proteins prevent protein aggregation due to water stress. *Biochemical Journal* 388: 151–157.
- Guidetti, R. & K. I. Jönsson, 2002. Long-term anhydrobiotic survival in semi-terrestrial micrometazoans. *Journal of Zoology* 257: 181–187.
- Guidetti, R. & N. Møbjerg, 2019. Environmental adaptations: encystment and cyclomorphosis. In Schill, R. O. (ed.), *Water Bears: The Biology of Tardigrades*, Vol. 9., *Zoological Monographs* Springer, Berlin: 249–271.
- Guidetti, R., D. Boschini, L. Rebecchi & R. Bertolani, 2006. Encystment processes and “Matrioshka-like stage” cyst in a moss-dwelling and in a limnic species of eutardigrades (Tardigrada). *Hydrobiologia* 558: 9–21.
- Guidetti, R., C. Colavita, T. Altiero, R. Bertolani & L. Rebecchi, 2007. Energy allocation in two species of Eutardigrada. *Journal of Limnology* 66(Suppl. 1): 111–118.
- Guidetti, R., D. Boschini, T. Altiero, R. Bertolani & L. Rebecchi, 2008. Diapause in tardigrades: a study of factors involved in encystment. *Journal of Experimental Biology* 211: 2296–2302.
- Guidetti, R., T. Altiero & L. Rebecchi, 2011a. On dormancy strategies in tardigrades. *Journal of Insect Physiology* 57: 567–576.
- Guidetti, R., T. Altiero, R. Bertolani, P. Grazioso & L. Rebecchi, 2011b. Survival of freezing by hydrated tardigrades inhabiting terrestrial and freshwater habitats. *Zoology* 114: 123–128.
- Guidetti, R., A. M. Rizzo, T. Altiero & L. Rebecchi, 2012. What can we learn from the toughest animals of the Earth? Water bears (tardigrades) as multicellular model organisms in order to perform scientific preparations for lunar exploration. *Planetary and Space Science* 74: 97–102.
- Guidetti, R., M. Cesari, R. Bertolani, T. Altiero & L. Rebecchi, 2019. High diversity in species, reproductive modes and distribution within the *Paramacrobiotus richtersi* complex (Eutardigrada, Macrobiotidae). *Zoological Letters* 5: 1.
- Hand, S. C., 1991. Metabolic dormancy in aquatic invertebrates. *Advances in Comparative and Environmental Physiology* 8. Springer, Berlin: 1–50.
- Hand, S. C., D. L. Denlinger, J. E. Podrabsky & R. Roy, 2016. Mechanisms of animal diapause: recent developments from nematodes, crustaceans, insects, and fish. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 310: R1193–R1211.
- Hansen, J. G. & A. K. Katholm, 2002. A study of the genus *Amphibolus* from Disko Island with special attention on the life cycle of *Amphibolus nebulosus* (Eutardigrada: Eohypsibiidae). In Hansen, J. G. (ed.), *Arctic Biology Field Course*, Godhavn, Zoological Museum. University of Copenhagen, H.C.Ø. TRYK, Copenhagen: 129–163.
- Hanson, S. J., C.-P. Stelzer, D. B. Mark Welch & J. M. Logsdon Jr., 2013. Comparative transcriptome analysis of obligately asexual and cyclically sexual rotifers reveals genes with putative functions in sexual reproduction, dormancy, and asexual egg production. *BMC Genomics* 14: 412.
- Hashimoto, T. & T. Kunieda, 2017. DNA protection protein, a novel mechanism of radiation tolerance: lessons from tardigrades. *Life* 7(2): 26.
- Hashimoto, T., D. D. Horikawa, Y. Saito, H. Kuwahara, H. Kozuka-Hata, T. Shin, Y. Minakuchi, K. Ohishi, A. Motoyama, T. Aizu & A. Enomoto, 2016. Extremotolerant tardigrade genome and improved radiotolerance of human cultured cells by tardigrade-unique protein. *Nature Communications* 7: 12808.
- Hengherr, S. & R. O. Schill, 2018. Environmental adaptations: cryobiosis. In Schill, R. O. (ed.), *Water Bears: The Biology of Tardigrades*, Vol. 2., *Ecological Monographs* Springer, Berlin: 295–310.
- Hengherr, S., F. Brümmer & R. O. Schill, 2008a. Anhydrobiosis in tardigrades and its effects on longevity traits. *Journal of Zoology* 275: 216–220.
- Hengherr, S., A. G. Heyer, H.-R. Köhler & R. O. Schill, 2008b. Trehalose and anhydrobiosis in tardigrades – evidence for divergence in responses to dehydration. *FEBS Journal* 275: 281–288.
- Hengherr, S., M. R. Worland, A. Reuner, F. Brümmer & R. O. Schill, 2009. High-temperature tolerance in anhydrobiotic tardigrades is limited by glass transition. *Physiological and Biochemical Zoology* 82(6): 749–755.
- Hespeels, B., M. Knapen, D. Hanot-Mambres, A. C. Heuskin, F. Pineux, S. Lucas, R. Koszul & K. Van Doninck, 2014. Gateway to genetic exchange? DNA double-strand breaks in the bdelloid rotifer *Adineta vaga* submitted to desiccation. *Journal of Evolutionary Biology* 27: 1334–1345.
- Hespeels, B., X. Li, J.-F. Flot, L.-M. Pigneur, J. Malaisse, C. Da Silva & K. Van Doninck, 2015. Against all odds: trehalose-6-phosphate synthase and trehalase genes in the bdelloid rotifer *Adineta vaga* were acquired by horizontal gene transfer and are upregulated during desiccation. *PLoS ONE* 10(7): e0131313.
- Horikawa, D. D., T. Kunieda, W. Abe, M. Watanabe, Y. Nakahara, F. Yukuhiro, T. Sakashita, N. Hamada, S. Wada, T. Funayama, C. Katagiri, Y. Kobayashi, S. Higashi & T. Okuda, 2008. Establishment of a rearing system of the extremotolerant tardigrade *Ramazzottius variegatus*: a



- new model animal for astrobiology. *Astrobiology* 8: 549–556.
- Hughes, S. E., K. Evason, C. Xiong & K. Kornfeld, 2007. Genetic and pharmacological factors that influence reproductive aging in nematodes. *PLoS Genetics* 3(2): e25.
- Irons, J. G., L. K. Miller & M. W. Oswood, 1993. Ecological adaptations of aquatic macroinvertebrates to overwintering in interior Alaska (USA) subarctic streams. *Canadian Journal of Zoology* 71(1): 98–108.
- Ivarsson, H. & K. I. Jönsson, 2004. Aggregation effects on anhydrobiotic survival in the tardigrade *Richtersius coronifer*. *Journal of Experimental Zoology Part A: Comparative Experimental Biology* 301(2): 195–199.
- Jönsson, K. I., 2005. The evolution of life histories in holo-anhydrobiotic animals: a first approach. *Integrative and Comparative Biology* 45(5): 764–770.
- Jonsson, K. I., 2007. Tardigrades as a potential model organisms in space research. *Astrobiology* 7: 757–766.
- Jönsson, K. I. & R. Bertolani, 2001. Facts and fiction about long-term survival in tardigrades. *Journal of Zoology (London)* 255: 121–123.
- Jönsson, K. I. & J. Järemo, 2003. A model on the evolution of cryptobiosis. *Annales Zoologici Fennici* 40: 331–340.
- Jönsson, K. I. & O. Persson, 2010. Trehalose in three species of desiccation tolerant tardigrades. *The Open Zoology Journal* 3: 1–5.
- Jönsson, K. J. & L. Rebecchi, 2002. Experimentally induced anhydrobiosis in the tardigrade *Richtersius coronifer*: phenotypic factors affecting survival. *Journal of Experimental Zoology* 293: 578–584.
- Jönsson, K. I. & R. O. Schill, 2007. Induction of Hsp70 by desiccation, ionizing radiation and heat-shock in the eutardigrade *Richtersius coronifer*. *Comparative Biochemistry and Physiology* 146B: 456–460.
- Jönsson, K. J. & A. Wojcik, 2017. Tolerance to X-rays and heavy ions (Fe, He) in the tardigrade *Richtersius coronifer* and the bdelloid rotifer *Mniobia russeola*. *Astrobiology* 17: 163–167.
- Jönsson, K. I., S. Borsari & L. Rebecchi, 2001. Anhydrobiotic survival in populations of the tardigrades *Richtersius coronifer* and *Ramazzottius oberhaeuseri* from Italy and Sweden. *Zoologischer Anzeiger* 240: 419–423.
- Jönsson, K. I., M. Harms-Ringdahl & J. Torudd, 2005. Radiation tolerance in the eutardigrade *Richtersius coronifer*. *International Journal of Radiation Biology* 81: 649–656.
- Jönsson, K. I., E. Rabbow, R. O. Schill, M. Harms-Ringdahl & P. Rettberg, 2008. Tardigrades survive exposure to space in low Earth orbit. *Current Biology* 18: R729–R731.
- Jönsson, K. I., R. A. Beltrán-Pardo, S. Haghdoost, A. Wojcik, R. M. Bermúdez-Cruz, J. Bernal Villegas & M. Harms-Ringdahl, 2013. Tolerance to gamma-irradiation in eggs of the tardigrade *Richtersius coronifer* depends on stage of development. *Journal of Limnology* 72(s1): 73–79.
- Jørgensen, A., N. Møbjerg & R. M. Kristensen, 2007. A molecular study of the tardigrade *Echiniscus testudo* (Echiniscidae) reveals low DNA sequence diversity over a large geographical area. *Journal of Limnology* 66: 77–83.
- Kaczmarek, Ł., M. Roszkowska, D. Fontaneto, M. Jezierska, B. Pietrzak, R. Wiczorek, I. Poprawa, J. Z. Kosicki, A. Karachitos & H. Kmita, 2019. Staying young and fit? Ontogenetic and phylogenetic consequences of animal anhydrobiosis. *Journal of Zoology (London)*. <https://doi.org/10.1111/jzo.12677>.
- Kamilari, M., A. Jørgensen, M. Schiøtt & N. Møbjerg, 2019. Comparative transcriptomics suggest unique molecular adaptations within tardigrade lineages. *BMC Genomics* 20: 607.
- Kaneko, G., T. Yoshinaga, Y. Yanagawa, S. Kinoshita, K. Tsukamoto & S. Watabe, 2005. Molecular characterization of Mn-superoxide dismutase and gene expression studies in dietary restricted *Brachionus plicatilis* rotifers. *Hydrobiologia* 546: 117–123.
- Keilin, D., 1959. The Leewenhoek lecture. The problem of anabiosis or latent life: history and current concept. *Proceedings of the Royal Society of London B: Biological Sciences* 150: 149–191.
- Koutsovoulos, G., S. Kumar, D. R. Laetsch, L. Stevens, J. Daub, C. Conlon, H. Maroon, F. Thomas, A. A. Aboobaker & M. Blaxter, 2016. No evidence for extensive horizontal gene transfer in the genome of the tardigrade *Hypsibius dujardini*. *Proceedings of the National Academy of Sciences of the United States of America* 113(18): 5053–5058.
- Krisko, A. & M. Radman, 2019. Protein damage, ageing and age-related diseases. *Open Biology* 9: 180249.
- Krisko, A., M. Leroy, M. Radman & M. Meselson, 2012. Extreme anti-oxidant protection against ionizing radiation in bdelloid rotifers. *Proceedings of the National Academy of Sciences of the United States of America* 109(7): 2354–2357.
- Kristensen, R. M., 1982. New aberrant eutardigrades from homothermic springs on Disko Island, West Greenland. In D.R. Nelson (ed.), *Proceedings of the Third International Symposium on Tardigrada*. East Tennessee State University Press, Johnson City, Tennessee: 203–220.
- Lapinski, J. & A. Tunnacliffe, 2003. Anhydrobiosis with trehalose in bdelloid rotifers. *FEBS Letters* 553: 387–390.
- Leandro, L. J., N. J. Szewczyk, A. Benguria, R. Herranz, D. Laván, F. J. Medina, G. Gasset, J. J. W. A. van Loon, C. A. Conley & R. Marco, 2007. Comparative analysis of *Drosophila melanogaster* and *Caenorhabditis elegans* gene expression experiments in the European Soyuz flights to the International Space Station. *Advances in Space Research* 40(4): 506–512.
- Lee, D. L., 2002. *The Biology of Nematodes*. CRC Press, Taylor and Francis group, New York.
- Mack, H. I., T. Heimbucher & C. T. Murphy, 2018. The nematode *Caenorhabditis elegans* as a model for aging research. *Drug Discovery Today: Disease Models* 27: 3–13.
- Madin, K. A. C. & J. H. Crowe, 1975. Anhydrobiosis in nematodes: carbohydrate and lipid metabolism during dehydration. *Journal of Experimental Zoology* 193: 335–342.
- Mali, B., M. A. Grohme, F. Förster, T. Dandekar, M. Schnölzer, D. Reuter, W. Welnicz, R. O. Schill & M. Frohme, 2010. Transcriptome survey of the anhydrobiotic tardigrade *Milnesium tardigradum* in comparison with *Hypsibius dujardini* and *Richtersius coronifer*. *BMC Genomics* 11: 168.
- Mark Welch, D. B. & M. Meselson, 2000. Evidence for the evolution of bdelloid rotifers without sexual reproduction or genetic exchange. *Science* 288: 1211–1215.

- Marotta, R., F. Leasi, A. Uggetti, C. Ricci & G. Melone, 2010. Dry and survive: morphological changes during anhydrobiosis in a bdelloid rotifer. *Journal of Structural Biology* 171(1): 11–17.
- Mattimore, V. & J. R. Battista, 1996. Radioresistance of *Deinococcus radiodurans*: functions necessary to survive ionizing radiation are also necessary to survive prolonged desiccation. *Journal of Bacteriology* 178: 633–637.
- Melone, G., C. Ricci, H. Segers & R. L. Wallace, 1998. Phylogenetic relationships of phylum Rotifera with emphasis on the families of Bdelloidea. *Hydrobiologia* 387(388): 101–107.
- Mills, S., A. Gómez & D. H. Lunt, 2017. Global isolation by distance despite strong regional phylogeography in a small metazoan. *BMC Evolutionary Biology* 7: 22.
- Møbjerg, N., K. A. Halberg, A. Jørgensen, D. Persson, M. Bjørn, H. Ramløv & R. M. Kristensen, 2011. Survival in extreme environments – on the current knowledge of adaptations in tardigrades. *Acta Physiologica* 202: 409–420.
- Mogle, M. J., S. A. Kimball, W. R. Miller & R. D. McKown, 2018. Evidence of avian-mediated long distance dispersal in American tardigrades. *PeerJ* 6: e5035.
- Nelson, D., R. Guidetti & L. Rebecchi, 2015. Phylum Tardigrada. 17. In Thorp, J. H. & D. C. Rogers (eds), *Ecology and General Biology: Thorp and Covich's Freshwater Invertebrates*. Academic Press, New York: 347–380.
- Nelson, D. R., P. J. Bartels & N. Guil, 2018. Ch 7. Tardigrade ecology. In Schill, R. O. (ed.), *Water Bears: The Biology of Tardigrades*, Vol. 2., Ecological Monographs Springer, Berlin: 163–210.
- Neumann, S., A. Reuner, F. Brummer & R. O. Schill, 2009. DNA damage in storage cells of anhydrobiotic tardigrades. *Comparative Biochemistry and Physiology – Part A: Molecular & Integrative Physiology* 153: 425–429.
- Nkem, J. N., D. H. Wall, R. A. Virginia, J. E. Barrett, E. J. Broos, D. L. Porazinska & B. J. Adams, 2006. Wind dispersal of soil invertebrates in the McMurdo Dry Valleys, Antarctica. *Polar Biology* 29: 346–352.
- Nowell, R. W., P. Almeida, C. G. Wilson, T. P. Smith, D. Fontaneto, A. Crisp, G. Micklem, A. Tunnacliffe, C. Boschetti & T. G. Barraclough, 2018. Comparative genomics of bdelloid rotifers: insights from desiccating and nondesiccating species. *PLoS Biology* 16(4): e2004830.
- Örstan, A., 1995. Desiccation survival of the eggs of the rotifer *Adineta vaga* (Davis, 1873). *Hydrobiologia* 313(314): 373–375.
- Örstan, A., 1998. Factors affecting long-term survival of dry bdelloid rotifers: a preliminary study. *Hydrobiologia* 387(388): 327–331.
- Persson, D., K. A. Halberg, A. Jørgensen, C. Ricci, N. Møbjerg & R. M. Kristensen, 2011. Extreme stress tolerance in tardigrades: surviving space conditions in low earth orbit. *Journal of Zoological Systematics and Evolutionary Research* 49(Suppl. 1): 90–97.
- Pilato, G., 1979. Correlations between cryptobiosis and other biological characteristics in some soil animals. *Bollettino di Zoologia* 46: 319–332.
- Pouchkina-Stantcheva, N. N., B. M. McGee, C. Boschetti, D. Tolleter, S. Chakrabortee, A. V. Popova, F. Meersman, D. Macherel, D. K. Hinch & A. Tunnacliffe, 2007. Functional divergence of former alleles in an ancient asexual invertebrate. *Science* 318(5848): 268–271.
- Ratnakumar, S. & A. Tunnacliffe, 2006. Intracellular trehalose is neither necessary nor sufficient for desiccation tolerance in yeast. *FEMS Yeast Research* 6(6): 902–913.
- Rebecchi, L., 2013. Dry up and survive: the role of antioxidant defences in anhydrobiotic organisms. *Journal of Limnology* 72: 62–72.
- Rebecchi, L. & R. Bertolani, 1994. Maturative pattern of ovary and testis in eutardigrades of freshwater and terrestrial habitats. *Invertebrate Reproduction and Development* 26: 107–118.
- Rebecchi, L., R. Guidetti, S. Borsari, T. Altiero & R. Bertolani, 2006. Dynamics of long-term anhydrobiotic survival of lichen-dwelling tardigrades. *Hydrobiologia* 558: 23–30.
- Rebecchi, L., T. Altiero & R. Guidetti, 2007. Anhydrobiosis: the extreme limit of desiccation tolerance. *Invertebrate Survival Journal* 4: 65–81.
- Rebecchi, L., M. Cesari, T. Altiero, A. Frigieri & R. Guidetti, 2009a. Survival and DNA degradation in anhydrobiotic tardigrades. *Journal of Experimental Biology* 212(24): 4033–4039.
- Rebecchi, L., T. Altiero, R. Guidetti, M. Cesari, R. Bertolani, M. Negroni & A. M. Rizzo, 2009b. Tardigrade resistance to space effects: first results of the experiment of LIFE – TARSE mission on FOTON-M3 (September 2007). *Astrobiology* 9: 581–591.
- Rebecchi, L., T. Altiero, M. Cesari, R. Bertolani, A. M. Rizzo, P. Corsetto & R. Guidetti, 2011. Resistance of the anhydrobiotic eutardigrade *Paramacrobiotus richtersi* to space flight (LIFE–TARSE mission on FOTON-M3). *Journal of Zoological Systematics and Evolutionary Research* 49(Suppl. 1): 98–103.
- Ricci, C., 1987. Ecology of bdelloids: how to be successful. *Hydrobiologia* 147: 117–127.
- Ricci, C., 1998. Anhydrobiotic capabilities of bdelloid rotifers. *Hydrobiologia* 387: 321–326.
- Ricci, C., 2001. Dormancy patterns of rotifers. *Hydrobiologia* 446(447): 1–11.
- Ricci, C. & C. Boschetti, 2003. Bdelloid rotifers as model system to study developmental biology in space. *Advances in Space Biology and Medicine* 9: 25–39.
- Ricci, C. & M. Caprioli, 2001. Recipes for successful anhydrobiosis in bdelloid rotifers. *Hydrobiologia* 446(447): 13–17.
- Ricci, C. & M. Caprioli, 2005. Anhydrobiosis in bdelloid species, populations and individuals. *Integrative and Comparative Biology* 45(5): 759–763.
- Ricci, C. & C. Covino, 2005. Anhydrobiosis of *Adineta ricciae*: costs and benefits. *Hydrobiologia* 546: 307–314.
- Ricci, C. & G. Melone, 2000. Key to the identification of the genera of bdelloid rotifers. *Hydrobiologia* 418: 73–80.
- Ricci, C. & M. Pagani, 1997. Desiccation of *Panagrolaimus rigidus* (Nematoda): survival, reproduction and the influence on the internal clock. *Hydrobiologia* 347(1–3): 1–13.
- Ricci, C. & F. Perletti, 2006. Starve and survive: stress tolerance and life-history traits of a bdelloid rotifer. *Functional Ecology* 20: 340–346.
- Ricci, C., G. Melone, N. Santo & M. Caprioli, 2003. Morphological response of a bdelloid rotifer to desiccation. *Journal of Morphology* 257(2): 246–253.

- Ricci, C., M. Caprioli, C. Boschetti & N. Santo, 2005. *Macrotrachela quadricornifera* featured in a space experiment. *Hydrobiologia* 534(1–3): 239–244.
- Ricci, C., M. Caprioli & D. Fontaneto, 2007. Stress and fitness in parthenogens: is dormancy a key feature for bdelloid rotifers? *BMC Evolutionary Biology* 7(2): S9.
- Ricci, C. & D. Fontaneto, 2009. The importance of being a bdelloid: ecological and evolutionary consequences of dormancy. *Italian Journal of Zoology* 76(3): 240–249.
- Rivas Jr., J. A., T. Schröder, T. E. Gill, R. L. Wallace & E. J. Walsh, 2019. Anemochory of diapausing stages of microinvertebrates in North American drylands. *Freshwater Biology* 64(7): 1303–1314.
- Rizzo, A. M., M. Negroni, T. Altiero, G. Montorfano, P. Corsetto, P. Berselli, B. Berra, R. Guidetti & L. Rebecchi, 2010. Antioxidant defences in hydrated and desiccated states of the tardigrade *Paramacrobiotus richtersi*. *Comparative Biochemistry and Physiology, Part B* 156: 115–121.
- Rozema, E., S. Kierszniowska, O. Almog-Gabai, E. G. Wilson, Y. H. Choi, R. Verpoorte, R. Hamo, V. Chalifa-Caspi, Y. G. Assaraf & E. Lubzens, 2019. Metabolomics reveals novel insight on dormancy of aquatic invertebrate encysted embryos. *Scientific Reports* 9: 8878.
- Rundle, S. D., A. L. Robertson & J. M. Schmid-Araya (eds), 2002. *Freshwater Meiofauna: Biology and Ecology*. Backhuys Publishers, Leiden, The Netherlands.
- Saragusty, J. & P. Loi, 2019. Exploring dry storage as an alternative biobanking strategy inspired by nature. *Theriogenology* 126: 17–27.
- Schaffitzel, E. & M. Hertweck, 2006. Recent aging research in *Caenorhabditis elegans*. *Experimental Gerontology* 41(6): 557–563.
- Schill, R. O. & S. Hengherr, 2018. Ch 10. Environmental adaptations: desiccation tolerance. In Schill, R. O. (ed.), *Water Bears: The Biology of Tardigrades*, Vol. 2., *Ecological Monographs* Springer, Berlin: 273–293.
- Schill, R. O., G. H. B. Steinbrück & H.-R. Köhler, 2004. Stress gene (hsp70) sequences and quantitative expression in *Milnesium tardigradum* (Tardigrada) during active and cryptobiotic stages. *The Journal of Experimental Biology* 207: 1607–1613.
- Schill, R. O., B. Mali, T. Dandekar, M. Schnölzer, D. Reuter & M. Frohme, 2009. Molecular mechanisms of tolerance in tardigrades: new perspectives for preservation and stabilization of biological material. *Biotechnology Advances* 27(4): 348–352.
- Schokraie, E., A. Hotz-Wagenblatt, U. Warnken, B. Mali, M. Frohme, F. Förster, T. Dandekar, S. Hengherr, R. O. Schill & M. Schnölzer, 2010. Proteomic analysis of tardigrades: towards a better understanding of molecular mechanisms by anhydrobiotic organisms. *PLoS ONE* 5(3): e9502.
- Schröder, T., 2005. Diapause in monogonont rotifers. *Hydrobiologia* 546: 291–306.
- Segers, H., 2007. Annotated checklist of the rotifers (Phylum Rotifera) with notes on nomenclature, taxonomy and distribution. *Zootaxa* 1564: 1–104.
- Selch, F., A. Higashibata, M. Imamizo-Sato, A. Higashitani, N. Ishioka, N. J. Szewczyk & C. A. Conley, 2008. Genomic response of the nematode *Caenorhabditis elegans* to spaceflight. *Advances in Space Research* 41(5): 807–815.
- Shannon, A. J., J. A. Browne, J. Boyd, D. A. Fitzpatrick & A. M. Burnell, 2005. The anhydrobiotic potential and molecular phylogenetics of species and strains of *Panagrolaimus* (Nematoda, Panagrolaimidae). *The Journal of Experimental Biology* 208: 2433–2445.
- Sielaff, M., H. Schmidt, T. H. Struck, D. Rosenkranz, D. B. Mark Welch, T. Hankeln & H. Herlyn, 2016. Phylogeny of Syndermata (syn. Rotifera): mitochondrial gene order verifies epizoic Seisonidea as sister to endoparasitic Acanthocephala within monophyletic Hemirotifera. *Molecular Phylogenetics and Evolution* 96: 79–92.
- Snare, D. J., A. M. Fields, T. W. Snell & J. Kubanek, 2013. Lifespan extension of rotifers by treatment with red algal extracts. *Experimental Gerontology* 48: 1420–1427.
- Snell, T. W., R. K. Johnson, B. Rabeneck, C. Zipperer & S. Teat, 2014. Joint inhibition of TOR and JNK pathways interacts to extend the lifespan of *Brachionus manjavacas* (Rotifera). *Experimental Gerontology* 52: 55–69.
- Snell, T. W., R. K. Johnson, K. E. Gribble & D. B. Mark Welch, 2015. Rotifers as experimental tools for investigating aging. *Invertebrate Reproduction & Development* 59(S1): 5–10.
- Sømme, L., 1996. Anhydrobiosis and cold tolerance in tardigrades. *European Journal of Entomology* 93: 349–357.
- Sommer, S., D. Fontaneto & A. Ozgul, 2019. Demographic processes underlying fitness restoration in bdelloid rotifers emerging from dehydration. *Freshwater Biology* 64: 1295–1302.
- Sommerville, R. I. & K. G. Davey, 2002. Diapause in parasitic nematodes. *Canadian Journal of Zoology* 80: 1817–1840.
- Sørensen, M. V., P. Funch & H. Segers, 2005. On a new *Seison* Grube, 1861, from coastal waters of Kenya, with a reappraisal of the classification of Seisonida (Rotifera). *Zoological Studies* 44: 34–43.
- Steiner, G. & F. E. Albin, 1946. Resuscitation of the nematode *Tylenchus polyhypnus* n. sp. after almost 39 years' dormancy. *Journal of the Washington Academy of Science* 36: 97–99.
- Swanstrom, J., K. Chen, K. Castillo, T. G. Barraclough & D. Fontaneto, 2011. Testing for evidence of inefficient selection in bdelloid rotifers: do sample size and habitat differences matter? *Hydrobiologia* 662: 19–25.
- Szydlowski, L., C. Boschetti, A. Crisp, E. G. G. Barbosa & A. Tunnacliffe, 2015. Multiple horizontally acquired genes from fungal and prokaryotic donors encode cellulolytic enzymes in the bdelloid rotifer *Adineta ricciae*. *Gene* 566: 125–137.
- Szymańska, B., 1995. Encystment in the tardigrade *Dactylobiotus dispar* (Murray, 1907) (Tardigrada: Eutardigrada) Part I: observation of leaving animals and structure of the cyst. *Zoologica Poloniae* 40: 91–102.
- Tanaka, S., J. Tanaka, Y. Miwa, D. D. Horikawa, T. Katayama, K. Arakawa, A. Toyoda, T. Kubo & T. Kunieda, 2015. Novel mitochondria-targeted heat-soluble proteins identified in the anhydrobiotic tardigrade improve osmotic tolerance of human cells. *PLoS ONE* 10(2): e0118272.
- Tapia, H., L. Young, D. Fox, C. R. Bertozzi & D. Koshland, 2015. Increasing intracellular trehalose is sufficient to confer desiccation tolerance to *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences of the United States of America* 112(19): 6122–6127.

- Teramoto, N., N. D. Sachinvala & M. Shibata, 2008. Trehalose and Trehalose-based polymers for environmentally benign, biocompatible and bioactive materials. *Molecules* 13(8): 1773–1816.
- Tomba, P., 2002. Intrinsically unstructured proteins. *Trends in Biochemical Sciences* 27(10): 527–533.
- Tunnacliffe, A. & J. Lapinski, 2003. Resurrecting Leuwenhoek's rotifers: a reappraisal of the role of disaccharides in anhydrobiosis. *Philosophical Transactions of the Royal Society B: Biological Sciences* 358: 1755–1771.
- Tunnacliffe, A. & M. J. Wise, 2007. The continuing conundrum of the LEA proteins. *Naturwissenschaften* 94: 791–812.
- Tunnacliffe, A., J. Lapinski & B. McGee, 2005. A putative LEA protein, but no trehalose, is present in anhydrobiotic bdelloid rotifers. *Hydrobiologia* 546: 315–321.
- Tunnacliffe, A., D. K. Hinch, O. Leprince & D. Macherel, 2010. LEA proteins: versatility of form and function. In Lubzens, E., et al. (eds), *Dormancy and Resistance in Harsh Environments – Topics in Current Genetics*, Vol. 21. Springer, Berlin: 91–108.
- Tyson, T., W. Reardon, J. A. Browne & A. M. Burnell, 2007. Gene induction by desiccation stress in the entomopathogenic nematode *Steinernema carpocapsae* reveals parallels with drought tolerance mechanisms in plants. *International Journal of Parasitology* 37: 763–776.
- Tyson, T., G. O. M. Zamora, S. Wong, M. Skelton, B. Daly, J. T. Jones, E. D. Mulvihill, B. Elsworth, M. Phillips, M. Blaxter & A. M. Burnell, 2012. A molecular analysis of desiccation tolerance mechanisms in the anhydrobiotic nematode *Panagrolaimus superbus* using expressed sequenced tags. *BMC Research Notes* 5(1): 68.
- Vukich, M., P. L. Ganga, D. Cavalieri, D. Rivero, S. Pollastri, S. Mugnai, S. Mancuso, S. Pastorelli, M. Lambreva, A. Antonacci, A. Margonelli, I. Bertalan, U. Johannmeier, M. T. Giardi, G. Rea, M. Pugliese, M. Quarto, V. Roca, A. Zanin, O. Borla, L. Rebecchi, T. Altiero, R. Guidetti, M. Cesari, T. Marchioro, R. Bertolani, E. Pace, A. De Sio, M. Casarosa, L. Tozzetti, S. Branciamore, E. Gallori, M. Scorigella, M. Bruzzi, M. Bucciolini, C. Talamonti, A. Donati & V. Zolesi, 2012. BLOKIS: a model payload for multidisciplinary experiments in microgravity. *Microgravity Science and Technology* 24: 397–409.
- Wallace, R. L. & T. W. Snell, 1991. Rotifera. In Thorpe, J. & A. Covich (eds), *Ecology and Classification of North American Freshwater Invertebrates*. Academic Press, New York: 187–248.
- Walsh, E. J., H. A. Smith & R. L. Wallace, 2014. Rotifers of temporary waters. *International Review of Hydrobiology* 99: 3–19.
- Wang, C., M. A. Grohme, B. Mali, R. O. Schill & M. Frohme, 2014. Towards decrypting cryptobiosis – Analyzing anhydrobiosis in the tardigrade *Milnesium tardigradum* using transcriptome sequencing. *PLoS ONE* 9(3): e92663.
- Watanabe, M., T. Kikawada, A. Fujita, E. Forczek, T. Adati & T. Okuda, 2004. Physiological traits of invertebrates entering cryptobiosis in a post-embryonic stage. *European Journal of Entomology* 101: 439–444.
- Watanabe, M., T. Sakashita, A. Fujita, T. Kikawada, D. D. Horikawa, Y. Nakahara, S. Wada, T. Funayama, N. Hamada, Y. Kobayashi & T. Okuda, 2006. Biological effects of anhydrobiosis in an African chironomid, *Polypedilum vanderplanki* on radiation tolerance. *International Journal of Radiation Biology* 82: 587–592.
- Weinicz, W., M. A. Grohme, L. Kaczmarek, R. O. Schill & M. Frohme, 2011. Anhydrobiosis in tardigrades – the last decade. *Journal of Insect Physiology* 57: 577–583.
- Westh, P. & R. M. Kristensen, 1992. Ice formation in the freeze-tolerant eutardigrades *Adorybiotus coronifer* and *Amphibolus nebulosus* studied by differential scanning calorimetry. *Polar Biology* 12: 693–699.
- Westh, P. & H. Ramløv, 1991. Trehalose accumulation in the tardigrade *Adorybiotus coronifer* during anhydrobiosis. *Journal of Experimental Zoology* 258: 303–311.
- Wharton, D. A., 1996. Water loss and morphological changes during desiccation of the anhydrobiotic nematode *Ditylenchus dipsaci*. *Journal of Experimental Biology* 199(5): 1085–1093.
- Wharton, D. A., 2003. The environmental physiology of Antarctic terrestrial nematodes: a review. *Journal of Comparative Physiology B: Biochemical, Systems, and Environmental Physiology* 173: 621–628.
- Wharton, D. A., G. Goodall & C. J. Marshall, 2003. Freezing survival and cryoprotective dehydration as cold tolerance mechanisms in the Antarctic nematode *Panagrolaimus davidi*. *Journal of Experimental Biology* 206(2): 215–221.
- Wharton, D. A., L. Petrone, A. Duncan & A. J. McQuillan, 2008. A surface lipid may control the permeability slump associated with entry into anhydrobiosis in the plant parasitic nematode *Ditylenchus dipsaci*. *Journal of Experimental Biology* 211(18): 2901–2908.
- Wilhelm, T. & H. Richly, 2018. Autophagy during ageing – from Dr Jekyll to Mr Hyde. *The FEBS Journal* 285(13): 2367–2376.
- Wilson, C. G. & P. W. Sherman, 2013. Spatial and temporal escape from fungal parasitism in natural communities of anciently asexual bdelloid rotifers. *Proceedings of the Royal Society of London B* 280: 20131255.
- Womersley, C., 1987. A re-evaluation of strategies employed by nematode anhydrobiotes in relation to their natural environment. In Veech, J. & D. W. Dickson (eds), *Vistas on Nematology*. Society of Nematologists, Hyattsville, MD: 165–173.
- Wright, J. C., 1988a. Structural correlates of permeability and tun formation in tardigrade cuticle: an image analysis study. *Journal of Ultrastructural and Molecular Structural Research* 101: 23–38.
- Wright, J. C., 1988b. The tardigrade cuticle. I. Fine structure and the distribution of lipids. *Tissue & Cell* 20: 745–758.
- Wright, J. C., 1989a. Desiccation tolerance and water-retentive mechanisms in tardigrades. *Journal of Experimental Biology* 142: 267–292.
- Wright, J. C., 1989b. The tardigrade cuticle II. Evidence for a dehydration-dependent permeability barrier in the intracuticle. *Tissue and Cell* 21: 263–279.
- Wright, J. C., 1989c. The significance of four xeric parameters in the ecology of terrestrial tardigrades. *Journal of Zoology, London* 224: 59–77.
- Wright, J. C., 2001. Cryptobiosis 300 years on from van Leeuwenhoek: what have we learned about tardigrades. *Zoologischer Anzeiger* 240: 563–582.
- Wright, J. C., P. Westh & H. Ramløv, 1992. Cryptobiosis in Tardigrada. *Biological Reviews* 67(1): 1–29.

- Wurdak, E. S., J. J. Gilbert & R. Jagels, 1978. Fine structure of resting eggs of rotifers *Brachionus calyciflorus* and *Asplanchna sieboldi*. *Transactions of the American Microscopy Society* 97: 49–72.
- Yamaguchi, A., S. Tanaka, S. Yamaguchi, H. Kuwahara, C. Takamura, S. Imajoh-Ohmi, D. D. Horikawa, A. Toyoda, T. Katayama, K. Arakawa, A. Fujiyama, T. Kubo & T. Kunieda, 2012. Two heat-soluble protein families abundantly expressed in an anhydrobiotic tardigrade. *PLoS ONE* 7: e44209.
- Yoshida, Y., G. Koutsovoulos, D. R. Laetsch, L. Stevens, S. Kumar, D. D. Horikawa, K. Ishino, S. Komine, T. Kunieda, M. Tomita, M. Blaxter & K. Arakawa, 2017. Comparative genomics of the tardigrades *Hypsibius dujardini* and *Ramazzottius varieornatus*. *PLoS Biology* 15(7): e2002266.
- Ziv, T., V. Chalifa-Caspi, N. Denekamp, I. Plaschkes, S. Kierszniowska, I. Blais, A. Admon & E. Lubzens, 2017. Dormancy in embryos: insight from hydrated encysted embryos of an aquatic invertebrate. *Molecular and Cellular Proteomics* 16(10): 1746–1769.
- Zullini, A., 2018. Cosmopolitanism and endemism in free living nematodes. *Biogeographia – The Journal of Integrative Biogeography* 33: 33–39.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.