

Late-onset Hypogonadism: Bone health

Vincenzo Rochira^{1,2} 

¹Unit of Endocrinology, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy

²Department of Medical Specialties, Azienda Ospedaliero-Universitaria di Modena, Ospedale Civile di Baggiovara, Modena, Italy

Correspondence

Vincenzo Rochira, Unit of Endocrinology, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy, Via Giardini, 1355, 41126 Modena, Italy.
Email: vincenzo.rochira@unimore.it

Abstract

Background: Bone health is underdiagnosed and undermanaged in men. Bone loss occurs in men with hypogonadism and in aging men. Thus, patients with a diagnosis of late-onset hypogonadism (LOH) are at risk of osteoporosis and osteoporotic fractures.

Objectives: To provide an update on research data and clinical implications regarding bone health in men with LOH by reviewing literature articles on this issue.

Materials and Methods: A thorough search of listed publications in PubMed on bone health in older men with hypogonadism was performed, and other articles derived from these publications were further identified.

Results: Late-onset Hypogonadism may be associated with reduced bone mineral density (BMD). In a pathophysiological perspective, the detrimental effects of testosterone (T) deficiency on BMD are partly ascribed to relative estrogen deficiency and both serum T and serum estradiol (E2) need to be above 200 ng/dL and 20 pg/mL to prevent bone loss. The effects of exogenous T on BMD are controversial, but most of the studies confirm that testosterone replacement therapy (TRT) increases BMD and prevents further bone loss in men with hypogonadism. No data are available on TRT and the prevention of fractures.

Discussion and Conclusion: In men with documented LOH, a specific clinical workup should be addressed to the diagnosis of osteoporosis in order to program subsequent follow-up and consider specific bone active therapy. TRT should be started according to guidelines of male hypogonadism while keeping in mind that it may also have positive effects also on bone health in men with LOH.

1 | INTRODUCTION

Bone health in men is a partially neglected health issue as it is less investigated than in the female counterpart.¹⁻³ Even the clinical trials investigating the efficacy of bone active drugs involve a smaller number of men than females, and not all of these drugs, which are available on the market for the treatment of female osteoporosis, have been approved also for men by regulatory agencies.^{3,4} At present, evidence from several studies has unequivocally proved that male osteoporosis (a) is often secondary to other clinical

conditions,^{3,5} (b) occurs later in life compared with women in the majority of cases,³ and (c) osteoporotic fractures are associated with higher morbidity and mortality in elderly men compared with women.^{1,6,7} Among secondary osteoporosis, male hypogonadism is one of the most important risk factors and accounts for the progressive bone loss in aging men, especially in case of a diagnosis of late-onset hypogonadism (LOH).^{1,8} Serum testosterone (T), in fact, decreases with advancing age in elderly men, and the amount of serum estradiol (E2) tends to decrease accordingly⁹; the same occurs for bone loss during aging.¹⁻³ Traditionally, T was considered

the main sex steroid acting on male bone, but starting from the nineties the pivotal role exerted by estrogens started to rise¹⁰⁻¹³ thanks to the description of the first cases of men with congenital estrogen deficiency due to estrogen resistance¹⁴ and aromatase deficiency.^{15,16} The observation that a condition of severe estrogen deficiency was constantly associated with the arrest of skeletal maturation and with severe bone loss resulting in osteopenia or osteoporosis in adult men with these rare diseases^{17,18} opened the way to fully understand the role of estrogens on bone as well as on other male physiological processes.^{12,19-22}

2 | PATHOPHYSIOLOGY OF T DEFICIENCY AND ITS METABOLITES IN BONE

Sex steroids, both androgens and estrogens, exert direct and indirect effects on bone tissue and regulate bone homeostasis.²³⁻²⁵ Estrogens derive from androgens after the aromatization of the A ring of androgens through the activity of the CYP19A1 enzyme, named aromatase, which is expressed in many male tissues.^{11,12,26}

2.1 | Effects of T on bone

T exerts direct and indirect effects on bone.²⁷⁻³⁰ Direct effects of T involve several cells within the bone; among them, human mesenchymal stem cells, osteoblasts, and osteocytes express the androgen receptor and are target cells for T³¹; vice versa, osteoclasts are not target cells for direct action of T and androgens regulate osteoclast proliferation and activity indirectly through the modulation of the receptor activator of nuclear factor κ -B (RANK ligand).²⁷ T and its metabolite dihydrotestosterone (DHT) exert anabolic action on osteocytes and osteoblasts by promoting cell proliferation and probably also their differentiation,³² but the latter effect is less clear.^{27,28} Furthermore, T inhibits apoptosis of osteoblasts.³³ Locally produced androgens (ie, DHT) within the bone contribute together with the circulating quote to support the direct effect of androgens on bone and account for different percentages of sex steroids within the tissue compared with blood concentration.²⁹

Indirect effect of androgens on bone cells may be mediated by several cytokines and by the local production of growth hormone (GH) and insulin-like growth factor 1 (IGF-1) that are known to be under the control of T.²⁷ Other indirect effects on bone are mediated by the mechanical load exerted by the muscle masses surrounding the bone. As muscle mass depends on serum T and hypogonadism leads to sarcopenia, this indirect effect is of great relevance for bone homeostasis.^{1,34,35} Both hypogonadism^{19,36} and aging³⁷ lead to physiological reduction in body muscle masses and sarcopenia in older men, thus promoting bone loss due to mechanical strain³⁸ and postural instability, thus increasing the risk of fracture due to osteoporosis or increased incidence of falls^{38,39} due to altered muscle–bone cross talk.⁴⁰

2.2 | Effects of E2 on bone

Estrogens in men exert their action through the binding to the ERs. Nuclear ER alpha and ER beta and the transmembrane G protein-coupled receptor GPR30 (GPER30) are expressed in human male tissue. The nuclear receptors account for genomic effects of estrogens, while the GPR30 transmembrane receptor accounts for non-genomic, rapid effects of estrogens.^{11,12,26}

In men, locally produced estrogens in bone come from androgens thanks to aromatase that is expressed in fibroblasts and other bone cells (ie, osteoblasts and osteoclasts)^{12,41}; the same occurs in tissues surrounding the bone such as the adipose tissue and the bone marrow.²⁶ Circulating estrogens as well as locally produced estrogens exert their effect on bone through the binding to both ER alpha and ER beta that are expressed in the following bone cells: osteoblasts, osteoclasts, and osteocytes,⁴² all these cell types express also GPER30.^{43,44} Furthermore, estrogens increase osteocytes vitality and inhibit their apoptosis,⁴⁵ induce apoptosis of osteoclasts,⁴⁶ and inhibit their differentiation through the modulation of RANKL.⁴² The final result is the decrease in bone resorption operated by osteoclasts.⁴⁷ On the contrary, estrogens promote/maintain bone formation by osteoblasts through an antiapoptotic effect⁴⁸ and by stimulating their differentiation.⁴⁹ Apart direct effects, estrogens modulate other hormones and cytokines involved in bone physiology (indirect effects), among them the GH/IGF-1 network being one of the most important.⁵⁰ Finally, estrogens positively modulate the bone response to mechanical strain^{51,52} similarly to what androgens^{38,40} do.

2.3 | Late-onset Hypogonadism, relative estrogen deficiency, and osteoporosis in men

In men, serum T progressively declines with aging⁵³ according to the decrease in testicular function, a condition which is common in elderly men.⁵⁴ On average, serum T is below the lower end of the normal range in about 20% of the older men (>60 years)⁵⁵ while remains within the normal range in the majority of them.⁵⁴ There is, however, a great variability in the decline of both total serum T and free serum T, accounting for interindividual differences and diverse trajectories of T decline over time.⁵⁶

The majority of large, prospective studies involving aging men such as the Baltimore Longitudinal Study of Aging (BLSA), the Massachusetts Male Aging Study (MMAS), the Osteoporotic Fractures in Men Study (MrOS), the European Male Aging Study (EMAS), the Rancho Bernardo Study (RBS), the InCHIANTI Study, the Tromsø Study, the Concord Health and Ageing in Men Project (CHAMP), and the Health in Men Study (HIMS) found, on average, a decline of total serum T around 3 ng/dL per year^{57,58} in aging men,⁹ which involves also both calculated serum-free T and serum-free T assayed by equilibrium dialysis.⁹ This finding results in an increase in the prevalence of biochemical hypogonadism with advancing age.^{9,54} The diagnosis of LOH, however, involves only a minority

of cases of older men with low serum T, depending on the severity of serum T deficiency, the presence of symptoms of hypogonadism, and concomitant comorbidities that may lead to functional hypogonadism.^{9,54,59,60}

In men with LOH, the reduced amount of T precursor available for aromatization into E2 may result in relative estrogen deficiency,⁶¹ which may prompt the acceleration of the physiological process of bone loss occurring in aging.^{34,62,63} By an elegant study design, Finkelstein *et al* proved that the pharmacologically induced serum T decrease below 200 ng/dL is constantly associated with a corresponding serum E2 fall below 10 pg/mL and with significant changes in both bone turnover markers and BMD²⁰ (Figure 1). This study proves that relative estrogen deficiency occurs in hypogonadal men and that it is responsible to a great extent for bone loss, whereas T deficiency has a minor direct role.²⁰ Outside the context of experimental design of pharmacologically induced hypogonadism, both serum T and serum E2 decline with advancing age⁶⁴⁻⁶⁶ as demonstrated by both the MrOS and the EMAS. The occurrence of relative estrogen deficiency in aging men with LOH may depend on several factors.⁶¹ First, the severity of hypogonadism obviously influences the amount of circulating serum E2. Second, several associated clinical conditions may modulate aromatase expression and estrogen production.^{61,67-69} As an example, obesity may boost estrogen production even in the presence of low serum T in hypogonadal men through an increased rate of aromatization of androgens, thus preventing both relative estrogen deficiency and bone loss.⁶⁸ Third, individual genetic differences may predispose to relative estrogen deficiency on equal serum T levels.¹ Both polymorphisms of the ER-alpha gene^{70,71} and of the gene encoding for the aromatase enzyme^{72,73} are associated with bone loss in men. Individual genetic differences influencing the expression and activity of the aromatase enzyme have been well-characterized in recent years^{61,68,69,72,73} (Figure 1). By using a genome-wide association study and Mendelian randomization analysis on a large number of men, several genetic variants of the aromatase enzyme resulted directly related to the amount of circulating E2 and BMD as the genetically determined increase of 1 pg/mL of serum E2 corresponded to a BMD increase of 0.048 standard deviation at lumbar spine.⁷⁴ The same experimental

design proved that serum E2 but not T has a causal effect also on bone fractures.⁷⁵ The importance of relative estrogen deficiency as pathophysiological process explaining several changes occurring in hypogonadal men as well as the need of keeping serum T and serum E2 above a certain threshold was strengthened thanks to these well-designed studies (Figure 1). Estrogen-related bone loss in men occurs especially when serum E2 is very low, similarly to what happens to post-menopausal women.⁹ The first evidence came from cases of severe aromatase deficiency where characterized by serum E2 is undetectable leading to bone loss,⁷⁶ and exogenous E2 administration results in a dose-dependent increase in BMD.⁶² The threshold for serum E2 below which the loss of BMD is relevant and the risk of osteoporosis and osteoporotic fractures increases has been estimated to be comprised between 15 and 20 pg/mL^{20,23,34,47,77-83} (Figure 1). In older men with hypogonadism, serum E2 remains higher if compared to the quite undetectable levels in post-menopausal older women (below 5 pg/mL),⁹ but it may fall below 20 pg/mL,⁷⁹ thus accounting for the development of osteopenia and osteoporosis. The threshold for serum T ensuring bone mass preservation seems to be 200 ng/dL^{20,79} (Figure 1).

3 | LOH AND BONE MINERAL DENSITY (BMD)

It is well known that mild-to-severe hypogonadism in young adult men is almost constantly associated with reduced BMD and osteopenia/osteoporosis.^{4,84,85} The effects of T on BMD remain, however, conflicting in elderly men who often display only slight reduction in circulating T that often is due to functional hypogonadism.^{60,86} Several large cohort studies investigating sex steroids in aging men, such as the BLSA, the MMAS, the InCHIANTI, and the HIMS, did not record data about bone health; in particular, BMD and osteoporotic fractures were not investigated.^{9,87} Conversely, other large cohort studies have investigated the relationships between sex steroids and BMD in aging men^{1,2,8,9} showing that low serum T is associated with reduced BMD in older men with overt hypogonadism (total T < 230 ng/dL; ie, 8 nml/L) compared with eugonadal older men




FIGURE 1 Circulating sex steroids and bone pathophysiology in men: relationship between serum T, serum E2, individual factors involved in the conversion of T into E2, putative thresholds for serum T^{20,79} and serum E2,^{20,34,47,80-82} and bone health




FIGURE 2 Information and issues to be covered during medical interview and at each visit

(EMAS)^{35,66,88} and that serum bioavailable T is directly correlated with BMD (RBS and the Tromsø Studies).⁸⁹⁻⁹¹ Only few studies such as the CHAMP did not find any association among serum T, serum E2, SHBG, and bone health.^{92,93}

Studies on the relationships between circulating sex steroids and osteoporotic fractures in aging men are scanty and provide conflicting results⁹ varying from no association (CHAMP and the Tromsø) between circulating sex steroids (total and free serum T, and serum E2) and fracture risk,^{90,93,94} to the association of an increased risk of non-vertebral osteoporotic fractures with both serum T and serum E2 such as in the MrOS.⁹⁵ Thanks to studies performed in the last 20 years using immunometric assays^{1,2,34,96} and those most recent using the gold standard liquid chromatography-tandem mass spectrometry (LC-MS/MS)⁹⁷⁻⁹⁹ for serum E2 measurement the concept that E2 is the main sex steroid involved in the control of BMD maintenance in adult and older men is now textbook knowledge^{21,34,44} (see above the paragraph on pathophysiology for more details). Accordingly, both cross-sectional^{70,89,100-103} and longitudinal studies^{77,104} have disclosed a direct relationship between serum E2, especially the bioavailable fraction, and BMD in men pointing out that relative estrogen deficiency rather than serum T decline per se is involved in bone loss.^{1,34,44} Among them, the most important are the EMAS,⁸⁸ RBS,^{89,105} and the MrOS.⁶⁴ In addition, the latter study showed also a direct role of serum E2 on fracture risk.⁹⁵

In clinical practice, serum T, serum E2, and SHBG should be considered, however weak predictors of vertebral fractures in older men.¹⁰⁶ It should be considered, in fact, that most of these large cohort studies were not specifically addressed to investigate the relationships between sex steroids and bone as their primary endpoint, thus often failed to find an association.⁹ Almost all the studies that had as primary endpoint the investigation of the role of sex steroids on bone in men have clearly demonstrated that low-circulating T is associated with low BMD.^{77,102,107,108}

4 | BONE HEALTH: CLINICAL APPROACH IN MEN WITH LOH

Andrological consultation is a good opportunity to check general health status in men, including bone.⁴

In the presence of a diagnosis of LOH, the investigation of bone health becomes mandatory.^{4,5} The first step is represented by the clinical interview, which should keep information on past medical history with particular concern to general health and physical changes, endocrinological issues, lifestyle, ongoing or previous medications, and additional risk factors for osteoporosis (ie, comorbidities) other than hypogonadism⁴ (Figure 2). Physical examination should check the presence of skeletal features associated with hypogonadism, such as eunuchoid skeletal proportions and the body composition

paying particular attention to the presence of reduced muscle mass (sarcopenia), which is common in hypogonadal men.^{19,40,109} Furthermore, specific features related to severe osteoporosis should be ruled out. Among them, height changes, kyphosis, wall-to-occiput distance (>0 cm), rib-to-pelvis distance (<3 fingers), and tooth loss¹¹⁰ are simple signs that may suggest osteoporosis and possible occult vertebral fractures.^{4,111}

4.1 | Clinical examinations

In order to investigate bone health status in men with LOH, further biochemical, hormonal, and imaging examination may be performed.

4.2 | Sex steroid measurements

Biochemical examinations should be addressed to gonadotropins, T, and SHBG for the diagnosis of LOH, according to available guidelines.^{60,112}

As previously outlined, relative estrogen deficiency is clinically relevant for the development of osteopenia/osteoporosis in older men with LOH,^{1,20,61,68} but the measurement of serum E2 is not recommended in clinical practice as the commercially available assays are not accurate for measuring low values typical of men (20–40 pg/mL),^{4,13,61,113} thus leaving the diagnosis of relative estrogen deficiency in men with hypogonadism still challenging.⁶¹ Having serum E2 measured at baseline may help, theoretically, to better stratify the risk of fracture and the effectiveness of TRT in normalizing not only serum T but also serum E2 (Figure 1), higher BMD being associated with higher serum E2 within the normal range for men.⁶⁸ Hopefully, the widespread diffusion of LC-MS/MS, which is the only accurate and reliable methodology,^{114,115} among clinical laboratories^{116,117} together with the standardization of LC-MS/MS reference ranges for serum sex steroids,¹¹⁷ will probably lead to the validation of this technique for the measurement of serum E2 in men also for clinical purposes in the next future.

At present, the measurement of serum E2 is not recommended in men in the diagnostic workup of osteoporosis,^{4,112} hypogonadism,^{60,112,118,119} and in general in the routine, clinical assessment of the male patient⁶¹ remaining confined to research or clinical settings for which the LC-MS/MS measurement is available or in rare cases in which one of the most accurate immunometric assays is available.¹²⁰ Only in this setting, serum E2 may be useful to rule out/rule in relative estrogen deficiency in a man with LOH and as a reliable target of TRT.^{1,34,61}

4.3 | Biochemical evaluation of bone metabolism

In men with LOH and documented low BMD (osteopenia or osteoporosis), serum calcium, phosphorous, vitamin D, and parathyroid hormone (PTH) should be checked in order to rule out other disorders of calcium/phosphorous metabolism.^{4,121,122}

Measuring markers of bone turnover are not mandatory, and their use should be confined to the monitoring of the effects of bone active therapy even though no clear evidence about their usefulness in the clinic is available.^{4,123}

4.4 | Imaging

The gold standard for obtaining information on bone health in men is dual-energy X-ray absorptiometry (DXA).^{4,111} This simple, safe, and rapid tool measures BMD at different skeletal sites (lumbar spine and femoral neck) and allows the diagnosis of osteopenia or osteoporosis when the t-score is between -1.0 and 2.5 or below -2.5 , respectively.^{124,125} A DXA examination at both lumbar and femoral sites is mandatory for men with a long history of hypogonadism¹ and for older men with LOH and a serum T below 200 ng/dL as in these cases an impairment of BMD is highly probable^{4,5,20,79} (Figure 1). In men with LOH and serum T between 200 and 300 ng/dL, DXA at both lumbar and femoral sites is indicated in the presence of symptoms of hypogonadism or if other risk factors for osteoporosis are present (Figure 2). Pitfalls of DXA measurement should be taken into account in order to minimize errors in the diagnostic interpretation of DXA report. Among them, the most common are the coexistence of osteomalacia, previous fractures resulting in compacted bone tissue, severe scoliosis, vertebral deformities (eg, osteophytes), and inadequate operating procedures (eg, machine calibration, patient's position).¹²⁶

Other techniques for the measurement of BMD such as ultrasound and quantitative computed tomography (QCT) are not validated in clinical practice and are only useful in research settings.^{4,5,111,125,127}

X-ray of the spine is a simple and inexpensive examination, which is mandatory for patients with back pain or with documented signs or symptoms of reduction in height of the spine, such as height or abnormal wall-to-occiput distance^{4,110} (Figure 2). X-ray of the spine allows confirming the diagnosis of severe osteoporosis while it is not useful for the diagnosis of osteopenia or osteoporosis.⁴ Furthermore, X-ray of the spine allows better characterizing the type and severity of fracture according to the Genant classification of vertebral fractures¹²⁸ (<https://www.iofbonehealth.org/radiological-assessment-and-bone-turnover-markers>).

4.5 | Fracture risk in men with LOH

Fracture risk assessment needs a careful collection of information from the male patient with a diagnosis of LOH in order to rule in/rule out additional risk factors that may modify the risk stratification (Figure 2).

All available data may be pooled all together in order to calculate the cumulative risk of fracture by using the Fracture Risk Assessment Tool (FRAX)[®] (<https://www.sheffield.ac.uk/FRAX/>).¹²⁹ At the condition that the patient's interview is

well-conducted and all information is available (Figure 2), FRAX is a simple and effective tool able to predict fracture risk in men both when BMD values are included¹³⁰ or not¹³¹ allowing to identify patients at high risk to develop fractures¹³² and to choose adequate therapeutic options.¹²⁵

BMD measurement by DXA is also useful to establish the risk of fractures in men as low BMD is associated with an increased prevalence of future osteoporotic fractures at all sites in men.^{125,133,134} Besides, change from baseline over time is able to predict future fractures in older men.¹³⁵ Accordingly, a 10-year risk of hip and major osteoporotic fractures obtained by FRAX $\geq 3\%$ $\geq 20\%$, respectively, is a good threshold useful to decide starting bone active therapy in men older than 50 years.¹²⁵

5 | EFFECTS OF T TREATMENT ON BONE IN LOH PATIENTS

During adulthood, bone mass remains the same in men after the achievement of the peak of bone mass and begins to decline only after the age of 50, with a significant bone loss starting around the age of 70.^{144,61,136} In young hypogonadal men, T replacement therapy (TRT) prevents further bone loss and ensures the achievement of a physiological peak of bone mass.¹³⁷ The role of TRT in older men is, however, more controversial. In uncontrolled studies, long-term TRT is able to improve BMD at both lumbar and femoral sites¹³⁸⁻¹⁴¹ or at the femoral site only in other research settings¹⁴² and to decrease markers of bone turnover.¹⁴⁰ In older men with osteopenia or osteoporosis and concomitant documented hypogonadism, TRT leads to a greater increase in spine BMD in the treated group compared with controls.¹⁴³ The effects of TRT seem to be dose-dependent,¹⁴⁰ and there is evidence that even T administration to eugonadal men is able to increase BMD by bringing serum T to levels near the upper end of the normal range.¹⁴⁴ Conversely, several studies did not find any significant effect of TRT on BMD¹⁴⁵⁻¹⁴⁷ and on bone turnover markers.¹⁴⁸

Randomized controlled studies provided conflicting results. TRT is able to significantly decrease bone turnover markers compared with placebo,¹⁴⁷ but no difference¹⁴⁷ or only a small improvement¹⁴⁹ was found in BMD in some studies. On the contrary, other randomized, controlled studies found a significant increase in vertebral and femoral BMD in the TRT group compared with placebo^{150,151} coupled with changes in bone markers.¹⁵⁰

Meta-analyses of clinical studies/trials confirm that TRT is effective in increasing BMD at the lumbar spine^{152,153} and in improving markers of bone turnover.¹⁵² The effect of TRT at the femoral site remains doubtful even though a not significant positive trend has been shown.¹⁵³ The effect of TRT on BMD remains at least doubtful as the patients enrolled in various studies were heterogeneous regarding the presence/absence and the severity of T deficiency. Some studies enrolled patients with low-normal serum T above the lower end of the normal range (300 ng/dL)^{145,146,148} or patients with various degree of T deficiency ranging from severe

(<100 ng/dL)^{138,154,155} to mild hypogonadism (between 200 and 300 ng/dL).^{139-142,146,147,149,150,156} However, the heterogeneity of serum T at baseline among studies and among subjects enrolled in each study together with different study designs (duration, endpoints, methodologies) does not help to reach robust evidence on the effects of TRT on bone and further studies are needed. T trial performed on men older than 65 and documented serum T below 275 ng/dL have found a significant effect of T on spine BMD and hip BMD.¹⁵¹ The efficacy of TRT on BMD increase probably is proportional to the severity of hypogonadism and serum T at baseline and a greater effect on bone should be expected in men with very low serum T at baseline.

Antiestrogens (ie, aromatase inhibitors and estrogen receptor blockers) have been suggested as a possible therapeutic alternative to TRT in men with LOH, but their efficacy has not been proved to be equal to that of TRT, especially in long-term treatment.¹⁵⁷ Furthermore, the number of studies with antiestrogens is limited as well as safety data. For these reasons, antiestrogens remain off-label and are not recommended for the treatment of hypogonadism, including LOH.^{60,157} In men with LOH and concomitant osteopenia or osteoporosis, the use of antiestrogens may unbalance the E/T ratio and cause relative estrogen deficiency and is not considered safe for preventing bone loss and ensure bone health.^{34,61} Accordingly, TRT increases BMD, but the aromatase inhibitor anastrozole did not in a study involving older men with low serum T¹⁵⁸ and there is evidence that aromatase inhibitors can cause bone loss.¹⁵⁷⁻¹⁵⁹

For all these reasons, TRT alone must not be considered as a treatment for osteoporosis in older men with low serum T.^{4,60,160} In clinical practice, TRT should be offered to older men with a documented diagnosis of LOH according to available advice from clinical guidelines.^{60,112} TRT should be started after having informed and discussed with the patients about possible risk and benefits and expected outcomes of TRT.^{59,60,112,119,161} In the clinical context of LOH, TRT may have beneficial effects on bone health too, but an impaired bone status should not lead the decision to start TRT as the latter must be based on the complex of both signs and symptoms (not solely the impaired BMD) of hypogonadism together with the evaluation of potential risks and benefits.^{4,60,84,112} TRT in men with LOH and osteoporosis may help preventing further bone loss and may increase BMD but to a lesser extent than bone active drugs.¹⁶² Of note, TRT has no documented effect on the prevention of fractures and for this reason cannot be considered a therapy for the treatment of osteoporosis. All formulations of exogenous T such as transdermal,^{138,156} oral,¹⁵⁴ and intramuscular are able to act on bone.^{60,148} Men with LOH treated with TRT should expect that TRT may help in preventing further bone loss and increase BMD especially in men with very low serum T at baseline.¹⁴⁶ In order to have beneficial effects on bone, TRT should ensure serum T levels at least above 200 ng/dL^{20,79} and, if accurate assays for estradiol are available, a serum E2 above 20 pg/mL^{1,34,61,68,69} in order to exert a protective role on bone (Figure 1). In this perspective, serum T and serum E2 may be considered reliable targets of TRT for bone health.^{34,61,68}

6 | FOLLOW-UP

Follow-up of men with LOH should also be addressed to bone health in men with no osteopenia or osteoporosis at first or previous visits, and interview focusing on bone health should be repeated at each visit (Figure 2). Furthermore, DXA remains the gold standard to monitor BMD over time and may be repeated every two to three years, depending on patient's health status and risk factors for osteoporosis.

In patients on treatment with active bone drugs, it is important to check about patient's compliance to therapy. Clinical investigation should include regular measurement of calcium, vitamin D, and kidney function. Bone turnover markers may be useful for checking efficacy of bone active therapy even if they are not mandatory.

7 | PHYSICAL ACTIVITY AND PREVENTION OF FURTHER BONE LOSS

Among lifestyle changes that may be beneficial for bone health (Figure 2), physical activity merits to be considered in a comprehensive way in men with LOH for its beneficial effects on muscle mass, skeletal trophism (see above the paragraph on pathophysiology for details), and its direct effects on T secretion. Accordingly, physical activity acts in a synergic fashion together with T by increasing and ensuring muscle mass and both the muscle contraction during physical activity and the muscle mass itself exert an anabolic action on bone through enhanced mechanical load.¹⁶³⁻¹⁶⁵ Besides, physical exercise is able to increase serum T after short time,^{166,167} thus supporting the concept that physical activity and serum T take part to a virtuous circle able to prevent bone loss.¹⁶⁵ In addition, physical activity is effective in preventing falls in older patients.¹⁶⁵ For all these reasons, the prescription of physical activity and exercise should be part of the therapeutic approach to the older man with LOH in order to ensure bone health and prevent fractures.^{4,165,168}

8 | TREATMENT OF OSTEOPENIA/OSTEOPOROSIS IN MEN WITH LOH

After the diagnosis of osteoporosis, a man with concomitant LOH requires a specific therapeutic approach which needs to be tailored according to patient's clinical data. In men with LOH and a diagnosis of osteoporosis, a specific treatment with a bone active drug is indicated, according to guidelines and expert advices for the treatment of male osteoporosis.^{3-5,160} Among them, bisphosphonates (zoledronic acid, alendronate, risedronate, and pamidronate), teriparatide or PTH analogues, and denosumab are all approved by regulatory agencies for the treatment of osteoporosis in men.^{4,160} These drugs should be prescribed in association with calcium plus vitamin D supplementation. In men, the treatment with calcium and vitamin D is indicated also in case of osteopenia when active bone therapy is not

yet necessary.^{4,160,169} Lifestyle changes and behavioral advice aiming to prevent falls are also indicated in men with osteoporosis.³⁻⁵

9 | UNRESOLVED ISSUES

The interpretation of DXA measurement in men suffers from the absence of normative values obtained from male populations. Most of the data used for obtaining reference ranges and for the calculation of t-scores come from female populations; thus, there is no consensus on BMD-based definition of osteoporosis.^{1,170} Future studies are needed in order to consider the risk of fracture based on BMD values by considering also sexual dimorphism of the skeleton^{23,44} and avoiding to transpose to men what we have learnt from women without considering differences between the two sexes in terms of bone pathophysiology.¹⁷¹ To do this, more stringent criteria based on validated male reference ranges for BMD are required in order to appropriately diagnose and manage osteoporosis in older men.

Concerning the role of TRT on the prevention of fractures, there are no available data. However, TRT is effective in ensuring bone anabolism, in preventing sarcopenia, and in improving physical performance and all these effects defend against both frailty and falls, that are known factors involved in the occurrence of fractures, especially the femoral one.⁴⁰

There is a lack of data concerning both the possible use of estrogen treatment in men and in particular in men with hypogonadism.⁶¹ At present, we have indirect information on estrogen effects on bone from rare cases of men with aromatase deficiency showing that physiological doses of exogenous E2 are able to increase BMD in a dose-dependent way.^{62,76} Little is known about the efficacy of selective ER modulators (SERMs) in men and suggests that raloxifene is not so effective as E2 in men with aromatase deficiency.¹⁷²

10 | CONCLUSIONS

The assessment of bone health should be included in the clinical workup of men with LOH bearing in mind that osteoporosis is undermanaged and underdiagnosed in older men and remains less investigated by researchers than in the female counterpart. Bone health, in fact, may be considered a rare case of gender health inequality favoring women rather than men.^{2,3} Normative data obtained by studies on male population are urgently need for both DXA reference values and FRAX risk calculation in order to better tailor the diagnosis and the follow-up of men with osteoporosis. Andrological consultation performed in men with LOH represents an opportunity for male health¹⁷³ including bone.⁴

ACKNOWLEDGMENTS

This study was supported by the Italian Ministry of University and Research by the "Departments of Excellence Programme" granted to the Department of Biomedical, Metabolic and Neural Sciences (University of Modena and Reggio Emilia (Italy).

CONFLICT OF INTEREST

The author has nothing to disclose.

ORCID

Vincenzo Rochira  <https://orcid.org/0000-0001-8169-0696>

REFERENCES

- Rochira V, Balestrieri A, Madeo B, Zirilli L, Granata AR, Carani C. Osteoporosis and male age-related hypogonadism: role of sex steroids on bone (patho)physiology. *Eur J Endocrinol*. 2006;154(2):175-185.
- Madeo B, Zirilli L, Caffagni G, et al. The osteoporotic male: overlooked and undermanaged? *Clin Interv Aging*. 2007;2(3):305-312.
- Adler RA. Update on osteoporosis in men. *Best Pract Res Clin Endocrinol Metab*. 2018;32(5):759-772.
- Rochira V, Antonio L, Vanderschueren D. EAA clinical guideline on management of bone health in the andrological outpatient clinic. *Andrology*. 2018;6(2):272-285.
- Watts NB, Adler RA, Bilezikian JP, et al. Osteoporosis in men: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2012;97(6):1802-1822.
- Cree MW, Juby AG, Carriere KC. Mortality and morbidity associated with osteoporosis drug treatment following hip fracture. *Osteoporosis Int*. 2003;14(9):722-727.
- Sattui SE, Saag KG. Fracture mortality: associations with epidemiology and osteoporosis treatment. *Nat Rev Endocrinol*. 2014;10(10):592-602.
- Rochira V, Balestrieri A, Madeo B, Zirilli L, Granata AR, Carani C. Bone loss, sex steroids and male age related hypogonadism. *J Endocrinol Invest*. 2005;28(suppl 11):46-48.
- Decaroli MC, Rochira V. Aging and sex hormones in males. *Virulence*. 2017;8(5):545-570.
- Rochira V, Granata AR, Madeo B, Zirilli L, Rossi G, Carani C. Estrogens in males: what have we learned in the last 10 years? *Asian J Androl*. 2005;7(1):3-20.
- Rochira V, Madeo B, Diazzi C, et al. Estrogens and male reproduction. In: De Groot LJ, Chrousos G, Dungan K, eds. *Endotext*. South Dartmouth MA: MDText.com, Inc.; 2000.
- Cooke PS, Nanjappa MK, Ko C, Prins GS, Hess RA. Estrogens in Male Physiology. *Physiol Rev*. 2017;97(3):995-1043.
- Santen R, Simpson E. History of estrogen: its purification, structure, synthesis, biologic actions and clinical implications. *Endocrinology*. 2019;160(3):605-625.
- Smith EP, Boyd J, Frank GR, et al. Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. *N Engl J Med*. 1994;331(16):1056-1061.
- Morishima A, Grumbach MM, Simpson ER, Fisher C, Qin K. Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. *J Clin Endocrinol Metab*. 1995;80(12):3689-3698.
- Carani C, Qin K, Simoni M, et al. Effect of testosterone and estradiol in a man with aromatase deficiency. *N Engl J Med*. 1997;337(2):91-95.
- Grumbach MM, Auchus RJ. Estrogen: consequences and implications of human mutations in synthesis and action. *J Clin Endocrinol Metab*. 1999;84(12):4677-4694.
- Faustini-Fustini M, Rochira V, Carani C. Oestrogen deficiency in men: where are we today? *Eur J Endocrinol*. 1999;140(2):111-129.
- Finkelstein JS, Lee H, Burnett-Bowie SA, et al. Gonadal steroids and body composition, strength, and sexual function in men. *N Engl J Med*. 2013;369(11):1011-1022.
- Finkelstein JS, Lee H, Leder BZ, et al. Gonadal steroid-dependent effects on bone turnover and bone mineral density in men. *J Clin Invest*. 2016;126(3):1114-1125.
- Russell N, Grossmann M. Mechanisms in Endocrinology: estradiol as a male hormone. *Eur J Endocrinol*. 2019;181(1):R23-R43.
- Hammes SR, Levin ER. Impact of estrogens in males and androgens in females. *J Clin Invest*. 2019;129(5):1818-1826.
- Laurent M, Antonio L, Sinnesael M, et al. Androgens and estrogens in skeletal sexual dimorphism. *Asian J Androl*. 2014;16(2):213-222.
- Vanderschueren D, Laurent MR, Claessens F, et al. Sex steroid actions in male bone. *Endocr Rev*. 2014;35(6):906-960.
- Farr JN, Khosla S. Skeletal changes through the lifespan—from growth to senescence. *Nat Rev Endocrinol*. 2015;11(9):513-521.
- Simpson ER, Mahendroo MS, Means GD, et al. Aromatase cytochrome P450, the enzyme responsible for estrogen biosynthesis. *Endocr Rev*. 1994;15(3):342-355.
- Vanderschueren D, Vandenput L, Boonen S, Lindberg MK, Bouillon R, Ohlsson C. Androgens and bone. *Endocr Rev*. 2004;25(3):389-425.
- Clarke BL, Khosla S. Androgens and bone. *Steroids*. 2009;74(3):296-305.
- Yarrow JF, Wronski TJ, Borst SE. Testosterone and adult male bone: actions independent of 5 α -reductase and aromatase. *Exerc Sport Sci Rev*. 2015;43(4):222-230.
- Mohamad NV, Soelaiman IN, Chin KY. A concise review of testosterone and bone health. *Clin Interv Aging*. 2016;11:1317-1324.
- Kasperk C, Helmboldt A, Borscok I, et al. Skeletal site-dependent expression of the androgen receptor in human osteoblastic cell populations. *Calcif Tissue Int*. 1997;61(6):464-473.
- Kasperk CH, Wakley GK, Hierl T, Ziegler R. Gonadal and adrenal androgens are potent regulators of human bone cell metabolism in vitro. *J Bone Miner Res*. 1997;12(3):464-471.
- Wiren KM, Toombs AR, Semirale AA, Zhang X. Osteoblast and osteocyte apoptosis associated with androgen action in bone: requirement of increased Bax/Bcl-2 ratio. *Bone*. 2006;38(5):637-651.
- Rochira V, Kara E, Carani C. The endocrine role of estrogens on human male skeleton. *Int J Endocrinol*. 2015;2015:165215.
- Ward KA, Pye SR, Adams JE, et al. Influence of age and sex steroids on bone density and geometry in middle-aged and elderly European men. *Osteoporosis Int*. 2011;22(5):1513-1523.
- McKee A, Morley JE, Matsumoto AM, Vinik A. Sarcopenia: an endocrine disorder? *Endocrine Pract*. 2017;23(9):1143-1152.
- Larsson L, Degens H, Li M, et al. Sarcopenia: aging-related loss of muscle mass and function. *Physiol Rev*. 2019;99(1):427-511.
- Laurent MR, Dedeyne L, Dupont J, Mellaerts B, Dejaeger M, Gielen E. Age-related bone loss and sarcopenia in men. *Maturitas*. 2019;122:51-56.
- Vandenput L, Mellstrom D, Laughlin GA, et al. Low testosterone, but not estradiol, is associated with incident falls in older men: the international MrOS study. *J Bone Miner Res*. 2017;32(6):1174-1181.
- Greco EA, Pietschmann P, Migliaccio S. Osteoporosis and sarcopenia increase frailty syndrome in the elderly. *Front Endocrinol (Lausanne)*. 2019;10:255.
- van der Eerden BC, Lowik CW, Wit JM, Karperien M. Expression of estrogen receptors and enzymes involved in sex steroid metabolism in the rat tibia during sexual maturation. *J Endocrinol*. 2004;180(3):457-467.
- Khalid AB, Krum SA. Estrogen receptors alpha and beta in bone. *Bone*. 2016;87:130-135.
- Smith EP, Specker B, Korach KS. Recent experimental and clinical findings in the skeleton associated with loss of estrogen hormone or estrogen receptor activity. *J Steroid Biochem Mol Biol*. 2010;118(4-5):264-272.
- Almeida M, Laurent MR, Dubois V, et al. Estrogens and androgens in skeletal physiology and pathophysiology. *Physiol Rev*. 2017;97(1):135-187.

45. Tomkinson A, Reeve J, Shaw RW, Noble BS. The death of osteocytes via apoptosis accompanies estrogen withdrawal in human bone. *J Clin Endocrinol Metab.* 1997;82(9):3128-3135.
46. Martin-Millan M, Almeida M, Ambrogini E, et al. The estrogen receptor- α in osteoclasts mediates the protective effects of estrogens on cancellous but not cortical bone. *Mol Endocrinol.* 2010;24(2):323-334.
47. Khosla S, Melton 3rd LJ, Riggs BL. The unitary model for estrogen deficiency and the pathogenesis of osteoporosis: is a revision needed? *J Bone Miner Res.* 2011;26(3):441-451.
48. Pantschenko AG, Zhang W, Nahounou M, et al. Effect of osteoblast-targeted expression of bcl-2 in bone: differential response in male and female mice. *J Bone Miner Res.* 2005;20(8):1414-1429.
49. Zhang W, Pantschenko AG, McCarthy MB, Gronowicz G. Bone-targeted overexpression of Bcl-2 increases osteoblast adhesion and differentiation and inhibits mineralization in vitro. *Calcif Tissue Int.* 2007;80(2):111-122.
50. Birzniece V, McLean M, Reddy N, Ho KKY. Disparate effect of aromatization on the central regulation of GH secretion by estrogens in men and postmenopausal women. *J Clin Endocrinol Metab.* 2019;104(7):2978-2984.
51. Miyazaki T, Zhao Z, Ichihara Y, et al. Mechanical regulation of bone homeostasis through p130Cas-mediated alleviation of NF- κ B activity. *Sci Adv.* 2019;5(9):eaau7802.
52. Swift SN, Swift JM, Bloomfield SA. Mechanical loading increases detection of estrogen receptor- α in osteocytes and osteoblasts despite chronic energy restriction. *J Appl Physiol (1985).* 2014;117(11):1349-1355.
53. Veldhuis JD. Changes in pituitary function with ageing and implications for patient care. *Nat Rev Endocrinol.* 2013;9(4):205-215.
54. Wu FC, Tajar A, Beynon JM, et al. Identification of late-onset hypogonadism in middle-aged and elderly men. *N Engl J Med.* 2010;363(2):123-135.
55. Feldman HA, Longcope C, Derby CA, et al. Age trends in the level of serum testosterone and other hormones in middle-aged men: longitudinal results from the Massachusetts male aging study. *J Clin Endocrinol Metab.* 2002;87(2):589-598.
56. Kaufman JM, Vermeulen A. The decline of androgen levels in elderly men and its clinical and therapeutic implications. *Endocr Rev.* 2005;26(6):833-876.
57. Harman SM, Metter EJ, Tobin JD, Pearson J, Blackman MR. Longitudinal effects of aging on serum total and free testosterone levels in healthy men. Baltimore Longitudinal Study of Aging. *J Clin Endocrinol Metab.* 2001;86(2):724-731.
58. Fabbri E, An Y, Gonzalez-Freire M, et al. Bioavailable testosterone linearly declines over a wide age spectrum in men and women from the Baltimore longitudinal study of aging. *J Gerontol Series A Biol Sci Med Sci.* 2016;71(9):1202-1209.
59. Wang C, Nieschlag E, Swerdloff R, et al. Investigation, treatment, and monitoring of late-onset hypogonadism in males: ISA, ISSAM, EAU, EAA, and ASA recommendations. *J Androl.* 2009;30(1):1-9.
60. Corona G, Goulis DG, Huhtaniemi I, et al. European Academy of Andrology (EAA) guidelines* on investigation, treatment and monitoring of functional hypogonadism in males. *Andrology.* 2020.
61. Rochira V, Carani C. Estrogen Deficiency in Men. In: Simoni M, Huhtaniemi IT, eds. *Endocrinology of the Testis and Male Reproduction.* Cham, Switzerland: Springer International Publishing; 2017:797-828.
62. Rochira V, Faustini-Fustini M, Balestrieri A, Carani C. Estrogen replacement therapy in a man with congenital aromatase deficiency: effects of different doses of transdermal estradiol on bone mineral density and hormonal parameters. *J Clin Endocrinol Metab.* 2000;85(5):1841-1845.
63. Santi D, Madeo B, Carli F, et al. Serum total estradiol, but not testosterone is associated with reduced bone mineral density (BMD) in HIV-infected men: a cross-sectional, observational study. *Osteoporosis Int.* 2016;27(3):1103-1114.
64. Mellstrom D, Johnell O, Ljunggren O, et al. Free testosterone is an independent predictor of BMD and prevalent fractures in elderly men: MrOS Sweden. *J Bone Miner Res.* 2006;21(4):529-535.
65. Camacho EM, Huhtaniemi IT, O'Neill TW, et al. Age-associated changes in hypothalamic-pituitary-testicular function in middle-aged and older men are modified by weight change and lifestyle factors: longitudinal results from the European Male Ageing Study. *Eur J Endocrinol.* 2013;168(3):445-455.
66. Antonio L, Wu Frederick CW, O'Neill Terence W, et al. Low free testosterone is associated with hypogonadal signs and symptoms in men with normal total testosterone. *J Clin Endocrinol Metab.* 2016;101(7):2647-2657. <https://doi.org/10.1210/jc.2015-4106>
67. Rochira V, Zirilli L, Orlando G, et al. Premature decline of serum total testosterone in HIV-infected men in the HAART-era. *PLoS One.* 2011;6(12):e28512.
68. Aguirre LE, Colleluori G, Fowler KE, et al. High aromatase activity in hypogonadal men is associated with higher spine bone mineral density, increased truncal fat and reduced lean mass. *Eur J Endocrinol.* 2015;173(2):167-174.
69. Aguirre LE, Colleluori G, Robbins D, et al. Bone and body composition response to testosterone therapy vary according to polymorphisms in the CYP19A1 gene. *Endocrine.* 2019;65(3):692-706.
70. Ongphiphadhanakul B, Rajatanavin R, Chanprasertyothin S, Piaseu N, Chailurkit L. Serum oestradiol and oestrogen-receptor gene polymorphism are associated with bone mineral density independently of serum testosterone in normal males. *Clinical Endocrinol.* 1998;49(6):803-809.
71. Khosla S, Riggs BL, Atkinson EJ, et al. Relationship of estrogen receptor genotypes to bone mineral density and to rates of bone loss in men. *J Clin Endocrinol Metab.* 2004;89(4):1808-1816.
72. Van Pottelbergh I, Goemaere S, Kaufman JM. Bioavailable estradiol and an aromatase gene polymorphism are determinants of bone mineral density changes in men over 70 years of age. *J Clin Endocrinol Metab.* 2003;88(7):3075-3081.
73. Gennari L, Masi L, Merlotti D, et al. A polymorphic CYP19 TTTA repeat influences aromatase activity and estrogen levels in elderly men: effects on bone metabolism. *J Clin Endocrinol Metab.* 2004;89(6):2803-2810.
74. Eriksson AL, Perry JRB, Coviello AD, et al. Genetic determinants of circulating estrogen levels and evidence of a causal effect of estradiol on bone density in Men. *J Clin Endocrinol Metab.* 2018;103(3):991-1004.
75. Nethander M, Vandenput L, Eriksson AL, Windahl S, Funck-Brentano T, Ohlsson C. Evidence of a causal effect of estradiol on fracture risk in men. *J Clin Endocrinol Metab.* 2019;104(2):433-442.
76. Rochira V, Carani C. Aromatase deficiency in men: a clinical perspective. *Nat Rev Endocrinol.* 2009;5(10):559-568.
77. Khosla S, Melton LJ 3rd, Atkinson EJ, O'Fallon WM. Relationship of serum sex steroid levels to longitudinal changes in bone density in young versus elderly men. *J Clin Endocrinol Metab.* 2001;86(8):3555-3561.
78. Khosla S, Melton 3rd LJ, Robb RA, et al. Relationship of volumetric BMD and structural parameters at different skeletal sites to sex steroid levels in men. *J Bone Miner Res.* 2005;20(5):730-740.
79. Fink HA, Ewing SK, Ensrud KE, et al. Association of testosterone and estradiol deficiency with osteoporosis and rapid bone loss in older men. *J Clin Endocrinol Metab.* 2006;91(10):3908-3915.
80. Lanfranco F, Zirilli L, Baldi M, et al. A novel mutation in the human aromatase gene: insights on the relationship among serum estradiol, longitudinal growth and bone mineral density

- in an adult man under estrogen replacement treatment. *Bone*. 2008;43(3):628-635.
81. Khosla S. Estrogen and bone: insights from estrogen-resistant, aromatase-deficient, and normal men. *Bone*. 2008;43(3):414-417.
 82. Mellstrom D, Vandenput L, Mallmin H, et al. Older men with low serum estradiol and high serum SHBG have an increased risk of fractures. *J Bone Miner Res*. 2008;23(10):1552-1560.
 83. Vandenput L, Ohlsson C. Sex steroid metabolism in the regulation of bone health in men. *J Steroid Biochem Mol Biol*. 2010;121(3-5):582-588.
 84. Antonio L, Caerels S, Jardi F, Delaunay E, Vanderschueren D. Testosterone replacement in congenital hypogonadotropic hypogonadism maintains bone density but has only limited osteoanabolic effects. *Andrology*. 2019;7(3):302-306.
 85. Cangiano B, Swee DS, Quinton R, Bonomi M. Genetics of congenital hypogonadotropic hypogonadism: peculiarities and phenotype of an oligogenic disease. *Hum Genet*. 2020. <http://dx.doi.org/10.1007/s00439-020-02147-1>
 86. Grossmann M, Matsumoto AM. A perspective on middle-aged and older men with functional hypogonadism: focus on holistic management. *J Clin Endocrinol Metab*. 2017;102(3):1067-1075.
 87. Barrett-Connor E, Laughlin GA, Li H, et al. The association of concurrent vitamin D and sex hormone deficiency with bone loss and fracture risk in older men: the osteoporotic fractures in men (MrOS) study. *J Bone Miner Res*. 2012;27(11):2306-2313.
 88. Vanderschueren D, Pye SR, Venken K, et al. Gonadal sex steroid status and bone health in middle-aged and elderly European men. *Osteoporosis Int*. 2010;21(8):1331-1339.
 89. Greendale GA, Edelstein S, Barrett-Connor E. Endogenous sex steroids and bone mineral density in older women and men: the Rancho Bernardo Study. *J Bone Miner Res*. 1997;12(11):1833-1843.
 90. Bjornerem A, Ahmed LA, Joakimsen RM, et al. A prospective study of sex steroids, sex hormone-binding globulin, and non-vertebral fractures in women and men: the Tromso Study. *Eur J Endocrinol*. 2007;157(1):119-125.
 91. Bjornerem A, Emaus N, Berntsen GK, et al. Circulating sex steroids, sex hormone-binding globulin, and longitudinal changes in forearm bone mineral density in postmenopausal women and men: the Tromso study. *Calcif Tissue Int*. 2007;81(2):65-72.
 92. Hsu B, Cumming RG, Seibel MJ, et al. Reproductive hormones and longitudinal change in bone mineral density and incident fracture risk in older men: the concord health and aging in men project. *J Bone Miner Res*. 2015;30(9):1701-1708.
 93. Hsu B, Seibel MJ, Cumming RG, et al. Progressive temporal change in serum shbg, but not in serum testosterone or estradiol, is associated with bone loss and incident fractures in older men: the concord health and ageing in men project. *J Bone Miner Res*. 2016;31(12):2115-2122.
 94. Hsu B, Cumming RG, Blyth FM, et al. The longitudinal relationship of sexual function and androgen status in older men: the Concord Health and Ageing in Men Project. *J Clin Endocrinol Metab*. 2015;100(4):1350-1358.
 95. LeBlanc ES, Nielson CM, Marshall LM, et al. The effects of serum testosterone, estradiol, and sex hormone binding globulin levels on fracture risk in older men. *J Clin Endocrinol Metab*. 2009;94(9):3337-3346.
 96. Rochira V, Balestrieri A, Faustini-Fustini M, Carani C. Role of estrogen on bone in the human male: insights from the natural models of congenital estrogen deficiency. *Mol Cell Endocrinol*. 2001;178(1-2):215-220.
 97. Orwoll E, Blank JB, Barrett-Connor E, et al. Design and baseline characteristics of the osteoporotic fractures in men (MrOS) study—a large observational study of the determinants of fracture in older men. *Contemp Clin Trials*. 2005;26(5):569-585.
 98. Lee DM, O'Neill TW, Pye SR, et al. The European Male Ageing Study (EMAS): design, methods and recruitment. *Int J Androl*. 2009;32(1):11-24.
 99. Lee DM, Pye SR, Tajar A, et al. Cohort profile: the European Male Ageing Study. *Int J Epidemiol*. 2013;42(2):391-401.
 100. Slemenda CW, Longcope C, Zhou L, Hui SL, Peacock M, Johnston CC. Sex steroids and bone mass in older men. Positive associations with serum estrogens and negative associations with androgens. *J Clin Invest*. 1997;100(7):1755-1759.
 101. Center JR, Nguyen TV, Sambrook PN, Eisman JA. Hormonal and biochemical parameters in the determination of osteoporosis in elderly men. *J Clin Endocrinol Metab*. 1999;84(10):3626-3635.
 102. Amin S, Zhang Y, Sawin CT, et al. Association of hypogonadism and estradiol levels with bone mineral density in elderly men from the Framingham study. *Ann Intern Med*. 2000;133(12):951-963.
 103. Szulc P, Munoz F, Claustrat B, et al. Bioavailable estradiol may be an important determinant of osteoporosis in men: the MINOS study. *J Clin Endocrinol Metab*. 2001;86(1):192-199.
 104. Cauley JA, Ewing SK, Taylor BC, et al. Sex steroid hormones in older men: longitudinal associations with 4.5-year change in hip bone mineral density—the osteoporotic fractures in men study. *J Clin Endocrinol Metab*. 2010;95(9):4314-4323.
 105. Laughlin GA, Barrett-Connor E, Bergstrom J. Low serum testosterone and mortality in older men. *J Clin Endocrinol Metab*. 2008;93(1):68-75.
 106. Orwoll ES, Lapidus J, Wang PY, et al. The limited clinical utility of testosterone, estradiol, and sex hormone binding globulin measurements in the prediction of fracture risk and bone loss in older men. *J Bone Miner Res*. 2017;32(3):633-640.
 107. Khosla S, Melton 3rd LJ, Atkinson EJ, O'Fallon WM, Klee GG, Riggs BL. Relationship of serum sex steroid levels and bone turnover markers with bone mineral density in men and women: a key role for bioavailable estrogen. *J Clin Endocrinol Metab*. 1998;83(7):2266-2274.
 108. Falahati-Nini A, Riggs BL, Atkinson EJ, O'Fallon WM, Eastell R, Khosla S. Relative contributions of testosterone and estrogen in regulating bone resorption and formation in normal elderly men. *J Clin Invest*. 2000;106(12):1553-1560.
 109. Nieschlag E. Late-onset hypogonadism: a concept comes of age. *Andrology*. 2019. <https://doi.org/10.1111/andr.12719>
 110. Green AD, Colon-Emeric CS, Bastian L, Drake MT, Lyles KW. Does this woman have osteoporosis? *JAMA*. 2004;292(23):2890-2900.
 111. Lewiecki EM, et al. Osteoporosis: Clinical Evaluation. In: De Groot LJ, Chrousos G, Dungan K, eds. *Endotext*. South Dartmouth MA: MDTText.com, Inc.; 2015.
 112. Bhasin S, Brito JP, Cunningham GR, et al. Testosterone therapy in men with hypogonadism: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab*. 2018;103(5):1715-1744.
 113. Santen RJ, Demers L, Ohorodnik S, et al. Superiority of gas chromatography/tandem mass spectrometry assay (GC/MS/MS) for estradiol for monitoring of aromatase inhibitor therapy. *Steroids*. 2007;72(8):666-671.
 114. Rosner W, Hankinson SE, Sluss PM, Vesper HW, Wierman ME. Challenges to the measurement of estradiol: an endocrine society position statement. *J Clin Endocrinol Metab*. 2013;98(4):1376-1387.
 115. Handelsman DJ, Newman JD, Jimenez M, McLachlan R, Sartorius G, Jones GR. Performance of direct estradiol immunoassays with human male serum samples. *Clin Chem*. 2014;60(3):510-517.
 116. Pagotto U, Fanelli F, Granata AR. Hormonal Laboratory Examination. In: Simoni M, Huhtaniemi IT, eds. *Endocrinology of the Testis and Male Reproduction*. Cham, Switzerland: Springer International Publishing; 2017:495-516.
 117. Fanelli F, Baronio F, Ortolano R, et al. Normative basal values of hormones and proteins of gonadal and adrenal functions from birth to adulthood. *Sex Dev*. 2018;12(1-3):50-94.

118. Dean JD, McMahon CG, Guay AT, et al. The international society for sexual medicine's process of care for the assessment and management of testosterone deficiency in adult men. *J Sex Med.* 2015;12(8):1660-1686.
119. Huhtaniemi I. Late-onset hypogonadism: current concepts and controversies of pathogenesis, diagnosis and treatment. *Asian J Androl.* 2014;16(2):192-202.
120. Ketha H, Girtman A, Singh RJ. Estradiol assays-The path ahead. *Steroids.* 2015;99(Pt A):39-44.
121. Madeo B, De Vincentis S, Kara E, et al. Reliability of calcium-phosphorus (Ca/P) ratio as a new, accurate and inexpensive tool in the diagnosis of some Ca-P disorders. *J Endocrinol Invest.* 2019;42(9):1041-1049.
122. Madeo B, De Vincentis S, Repaci A, et al. The calcium-to-phosphorous (Ca/P) ratio in the diagnosis of primary hyperparathyroidism and hypoparathyroidism: a multicentric study. *Endocrine.* 2020. <http://dx.doi.org/10.1007/s12020-020-02276-7>
123. Jain S, Camacho P. Use of bone turnover markers in the management of osteoporosis. *Curr Opin Endocrinol Diabetes Obes.* 2018;25(6):366-372.
124. Kanis JA, Melton 3rd LJ, Christiansen C, Johnston CC, Khaltaev N. The diagnosis of osteoporosis. *J Bone Miner Res.* 1994;9(8):1137-1141.
125. Cosman F, de Beur SJ, LeBoff MS, et al. Clinician's guide to prevention and treatment of osteoporosis. *Osteoporosis Int.* 2014;25(10):2359-2381.
126. Kanis JA, Delmas P, Burckhardt P, Cooper C, Torgerson D. Guidelines for diagnosis and management of osteoporosis. The European Foundation for Osteoporosis and Bone Disease. *Osteoporosis Int.* 1997;7(4):390-406.
127. Lewiecki EM. Imaging technologies for assessment of skeletal health in men. *Curr Osteoporos Rep.* 2013;11(1):1-10.
128. Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res.* 1993;8(9):1137-1148.
129. Kanis JA, Hans D, Cooper C, et al. Interpretation and use of FRAX in clinical practice. *Osteoporosis Int.* 2011;22(9):2395-2411.
130. Harvey NC, Johansson H, Oden A, et al. FRAX predicts incident falls in elderly men: findings from MrOs Sweden. *Osteoporosis Int.* 2016;27(1):267-274.
131. Ettinger B, Ensrud KE, Blackwell T, Curtis JR, Lapidus JA, Orwoll ES. Performance of FRAX in a cohort of community-dwelling, ambulatory older men: the Osteoporotic Fractures in Men (MrOS) study. *Osteoporosis Int.* 2013;24(4):1185-1193.
132. Kanis JA, Harvey NC, Cooper C, Johansson H, Oden A, McCloskey EV. A systematic review of intervention thresholds based on FRAX: A report prepared for the National Osteoporosis Guideline Group and the International Osteoporosis Foundation. *Arch Osteoporos.* 2016;11(1):25.
133. Cummings SR, Cawthon PM, Ensrud KE, Cauley JA, Fink HA, Orwoll ES. BMD and risk of hip and nonvertebral fractures in older men: a prospective study and comparison with older women. *J Bone Mineral Res.* 2006;21(10):1550-1556.
134. Pasco JA, Lane SE, Brennan SL, et al. Fracture risk among older men: osteopenia and osteoporosis defined using cut-points derived from female versus male reference data. *Osteoporosis Int.* 2014;25(3):857-862.
135. Cawthon PM, Ewing SK, Mackey DC, et al. Change in hip bone mineral density and risk of subsequent fractures in older men. *J Bone Miner Res.* 2012;27(10):2179-2188.
136. Weaver CM, Gordon CM, Janz KF, et al. The National Osteoporosis Foundation's position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. *Osteoporosis Int.* 2016;27(4):1281-1386.
137. Laitinen EM, Hero M, Vaaralahti K, Tommiska J, Raivio T. Bone mineral density, body composition and bone turnover in patients with congenital hypogonadotropic hypogonadism. *Int J Androl.* 2012;35(4):534-540.
138. Behre HM, Kliesch S, Leifke E, Link TM, Nieschlag E. Long-term effect of testosterone therapy on bone mineral density in hypogonadal men. *J Clin Endocrinol Metab.* 1997;82(8):2386-2390.
139. Aversa A, Bruzziches R, Francomano D, et al. Effects of long-acting testosterone undecanoate on bone mineral density in middle-aged men with late-onset hypogonadism and metabolic syndrome: results from a 36 months controlled study. *Aging Male.* 2012;15(2):96-102.
140. Bouloux PM, Legros JJ, Elbers JM, et al. Effects of oral testosterone undecanoate therapy on bone mineral density and body composition in 322 aging men with symptomatic testosterone deficiency: a 1-year, randomized, placebo-controlled, dose-ranging study. *Aging Male.* 2013;16(2):38-47.
141. Rodriguez-Tolra J, Torremade J, di Gregorio S, Del Rio L, Franco E. Effects of testosterone treatment on bone mineral density in men with testosterone deficiency syndrome. *Andrology.* 2013;1(4):570-575.
142. Basurto L, Zarate A, Gomez R, Vargas C, Saucedo R, Galvan R. Effect of testosterone therapy on lumbar spine and hip mineral density in elderly men. *Aging Male.* 2008;11(3):140-145.
143. Shigehara K, Konaka H, Koh E, et al. Effects of testosterone replacement therapy on hypogonadal men with osteopenia or osteoporosis: a subanalysis of a prospective randomized controlled study in Japan (EARTH study). *Aging Male.* 2017;20(3):139-145.
144. Anderson FH, Francis RM, Peaston RT, Wastell HJ. Androgen supplementation in eugonadal men with osteoporosis: effects of six months' treatment on markers of bone formation and resorption. *J Bone Miner Res.* 1997;12(3):472-478.
145. Kenny AM, Prestwood KM, Gruman CA, Marcello KM, Raisz LG. Effects of transdermal testosterone on bone and muscle in older men with low bioavailable testosterone levels. *J Gerontol Series A Biol Sci Med Sci.* 2001;56(5):M266-272.
146. Snyder PJ, Peachey H, Hannoush P, et al. Effect of testosterone treatment on bone mineral density in men over 65 years of age. *J Clin Endocrinol Metab.* 1999;84(6):1966-1972.
147. Merza Z, Blumsohn A, Mah PM, et al. Double-blind placebo-controlled study of testosterone patch therapy on bone turnover in men with borderline hypogonadism. *Int J Androl.* 2006;29(3):381-391.
148. Kenny AM, Prestwood KM, Raisz LG. Short-term effects of intramuscular and transdermal testosterone on bone turnover, prostate symptoms, cholesterol, and hematocrit in men over age 70 with low testosterone levels. *Endocr Res.* 2000;26(2):153-168.
149. Kenny AM, Kleppinger A, Annis K, et al. Effects of transdermal testosterone on bone and muscle in older men with low bioavailable testosterone levels, low bone mass, and physical frailty. *J Am Geriatr Soc.* 2010;58(6):1134-1143.
150. Amory JK, Watts NB, Easley KA, et al. Exogenous testosterone or testosterone with finasteride increases bone mineral density in older men with low serum testosterone. *J Clin Endocrinol Metab.* 2004;89(2):503-510.
151. Snyder PJ, Kopperdahl DL, Stephens-Shields AJ, et al. Effect of testosterone treatment on volumetric bone density and strength in older men with low testosterone: a controlled clinical trial. *JAMA Intern Med.* 2017;177(4):471-479.
152. Isidori AM, Giannetta E, Greco EA, et al. Effects of testosterone on body composition, bone metabolism and serum lipid profile in middle-aged men: a meta-analysis. *Clinical Endocrinol.* 2005;63(3):280-293.
153. Tracz MJ, Sideras K, Bolona ER, et al. Testosterone use in men and its effects on bone health. A systematic review and meta-analysis

- of randomized placebo-controlled trials. *J Clin Endocrinol Metab.* 2006;91(6):2011-2016.
154. Wang C, Eyre DR, Clark R, et al. Sublingual testosterone replacement improves muscle mass and strength, decreases bone resorption, and increases bone formation markers in hypogonadal men—a clinical research center study. *J Clin Endocrinol Metab.* 1996;81(10):3654-3662.
 155. Katznelson L, Finkelstein JS, Schoenfeld DA, Rosenthal DI, Anderson EJ, Klibanski A. Increase in bone density and lean body mass during testosterone administration in men with acquired hypogonadism. *J Clin Endocrinol Metab.* 1996;81(12):4358-4365.
 156. Wang C, Swerdloff RS, Iranmanesh A, et al. Effects of transdermal testosterone gel on bone turnover markers and bone mineral density in hypogonadal men. *Clin Endocrinol.* 2001;54(6):739-750.
 157. Awouters M, Vanderschueren D, Antonio L. Aromatase inhibitors and selective estrogen receptor modulators: Unconventional therapies for functional hypogonadism?. *Andrology* 2020;8: 1590-1597.
 158. Dias JP, Melvin D, Simonsick EM, et al. Effects of aromatase inhibition vs. testosterone in older men with low testosterone: randomized-controlled trial. *Andrology.* 2016;4(1):33-40.
 159. Burnett-Bowie SA, McKay EA, Lee H, Leder BZ. Effects of aromatase inhibition on bone mineral density and bone turnover in older men with low testosterone levels. *J Clin Endocrinol Metab.* 2009;94(12):4785-4792.
 160. Vescini F, Attanasio R, Balestrieri A, et al. Italian association of clinical endocrinologists (AME) position statement: drug therapy of osteoporosis. *J Endocrinol Invest.* 2016;39(7):807-834.
 161. Petak SM, Nankin HR, Spark RF, Swerdloff RS, Rodriguez-Rigau LJ. American Association of Clinical Endocrinologists Medical Guidelines for clinical practice for the evaluation and treatment of hypogonadism in adult male patients—2002 update. *Endocr Pract.* 2002;8(6):440-456.
 162. Welch BJ, Denke MA, Kermani A, Adams-Huet B, Gazmen NM, Gruntmanis U. Comparison of testosterone, alendronate, and a combination of both therapies in men with low bone mineral density. *J Investig Med.* 2007;55(4):168-173.
 163. Kemmler W, von Stengel S, Kohl M. Exercise frequency and fracture risk in older adults—how often is enough? *Curr Osteoporos Rep.* 2017;15(6):564-570.
 164. Herrmann M, Engelke K, Ebert R, et al. Interactions between muscle and bone—where physics meets biology. *Biomolecules.* 2020;10(3):432.
 165. Cauley JA, Giangregorio L. Physical activity and skeletal health in adults. *Lancet Diabetes Endocrinol.* 2020;8(2):150-162.
 166. D'Andrea S, Spaggiari G, Barbonetti A, Santi D. Endogenous transient doping: physical exercise acutely increases testosterone levels—results from a meta-analysis. *J Endocrinol Invest.* 2020. <https://doi.org/10.1007/s40618-020-01251-3>
 167. Cano Sokoloff N, Misra M, Ackerman KE. Exercise, training, and the hypothalamic-pituitary-gonadal axis in men and women. *Front Horm Res.* 2016;47:27-43.
 168. Beck BR, Daly RM, Singh MA, Taaffe DR. Exercise and Sports Science Australia (ESSA) position statement on exercise prescription for the prevention and management of osteoporosis. *J Sci Med Sport.* 2017;20(5):438-445.
 169. Bouillon R, Van Schoor NM, Gielen E, et al. Optimal vitamin D status: a critical analysis on the basis of evidence-based medicine. *J Clin Endocrinol Metab.* 2013;98(8):E1283-1304.
 170. Kaufman JM, Goemaere S. Osteoporosis in men. *Best Pract Res Clin Endocrinol Metab.* 2008;22(5):787-812.
 171. Briot K, Cortet B, Tremollieres F, et al. Male osteoporosis: diagnosis and fracture risk evaluation. *Joint Bone Spine.* 2009;76(2):129-133.
 172. Zirilli L, Maffei L, Meunier PJ, Chavassieux P, Carani C, Rochira V. The effects of long-term raloxifene and estradiol treatments on bone in a patient with congenital aromatase deficiency. *Bone.* 2009;45(5):827-832.
 173. Foresta C, Ferlin A, Lenzi A, Montorsi P. The great opportunity of the andrological patient: cardiovascular and metabolic risk assessment and prevention. *Andrology.* 2017;5(3):408-413.

How to cite this article: Rochira V. Late-onset Hypogonadism: Bone health. *Andrology.* 2020;8:1539–1550. <https://doi.org/10.1111/andr.12827>