

This is the peer reviewed version of the following article:

Effect of ripening and in vitro digestion on the evolution and fate of bioactive peptides in Parmigiano-Reggiano cheese / Martini, Serena; Conte, Angela; Tagliazucchi, Davide. - In: INTERNATIONAL DAIRY JOURNAL. - ISSN 0958-6946. - 105:(2020), pp. 1-9. [10.1016/j.idairyj.2020.104668]

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

19/12/2025 03:02

# **Effect of ripening and *in vitro* digestion on the evolution and fate of bioactive peptides in Parmigiano-Reggiano cheese**

Serena Martini, Angela Conte and Davide Tagliazucchi\*

*Department of Life Sciences, University of Modena and Reggio Emilia, Via Amendola, 2 - Pad. Besta, 42100 Reggio Emilia, Italy*

\* Corresponding author. Tel.: +39-0522-522060; fax: +39-0522-522053

*E-mail address:* [davide.tagliazucchi@unimore.it](mailto:davide.tagliazucchi@unimore.it) (D. Tagliazucchi)

## 1    **ABSTRACT**

2    The influence of ripening (12, 18 and 24 months) and *in vitro* digestion on the peptidomic profile of  
3    Parmigiano-Reggiano (PR) cheeses were investigated. Ripening and *in vitro* digestion thoroughly  
4    modified the peptidomic profile of the three cheeses. Twenty-six bioactive peptides were identified  
5    in undigested PR. Some peptides were degraded and others released during ripening. After  
6    digestion, 52 bioactive peptides were identified. Semi-quantitative data suggested that bioactive  
7    peptides released after digestion can be clustered in 5 groups according to the ripening time. VPP  
8    and IPP peptide levels in undigested samples were in the range of 4.52-11.34 and 0.66-4.24 mg Kg<sup>-1</sup>  
9    <sup>1</sup>, with the highest amounts found in 18-month ripened PR. YPFPGPI peptide was absent in  
10    undigested PRs but was released after digestion, especially in the 12-month-old sample (20.18 mg  
11    Kg<sup>-1</sup>). The present study suggests possible differences in bioactive peptide levels after digestion as a  
12    function of the duration of ripening of PR cheese.

## 13    **1. Introduction**

14    Cheese ripening is characterized by a complex chain of events that entails an intricate set of  
15    biochemical reactions. Among the different biochemical events occurring during cheese ripening,  
16    proteolysis is undeniably one of the most important. Several enzymatic activities originating from  
17    various sources are involved in the proteolysis process during cheese ripening. Curd and  
18    endogenous milk proteolytic enzymes (such as plasmin) initially hydrolyze caseins, generating large  
19    or intermediate-size peptides. The released peptides are further cleaved by the action of proteinases  
20    and peptidases coming from starter (S-LAB) and nonstarter (NS-LAB) lactic acid bacteria (Sforza  
21    et al., 2012).

22    Bioactive peptides can be defined as short amino acid sequences, originally encrypted within the  
23    sequence of the parent protein, which can be released after proteolysis and may have a positive  
24    impact on human health (Rizzello et al., 2016). Numerous bioactive peptides have been identified  
25    and characterized after hydrolysis of different food proteins or in fermented dairy products,  
26    presenting different functional activities including antimicrobial, antioxidative, dipeptidyl  
27    peptidase-IV (DPP-IV) and angiotensin-converting enzyme (ACE) inhibition, antihypertensive,  
28    immunomodulatory and opioid activities (Nongonierma & FitzGerald, 2015; Rizzello et al., 2016).

29    In cheese, the presence of bioactive peptides is the result of a sensitive equilibrium between their  
30    release and their degradation by the activity of lactic acid bacteria proteinases and peptidases during  
31    cheese ripening (Sforza et al., 2004; Sforza et al., 2012). Numerous bioactive peptides, especially  
32    ACE-inhibitory and anti-hypertensive peptides, have been identified in various cheeses (Sieber et  
33    al., 2010; Lu, Govindasamy-Lucey, & Lucey, 2015; Stuknyte, Cattaneo, Masotti, & De Noni, 2015;  
34    Basiricò et al., 2015). Meyer, Bütikofer, Walther, Wechsler, & Sieber (2009) investigated the  
35    changes in concentration of the lactotripeptides VPP and IPP in different Swiss cheese varieties  
36    during the ripening; they found that the concentration of VPP and IPP increased in semi-hard  
37    cheeses according to the ripening time whereas, in hard cheeses, the behavior was dependent on the  
38    cheese varieties. Gómez-Ruiz, Ramos, & Recio (2004) and Ong, & Shah (2008) investigated the

39 release of 5 and 6 ACE-inhibitory peptides during ripening of Manchego and Cheddar cheeses,  
40 respectively. In both cases, the authors did not find a general trend for the release of those peptides  
41 during cheese ripening.

42 Parmigiano-Reggiano is a long ripened, hard cheese made from raw cow milk and whey starter as  
43 sources of fermenting microorganisms (Solieri, Bianchi, & Giudici, 2012). Parmigiano-Reggiano is  
44 characterized by positive nutritional qualities, being an important source of essential nutrients, such  
45 as proteins, fat, vitamins and minerals, and is considered a functional food due to the presence of  
46 different compounds with particular biological activities (Summer et al., 2017; Godos et al., 2019).  
47 Very few studies have investigated the presence of bioactive peptides and their fate during ripening  
48 in Parmigiano-Reggiano. Sforza et al. (2012) gave the most detailed scenario of the evolution of  
49 peptides during Parmigiano-Reggiano ripening. Several bioactive peptides were found to be present  
50 in Parmigiano-Reggiano, such as the antimicrobial peptide isracidin ( $\alpha_{S1}$ -casein fragment 1-23), the  
51 multifunctional bioactive peptides YQEPVLGPVRGPFPIIV ( $\beta$ -casein fragment 193-209) and some  
52 caseinophosphopeptides (Sforza et al., 2012). Basiricò et al. (2015) identified and quantified 4 anti-  
53 hypertensive peptides (VPP, IPP, HLPLP and LHLPLP) in the water-soluble extract of 12-months  
54 ripened Parmigiano-Reggiano. However, a detailed picture of the presence and evolution of such  
55 bioactive peptides during cheese ripening is still lacking.

56 In addition, bioactive peptides might be degraded by gastro-intestinal proteases after ingestion.  
57 Nevertheless, new sequences could be released from inactive or less active precursors after  
58 digestion. Stuknite et al. (2015) identified and quantified 8 ACE-inhibitory peptides in different  
59 types of cheeses, which amounts were variably influenced by *in vitro* gastro-intestinal digestion. In  
60 another study, Sánchez-Rivera et al. (2014) investigated the influence of *in vitro* gastro-intestinal  
61 digestion on the peptidomic profile of Spanish blue cheese. They found that some peptides were  
62 degraded during cheese digestion, whereas some others were newly released by gastro-intestinal  
63 proteases. In this way, at the end of the digestion, a higher number of bioactive peptides were  
64 found. Basiricò et al. (2015) studied the fate of 8 anti-hypertensive peptides during *in vitro* gastro-

65 intestinal digestion of Parmigiano-Reggiano. The concentration of some peptides such as VPP and  
66 IPP was mostly un-affected by the *in vitro* digestion, whereas HLPLP and LHLPLP levels greatly  
67 increased after the digestive process. Some other peptides, such as AYFYPE and AYFYPEL, were  
68 not found in the Parmigiano-Reggiano samples but they were released during *in vitro* digestion.  
69 These results suggest that *in vitro* digestion greatly influences the peptidomic profile of cheese.  
70 The present study was designed to compare the peptidomic profile of Parmigiano-Reggiano cheese  
71 at different times of ripening as well as the influence of *in vitro* gastro-intestinal digestion.  
72 Bioactive peptides were identified, relatively quantified (by integration of the peak area of  
73 individual peptides) and their fate was followed during ripening and *in vitro* digestion. Finally, three  
74 well-known bioactive peptides, namely VPP, IPP and YPFPGPI, were quantified in the different  
75 samples before and after *in vitro* digestion.

76

## 77 **2. Materials and methods**

### 78 *2.1. Materials*

79 All MS/MS reagents were from Bio-Rad (Hercules, CA, U.S.A.), whereas the chemicals and  
80 enzymes for the digestion procedure and hydrolysis degree determination were purchased from  
81 Sigma-Aldrich (Milan, Italy). Amicon Ultra-4 regenerated cellulose filters with a molecular weight  
82 cut-off of 3 kDa were supplied by Millipore (Milan, Italy). Parmigiano-Reggiano cheese samples at  
83 12 (PR12), 18 (PR18) and 24 (PR24) months of ripening were withdrawn from the same cheese  
84 factory in the province of Reggio Emilia (Italy). Three different Parmigiano-Reggiano cheese  
85 samples for each time-point were analysed; these samples were collected the same day from  
86 different batches. Cheese production and ripening were carried out according to the Production  
87 Specification regulated by the Safeguarding Consortium. The analysed cheese sample were  
88 registered as a Protected Designation of Origin (PDO) cheese. A detailed description of the  
89 Parmigiano-Reggiano cheese production and maturation according to the PDO can be found in

90 Sforza et al. (2012). VPP, IPP and YPFPGPI (95% purity) were synthesized by Bio-Fab research  
91 (Rome, Italy). All the other reagents were from Carlo Erba (Milan, Italy).

92

## 93 *2.2. Preparation of water-soluble peptides extract (WSPE) from Parmigiano-Reggiano samples*

94 Water-soluble peptides extracts were obtained as described by Sforza et al. (2012) with slight  
95 modifications. Five grams of cheese samples were mixed with 45 mL of 0.1 mmol L<sup>-1</sup> HCl and  
96 homogenized for 1 min (3 cycles) using an Ultra-Turrax homogenizer. The samples were then  
97 centrifuged at 4000g for 40 min at 4°C. At the end of the centrifugation, the supernatants were  
98 collected and filtered through Whatman filters paper 4 (Maidstone, Kent, UK).

99

## 100 *2.3. In vitro gastro-intestinal digestion of Parmigiano-Reggiano samples using the harmonized* 101 *protocol*

102 The *in vitro* digestion of Parmigiano-Reggiano samples was carried out by following the protocol  
103 previously developed within the COST Action INFOGEST (Minekus et al., 2014). Simulated  
104 salivary, gastric and intestinal fluids (SSF, SGF and SIF) were prepared exactly as described by  
105 Minekus et al. (2014). Cheese samples (5 g) were mixed with 5 mL of SSF (containing 150 U mL<sup>-1</sup>  
106 of salivary  $\alpha$ -amylase), ground and incubated for 5 min at 37°C to reproduce mastication. The bolus  
107 was then mixed with 10 mL of SGF (containing 4000 U mL<sup>-1</sup> of porcine pepsin) and the pH  
108 adjusted to 2.0 with 6 mol L<sup>-1</sup> of HCl. After 2 h of incubation at 37°C, the final intestinal step was  
109 carried out by adding 20 mL of SIF (containing pancreatin 200 U mL<sup>-1</sup> based on trypsin activity).  
110 Then, the pH was adjusted to 7.0 and the samples were further incubated at 37°C for 2 h. All  
111 samples were immediately cooled on ice, centrifuged at 10000 g for 20 min at 4°C and frozen at -  
112 80°C for further analysis. The digestions were performed in triplicate. In addition, a control  
113 digestion, which included only the gastro-intestinal juices and enzymes, and water in place of  
114 cheese, was carried out to consider the possible impact of the digestive enzymes in the subsequent

115 analysis. For each digestion, aliquots were taken at the end of the gastric and intestinal phases of  
116 digestion.

117

#### 118 *2.4. Preparation of the peptide fractions and determination of peptides concentration*

119 Low molecular weight peptides from WSPE and digested samples were extracted by ultrafiltration  
120 (cut-off 3 kDa) as described in Tagliazucchi et al. (2017). The peptide content in these peptide  
121 fractions was determined by measuring the amount of released amino groups using the 2,4,6-  
122 trinitrobenzenesulfonic acid (TNBS) assay and leucine as standard (Adler-Nissen, 1979). The  
123 obtained raw data from the digested samples were corrected by the contribution of the control  
124 digestion. Data are expressed as mmol leucine equivalent g<sup>-1</sup> of cheese.

125

#### 126 *2.5. Identification of low molecular weight peptides by ultra high performance liquid* 127 *chromatography/high resolution mass spectrometry (UHPLC/HR-MS)*

128 The peptide fractions from WSPE and digested samples were subjected to UHPLC/HR-MS analysis  
129 for peptide identification. UHPLC/MS and tandem MS experiments were carried out on an UHPLC  
130 Ultimate 3000 separation module interfaced with a Q Exactive Hybrid Quadrupole-Orbitrap Mass  
131 Spectrometer (Thermo Scientific, San Jose, CA, USA) using a C18 column (Zorbax SB-C18  
132 reversed-phase, 2.1 × 50 mm, 1.8 µm particle size, Agilent Technologies, Santa Clara, CA, USA).  
133 The mobile phase consisted of (A) H<sub>2</sub>O/formic acid (99.9:0.1, v/v) and (B) acetonitrile. The sample  
134 (10 µL, 100-fold diluted) was loaded into the column at a flow rate of 0.3 mL/min. The gradient  
135 started at 2% B, and grew to 3% B in 2 min. The mobile phase composition was raised to 27% B in  
136 19 min and then to 90% in 4 min. The mass spectrometer was set as follow: spray voltage 3.5 kV,  
137 capillary temperature 320°C, sheath gas 40 and auxiliary gas 30. Full MS parameters were:  
138 resolution 70000, AGC target 3e6, maximum IT 333 ms and scan range 200 to 2000 *m/z*. MS/MS  
139 parameters were: resolution 17500, AGC target 1e5, maximum IT 120 ms and isolation window 3  
140 *m/z*.



141 The MS/MS spectra were then converted to .mgf files and the peptides were identified by using the  
142 Swiss-Prot database through MASCOT (Matrix Science, Boston, MA, USA) protein identification  
143 software. The following parameters were considered: enzyme, none; peptide mass tolerance,  $\pm 5$   
144 ppm; fragment mass tolerance,  $\pm 0.12$  Da; variable modification, oxidation (M) and  
145 phosphorylation (ST); maximal number of post-translational modifications permitted in a single  
146 peptide, 4. The assignment process was validated by the manual inspection of MS/MS spectra.

147

## 148 *2.6. Identification of bioactive peptides*

149 Peptides identified in the peptide fractions from WSPE and digested samples were investigated in  
150 relation to bioactive peptides previously identified in the literature using the Milk Bioactive  
151 Peptides Database (MBPDB) (Nielsen, Beverly, Qu, & Dallas, 2017). Only peptides with 100%  
152 homology to acknowledged functional peptides were considered as bioactive peptides. The relative  
153 amount of the bioactive peptides was estimated by integrating the area under the peak (AUP). AUP  
154 was measured from the extracted ion chromatograms (EIC) obtained for each peptide (tolerance  $\pm 5$   
155 ppm). Data are expressed as AUP g<sup>-1</sup> of cheese.

156

## 157 *2.7. Quantification of VPP, IPP and YPFPGPI by parallel reaction monitoring (PRM)*

158 Synthetic peptides were dissolved in solvent A (H<sub>2</sub>O/formic acid; 99.9:0.1, v/v) at a concentration  
159 of 5 mg mL<sup>-1</sup>. The selected analytes were quantified by standard addition method spiking known  
160 amounts of standard solutions directly to the analyzed samples. For each sample, linear range for  
161 VPP and IPP standards were generated by using 0, 4, 8, 16 and 32  $\mu\text{g L}^{-1}$  of standard (final  
162 concentrations in the samples). For YPFPGPI, the linear range was obtained at 0, 5, 15, 30, 50  $\mu\text{g L}^{-1}$   
163 of standard (final concentrations in the samples).

164 The samples (10  $\mu\text{L}$ ; 100-fold diluted) were then injected in the same UHPLC/HR-MS instrument  
165 as describe above. Each sample was analyzed two times. Mobile phase A was 0.1% formic acid in  
166 water and mobile phase B was acetonitrile. The elution gradient started with 2% B, was maintained

for 2 min, and then increased to 15% B between 2 and 6 min. The mobile phase composition was then increased to 27% B in 15 min and further raised to 90% B in 4 min. The flow rate was set at 0.4 mL min<sup>-1</sup>. Ion source parameters was as follow: spray voltage 4 kV, capillary temperature 320 °C, sheath gas 50 and auxiliary gas 25. PRM parameters were as follow: resolution 17500, AGC target 5e5, max IT 150 ms, MSX count 1 and isolation window 3.0 m/z. The precursor ions selected for VPP, IPP and YPFPGPI were [M + H]<sup>+</sup> *m/z* 312.1918, 326.2074 and 790.4134, respectively. The product ion *y*<sub>2</sub><sup>+</sup> at *m/z* 213.1234 was selected for quantitation of VPP and IPP. The product ion *y*<sub>2</sub><sup>+</sup> at *m/z* 229.1547 was selected for quantitation of YPFPGPI. Peaks were integrated by using the Genesis algorithm function in the Thermo Xcalibur Quantitative Browser, and 5 ppm mass tolerance was applied for the extraction of target product ions. For each sample analyzed, three calibration curves were generated by linear regression analysis and the concentration of each peptide in the sample was calculated by determining the value of the intercept in the *x*-axis, which represent the initial analyte concentration in the sample.

181

## 182 2.8. Statistical analysis

183 All data are presented as mean ± standard deviation (SD) for three replicates for each prepared  
184 sample. Univariate analysis of variance (ANOVA) with Tukey post-hoc test was applied using  
185 GraphPad Prism 6.0 (GraphPad Software, San Diego, CA, USA). The differences were considered  
186 significant with *P* < 0.05.

### 187 3. Results and discussion

188

#### 189 3.1. Total peptides quantification in the peptide fractions of Parmigiano-Reggiano (PR) WSPE and 190 digested samples

191 During cheese ripening, caseins can be hydrolyzed by the activity of proteases and peptidases  
192 mainly derived from S-LAB and NS-LAB (Pangallo et al., 2019). Generally, LAB possess a  
193 complex proteolytic system, which is able to hydrolyze milk caseins to short peptides and amino  
194 acids to fulfill their amino acid requirements (Tagliazucchi, Martini, & Solieri, 2019). LAB cell-  
195 envelope proteases (CEPs) break down caseins into protein fragments (mainly oligopeptides of  
196 about 5-30 amino acids) (Solieri, De Vero, & Tagliazucchi, 2018). These peptides can be  
197 transported into the cell and further hydrolyzed by cytoplasmic peptidases into smaller peptides and  
198 amino acids (Tagliazucchi et al., 2019). As shown in Figure 1, ripening affected the extent of  
199 proteolysis in PR samples, as determined by the TNBS assay. The amount of water-soluble low  
200 molecular weight peptides did not differ between PR12 and PR18 samples whereas a significant  
201 increase was observed in PR24 sample ( $P < 0.05$ ).

202 Previous studies already confirmed an increase in proteolysis as a function of cheese ripening  
203 (Bütikofer, Meyer, Sieber, Walther, & Wechsler, 2008; Stuknite et al., 2015). Gaiaschi et al. (2001)  
204 reported that the extent of proteolysis in Grana Padano cheese was high in the first 12 months of  
205 ripening, reaching a plateau and further increased after 22 months of ripening.

206 An increase in the level of low molecular weight peptides was observed for PR cheeses at different  
207 ripening time-points after gastric digestion (Figure 1). The amount of peptides released from PR24  
208 after gastric digestion was significantly higher ( $P < 0.001$ ) than that released from PR12 and PR18.  
209 No significant differences were observed between PR12 and PR18 after gastric digestion  
210 ( $P > 0.05$ ). Intestinal digestion brought about an increase in the amount of low molecular weight  
211 peptides in all of the samples. Once again, the concentration of peptides detected after the intestinal  
212 digestion step of PR24 was significantly higher than that measured in PR12 and PR18 ( $P < 0.05$ ;

Fig. 1). No significant differences ( $P > 0.05$ ) were found between peptide concentrations in PR12 and PR18.

### 3.2. Effect of ripening on the peptidomic profile of Parmigiano-Reggiano (PR) WSPE peptide fractions

Overall, 278 unique peptides were identified in the three PR WSPE peptide fractions at different ripening time-points (Table S1). According to the TNBS assay data, the PR24 sample contained the highest amount of peptides (257 peptides), whereas the amount of peptides identified in PR12 and PR18 samples was similar (84 and 72 peptides, respectively) and lower compared to the number observed in the PR24 sample (Fig. 2A). The majority of the peptides identified in PR12 and PR18 samples were from  $\beta$ -casein (63.1% and 58.3%, respectively), whereas the remaining identified peptides were from  $\alpha_{S1}$ -casein (Fig. 2A). PR24 sample also contained peptides released from  $\alpha_{S2}$ -casein (13.6% of total peptides). The Venn diagram (Fig. 3A) shows that 64 peptides (23% of total peptides) were commonly found in all the PR samples. Five peptides (1.8% of total peptides) were in common between PR12 and PR18, whereas the PR24 sample contained 191 (68.7% of total peptides) unique peptides. Among the 84 peptides identified in PR12, 15 of them (~18% of peptides identified in PR12) were found only in this sample.

The peptidomic profile of the three PR samples highlighted the paramount importance of LAB CEPs in the proteolysis of cheese caseins, as indicated by the cleavage sites in the N-terminal region of  $\alpha_{S1}$ -casein. Peptidic bonds at the H<sub>8</sub>–Q<sub>9</sub>, Q<sub>9</sub>–G<sub>10</sub>, Q<sub>13</sub>–E<sub>14</sub>, E<sub>14</sub>–V<sub>15</sub>, L<sub>16</sub>–N<sub>17</sub> and F<sub>23</sub>–F<sub>24</sub> positions are well-known cleavage sites for LAB CEPs (Solieri et al., 2018; Jensen, Vogensen, & Ardö, 2009; Hebert et al., 2008). Another trait indicating the action of CEPs in the production of these peptides is that the majority of cleavage sites in  $\beta$ -casein (61.8%, 63.8% and 74.0% in the PR12, PR18 and PR24, respectively) have been previously reported to be typical for several CEPs from *Lactobacillus casei*, *Lactobacillus rhamnosus*, *Lactobacillus delbrueckii*, *Lactobacillus paracasei*, *Lactobacillus lactis* and *Lactobacillus helveticus* (Solieri et al., 2018; Hebert et al., 2008;

239 Lozo et al., 2011; Juillard et al., 1995; Miyamoto et al., 2015). It is important to emphasize that a  
 240 complex and dynamic population of LAB, which thoroughly changes during ripening, is  
 241 characteristic of PR cheeses (Solieri et al., 2012).  
 242 The action of the extracellular proteinase and intracellular peptidases in *Lactobacillus* can explain  
 243 the presence of the high number of unique peptides in PR24. Several studies showed that the LAB  
 244 population decreases as the ripening of PR proceeds (Solieri et al., 2012; Coppola et al., 1999).  
 245 LAB cell lysis may release cell envelope-proteases and intracellular peptidases in the matrices,  
 246 which can then enhance the proteolysis of caseins or caseins peptides. Indeed, the bacterial lysis  
 247 decreases the number of vital LAB and therefore the amount of peptides translocated into the cells.  
 248  
 249 *3.3. Effect of in vitro digestion on the peptidomic profile of digested Parmigiano-Reggiano (PR)*  
 250 *peptide fractions*  
 251 UHPLC/HR-MS analysis revealed different peptide profiles for the peptide fractions from the PR  
 252 samples after *in vitro* gastro-intestinal digestion (Figs. 2B and 3B). In each sample, the majority of  
 253 the peptides were from  $\beta$ -caseins (47.8, 40.4 and 38.2% of the total identified peptides in digested  
 254 PR12, PR18 and PR24, respectively) followed by  $\alpha_{s1}$ -casein (29.3, 30.9 and 32.6% of the total  
 255 identified peptides in digested PR12, PR18 and PR24, respectively) (Fig. 2B).  
 256 The Venn diagram indicated that 158 identified peptides (corresponding to the 33.7% of total  
 257 identified peptides) co-existed in the three PR peptide fractions after *in vitro* digestion (Fig. 3B).  
 258 There were 115, 31 and 24 peptides exclusively found in digested PR12, PR18 and PR24,  
 259 respectively. The highest similarity in peptide profiles was found between digested PR18 and PR24  
 260 samples, with 258 common peptides.  
 261 Only 22, 21 and 37 peptides were commonly found in undigested and digested samples from PR12,  
 262 PR18 and PR24 peptide fractions, respectively (supplementary Table S1). Among these peptides,  
 263 14 (30.4% of total peptides) were commonly found in each of the undigested and digested samples  
 264 (supplementary Fig. S1). The peptides RELEEL, ELEEL, DKIHPP, LVYPFP, EMPFPK,

265 SLVYFPFGPIP, LVYFPFGPIP, YFPFGPIP, VLPVPQK and AVPYYPQR were from  $\beta$ -casein  
266 whereas the peptides FVAPFPE, VAPFPE, EIVPN and YKVPQ were from  $\alpha_{S1}$ -casein. It is not  
267 surprising that most of these peptides contain a PXP sequence or a proline residue near to the  
268 carboxylic end. These peptide structural motifs increase the resistance to gastro-pancreatic proteases  
269 action, which do not readily hydrolyze proline-containing peptides (Tagliazucchi, Helal, Verzelloni,  
270 Bellesia, & Conte, 2016). Indeed, most of these PXP-containing peptides have been found after *in*  
271 *vivo* or *in vitro* gastro-intestinal digestion of milk or milk proteins (Tagliazucchi et al., 2016;  
272 Tagliazucchi, Martini, Shamsia, Helal, & Conte, 2018; Boutrou et al., 2013; Boutrou, Henry, &  
273 Sanchez-Rivera, 2015). The peptides found only in undigested cheeses were likely degraded during  
274 *in vitro* gastro-intestinal digestion.

275

#### 276 3.4. Effect of ripening on the evolution and fate of bioactive peptides in Parmigiano-Reggiano 277 cheeses

278 Peptides in the undigested PR peptide fractions from WSPE were compared for sequence matches  
279 with the milk bioactive peptide database MBPDB (Nielsen et al., 2017). Across the categories, 26  
280 peptides in undigested PR samples (Table 1) shared the same sequence (100% of homology) with  
281 functional peptides previously reported to have various bioactivities.

282 Among the peptide fractions from undigested WSPE, the PR24 sample contained the highest  
283 amount of bioactive peptides (26 peptides) respect to PR18 and PR12 samples (12 and 11 peptides,  
284 respectively) (Table 1 and supplementary Figure S2A). The Venn diagram (supplementary Fig.  
285 S2A) shows that 11 bioactive peptides (42.3% of total peptides) were commonly found in all the PR  
286 samples, whereas 14 bioactive peptides were uniquely present in the PR24 sample (Table 1). Nine  
287 of the identified bioactive peptides were angiotensin converting-enzyme (ACE) inhibitors, 5  
288 peptides were anti-microbial, 1 was a di-peptidyl-peptidase IV (DPPIV) inhibitor, 1 was anxiolytic,  
289 1 was antioxidant and 9 were multifunctional bioactive peptides. Considering also the  
290 multifunctional bioactive peptides, 9 were anti-microbial essentially active against pathogenic

Gram-negative bacteria such as *Escherichia coli*, *Cronobacter sakazakii* and *Staphylococcus aureus* (Sedaghati, Ezzattpanah, Mashhadi Akbar Boojar, Tajabadi Ebrahimi, & Kobarfard, 2015; Birkemo, O'Sullivan, Ross, & Hill, 2009; Kent et al., 2012). The ability of LAB present in PR cheese to produce anti-microbial peptides from hydrolysis of milk proteins may confer a competitive advantage, thus decreasing the risk of the growth and survival of food-borne pathogens (Settanni, & Moschetti, 2010).

Of the 11 commonly identified peptides, the relative abundance of 10 was significantly higher in P24 sample respect to P12 and P18 samples ( $P<0.05$ ). In contrast, the ACE-inhibitory peptide FFVAPFPEVFGK displayed a decreasing trend during ripening with the highest relative abundance found in P12 sample ( $P<0.05$ ). The ACE-inhibitory peptide SKVLPVPQ was not detected in sample P12 but showed an increasing trend during ripening with the highest relative abundance found in sample P24 ( $P<0.05$ ).

### 3.5. Effect of *in vitro* digestion on the evolution and fate of bioactive peptides in Parmigiano-Reggiano cheeses

The bioactive peptide profile varied in the PR samples after *in vitro* gastro-intestinal digestion (Table 2). Globally, 52 peptides with 100% of homology with previously reported functional peptides were identified in digested PR samples (Table 2). The majority (75%) of total identified bioactive peptides (39 peptides) were commonly found in the three digested PR samples (Table 2 and supplementary Fig. S2B). Most of the identified bioactive peptides were ACE-inhibitors (17 peptides) and multifunctional peptides (16 peptides). The other identified bioactive peptides were anti-microbial (7 peptides), DPPIV-inhibitors (4 peptides), antioxidant (2 peptides), opioid (2 peptides), cathepsin B-inhibitors (2 peptides), prolyl-endopeptidase-inhibitor (1 peptide) and immunomodulatory (1 peptide). Three bioactive peptides (LHLPLP, HLPLP and AYPYPEL) were already reported after *in vitro* digestion of PR cheese at 12 months of ripening (Basiricò et al.,

2015), whereas the other bioactive peptides were identified in digested PR cheeses for the first time in this study.

The resulting data from semi-quantitative analysis demonstrated that the majority of identified bioactive peptides were not present at a constant level after digestion with respect to the ripening time, but each peptide showed a characteristic trend. Bioactive peptides identified after *in vitro* digestion can be clustered into 5 different groups as a function of the evolutive trend respect to the ripening time (Table 2).

The first group was represented by bioactive peptides whose release after *in vitro* digestion continuously increased according to the ripening time (Table 2). This group was mainly characterized by the presence of ACE-inhibitory peptides (10 bioactive peptides out of 13). Most of these peptides showed low or very low IC<sub>50</sub> values against ACE. The peptides LHLPLPL, LHLPLP and YKVPQL have been reported to reduce hypertension in spontaneously hypertensive rats (SHR) (Quirós et al., 2007; Maeno, Yamamoto, & Takano, 1996; Miguel, Recio, Ramos, Delgado, & Aleixandre, 2006). Peptides LHLPLP, YKVPQL and EMPFPK were also found intact in human gastro-intestinal tract (Boutrou et al., 2013).

Peptide LHLPLP was able to resist *in vitro* gastro-intestinal digestion but it was hydrolyzed to HLPLP by cellular peptidases prior to being transported across Caco-2 cells (Quirós et al., 2008; Tagliazucchi et al., 2006). The latter can actually be absorbed by intestinal cells and has been found in human plasma after oral administration (Van Platerink et al., 2006). It has been suggested that the peptide LHLPLP, released after *in vitro* gastro-intestinal digestion of Grana-Padana cheese, may be partially responsible for the blood pressure lowering effect observed *in vivo* after diet enrichment with Grana-Padana cheese (Stuknite et al., 2015; Crippa et al., 2018).

The second group was characterized by bioactive peptides whose release after *in vitro* digestion increased according to the ripening time reaching a plateau after 18 months of ripening (Table 2). This group contained the majority of anti-microbial peptides and some ACE-inhibitory peptides with demonstrated *in vivo* activity on spontaneously hypertensive rats (SHR) and low IC<sub>50</sub> values.



342 The peptide AVYPYQR was able to decrease the blood pressure in SHR and behaved as a  
343 multifunctional bioactive peptide also showing anti-microbial, anticoagulant and antioxidant  
344 activities (Karaki et al., 1990; Tonolo et al., 2018; Tu et al., 2019).

345 The third group was characterized by bioactive peptides the release of which after *in vitro* digestion  
346 increased according to the ripening time reaching a maximum value at 18 months of ripening (Table  
347 2). To this group belonged peptides with different biological activities. The peptide AYFYPEL  
348 presented a very low IC<sub>50</sub> value against ACE and was able to reduce blood pressure in SHR  
349 (Contreras, Carrón, Montero, Ramos, & Recio, 2009).

350 The fourth group was represented by bioactive peptides whose release after *in vitro* digestion  
351 decreased according to the ripening time (Table 2). This group was characterized for the presence of  
352 the peptide YPFPGPI (also known as  $\beta$ -casomorphin-7) and its precursors. Some ACE-inhibitory  
353 peptides were also found in this group but, with the exception of YPFPGPIP<sub>N</sub>, they displayed  
354 higher IC<sub>50</sub> values.

355 Finally, the last group contained peptides whose amount after *in vitro* digestion remained constant  
356 throughout ripening (Table 2).

357

### 358 3.5. Quantification of YPFPGPI, VPP and IPP in the peptide fractions of WSPE and digested 359 samples of Parmigiano-Reggiano (PR)

360 Three peptides, namely VPP, IPP and YPFPGPI with documented *in vivo* effect on humans were  
361 quantified in undigested and digested PR samples. The tripeptides VPP and IPP received particular  
362 consideration since several *in vivo* studies confirmed their antihypertensive effect on SHR and  
363 mildly hypertensive patients (Cicero, Fogacci, & Colletti, 2017; Fitzgerald, Murray, & Walsh,  
364 2004). Vice versa, different studies have suggested adverse effects of YPFPGPI ( $\beta$ -casomorphin-7)  
365 on human health, including cardiovascular diseases, diabetes and digestive disorders (Asledottir et  
366 al., 2018).

VPP and IPP have been detected in the WSPE peptide fractions from undigested PR samples at each ripening time (Table 3). The amount of VPP and IPP found in the 12-month ripened PR were  $6.87 \pm 0.68$  and  $1.63 \pm 0.82$  mg kg<sup>-1</sup>, respectively. These data are in accordance with the range reported by Basiricò et al. (2015) in PR sample at 12 months of ripening. The amount of VPP and IPP increased in the sample at 18 months of ripening, reaching a concentration of  $11.34 \pm 0.21$  and  $4.24 \pm 2.85$  mg kg<sup>-1</sup>, respectively. After that, we observed a strong decline in the concentration of VPP and IPP at 24 months of ripening ( $4.52 \pm 0.28$  and  $0.66 \pm 0.05$  mg kg<sup>-1</sup>, respectively). Peptide YPFPGPI, in contrast, was not detected in any undigested PR sample. This is consistent with the report of De Noni, & Cattaneo (2010), who did not observe YPFPGPI in Grana Padano cheese at 10, 17 or 25 months of ripening. These results suggested that LAB proteases and peptidases are not able to release  $\beta$ -casomorphin-7 during cheese ripening.

As shown in Table 3, at the end of the *in vitro* gastro-intestinal digestion, the VPP content of PR12 and PR18 remained almost unchanged. Moreover, the quantitative analysis mainly showed an increase in the content of IPP in both the samples ( $P < 0.05$ ). In contrast, we observed a significant decrease ( $P < 0.05$ ) in VPP concentrations after *in vitro* digestion in the PR24 sample, whereas the amount of IPP was unaltered.

$\beta$ -casomorphin-7, which was not present in the cheese WSPE, was released during *in vitro* digestion of PR samples (Table 3). Previous research highlighted the ability of gastro-intestinal proteases to release  $\beta$ -casomorphin-7 during *in vitro* digestion of milk  $\beta$ -casein variant A1 (Asledottir et al., 2018). The concentration of YPFPGPI after *in vitro* digestion decreased during ripening, with the highest concentration found in digested PR12 sample.

388

#### 389 4. Conclusion

390 According to the data reported in this study, ripening of PR cheese has an important influence in the  
 391 release of bioactive peptides and the *in vitro* digestion further increased their number in the PR  
 392 samples. Most of them were found in all of the samples, but in different amounts. Interestingly, they

393 can be clustered accordingly to ripening time and bioactivities. For example, the majority of the  
394 bioactive peptides showing an increasing trend after digestion, as a function of the ripening time,  
395 were potent ACE-inhibitory peptides. By contrast, most of the identified anti-microbial peptides  
396 reached a plateau after 18 months of ripening. Moreover, the opioid peptides  $\beta$ -casomorphin-7 and  
397 its precursor displayed a typical behavior with a decreasing trend after *in vitro* digestion as a  
398 function of the ripening time. The present study suggests possible differences in the biological  
399 effect after ingestion of PR cheese as a function of the ripening time. The major driving force for  
400 consumers to choose a cheese with different ripening times is the organoleptic characteristics but,  
401 nevertheless, the peptide profile and bioactivities may also change.

**Author contributions**

SM, AC and DT conceived and designed the study. SM performed the *in vitro* digestion and bioactivity experiments. SM and DT performed the peptidomic experiments and the bioinformatic analysis. DT wrote the manuscript. SM and AC critically revised the manuscript. All the authors read the manuscript and discussed the interpretation of results.

## **Acknowledgements**

The authors acknowledge the Fondazione Cassa di Risparmio di Modena for funding the HPLC-ESI-Q Exactive Hybrid Quadrupole-Orbitrap Mass Spectrometer system at the Centro Interdipartimentale Grandi Strumenti (CIGS).

**Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## **Declarations of interest**

None

## References

- Adler-Nissen, J. (1979). Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzensulfonic acid. *Journal of Agricultural and Food Chemistry*, 27, 1256-1262.
- Asledottir, T., Le, T. T., Poulsen, N. A., Devold, T. G., Larsen, L. B., & Vegarud, G. E. (2018). Release of  $\beta$ -casomorphin-7 from bovine milk of different  $\beta$ -casein variants after ex vivo gastrointestinal digestion. *International Dairy Journal*, 81, 8-11.
- Basiricò, L., Catalani, E., Morera, P., Cattaneo, S., Stuknyte, M., Bernabucci, U., De Noni, I., & Nardone, A. Release of angiotensin converting enzyme-inhibitor peptides during *in vitro* gastrointestinal digestion of Parmigiano-Reggiano PDO cheese and their absorption through an in vitro model of intestinal epithelium. *Journal of Dairy Science*, 98, 7595-7601.
- Birkemo, G. A., O'Sullivan, O., Ross, R. P., & Hill, C. (2009). Antimicrobial activity of two peptides casecidin 15 and 17, found naturally in bovine colostrum. *Journal of Applied Microbiology*, 106, 233-240.
- Boutrou, R., Gaudichon, C., Dupont, D., Jardin, J., Airinei, G., Marsset-Baglieri, A., et al. (2013). Sequential release of milk protein-derived bioactive peptides in the jejunum in healthy humans. *American Journal of Clinical Nutrition*, 97, 1314-1323.
- Boutrou, R., Henry, G., & Sanchez-Rivera, L. (2015). On the trail of milk bioactive peptides in human and animal intestinal tracts during digestion: A review. *Dairy Science and Technology*, 95, 815-829.
- Bütikofer, U., Meyer, J., Sieber, R., Walther, B., & Wechsler, D. (2008). Occurrence of the angiotensin-converting enzyme-inhibiting tripeptides Val-Pro-Pro and Ile-Pro-Pro in different cheese varieties of Swiss origin. *Journal of Dairy Science*, 91, 29-38.
- Cicero, A. F. G., Fogacci, F., & Colletti, A. (2017). Potential role of bioactive peptides in prevention and treatment of chronic diseases: a narrative review. *British Journal of Pharmacology*, 174, 1378-1394.



- Contreras, M. M., Carrón, R., Montero, M. J., Ramos, M., & Recio, I. (2009). Novel casein-derived peptides with antihypertensive activity. *International Dairy Journal*, 19, 566-573.
- Coppola, R., Nanni, M., Iorizzo, M., Sorrentino, A., Sorrentino, E., & Grazia, L. (1997). Survey of lactic acid bacteria isolated during the advanced stages of the ripening of Parmigiano-Reggiano cheese. *Journal of Dairy Research*, 64, 305-310.
- Crippa, G., Zabzuni, D., Bravi, E., Piga, G., De Noni, I., Bigli, E., & Rossi, F. (2018). Randomized, double blind placebo-controlled pilot study of the antihypertensive effects of Grana Padano D.O.P. cheese consumption in mild - moderate hypertensive subjects. *European Review for Medical and Pharmacological Sciences*, 22, 7573-7581.
- De Noni, I., & Cattaneo, S. (2010). Occurrence of  $\beta$ -casomorphins 5 and 7 in commercial dairy products and in their digests following *in vitro* simulated gastro-intestinal digestion. *Food Chemistry*, 119, 560-566.
- Fitzgerald, R. J., Murray, B. A., & Walsh, D. J. (2004). Hypotensive peptides from milk proteins. *Journal of Nutrition*, 134, 980S-988S.
- Gaiaschi, A., Beretta, B., Poiesi, C., Conti, A., Giuffrida, M. G., Galli, C. L., & Restani, P. (2001). Proteolysis of  $\beta$ -casein as a marker of Grana Padano cheese ripening. *Journal of Dairy Science*, 84, 60-65.
- Godos, J., Tieri, M., Ghelfi, F., Titta, L., Marventano, S., Lafranconi, A., Gambera, A., Alonzo, E., Sciacca, S., Buscemi, S., Ray, S., Del Rio, D., Galvano, F., & Grosso, G. (2019). Dairy foods and health: an umbrella review of observational studies. *International Journal of Food Sciences and Nutrition*, doi:10.1080/09637486.2019.1625035.
- Gómez-Ruiz, J. Á., Ramos, M., & Recio I. (2004). Identification and formation of angiotensin converting enzyme-inhibitory peptides in Manchego cheese by high-performance liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A*, 1054, 269-277.
- Hebert, E. M., Mamone, G., Picariello, G., Raya, R. R., Savoy, G., Ferranti, P., et al. (2008). Characterization of the pattern of  $\alpha$ S1- and  $\beta$ -casein breakdown and release of bioactive peptide

- by a cell envelope proteinase from *Lactobacillus delbrueckii* subsp. *lactis* CRL 581. *Applied and Environmental Microbiology*, 74, 3682-3689.
- Jensen, M. P., Vogensen, F. K., & Ardö, Y. (2009). Variation in caseinolytic properties of six cheese related *Lactobacillus helveticus* strains. *International Dairy Journal*, 19, 661–668.
- Juillard, V., Laan, H., Kunji, E. R. S., Jeronimus-Stratingh, C. M., Bruins, A. P., & Konings, W. N. (1995). The extracellular PI-type proteinase of *Lactococcus lactis* hydrolyzes  $\beta$ -casein into more than one hundred different oligopeptides. *Journal of Bacteriology*, 177, 3472-3478.
- Karaki, H., Doi, K., Sugano, S., Huchiwa, H., Sugai, R., Muramaki, U., & Takemoto, S. (1990). Antihypertensive effect of tryptic hydrolysate of milk casein in spontaneously hypertensive rats. *Comparative Biochemistry and Physiology*, 96, 367-371.
- Kent, R. M., Guinane, C. M., O'Connor, P. M., Fitzgerald, G. F., Hill, C., Stanton, C., & Ross, R. P. (2012). Production of the antimicrobial peptides Caseicin A and B by *Bacillus* isolates growing on sodium caseinate. *Letters of Applied Microbiology*, 55, 141-148.
- Lozo, J., Strahinic, I., Dalgarrondo, M., Chobert, J. M., Haertle, T., & Topisirovic, C. (2011). Comparative analysis of  $\beta$ -casein proteolysis by PrtP proteinase from *Lactobacillus paracasei* subsp. *paracasei* BGHN14, PrtR proteinase from *Lactobacillus rhamnosus* BGT10 and PrtH proteinase from *Lactobacillus helveticus* BGRA43. *International Dairy Journal*, 21, 863-868.
- Lu, Y., Govindasamy-Lucey, S., & Lucey, J. A. (2016). Angiotensin-I-converting enzyme-inhibitory peptides in commercial Wisconsin Cheddar cheeses of different ages. *Journal of Dairy Science*, 99, 41-52.
- Maeno, M., Yamamoto, N. & Takano, T. (1996) Identification of antihypertensive peptides from casein hydrolysate produced by a proteinase from *Lactobacillus helveticus* CP790. *Journal of Dairy Science*, 73, 1316-1321.
- Meyer, J., Bütikofer, U., Walther, B., Wechsler, D., & Sieber, R. (2009). Changes in angiotensin-converting enzyme inhibition and concentrations of the tripeptides Val-Pro-Pro and Ile-Pro-Pro during ripening of different Swiss cheese varieties. *Journal of Dairy Science*, 92, 826-836.

- Miguel, M., Recio, I., Ramos, M., Delgado, M. A., & Aleixandre, M. A. (2006). Antihypertensive effect of peptides obtained from *Enterococcus faecalis*-fermented milk in rats. *Journal of Dairy Science*, 89, 3352–3359.
- Minekus, M., Alming, M., Alvito, P., Ballance, S., Bohn, T., Bourlieu, C., et al. (2014). A standardised static in vitro digestion method suitable for food—an international consensus. *Food and Function*, 5, 1113 –1124.
- Miyamoto, M., Ueno, H. M., Watanabe, M., Tatsuma, Y., Seto, Y., Miyamoto, T., & Nakajima, H. (2015). Distinctive proteolytic activity of cell envelope proteinase of *Lactobacillus helveticus* isolated from airag, a traditional Mongolian fermented mare's milk. *International Journal of Food Microbiology*, 197, 65-71.
- Nielsen, S. D., Beverly, R. L., Qu, Y., & Dallas, D. C. (2017). Milk bioactive peptide database: A comprehensive database of milk protein-derived bioactive peptides and novel visualization. *Food Chemistry*, 232, 673–82.
- Nongonierma, A. B., & FitzGerald, R. J. (2015). The scientific evidence for the role of milk protein-derived bioactive peptides in humans: A Review. *Journal of Functional Foods*, 17, 640 - 656.
- Ong, L., & Shah N. P. (2008). Release and identification of angiotensin-converting enzyme inhibitory peptides as influenced by ripening temperatures and probiotic adjuncts in Cheddar cheeses. *LWT-Food Science and Technology*, 41, 1555–1566.
- Pangallo, D., Kraková, L., Puškárová, A., Šoltys, K., Bučková, M., Koreňová, J., Budiš, J., & Kuchta, T. (2019). Transcription activity of lactic acid bacterial proteolysis-related genes during cheese maturation. *Food Microbiology*, 82, 416-425.
- Quirós, A., Ramos, M., Muguerza, B., Delgado, M. A., Miguel, M., Aleixandre, A., & Recio, I. (2007). Identification of novel antihypertensive peptides in milk fermented with *Enterococcus faecalis*. *International Dairy Journal*, 17, 33-41.

- Quirós, A., Dávalos, A., Lasunción, M. A., Ramos, M., & Recio, I. (2008). Bioavailability of the antihypertensive peptide LHLPLP: Transepithelial flux of HLPLP. *International Dairy Journal*, 18, 279-286.
- Rizzello, C. G., Tagliazucchi, D., Babini, E., Rutella, G. S., Taneyo Saa, D. L., & Gianotti, A. (2016). Bioactive peptides from vegetable food matrices: Research trends and novel biotechnologies for synthesis and recovery. *Journal of Functional Foods*, 27, 549-569.
- Sánchez-Rivera, L., Diezhandino, I., Gómez-Ruiz, J. Á., Fresno, J. M., Miralles, B., & Recio, I. (2014). Peptidomic study of Spanish blue cheese (Valdeón) and changes after simulated gastrointestinal digestion. *Electrophoresis*, 35, 1627-1636.
- Sedaghati, M., Ezzatpanah, H., Mashhadi Akbar Boojari, M., Tajabadi Ebrahimi, M., & Kobarfard, F. (2015). Isolation and identification of some antibacterial peptides in the plasmin-digest of  $\beta$ -casein. *LWT-Food Science and Technology*, 68, 217-225.
- Settanni, L., & Moschetti, G. (2010). Non-starter lactic acid bacteria used to improve cheese quality and provide health benefits. *Food Microbiology*, 27, 691-697.
- Sforza, S., Galaverna, G., Neviani, E., Pinelli, C., Dossena, A., & Marchelli, M. (2004). Study of the oligopeptide fraction in Grana Padano and Parmigiano-Reggiano cheeses by liquid chromatography-electrospray ionization mass spectrometry. *European Journal of Mass Spectrometry*, 10, 421-427.
- Sforza, S., Cavatorta, V., Lambertini, F., Galaverna, G., Dossena, A., & Marchelli, R. (2012). Cheese peptidomics: a detailed study on the evolution of the oligopeptide fraction in Parmigiano-Reggiano cheese from curd to 24 months of aging. *Journal of Dairy Science*, 95, 3514-26.
- Sieber, R., Bütikofer, U., Egger, C., Portmann, R., Walther, B., & Wechsler, D. (2010). ACE-inhibitory activity and ACE-inhibiting peptides in different cheese varieties. *Dairy Science and Technology*, 90, 47-73.

- Solieri, L., Bianchi, A., & Giudici, P. (2012). Inventory of non starter lactic acid bacteria from ripened Parmigiano-Reggiano cheese as assessed by a culture dependent multiphasic approach. *Systematic and Applied Microbiology*, 35, 270-277.
- Solieri, L., De Vero, L., & Tagliazucchi, D. (2018). Peptidomic study of casein proteolysis in bovine milk by *Lactobacillus casei* PRA205 and *Lactobacillus rhamnosus* PRA331. *International Dairy Journal*, 85, 237-246.
- Stuknyte, M., Cattaneo, S., Masotti, F., & De Noni, I. (2015). Occurrence and fate of ACE-inhibitor peptides in cheeses and in their digestates following *in vitro* static gastrointestinal digestion. *Food Chemistry*, 168, 27-33.
- Summer, A., Formaggioni, P., Franceschi, P., Di Frangia, F., Righi, F., & Malacarne, M. (2017). Cheese as functional food: the example of Parmigiano-Reggiano and Grana Padano. *Food Technology and Biotechnology*, 55, 277-289.
- Tagliazucchi, D., Helal, A., Verzelloni, E., Bellesia, A., & Conte, A. (2016). Composition and properties of peptides that survive standardised *in vitro* gastro-pancreatic digestion of bovine milk. *International Dairy Journal*, 61, 196-204.
- Tagliazucchi, D., Shamsia, S., Helal, A., & Conte, A. (2017). Angiotensin-converting enzyme inhibitory peptides from goats' milk released by *in vitro* gastro-intestinal digestion. *International Dairy Journal*, 71, 6-16.
- Tagliazucchi, D., Martini, S., Shamsia, S., Helal, A., & Conte, A. (2018). Biological activity and peptidomic profile of *in vitro* digested cow, camel, goat and sheep milk. *International Dairy Journal*, 81, 19-27.
- Tagliazucchi, D., Martini, S. & Solieri, L. (2019). Bioprospecting for bioactive peptide production by lactic acid bacteria isolated from fermented dairy food. *Fermentation*, 5, 96.
- Tonolo, F., Sandre, M., Ferro, S., Folda, A., Scalcon, V., Scutari, G., Feller, E., Marin, O., Bindoli, A., & Rigobello, M. P. (2018). Milk-derived bioactive peptides protect against oxidative stress in a Caco-2 cell model. *Food and Function*, 9, 1245-1253.

- Tu, M., Liu, H., Cheng, S., Mao, F., Chen, H., Fan, F., Lu, W., & Du, M. (2019). Identification and characterization of a novel casein anticoagulant peptide derived from *in vivo* digestion. *Food and Function*, 10, 2552-2559.
- Van Platerink, C. J., Janssen, H. G. M., Horsten, R., & Haverkamp, J. (2006). Quantification of ACE inhibiting peptides in human plasma using high performance liquid chromatography–mass spectrometry. *Journal of Chromatography B*, 830, 151-157.

## Figure captions

**Fig. 1.** Effect of ripening time and *in vitro* digestion on the peptide concentrations in Parmigiano-Reggiano (PR). The total amount of peptides was expressed as mmol of leucine equivalent per g of cheese. WSPE means water-soluble peptide extracts and represent the peptide fractions obtained after ultrafiltration (< 3 kDa) of water-soluble peptides extracted from un-digested PR samples. PR12: Parmigiano-Reggiano at 12 months of ripening. PR18: Parmigiano-Reggiano at 18 months of ripening. PR24: Parmigiano-Reggiano at 24 months of ripening. Values are means of data from three independent digestions  $\pm$  standard deviation (SD). Different letters indicate significantly different values ( $P < 0.05$ )

**Fig. 2.** Number of unique peptides identified in the Parmigiano-Reggiano (PR) peptide fractions. (A) Number of peptides in un-digested PR peptide fractions at 12 (PR12), 18 (PR18) and 24 (PR24) months of ripening. (B) Number of peptides in *in vitro* digested PR peptide fractions at 12 (PR12), 18 (PR18) and 24 (PR24) months of ripening. The incidence of the different milk proteins on the released peptides is also shown.

**Fig. 3.** Venn diagrams of peptides obtained from Parmigiano-Reggiano (PR) peptide fractions. (A) Venn diagram created with all the identified peptides in un-digested PR peptide fractions at 12 (WSPE PR12), 18 (WSPE PR18) and 24 (WSPE PR24) months of ripening (see on line supplementary material Tables S1 for the peptide sequences). (B) Venn diagram created with all the identified peptides in digested PR peptide fractions at 12 (D PR12), 18 (D PR18) and 24 (D PR24) months of ripening (see on line supplementary material Tables S1 for the peptide sequences).

Table 1. Relative amount of bioactive peptides identified in water-soluble extract (WSPE) peptide fractions from Parmigiano Reggiano samples at 12 (PR12), 18 (PR18) and 24 (PR24) months of ripening with previously demonstrated bioactivity<sup>a</sup>.

<i>Sequence<sup>b</sup></i>	<i>Fragment</i>	<i>Relative amount in WSPE PR12<sup>c</sup></i>	<i>Relative amount in WSPE PR18<sup>c</sup></i>	<i>Relative amount in WSPE PR24<sup>c</sup></i>
<i>ACE-inhibitory</i>				
DKIHFP	βCN 47-52	9.02x10 <sup>5</sup> ± 7.75x10 <sup>4a</sup>	9.80x10 <sup>5</sup> ± 5.81x10 <sup>4a</sup>	1.41x10 <sup>6</sup> ± 7.43x10 <sup>4b</sup>
LVYPFP	βCN 58-63	1.11x10 <sup>5</sup> ± 1.04x10 <sup>4a</sup>	1.04x10 <sup>5</sup> ± 2.05x10 <sup>4a</sup>	4.01x10 <sup>5</sup> ± 9.12x10 <sup>4b</sup>
SQSKVLPVPQ	βCN 166-175	n.d.	n.d.	1.84x10 <sup>6</sup> ± 1.05x10 <sup>5</sup>
SKVLPVPQ	βCN 168-175	n.d.	1.01x10 <sup>4</sup> ± 1.63x10 <sup>2a</sup>	4.79x10 <sup>5</sup> ± 1.27x10 <sup>5b</sup>
VLPVPQK <sup>d,e</sup>	βCN 170-176	2.89x10 <sup>5</sup> ± 2.82x10 <sup>4a</sup>	2.44x10 <sup>5</sup> ± 5.50x10 <sup>4a</sup>	3.52x10 <sup>6</sup> ± 1.06x10 <sup>5b</sup>
RDMPIQAF	βCN 183-190	9.47x10 <sup>3</sup> ± 8.98x10 <sup>2a</sup>	1.09x10 <sup>4</sup> ± 6.11x10 <sup>2a</sup>	1.01x10 <sup>6</sup> ± 6.77x10 <sup>4b</sup>
YQEPVLGPVRGPFPIIV <sup>e,f</sup>	βCN 193-209	1.32x10 <sup>4</sup> ± 5.95x10 <sup>3a</sup>	4.18x10 <sup>4</sup> ± 3.18x10 <sup>3b</sup>	4.04x10 <sup>5</sup> ± 4.26x10 <sup>4b</sup>
QEPVLGPVRGPFPIIV	βCN 194-209	n.d.	n.d.	5.06x10 <sup>5</sup> ± 7.12x10 <sup>4</sup>
FALPQYLK	α <sub>S2</sub> CN 174-181	n.d.	n.d.	1.66x10 <sup>5</sup> ± 8.02x10 <sup>4</sup>
AMKPWIQPK	α <sub>S2</sub> CN 189-197	n.d.	n.d.	4.96x10 <sup>5</sup> ± 6.73x10 <sup>4</sup>
<i>Anti-hypertensive</i>				
VYPFPGPIPN <sup>g</sup>	βCN 59-68	n.d.	n.d.	8.99x10 <sup>5</sup> ± 2.44x10 <sup>4</sup>
YFPFPGPIPN <sup>g</sup>	βCN 60-68	6.64x10 <sup>4</sup> ± 5.57x10 <sup>3a</sup>	3.33x10 <sup>4</sup> ± 4.34x10 <sup>3b</sup>	2.49x10 <sup>5</sup> ± 2.32x10 <sup>4c</sup>
EMPFPG <sup>g</sup>	βCN 108-113	1.47x10 <sup>4</sup> ± 4.60x10 <sup>2a</sup>	1.77x10 <sup>4</sup> ± 1.44x10 <sup>3b</sup>	7.06x10 <sup>5</sup> ± 4.75x10 <sup>4c</sup>
AVPYPQR <sup>g</sup>	βCN 177-183	1.89x10 <sup>4</sup> ± 3.16x10 <sup>3a</sup>	1.66x10 <sup>4</sup> ± 1.12x10 <sup>3a</sup>	4.52x10 <sup>5</sup> ± 4.21x10 <sup>4b</sup>
LLYQEPVLGPVRGPFPIIV <sup>g</sup>	βCN 191-209	n.d.	n.d.	1.72x10 <sup>6</sup> ± 2.41x10 <sup>5</sup>
FFVAPFPEVFGK <sup>g</sup>	α <sub>S1</sub> CN 23-34	1.38x10 <sup>6</sup> ± 9.46x10 <sup>4a</sup>	1.19x10 <sup>6</sup> ± 5.61x10 <sup>4b</sup>	4.42x10 <sup>5</sup> ± 2.47x10 <sup>5c</sup>
<i>Anti-microbial</i>				
EAMAPK	βCN 100-105	n.d.	n.d.	2.27x10 <sup>5</sup> ± 1.77x10 <sup>5</sup>
VLPVPQKAVPYPQR	βCN 170-183	n.d.	n.d.	1.50x10 <sup>6</sup> ± 1.31x10 <sup>5</sup>
VLNENLLR	α <sub>S1</sub> CN 15-22	n.d.	n.d.	1.76x10 <sup>5</sup> ± 1.08x10 <sup>5</sup>
HIQKEDVPSELYLGYLEQLRLK	α <sub>S1</sub> CN 80-102	n.d.	n.d.	1.09x10 <sup>5</sup> ± 7.56x10 <sup>4</sup>
YLEQLLR	α <sub>S1</sub> CN 94-101	n.d.	n.d.	5.57x10 <sup>5</sup> ± 7.73x10 <sup>4</sup>
<i>Immunomodulatory</i>				
PGPIPN	βCN 63-68	2.08x10 <sup>5</sup> ± 1.20x10 <sup>4a</sup>	1.21x10 <sup>5</sup> ± 2.18x10 <sup>4b</sup>	2.70x10 <sup>5</sup> ± 3.04x10 <sup>4c</sup>
<i>Antioxidant</i>				
VKEAMAPK <sup>e</sup>	βCN 98-105	n.d.	n.d.	2.80x10 <sup>5</sup> ± 2.75x10 <sup>4</sup>
TQTPVVVPPFLQPE	βCN 78-91	n.d.	n.d.	5.57x10 <sup>4</sup> ± 1.64x10 <sup>4</sup>
<i>DPPIV-inhibitory</i>				



LPVPQ	$\beta$ CN 171-175	n.d.	n.d.	$2.25 \times 10^5 \pm 1.14 \times 10^5$
<i>Anxiolytic</i>				
YLGYLEQLLR	$\alpha_{S1}$ CN 91-101	$6.79 \times 10^5 \pm 4.29 \times 10^{4a}$	$6.64 \times 10^5 \pm 2.65 \times 10^{4a}$	$1.14 \times 10^6 \pm 4.95 \times 10^{3b}$

<sup>a</sup>Abbreviations are: ACE, angiotensin converting enzyme; DPPIV, dipeptidyl peptidase IV; CN, casein.

<sup>b</sup>One code letter was used for amino acid nomenclature. Potential bioactivities were achieved from MBPDB databases (Nielsen et al., 2017). Anti-hypertensive activity was measured on spontaneously anti-hypertensive rats.

Multifunctional peptides were labelled with superscript letters: <sup>d</sup>antioxidant activity; <sup>e</sup>antimicrobial activity;

<sup>f</sup>immunomodulator; <sup>g</sup>ACE-inhibitory activity.

<sup>c</sup>Relative amount was expressed as the area under the peak (AUP) g<sup>-1</sup> of cheese measured from the extracted ion chromatograms (EIC) obtained for each peptide. N.d. means peptide not detected in the sample. Different superscript letters within the same row indicate that the values are significantly different ( $P < 0.05$ )

Table 2. Relative amount of bioactive peptides identified in digested peptide fractions from Parmigiano Reggiano samples at 12 (PR12), 18 (PR18) and 24 (PR24) months of ripening with previously demonstrated bioactivity<sup>a</sup>

<i>Sequence<sup>b</sup></i>	<i>Fragment</i>	<i>Relative amount in digested PR12<sup>c</sup></i>	<i>Relative amount in digested PR18<sup>c</sup></i>	<i>Relative amount in digested PR24<sup>c</sup></i>
<i>Peptides with increasing trend according to the ripening time</i>				
LVYFPF <sup>d</sup>	βCN 58-63	1.80x10 <sup>5</sup> ± 1.15x10 <sup>4a</sup>	1.18x10 <sup>5</sup> ± 8.77x10 <sup>4a</sup>	7.60x10 <sup>5</sup> ± 4.66x10 <sup>4b</sup>
NIPPLTQTPVVVPPFLQ <sup>d</sup>	βCN 73-89	1.18x10 <sup>4</sup> ± 7.45x10 <sup>3a</sup>	6.91x10 <sup>5</sup> ± 5.74x10 <sup>4b</sup>	9.09x10 <sup>5</sup> ± 1.50x10 <sup>4c</sup>
EMFPFK <sup>d,e</sup>	βCN 108-113	7.28x10 <sup>6</sup> ± 1.02x10 <sup>5a</sup>	3.04x10 <sup>7</sup> ± 1.98x10 <sup>6b</sup>	7.90x10 <sup>8</sup> ± 3.90x10 <sup>7c</sup>
LHLPLP <sup>d,f</sup>	βCN 133-138	2.45x10 <sup>8</sup> ± 1.85x10 <sup>7a</sup>	4.33x10 <sup>8</sup> ± 3.88x10 <sup>7b</sup>	6.83x10 <sup>8</sup> ± 6.22x10 <sup>7c</sup>
LHLPLPL <sup>d,f</sup>	βCN 133-139	2.75x10 <sup>7</sup> ± 1.51x10 <sup>6a</sup>	6.95x10 <sup>7</sup> ± 6.87x10 <sup>6b</sup>	8.63x10 <sup>7</sup> ± 4.92x10 <sup>6c</sup>
YQEPVL <sup>d</sup>	βCN 193-198	5.87x10 <sup>4</sup> ± 4.25x10 <sup>4a</sup>	4.38x10 <sup>6</sup> ± 5.53x10 <sup>5b</sup>	1.98x10 <sup>7</sup> ± 2.40x10 <sup>5c</sup>
YKVPQL <sup>d,f</sup>	α <sub>S1</sub> CN 104-109	4.06x10 <sup>4</sup> ± 3.65x10 <sup>4a</sup>	6.25x10 <sup>7</sup> ± 7.97x10 <sup>6b</sup>	1.38x10 <sup>8</sup> ± 3.32x10 <sup>7c</sup>
NMAINPSK <sup>d</sup>	α <sub>S2</sub> CN 25-32	2.05x10 <sup>6</sup> ± 3.38x10 <sup>6a</sup>	3.83x10 <sup>7</sup> ± 2.50x10 <sup>6b</sup>	6.00x10 <sup>7</sup> ± 3.12x10 <sup>6c</sup>
SRYPST <sup>g</sup>	κCN 33-38	n.d.	4.81x10 <sup>6</sup> ± 5.04x10 <sup>5a</sup>	7.55x10 <sup>6</sup> ± 8.60x10 <sup>5b</sup>
INNQLFPYPY <sup>h</sup>	κCN 51-60	1.34x10 <sup>5</sup> ± 1.39x10 <sup>4a</sup>	1.10x10 <sup>8</sup> ± 1.26x10 <sup>7b</sup>	2.48x10 <sup>8</sup> ± 6.84x10 <sup>6c</sup>
LPYPY <sup>d,h</sup>	κCN 56-60	3.61x10 <sup>5</sup> ± 5.51x10 <sup>4a</sup>	2.19x10 <sup>7</sup> ± 1.72x10 <sup>6b</sup>	4.64x10 <sup>7</sup> ± 3.20x10 <sup>6c</sup>
IPAVF <sup>h</sup>	βLB 78-82	4.67x10 <sup>5</sup> ± 6.27x10 <sup>4a</sup>	1.88x10 <sup>6</sup> ± 2.90x10 <sup>5b</sup>	7.95x10 <sup>6</sup> ± 1.29x10 <sup>5c</sup>
VLDTDYK <sup>d</sup>	βLB 94-100	n.d.	2.61x10 <sup>6</sup> ± 1.92x10 <sup>5a</sup>	5.27x10 <sup>6</sup> ± 1.15x10 <sup>5b</sup>
<i>Peptides with increasing trend according to the ripening time reaching a plateau at 18 months</i>				
PVVVPPFLQPE <sup>e</sup>	βCN 81-91	n.d.	1.57x10 <sup>7</sup> ± 1.67x10 <sup>6a</sup>	1.32x10 <sup>7</sup> ± 3.94x10 <sup>6a</sup>
VENLHLPLPL <sup>d</sup>	βCN 130-140	n.d.	1.19x10 <sup>6</sup> ± 1.80x10 <sup>5a</sup>	1.44x10 <sup>6</sup> ± 2.64x10 <sup>5a</sup>
VLPVPQK <sup>d</sup>	βCN 170-176	2.22x10 <sup>7</sup> ± 3.68x10 <sup>6a</sup>	1.00x10 <sup>9</sup> ± 1.49x10 <sup>6b</sup>	1.08x10 <sup>9</sup> ± 5.94x10 <sup>6b</sup>
AVPYPQR <sup>d,e,f,i</sup>	β177-183	3.00x10 <sup>5</sup> ± 1.31x10 <sup>5a</sup>	5.48x10 <sup>8</sup> ± 5.31x10 <sup>7b</sup>	6.01x10 <sup>8</sup> ± 2.76x10 <sup>7b</sup>
DAYPSGAW <sup>d</sup>	α <sub>S1</sub> CN 157-164	7.32x10 <sup>5</sup> ± 1.43x10 <sup>5a</sup>	4.61x10 <sup>6</sup> ± 9.28x10 <sup>5b</sup>	5.11x10 <sup>6</sup> ± 9.02x10 <sup>5b</sup>
SDIPNPIGSENSEK <sup>e</sup>	α <sub>S1</sub> CN 180-193	3.79x10 <sup>6</sup> ± 5.49x10 <sup>5a</sup>	2.48x10 <sup>8</sup> ± 3.92x10 <sup>7b</sup>	2.69x10 <sup>8</sup> ± 6.55x10 <sup>7b</sup>
FFSDK <sup>e</sup>	κCN 17-21	3.65x10 <sup>6</sup> ± 3.09x10 <sup>5a</sup>	1.26x10 <sup>7</sup> ± 1.32x10 <sup>6b</sup>	1.71x10 <sup>7</sup> ± 2.43x10 <sup>6b</sup>
YIPIQY <sup>d</sup>	κCN 25-30	2.59x10 <sup>4</sup> ± 2.49x10 <sup>4a</sup>	9.80x10 <sup>6</sup> ± 1.45x10 <sup>5b</sup>	1.23x10 <sup>7</sup> ± 2.95x10 <sup>6b</sup>
GLDIQK <sup>d,l</sup>	βLB 9-14	n.d.	4.53x10 <sup>6</sup> ± 7.53x10 <sup>5a</sup>	3.95x10 <sup>6</sup> ± 3.39x10 <sup>5a</sup>
DAQSAPLR <sup>e</sup>	βLB 33-40	n.d.	1.34x10 <sup>6</sup> ± 1.17x10 <sup>5a</sup>	1.87x10 <sup>6</sup> ± 2.95x10 <sup>5a</sup>
IIAEK <sup>d,l</sup>	βLB 71-75	1.76x10 <sup>6</sup> ± 1.50x10 <sup>5a</sup>	5.46x10 <sup>6</sup> ± 6.79x10 <sup>5b</sup>	4.61x10 <sup>6</sup> ± 3.52x10 <sup>5b</sup>
IDALNENK <sup>e</sup>	βLB 84-91	n.d.	1.96x10 <sup>6</sup> ± 1.53x10 <sup>5a</sup>	2.06x10 <sup>6</sup> ± 1.82x10 <sup>5a</sup>
<i>Peptides with increasing trend according to the ripening time reaching a maximum at 18 months</i>				
TEDELQDKIHPPF <sup>e</sup>	βCN 41-52	1.80x10 <sup>5</sup> ± 1.93x10 <sup>4a</sup>	4.39x10 <sup>6</sup> ± 1.81x10 <sup>5b</sup>	3.80x10 <sup>6</sup> ± 3.45x10 <sup>4c</sup>
DKIHPPF <sup>d</sup>	βCN 47-52	7.34x10 <sup>5</sup> ± 5.30x10 <sup>4a</sup>	2.66x10 <sup>6</sup> ± 1.77x10 <sup>5b</sup>	1.28x10 <sup>6</sup> ± 5.57x10 <sup>4c</sup>
PGPIPN <sup>m</sup>	βCN 63-67	7.73x10 <sup>6</sup> ± 2.89x10 <sup>4a</sup>	8.89x10 <sup>6</sup> ± 4.46x10 <sup>4b</sup>	6.55x10 <sup>6</sup> ± 3.66x10 <sup>4c</sup>
EAMAPK <sup>e</sup>	βCN 100-105	1.61x10 <sup>8</sup> ± 2.06x10 <sup>7a</sup>	2.55x10 <sup>8</sup> ± 3.22x10 <sup>6b</sup>	1.61x10 <sup>8</sup> ± 3.29x10 <sup>7a</sup>
KVLPVPQK <sup>i</sup>	βCN 169-176	1.89x10 <sup>4</sup> ± 2.73x10 <sup>3a</sup>	5.85x10 <sup>6</sup> ± 2.34x10 <sup>4b</sup>	3.35x10 <sup>6</sup> ± 1.88x10 <sup>5c</sup>

QEPVL <sup>m</sup>	βCN 194-198	n.d.	1.17x10 <sup>7</sup> ± 2.08x10 <sup>5</sup>	n.d.
AYFYPEL <sup>d,f,g,i</sup>	α <sub>S1</sub> CN143-149	2.19x10 <sup>5</sup> ± 2.51x10 <sup>4a</sup>	5.54x10 <sup>6</sup> ± 6.29x10 <sup>5b</sup>	1.38x10 <sup>6</sup> ± 1.67x10 <sup>5c</sup>
FYPEL <sup>d,i</sup>	α <sub>S1</sub> CN145-149	5.89x10 <sup>8</sup> ± 5.16x10 <sup>7a</sup>	9.52x10 <sup>8</sup> ± 3.07x10 <sup>7b</sup>	6.43x10 <sup>8</sup> ± 6.22x10 <sup>7a</sup>
IPIQY <sup>h</sup>	κCN 26-30	n.d.	1.28x10 <sup>8</sup> ± 6.83x10 <sup>6a</sup>	6.56x10 <sup>7</sup> ± 1.23x10 <sup>7b</sup>

*Peptides with decreasing trend according to the ripening time*

HKEMPFPK <sup>e</sup>	βCN 106-113	5.39x10 <sup>6</sup> ± 4.36x10 <sup>5a</sup>	3.96x10 <sup>6</sup> ± 2.55x10 <sup>5b</sup>	n.d.
TQTPVVVPPFLQPE <sup>i</sup>	βCN 78-91	1.04x10 <sup>7</sup> ± 8.33x10 <sup>5</sup>	n.d.	n.d.
LVYFPFGPI <sup>d</sup>	βCN 58-66	3.32x10 <sup>7</sup> ± 7.09x10 <sup>5a</sup>	2.20x10 <sup>7</sup> ± 7.67x10 <sup>5b</sup>	1.55x10 <sup>7</sup> ± 3.63x10 <sup>5c</sup>
VYFPFGPI <sup>n</sup>	βCN 59-66	4.62x10 <sup>8</sup> ± 4.34x10 <sup>6a</sup>	3.75x10 <sup>8</sup> ± 7.67x10 <sup>6b</sup>	2.24x10 <sup>8</sup> ± 5.46x10 <sup>6c</sup>
VYFPFGPIPN <sup>d,i</sup>	βCN 59-68	5.35x10 <sup>8</sup> ± 1.86x10 <sup>7a</sup>	4.70x10 <sup>8</sup> ± 8.66x10 <sup>6b</sup>	3.20x10 <sup>8</sup> ± 4.84x10 <sup>6c</sup>
YFPFGPI <sup>d,g,m</sup>	βCN 60-66	6.87x10 <sup>6</sup> ± 1.29x10 <sup>5a</sup>	2.49x10 <sup>6</sup> ± 1.34x10 <sup>5b</sup>	1.79x10 <sup>6</sup> ± 5.68x10 <sup>4c</sup>
YFPFGPIPN <sup>d,f,h</sup>	βCN 60-67	1.13x10 <sup>7</sup> ± 3.80x10 <sup>5a</sup>	3.13x10 <sup>6</sup> ± 2.88x10 <sup>5b</sup>	2.66x10 <sup>6</sup> ± 1.02x10 <sup>5c</sup>
PFPFGPI <sup>o</sup>	βCN 61-66	9.65x10 <sup>7</sup> ± 1.02x10 <sup>6a</sup>	4.84x10 <sup>7</sup> ± 2.01x10 <sup>6b</sup>	3.56x10 <sup>7</sup> ± 1.36x10 <sup>6c</sup>
NIPPLTQTPV <sup>d</sup>	βCN 73-82	1.45x10 <sup>8</sup> ± 1.73x10 <sup>6</sup>	n.d.	n.d.
HLPLP <sup>d</sup>	βCN 134-138	2.27x10 <sup>8</sup> ± 1.06x10 <sup>7</sup>	n.d.	n.d.
LPVPQ <sup>h</sup>	βCN 171-175	1.25x10 <sup>6</sup> ± 1.55x10 <sup>5a</sup>	4.58x10 <sup>5</sup> ± 9.46x10 <sup>4b</sup>	2.07x10 <sup>5</sup> ± 1.29x10 <sup>4c</sup>
FVAPFPEVFG <sup>d</sup>	α <sub>S1</sub> CN 24-33	8.90x10 <sup>6</sup> ± 1.75x10 <sup>6a</sup>	3.21x10 <sup>5</sup> ± 2.94x10 <sup>5b</sup>	5.80x10 <sup>5</sup> ± 4.05x10 <sup>5c</sup>
YFYPE <sup>g</sup>	α <sub>S1</sub> CN 144-148	3.36x10 <sup>7</sup> ± 1.13x10 <sup>6a</sup>	1.59x10 <sup>6</sup> ± 9.18x10 <sup>5b</sup>	2.02x10 <sup>5</sup> ± 1.95x10 <sup>5c</sup>
AMKPW <sup>d</sup>	α <sub>S2</sub> CN 189-193	2.15x10 <sup>7</sup> ± 3.04x10 <sup>6a</sup>	1.32x10 <sup>7</sup> ± 1.08x10 <sup>6b</sup>	1.58x10 <sup>7</sup> ± 6.12x10 <sup>5c</sup>

*Peptides found in constant amount according to the ripening time*

YPVEFPG <sup>g,h</sup>	βCN 114-119	1.40x10 <sup>9</sup> ± 7.92x10 <sup>7ab</sup>	1.44x10 <sup>9</sup> ± 1.98x10 <sup>7a</sup>	1.38x10 <sup>9</sup> ± 2.99x10 <sup>7b</sup>
GPFPI <sup>o</sup>	βCN 203-207	1.96x10 <sup>9</sup> ± 4.74x10 <sup>7a</sup>	1.72x10 <sup>9</sup> ± 9.60x10 <sup>7a</sup>	1.82x10 <sup>9</sup> ± 3.61x10 <sup>7a</sup>
YFYPEL <sup>g,i</sup>	α <sub>S1</sub> CN 144-149	3.35x10 <sup>7</sup> ± 5.18x10 <sup>6a</sup>	4.54x10 <sup>7</sup> ± 7.41x10 <sup>6a</sup>	3.66x10 <sup>7</sup> ± 5.20x10 <sup>6a</sup>
TPEVDDEALEK <sup>e,h</sup>	βLB 125-135	1.69x10 <sup>7</sup> ± 1.93x10 <sup>6a</sup>	2.06x10 <sup>7</sup> ± 2.43x10 <sup>6a</sup>	1.85x10 <sup>7</sup> ± 2.97x10 <sup>6a</sup>

<sup>a</sup>Abbreviations are: CN, casein; LB, lactoglobulin.

<sup>b</sup>One code letter was used for amino acid nomenclature. Potential bioactivities were achieved from MBPDB databases (Nielsen et al., 2017). Anti-hypertensive activity was measured on spontaneously anti-hypertensive rats. Bioactive peptides are labelled as follow: <sup>d</sup>ACE-inhibitory activity; <sup>e</sup>anti-microbial activity; <sup>f</sup>anti-hypertensive activity; <sup>g</sup>opioid; <sup>h</sup>DPPIV-inhibitory activity; <sup>i</sup>antioxidant activity; <sup>h</sup>hypcholesterolemic; <sup>m</sup>immunomodulator; <sup>n</sup>PEP-inhibitory activity; <sup>o</sup>cathepsin B-inhibitory activity. Abbreviations are: ACE, angiotensin converting enzyme; DPPIV, dipeptidyl peptidase IV; PEP, prolyl endopeptidase.


<sup>c</sup>Relative amount was expressed as the area under the peak (AUP)/g of cheese measured from the extracted ion chromatograms (EIC) obtained for each peptide. N.d. means peptide not detected in the sample. Different superscript letters within the same row indicate that the values are significantly different ( $P < 0.05$ ).



Table 3. Amount of bioactive peptides in water-soluble extract (WSPE) and in digested peptidic fractions from Parmigiano Reggiano samples at 12, 18 and 24 months of ripening

<i>Sequence<sup>a</sup></i>	<i>WSPE PR12</i> <i>mg kg<sup>-1</sup></i>	<i>WSPE PR18</i> <i>mg kg<sup>-1</sup></i>	<i>WSPE PR24</i> <i>mg kg<sup>-1</sup></i>	<i>Digested PR12</i> <i>mg kg<sup>-1</sup></i>	<i>Digested PR18</i> <i>mg kg<sup>-1</sup></i>	<i>Digested PR24</i> <i>mg kg<sup>-1</sup></i>
VPP	6.87 ± 0.68 <sup>a</sup>	11.34 ± 1.01 <sup>b</sup>	4.52 ± 0.28 <sup>c</sup>	7.73 ± 0.91 <sup>a</sup>	12.46 ± 0.97 <sup>b</sup>	2.74 ± 0.03 <sup>d</sup>
IPP	1.63 ± 0.82 <sup>a</sup>	4.24 ± 0.85 <sup>b</sup>	0.66 ± 0.05 <sup>c</sup>	3.26 ± 0.21 <sup>b</sup>	5.64 ± 0.12 <sup>d</sup>	0.66 ± 0.23 <sup>ac</sup>
YPFPGPI	n.d.	n.d.	n.d.	20.18 ± 3.00 <sup>a</sup>	8.91 ± 0.74 <sup>b</sup>	6.38 ± 1.39 <sup>c</sup>

<sup>a</sup>One code letter was used for amino acid nomenclature.

N.d. means peptide not detected in the sample






**A**

**WSPE PR 12**

**WSPE PR 18**

**WSPE PR 24**





**B**

**D PR 12**

**D PR 18**

**D PR 24**








**A**

**WSPE PR 12**

**WSPE PR 18**

**WSPE PR 24**



**B**

**D PR 12**

**D PR 18**

**D PR 24**

