

This is the peer reviewed version of the following article:

Mediterranean diet vegetable foods protect meat lipids from oxidation during in vitro gastro-intestinal digestion / Martini, S.; Conte, A.; Bottazzi, S.; Tagliazucchi, D.. - In: INTERNATIONAL JOURNAL OF FOOD SCIENCES AND NUTRITION. - ISSN 0963-7486. - 71:4(2020), pp. 424-439. [10.1080/09637486.2019.1677570]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

05/01/2026 18:06

Mediterranean Diet vegetable foods protect meat lipids from oxidation during *in vitro* gastrointestinal digestion

Serena Martini, Angela Conte, Silvia Bottazzi, Davide Tagliazucchi*

Department of Life Sciences, University of Modena and Reggio Emilia, Via Amendola 2, 42100
Reggio Emilia, Italy

* Corresponding author. Tel.: +39-0522-522060; fax: +39-0522-522053

E-mail address: davide.tagliazucchi@unimore.it (D. Tagliazucchi)

1 **Abstract**

2 Meat lipids oxidation during digestion gives rise to a post-prandial oxidative stress condition, which
3 negatively affects human health. Mediterranean Diet vegetables contain high amount of phenolic
4 compounds, which potentially may reduce the oxidative phenomena during digestion. *In vitro* co-
5 digestion of turkey meat with a typical Mediterranean Diet salad containing tomato, onion, black
6 olives, extra-virgin olive oil (EVOO) and basil, dose-dependently reduced lipid peroxidation. Onion
7 and EVOO were more effective in limiting oxidation than the other foods, resulting in negligible
8 concentrations of lipid hydroperoxides after digestion. Specific phenolic classes dominated the
9 phenolic profile of the different foods, such as flavonols and anthocyanins in onion, phenolic acids
10 in tomato and basil, and tyrosol-derivatives in black olives and EVOO. The correlation between lipid
11 peroxidation inhibition, phenolic constituents and antioxidant properties was evaluated by principal
12 component analysis (PCA). Flavonols and anthocyanin were the major contributors to the bioactive
13 response of vegetable foods.

14 **Keywords:** onion, mass spectrometry, lipid hydroperoxides, flavonols, anthocyanins, antioxidant
15 activity

16 **1. Introduction**

17 Poly-unsaturated fatty acids peroxidation during gastro-intestinal digestion of foods is an oxidative
18 phenomenon, which may result in the generation of toxic compounds such as lipid hydroperoxides
19 and lipid oxidation end-products that might adversely affect human health (Nogueira et al. 2016;
20 Perše, 2013; Sies et al., 2005).

21 Meat is peculiarly vulnerable to lipid oxidation due to its content of poly-unsaturated fatty acids
22 such as linoleic, linolenic, arachidonic, and docosaheptaenoic acids and high concentrations of iron
23 catalysers (Tirosh, et al. 2015). In fact, high intakes of meat are usually associated to an increased
24 risk of colorectal cancer and cardiovascular diseases (Sasso and Latella 2018; Ferguson, 2010;
25 Micha et al. 2010). It has been supposed that this risk may be not caused by meat per se but a
26 consequence of high-fat intake, generation of carcinogens during meat processing as well as
27 oxidation of poly-unsaturated fatty acids that occurs during cooking and gastro-intestinal digestion
28 (Ferguson, 2010; Gorelik et al. 2013; Kanner and Lapidot 2001; Martini et al. 2018). In addition,
29 lipid peroxidation proceeds promptly when the raw meat structure is broken such as after cooking
30 and mastication (Papuc et al. 2017).

31 The formation of hydroxyl ($\text{HO}\bullet$) and perhydroxy ($\text{HOO}\bullet$) radicals as well as the generation of
32 perferrylmyoglobin-containing peptides are considered the main pathways to free radical chain
33 reaction initiation of lipid peroxidation during gastro-intestinal digestion of meat (Carlsen and
34 Skibsted 2004; Oueslati et al. 2016; Tagliazucchi et al. 2010; Martini et al. 2018). Ferrous iron and
35 dissolved oxygen may generate $\text{O}_2^{\bullet-}$, which at low pH such as found in the gastric milieu forms
36 $\text{HOO}\bullet$. Indeed, acidic disproportionation of $\text{O}_2^{\bullet-}$ may produce hydrogen peroxide (H_2O_2) and
37 oxygen (Oueslati et al. 2016). The formation of $\text{HO}\bullet$ is possible by H_2O_2 decomposition, catalysed
38 by ferrous iron (Fenton reaction), or by H_2O_2 reaction with $\text{O}_2^{\bullet-}$ (Haber–Weiss reaction) (Papuc et
39 al. 2017). These reactive species are able to initiate lipid peroxidation by subtracting a hydrogen
40 from lipids and generating a fatty acyl radical ($\text{L}\bullet$). The resulting radical may react with dissolved
41 oxygen to form a hydroperoxyl radical ($\text{LOO}\bullet$), which can further abstract a hydrogen atom from

42 another unsaturated fatty acyl group (LH) producing a new fatty acyl radical (L•) and a lipid
43 hydroperoxide (LOOH). In the presence of ferrous iron (Fe²⁺), lipid hydroperoxide can decompose
44 giving rise to a vast range of volatile and non-volatile compounds, collectively known as advanced
45 lipoxidation end-products (Papuc et al. 2017).

46 Since meat is considered the best dietary source of essential amino acids and contains an array of
47 important micronutrients such as iron, zinc, selenium, potassium and a range of B-vitamins, it is
48 essential for optimal health throughout the lifecycle. Due to this, the elimination of meat from the
49 diet does not seem to be a nutritionally concrete strategy (Binnie et al. 2014). A recent proposed
50 strategy suggests consuming meat with foods rich in antioxidant compounds typical of the
51 Mediterranean Diet to mitigate the production of lipid oxidation toxic compounds during meat
52 digestion (Gorelik et al. 2013; Kanner et al. 2017).

53 Recently, *in vitro* and *in vivo* studies have demonstrated that oxidation during digestion of various
54 type of meat can be reduced when is combined with Mediterranean Diet antioxidant-rich foods such
55 as red wine, herbs, spices and extra-virgin olive oil (Gorelik et al. 2008a; Gorelik et al. 2008b; Van
56 Hecke et al. 2017; Martini et al.,2018).

57 Although Mediterranean Diet pattern is often described as being low in meat intake, in the last
58 twenty years a general increased consumption of meat (especially in pork and poultry) has been
59 observed in Mediterranean countries (Leone et al. 2017; Chamorro et al. 2012). Nevertheless, in the
60 typical Mediterranean cuisine meat is consumed in combination with antioxidant-rich vegetable
61 foods such as tomatoes, onions, herbs and extra-virgin olive oil.

62 Therefore, this study was designed to understand if the combined consumption of a typical
63 Mediterranean Diet salad (containing tomatoes, onions, black olives, fresh basil and extra-virgin
64 olive oil) with grilled turkey meat could affect the oxidative phenomena during *in vitro* gastro-
65 intestinal digestion. Vegetable foods were also characterized for their phenolic profile by LC-ESI-
66 IT-MS/MS and for their antioxidant properties. Moreover, to gain more information about the role
67 of phenolic compounds, co-digestions between grilled turkey meat and extracted phenolic fractions

68 were carried out. Finally, multivariate analysis was applied to investigate the relationships between
69 the phenolic composition, the antioxidant properties and the lipid peroxidation inhibitory activity of
70 tested vegetable foods.

71 **2. Materials and methods**

72 **2.1. Materials**

73 All of the digestive enzymes (α -amylase from porcine pancreas, pepsin from porcine gastric mucosa
74 and pancreatin from porcine pancreas), phenolic standards and reagents for analytical determination
75 were obtained from Sigma-Aldrich (Milan, Italy). The mass spectrometry reagents and solvents for
76 phenolic compounds extraction were obtained from BioRad (Hercules, CA, USA). Turkey breast
77 meat (*pectoralis major*) and vegetables were purchased in a local supermarket (Reggio Emilia,
78 Italy).

79

80 **2.2. Preparation of the Mediterranean Diet salad**

81 The salad was prepared following the typical recipe from South Italy. The Mediterranean Diet salad
82 contained 200 g of tomato, 25 g of pink onion, 25 g of black olives, 10 g of extra-virgin olive oil
83 (EVOO) and 0.5 g of fresh basil. The above quantities were intended as a salad dish consumed with
84 100 g of cooked turkey meat. **Figure S1** shows a visual impact of the proportion of the single
85 vegetables in the Mediterranean Diet salad and of the salad/meat proportion.

86

87 **2.3. In vitro co-digestion of grilled turkey breast meat with the Mediterranean Diet salad and** 88 **determination of lipid hydroperoxides**

89 Turkey breast meat (average size of 10x15x0.4 cm) was grilled at 140°C for 5 min until complete
90 cooking was achieved. After cooking, the meat was cooled on ice and stored at -80°C overnight.
91 Mediterranean Diet salad was prepared by mixing the single fresh vegetables in the proportion
92 reported above. For the digestion, 10 g of frozen meat was homogenized in a laboratory blender
93 together with 26.05 g of the Mediterranean Diet salad. An aliquot of 5 g of the homogenate was
94 then used for the *in vitro* digestion experiments following the protocol previously developed within
95 the COST Action INFOGEST (Minekus et al. 2014). Simulated salivary, gastric, pancreatic and bile
96 fluids were prepared according to Minekus et al. (2014). To simulate the oral phase, 5 g of

97 homogenate were mixed with 5 mL of simulated salivary fluid containing 150 U/mL of porcine α -
98 amylase and incubated for 5 min at 37°C in a rotating wheel (10 rpm). The gastric phase was
99 carried out by adding 10 mL of simulated gastric fluid to the bolus. The pH was adjusted to 3.0 with
100 HCl 6 mol/L and supplemented with porcine pepsin (2000 U/mL of digest). The gastric bolus was
101 then incubated for 120 min at 37°C in a rotating wheel (10 rpm). The intestinal digestion was
102 carried out by adding 10 mL of pancreatic fluid (100 U trypsin activity/mL of digest) and 5 mL of
103 bile fluid (10 mmol/L in the total digest) to the gastric bolus and adjusting the pH to 7.0. The chyme
104 was further incubated for 120 min at 37°C in a rotating wheel (10 rpm).

105 At the end of the digestion, lipid hydroperoxides were extracted by 10-fold dilution in methanol
106 HPLC grade containing 4 mmol/L of butyl-hydroxytoluene (BHT) under slow stirring for 60 min
107 (Tagliazucchi et al. 2010). After centrifugation at 3000g for 15 min at 4°C, the hydroperoxides in
108 the supernatants were determined with the FOX assay at 560 nm adapted to a microplate reader
109 (Nourooz-Zadeh 1999; Martini et al. 2018). The FOX reagent contained 250 μ mol/L of ammonium
110 ferrous sulphate, 100 μ mol/L xylenol orange, 25 mmol/L H₂SO₄, and 4 mmol/L BHT in 90% (v/v)
111 methanol HPLC grade. For the assay, 60 μ L of extracted sample were added to 140 μ L of FOX
112 reagent and incubated for 30 minutes at room temperature. The hydroperoxides content was
113 expressed in nmol H₂O₂ equivalents per g of meat.

114 In the control digestion, 10 g of cooked meat were mixed with 26.05 g of distilled water (in place of
115 the salad) and homogenized as reported above. The *in vitro* digestion was carried out exactly as
116 reported above. At the end of the digestion lipid hydroperoxides were extracted and quantified,
117 representing the amount of lipid hydroperoxides generated during the digestion of meat without
118 vegetables.

119 The dose-response effect of the Mediterranean Diet salad was assessed by homogenising 10 g of
120 cooked meat with 13.025 g of salad (plus 13.025 g of water) and with 6.51 g of salad (plus 19.54 g
121 of water). After that, the homogenates were subjected to *in vitro* digestion and lipid hydroperoxides
122 quantification.

123 Finally, a blank digestion, which included only the gastro-intestinal juices and enzymes and water
124 in place of meat and salad, was carried out to consider the possible impact of the digestive enzymes
125 and fluids in the subsequent analysis.

126

127 ***2.4. Extraction of phenolic compounds from vegetables and extra-virgin olive oil***

128 Phenolic compounds from extra-virgin olive oil (EVOO) were extracted following the procedure
129 reported in Martini et al. (2018). Briefly, 15 grams of EVOO were mixed with 15 mL of a solution
130 of methanol/water (70:30, v/v) and incubated for 120 minutes at 30°C in a rotary wheel. After
131 incubation, the mixture was centrifuged at 3000g for 30 minutes at 4°C. When extraction was
132 completed, the samples were stored on freezer shelves at -20°C and allowed to stand overnight for
133 lipid precipitation and separation. The liquid supernatant containing phenolics was withdrawn and
134 stored at -20°C until analysis.

135 Phenolic compounds from vegetables were extracted adapting the procedure reported in Martini et
136 al. (2017). Vegetables (10 g) were homogenized with 20 mL of methanol/water solution (70:30,
137 v/v) and incubated for 30 min at 37°C. Homogenates were then centrifuged (6000g, 20 min, 4°C)
138 and the collected supernatant filtered on paper. The filtrates were concentrated by a rotary
139 evaporator and re-dissolved in 10 mL of water.

140

141 ***2.5. In vitro co-digestion of grilled turkey breast meat with single salad ingredients and*** 142 ***vegetables phenolic fractions***

143 In these co-digestion experiments, vegetable salad ingredients (tomatoes, onions, black olives,
144 EVOO and fresh basil) were added singularly to the grilled turkey breast meat in the same
145 proportions as found in the Mediterranean Diet salad. For the experiments, 10 g of cooked meat
146 were homogenized in presence of 20 g of tomato (plus 6.05 g of water) or 2.5 g of onion (plus 23.55
147 g of water) or 2.5 g of black olives (plus 23.55 g of water) or 1 g of EVOO (plus 25.05 g of water)
148 or 0.05 g of fresh basil (plus 26 g of water). The proportion meat/ingredients were 200% tomato,

149 25% onion or black olives, 10% EVOO and 0.5% basil respect to meat (w/w). After that, the *in*
150 *vitro* digestions were carried out as reported above.
151 Further experiments were carried out to gain more information about the effect of vegetables and
152 EVOO phenolic compounds on the oxidative phenomena during *in vitro* co-digestion with meat.
153 These co-digestions were carried out as reported above but replacing the vegetable foods or EVOO
154 with the corresponding amount of phenolic fraction.

155

156 ***2.6. Identification and quantification of phenolic compounds by liquid chromatography***
157 ***electrospray ionization ion trap mass spectrometer (LC-ESI-IT-MS)***

158 Phenolic fractions were analyzed on a HPLC Agilent 1200 Series system equipped with a C18
159 column (HxSil C18 Reversed phase, 250×4.6 mm, 5 µm particle size, Hamilton Company, Reno,
160 Nevada, USA) as reported in Martini et al. (2017). The mobile phase consisted of (A) H₂O/formic
161 acid (99:1, v/v) and (B) acetonitrile/formic acid (99:1, v/v). The gradient started at 4% B for 0.5
162 min then linearly ramped up to 30% B in 60 min. The mobile phase composition was raised up to
163 100% B in 1 min and maintained for 5 min in order to wash the column before returning to the
164 initial condition. The flow rate was set at 1 mL/min. After passing through the column, the eluate
165 was split and 0.3 mL/min was directed to an Agilent 6300 ion trap mass spectrometer. Two MS
166 experiments were performed, one in ESI negative ion mode and one using positive ESI ionization
167 (for anthocyanins), under the same chromatographic conditions. Identification of phenolic
168 compounds in all samples was carried out using full scan and data-dependent MS² scanning from
169 *m/z* 100 to 1500.

170 Phenolic compounds were quantified by using representative standards for each identified phenolic
171 class. Flavonols were quantified as quercetin-3-*O*-glucoside or quercetin-3-*O*-rutinoside
172 equivalents. Hydroxybenzoic acids were quantified in protocatechuic acid equivalents.
173 Hydroxycinnamic acids were quantified in coumaric or caffeic or ferulic acid equivalents.
174 Flavanones and flavones were quantified as naringenin-7-*O*-rutinoside equivalents. Tyrosol-

175 derivative were quantified in hydroxytyrosol equivalents. Anthocyanins were quantified as
176 cyanidin-3-*O*-glucoside equivalents.
177 ESI-MS parameters, limits of detection (LOD) and limits of quantification (LOQ) for the different
178 standards were the same as reported in Martini et al. (2017).
179 Quantitative results were expressed as mg of compounds per 100 g of vegetable or EVOO.
180

181 ***2.7. Antioxidant activity determination in vegetables and extra-virgin olive oil phenolic-rich*** 182 ***fraction***

183 The total antioxidant properties of phenolic fractions were analyzed by using five different assays.
184 The radical scavenging ability was assayed by using the ABTS assay according to Re et al. (1999).
185 For the determination of the Fe³⁺ reducing ability, a protocol based on the ferric
186 reducing/antioxidant power (FRAP) assay was utilized (Benzie and Strain 1999). The capacity to
187 scavenge hydroxyl radical and superoxide anion were evaluated according to the methods reported
188 by Martini et al. (2017). The results were expressed as μmol of ascorbic acid equivalent/mg of
189 phenolic compounds. The Fe²⁺-chelation ability of phenolic-rich fractions was evaluated by the
190 ferrozine assay (Karama and Pegg 2009).
191

192 ***2.8. Statistics***

193 All the digestions were carried out in triplicate and data are presented as mean ± SD for three
194 analytical replicates for each prepared sample. Univariate analysis of variance (ANOVA) with
195 Tukey's post-hoc test was applied using Graph Pad prism 6.0 (GraphPad Software, San Diego, CA,
196 U.S.A.) when multiple comparisons were performed. The differences were considered significant
197 with $P < 0.05$.

198 3. Result and discussion

199 3.1. Effect of Mediterranean Diet salad on lipid oxidation during co-digestion with turkey breast 200 meat

201 An eight-fold increase (from 33.9 ± 3.1 to 277.5 ± 16.1 nmol H₂O₂/g of meat) in the amount of lipid
202 hydroperoxides was observed after the *in vitro* gastro-intestinal digestion of turkey breast meat
203 without added vegetables. Whereas numerous studies determined the amount of lipid
204 hydroperoxides after *in vitro* gastric digestion (Kanner and Lapidot 2001; Gorelik et al. 2018a;
205 Tagliazucchi et al. 2010), very few of them measured their concentration after *in vitro* intestinal
206 digestion. However, a recent study by our research group showed a sharp increase in the generation
207 of lipid hydroperoxides during intestinal digestion of cooked turkey meat (Martini et al. 2018). This
208 increase could be a consequence of the bile salts emulsification and micellarization of fatty acids
209 present in turkey meat. Berton-Carabin et al. (2014) found that lipid peroxidation occurred much
210 faster in a water/oil system rather than in oil alone or in dispersion without emulsifier. This may be
211 due to several factors. First of all the formation of an interface between the aqueous phase and the
212 fat may favour the contact between the oxidants and the oxygen (dissolved in the aqueous phase)
213 and fatty acids (Berton-Carabin et al. 2014). Secondly, it could be ascribed to the solubilisation of
214 lipid hydroperoxides already formed in the micelles, which in turn may promote oxidation in the
215 micelles themselves (Donnelly et al. 1998). Finally, Sreejayan and von Ritter (1998) suggested that
216 bile salts, in the presence of iron, were able to favour the oxidation of arachidonic acid. The amount
217 of lipid hydroperoxides measured was about 23% lower than that found by Martini et al. (2018) at
218 the end of the intestinal digestion.

219 Data in **Figure 1** show that lipid hydroperoxides production from turkey meat after gastro-intestinal
220 digestion was greatly reduced by including increasing amounts of the Mediterranean Diet salad
221 mixture and was totally inhibited when meat and salad were co-digested in the original proportion
222 (260.5 g of salad/100 g of meat). Halving the amount of the Mediterranean Diet salad (130.3 g of

223 salad/100 g of meat) also resulted in a complete inhibition in the formation of lipid hydroperoxides
224 whereas further halving (65.2 g of salad/100 g of meat) produced an inhibition of 49.5% (**Figure 1**).
225 No previous data were found in literature about the inhibitory activity of food combination (e.g.
226 salads) on the generation of lipid hydroperoxides after gastro-intestinal digestion of meat. However,
227 in agreement with our results, Kanner and co-worker (2017) reported an inhibition of about 90% in
228 the formation of malondialdehyde after *in vitro* gastric digestion of meat with a Greek salad (274 g
229 of salad/200 g of meat) composed of tomato, cucumber, red pepper, green-cabbage, onion and black
230 olives.

231 Several authors demonstrated the correlation between lipid peroxidation during *in vitro* digestion of
232 meat, with or without phenolic-rich foods, and the concentrations of lipid hydroperoxides and
233 lipoxidation end-products in the plasma of human volunteers after consumption of the same test
234 meals (Natella et al. 2011; Kanner et al. 2001; Gorelik et al. 2008a; Sirota et al. 2013). Therefore,
235 the results of the present *in vitro* digestion study are likely to be relevant for the *in vivo* situation.

236

237 ***3.2. Effect of the single components of the Mediterranean Diet salad on lipid oxidation during*** 238 ***co-digestion with turkey breast meat***

239 To understand which component of the Mediterranean Diet salad was mainly responsible for the
240 observed inhibitory effect, we carried out co-digestion with turkey meat and each single
241 components of the salad in the same proportion as found in the Mediterranean Diet salad itself. As
242 reported in section 2.2, the Mediterranean Diet salad, related to 100 g of meat, consisted of 200 g of
243 tomato, 25 g of pink onion, 25 g of black olives, 10 g of extra-virgin olive oil (EVOO) and 0.5 g of
244 fresh basil. This means that, for example, in the co-digestion between turkey meat and tomato, the
245 proportion between meat and tomato was 100 g of meat and 200 g of tomato (200% of tomato
246 respect to meat; w/w). Basing on the same rationale, onion or black olives were added in the
247 proportion of 25% respect meat (w/w), EVOO in the proportion of 10% respect to meat (w/w) and
248 basil in the proportion of 0.5% respect to meat (w/w).

249 When turkey breast meat was co-digested with the single components of the Mediterranean Diet
250 salad, we observed a differential inhibition in lipid hydroperoxides generation (**Figure 2A**). Except
251 for fresh basil, the addition of all the vegetable components of the salad during co-digestion with
252 turkey meat resulted in a significantly lower amount of generated lipid hydroperoxides. Among the
253 different vegetables, digests of turkey meat with 25% onion and 10% EVOO had undetectable
254 levels of lipid hydroperoxides (100% inhibition). Tomato, although present in a higher amount (200
255 g per 100 g of meat) respect to onion and EVOO (25 g and 10 g per 100 g of meat, respectively),
256 showed a lower inhibitory effect ($P<0.05$) on the generation of lipid hydroperoxides during co-
257 digestion with turkey meat (~ 75% of inhibition). The inhibition in lipid hydroperoxides formation
258 was significantly lower ($P<0.05$) when 25% of black olives were added to the cooked turkey meat
259 in the digestion system (~ 62% of inhibition). Finally, no significant differences were found in the
260 amount of lipid hydroperoxides in the digests when 0.5% of fresh basil was added to turkey meat
261 ($P>0.05$).

262 Previous studies reported the ability of EVOO and onion to inhibit lipid peroxidation during *in vitro*
263 digestion of meat (Martini et al. 2018; Kanner et al. 2017; Tirosh et al. 2015). Kanner et al. (2017)
264 also found that tomato inhibited with less effectiveness lipid peroxidation during *in vitro* gastric
265 digestion of turkey meat respect to onion, black olives and EVOO. Differently from our study, they
266 observed a higher effectiveness of black olives respect to onion. However, they measured the lipid
267 peroxidation inhibitory activity at the end of the gastric phase of digestion and not after the
268 intestinal phase. Van Hecke et al. (2017) found that dried basil inhibited lipid peroxidation after *in*
269 *vitro* intestinal digestion of beef when added in the amounts of 0.5% or 1% respect to meat. Instead,
270 in this study we found that the addition of 0.5% of fresh basil had no effect on lipid peroxidation
271 during *in vitro* digestion of turkey meat. This difference was clearly related to the fact that the same
272 amount (0.5%) of dried basil delivered more antioxidant compounds to the digestive system respect
273 to the fresh herb (Henning et al. 2011).

274

275 **3.3. Effect of the phenolic fractions of single components of the Mediterranean Diet salad on**
276 **lipid oxidation during co-digestion with turkey breast meat**

277 Previous studies described a strong correlation between the concentration of total phenolic
278 compounds in fruit, beverages, vegetables and spices and the reduction in the lipid peroxidation
279 during *in vitro* digestion of meat (Kanner et al. 2017; Van Hecke et al. 2017; Martini et al. 2018).
280 Therefore, the phenolic fractions extracted from the different vegetables were co-digested with
281 turkey breast meat (**Figure 2B**).

282 Phenolic fractions extracted from onion and EVOO and co-digested with meat at the same
283 concentration as found in 25% onion and 10% EVOO totally inhibited the generation of lipid
284 hydroperoxides without any differences with the data obtained after the co-digestion of meat with
285 whole onion or EVOO ($P>0.05$). Similarly, co-digestion of turkey breast meat with phenolic
286 fractions extracted from black olives and fresh basil resulted in the same inhibitory potency as the
287 whole foods ($P>0.05$). However, in the case of tomato phenolic fraction, the effect was less
288 pronounced respect to that observed after co-digestion with whole food (58% vs 75% of inhibition;
289 $P<0.05$). Despite phenolic compounds, tomato is rich in other lipophilic antioxidants such as
290 carotenoids and in particular lycopene (Martí et al. 2016). Previous studies described lycopene as an
291 efficient inhibitor of lipid peroxidation both in meat products and in cell cultures (Rohlík et al.
292 2013; Chisté et al. 2014). Lycopene is highly hydrophobic and not extracted in the water/methanol
293 mixture used to prepare the phenolic fraction from tomato. Moreover, it can be released during the
294 intestinal phase of digestion and exert its anti-peroxidative effect (Tagliazucchi et al. 2012).

295

296 **3.4. Phenolic profile of vegetables and EVOO**

297 The phenolic profile of vegetables and EVOO was investigated using a non-targeted method
298 through LC-ESI-MS/MS experiments. The mass spectrum data along with peak assignments and
299 retention time for the identified phenolic compounds are described in **Table 1**. This approach
300 allowed the tentative identification of 132 compounds (**Table 1**). Seven compounds (compounds 4,

301 **23, 26, 40, 53, 95** and **99**) were identified by comparison with their respective authentic standards.
 302 The remaining compounds were tentatively identified based on the interpretation of the
 303 fragmentation patterns obtained from mass spectra and by comparing their mass spectral
 304 characteristics with the available literature. The interpretation of the mass spectra fragmentation
 305 patterns reported in the literature is not further discussed.
 306 The profile of individual phenolic compounds as well as the total phenolic compounds amount for
 307 each vegetable and EVOO are reported in **Table 2** and **Figure 3**. The highest phenolic content was
 308 found in EVOO > tomato > black olives > onion > fresh basil ($P < 0.05$). Each single ingredients
 309 was characterized for its specific phenolic profile (**Figure 3**). In tomato, hydroxycinnamic acids
 310 were the predominant class of phenolic compounds (94.4%) whereas in fresh basil hydroxybenzoic
 311 acids prevailed (63.3%) respect to hydroxycinnamic acids (30.3%). The phenolic profile of onion
 312 was mainly characterized by flavonols (58.3%) and anthocyanins (21.9%). EVOO and black olives
 313 were characterized for the presence of tyrosol-derivatives. Concerning the individual phenolic
 314 profile, sixty-five phenolic compounds were identified in tomato, which was characterized by the
 315 presence of relevant amounts of di-hydro-ferulic acid-*O*-hexoside (**47**), caffeic acid-*O*-hexoside
 316 (**25**) and 4- and 5-*O*-caffeoylquinic acids (**56** and **51**). Apart from hydroxycinnamic acids, modest
 317 amounts of hydroxybenzoic acids and flavonols were detected in tomato. Flavanones were present
 318 only in tomato but in very low concentrations. Thirty-five phenolic compounds were identified and
 319 quantified in onion (**Table 2**). Flavonols were the major group of phenolic compounds identified in
 320 onion. Quercetin-3-*O*-hexoside-4'-*O*-hexoside (**82**), quercetin-4'-*O*-hexoside (**119**) and cyanidin-3-
 321 *O*-malonylhexoside (**64**) were the main individual phenolics. With respect to black olives, a higher
 322 prevalence of tyrosol-derivatives was noted (**Table 2**). Twenty-seven phenolic compounds were
 323 quantified in black olives, with oleuropein aglycone (**70**) and hydroxytyrosol-*O*-hexoside isomers (**5**
 324 and **8**) present in high concentrations. Similar to what was reported for black olives, tyrosol-
 325 derivatives were the major group of phenolic compounds in EVOO, where oleuropein aglycone
 326 (**118**) and ligstroside aglycone (**64**) were the main individual phenolics (**Table 2**). Finally, twenty-

327 six individual phenolic compounds were identified and quantified in fresh basil with a prevalence of
328 phenolic acids. Syringic acid-4-*O*-pentoside (**36**), protocatechuic acid-*O*-hexoside-*O*-pentoside (**22**)
329 and ferulic acid-4-*O*-pentoside isomers (**73** and **74**) were the main phenolic acids.

330 331 **3.4. Antioxidant properties of vegetable and EVOO phenolic fractions**

332 Vegetables and EVOO phenolic fractions were characterized for their ability to scavenge
333 superoxide anions and hydroxyl radicals as well as for their total radical scavenging capacity
334 (ABTS assay). Moreover, their ability to chelate Fe²⁺ and their ferric reducing properties were
335 assessed (**Table 3**). Black olives and onion phenolic compounds showed the highest ABTS and
336 hydroxyl radical scavenging activities. Instead, tomato phenolic compounds displayed the highest
337 ability to scavenge superoxide anions and the highest ferric reducing ability. With respect to the
338 Fe²⁺-chelating ability, black olives and fresh basil phenolic compounds were the most active.
339 The different antioxidant properties of the phenolic fractions reflect differences in their phenolic
340 compositions (Martini et al. 2019). Onion and black olives were found to be particularly rich in
341 quercetin- and cyanidin-derivatives and hydroxytyrosol-derivatives, respectively. These compounds
342 share a 3',4'-dihydroxy structure in the B-ring (i.e. catechol moiety) which is considered of
343 paramount importance to determine the ABTS and hydroxyl radical scavenging properties (Rice-
344 Evans et al. 1999, Ozyürek et al. 2008, Zamora and Hidalgo 2016). Diversely, tomato and fresh
345 basil showed significantly lower ABTS and hydroxyl radical scavenging properties than onion and
346 black olives. Indeed, they were rich in phenolic acids, which showed the lowest hydroxyl radical
347 and ABTS scavenging activities among phenolic compounds (Rice-Evans et al. 1999, Ozyürek et al.
348 2008). Differences between black olives and EVOO hydroxyl radical and ABTS scavenging
349 activities may be related to the presence of additional antioxidant compounds in black olives. The
350 latter contain also non-phenolic compounds such as oleoside and its methyl- and dimethyl-
351 derivatives, which showed radical scavenging properties (Wang et al. 2000). Hydroxycinnamic
352 acids seemed to be the best superoxide anions (O₂•⁻) scavenging phenolic compounds since the

353 most active extract against $O_2^{\bullet-}$ was tomato which was mainly consisted of hydroxycinnamic acids
354 (**Figure 3**). Moreover, hydroxycinnamic acids were also the compounds with the highest ferric
355 reducing properties as suggested by the highest ferric reducing power of tomato and fresh basil
356 phenolic fractions. Previous works indicated that hydroxycinnamic and hydroxybenzoic acids
357 displayed higher ferric reducing activities than flavan-3-ols and glycosylated flavonols (Pulido et al.
358 2000; Martini et al. 2019). Finally, no clear relationship was found between the phenolic
359 composition and the Fe^{2+} -chelating ability of the phenolic fractions extracted from vegetables and
360 EVOO.

361

362 ***3.5. Relationship between the lipid peroxidation inhibitory activity, the phenolic profile and the*** 363 ***antioxidant properties of phenolic fractions extracted from vegetables and EVOO***

364 Principal component analysis (PCA) was performed as exploratory analysis allowing data
365 comprehension, clusters association and a quick network identification between phenolic
366 compounds determined by LC-MS/MS, the antioxidant properties and the lipid peroxidation
367 inhibitory activity of vegetables and EVOO. This approach can help to describe the variance
368 (information) in a set of multivariate data where the original variables (here: phenolic classes) may
369 be expressed as linear combination of orthogonal principal components (PCs).

370 Three principal components explained about 90.5% of total variance. In particular, a bidimensional
371 plot (PC1xPC2 biplot) was reported (**Figure 4**), recording the 63.3% cumulative percentage of the
372 total variance. **Figure 4** shows a clear separation of the phenolic-rich food ingredients, described by
373 the respective and representative phenolic classes. In fact, aiming to fully understand the causative
374 variables for the obtained distribution and the correlation between phenolic classes and
375 bioactivities, they were added to the bidimensional plot. ABTS and hydroxyl radical scavenging
376 activities and the inhibition of the lipid peroxidation displayed the same negative loading vectors on
377 PC1, positively correlated to the onion and its typical phenolic classes: anthocyanins and flavonols.
378 This reflects their higher effectiveness in antioxidant and lipid peroxidation inhibitory activities

379 than the other ingredients or phenolic classes. Regarding this, the orthogonal directions of
380 hydroxycinnamic and hydroxybenzoic acids did not suggest any kind of relationship. An inverse
381 relationship between ferric reducing power and the inhibition of the lipid peroxidation is depicted
382 by FRAP loading on PC1. The explanation could lay in the mechanisms of action of the used
383 antioxidant activity assays. According to the chemistry of the ABTS and hydroxyl radical
384 scavenging assays, their mechanisms may involve both the single electron transfer (SET) and
385 hydrogen atom transfer (HAT) (Prior et al. 2005); whereas FRAP assay is only characterized by
386 single electron transfer mechanism. Indeed, the capacity to reduce Fe^{3+} to Fe^{2+} may retain the
387 optimal conditions to maintain and stimulate the Fenton and Haber-Weiss reactions. Whereas, the
388 HAT mechanism might stop the lipid peroxidation reaction at several levels. Tyrosol- and
389 hydroxytyrosol-derivatives, describing the phenolic profile of black olives and EVOO, had the same
390 negative loadings on PC1 of the lipid peroxidation inhibition, reflecting their possible involvement
391 in the peroxidation phenomena. However, the negative loading vectors on PC2 could reflect their
392 intrinsic and paradoxical behaviour already investigated in Martini et al. (2018) outlining how
393 tyrosol- and hydroxytyrosol-derivatives peroxidation inhibitory activity is strictly related to their
394 final concentration.

395 4. Conclusions

396 This study provides evidence of a protective effect of a typical Mediterranean Diet salad on lipid
397 peroxidation during co-digestion of turkey breast meat. The co-digestion carried out with the single
398 ingredients and phenolic extracts of the Mediterranean Diet salad displayed differences in the lipid
399 peroxidation inhibitory effect. With the exception of tomato, there were not significant differences
400 between the inhibitory effect of the whole ingredients and the respective phenolic fractions,
401 implying that phenolic compounds were mainly responsible for the reported effect. Moreover, our
402 data suggested that the inhibitory effect was related to the different phenolic composition of the
403 tested ingredients and that some phenolic compounds, especially that with a B-ring catechol moiety
404 in their structure (i.e. flavonols and anthocyanins), were the most effective in reducing the oxidative
405 phenomena after co-digestion with meat. This effect was ascribed to the highest radical scavenging
406 and hydroxyl radical scavenging activities of these compounds. On the contrary, phenolic acids,
407 which showed the highest ability to reduce Fe^{3+} to Fe^{2+} , exhibited the lowest lipid peroxidation
408 inhibitory effect. This study gives strong evidence about the structure-activity relationship between
409 phenolic compounds and lipid peroxidation inhibitory activity. Therefore, it is of paramount
410 importance to profile the phenolic composition of antioxidant-rich foods used in this type of study
411 to predict their possible impact on lipid peroxidation during the digestion of meat. Indeed, our study
412 underlines the importance of consuming specific food combinations, in specific amounts to achieve
413 significant biological effects.

414 Lipid peroxidation inhibitory properties of phenolic compounds in the gastro-intestinal tract, during
415 a meal, may play a key role in the health effect of the Mediterranean Diet. The maintenance of the
416 right redox balance in the gastro-intestinal tract by phenolic-rich foods seems to be a concrete
417 nutritional strategy for healthy living.

418 **Disclosure Statement**

419 The authors report no conflict of interest.

Financial support

This work was supported by a grant from Department of Life Sciences, University of Modena and Reggio Emilia (research project FAR2016 “Dieta Mediterranea e salute: riduzione dei fenomeni ossidativi durante la digestione della carne”). The authors acknowledge the Fondazione Cassa di Risparmio di Modena for funding the HPLC-ESI-IT system at the Centro Interdipartimentale Grandi Strumenti (CIGS).

References

- Benzie IFF, Strain JJ. 1999. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol.* 299:15-27.
- Berton-Carabin CC, Ropers MH, Genot C. 2014. Lipid oxidation in oil-in-water emulsions: Involvement of the interfacial layer. *Compr Rev Food Sci Food Saf.* 13:945-977.
- Binnie MA, Barlow K, Johnson V, Harrison C. 2014. Red meats: time for a paradigm shift in dietary advice. *Meat Sci.* 98:445-451.
- Carlsen CU, Skibsted LH. 2004. Myoglobin species with enhanced prooxidative activity is formed during mild proteolysis by pepsin. *J Agric Food Chem.* 52:1675-1781.
- Chamorro A, Miranda FJ, Rubio S, Valero V. 2012. Innovations and trends in meat consumption: An application of the Delphi method in Spain. *Meat Sci.* 92:816-822.
- Chisté RC, Freitas M, Mercadante AZ, Fernandes E. 2014. Carotenoids inhibit lipid peroxidation and hemoglobin oxidation, but not the depletion of glutathione induced by ROS in human erythrocytes. *Life Sci.* 99:52-60.
- Donnelly JL, Decker EA, McClements DJ. 1998. Iron-catalyzed oxidation of Menhaden oil as affected by emulsifiers. *J Food Sci.* 63:997-1000.
- Ferguson LR. 2010. Meat and cancer. *Meat Sci.* 84:308-313.
- Gorelik S, Ligumsky M, Kohen R, Kanner J. 2008a. The stomach as a “bioreactor”: when red meat meets red wine. *J Agric Food Chem.* 56:5002-5007.
- Gorelik S, Ligumsky M, Kohen R, Kanner J. 2008b. A novel function of red wine polyphenols in humans: prevention of absorption of cytotoxic lipid peroxidation products. *FASEB J.* 22:41-46.
- Gorelik S, Kanner J, Schurr D, Kohen R. 2013. A rational approach to prevent postprandial modification of LDL by dietary polyphenols. *J Funct Foods.* 5:163-169.

- Henning SM, Zhang Y, Seeram NP, Lee RP, Wang P, Bowerman S, Heber D. 2011. Antioxidant capacity and phytochemical content of herbs and spices in dry, fresh and blended herb paste form. *Int J Food Sci Nutr.* 62:219-225.
- Kanner J, Lapidot T. 2001. The stomach as a bioreactor: Dietary lipid peroxidation in the gastric fluid and the effects of plant-derived antioxidants. *Free Radic Biol Med.* 31:1388-1395.
- Kanner J, Selhub J, Shpaizer A, Rabkin B, Schacham I, Tirosh O. 2017. Redox homeostasis in stomach medium by foods: The Postprandial Oxidative Stress Index (POSI) for balancing nutrition and human health. *Redox Biol.* 12:926-936.
- Karama M, Pegg RB. 2009. Limitations of the tetramethylmurexide assay for investigating the Fe(II) chelation activity of phenolic compounds. *J Agric Food Chem.* 57:6425-6431.
- Leone A, Battezzati A, De Amicis R, De Carlo G, Bertoli S. 2017. Trends of adherence to the Mediterranean dietary pattern in Northern Italy from 2010 to 2016. *Nutrients.* 11:E734.
- Martí R, Roselló S, Cebolla-Cornejo J. 2016. Tomato as a source of carotenoids and polyphenols targeted to cancer prevention. *Cancers.* 20:E58.
- Martini S, Conte A, Tagliazucchi D. 2017. Phenolic compounds profile and antioxidant properties of six sweet cherry (*Prunus avium*) cultivar. *Food Res Int.* 97:15-26.
- Martini S, Cavalchi M, Conte A, Tagliazucchi D. 2018. The paradoxical effect of extra-virgin olive oil on oxidative phenomena during *in vitro* co-digestion with meat. *Food Res Int.* 109:82-90.
- Martini S, Conte A, Tagliazucchi D. 2019. Bioactivity and cell metabolism of *in vitro* digested sweet cherry (*Prunus avium*) phenolic compounds. *Int J Food Sci Nutr.* 70:335-348.
- Micha R, Wallace SK, Mozaffarian D. 2010. Red and processed meat consumption and risk of incident coronary heart disease, stroke, and diabetes mellitus. *Circulation.* 121:2271-2283.
- Minekus M, Alming M, Alvito P, Ballance S, Bohn T, Bourlieu C, et al. 2014. A standardised static *in vitro* digestion method suitable for food – an international consensus. *Food Funct.* 5:1113-1124.

- Natella F, Macone A, Ramberti A, Forte M, Mattivi F, Matarese RM, Scaccini C. 2011. Red wine prevents the postprandial increase in plasma cholesterol oxidation products: A pilot study. *Br J Nutr.* 105:1718-1723.
- Nogueira MS, Kessuane MC, Lobo Ladd AA, Lobo Ladd FV, Cogliati B, Castro IA. 2016. Effect of long-term ingestion of weakly oxidised flaxseed oil on biomarkers of oxidative stress in LDL-receptor knockout mice. *Br J Nutr.* 116: 258-269.
- Nourooz-Zadeh J. 1999. Ferrous ion oxidation in the presence of xylenol orange for detection of lipid hydroperoxides in plasma. *Methods Enzymol.* 300:58-62.
- Oueslati K, de La Pomélie D, Santé-Lhoutellier V, Gatellier P. 2016. Impact of Fenton process in meat digestion as assessed using an *in vitro* gastro-intestinal model. *Food Chem.* 209:43-49.
- Ozyürek M, Bektaşoğlu B, Güçlü K, Apak R. 2008. Hydroxyl radical scavenging assay of phenolics and flavonoids with a modified cupric reducing antioxidant capacity (CUPRAC) method using catalase for hydrogen peroxide degradation. *Anal Chim Acta.* 616:196–206.
- Papuc C, Goran GV, Predescu CN, Nicorescu V. 2016. Mechanisms of oxidative processes in meat and toxicity induced by postprandial degradation products: A Review. *Compr Rev Food Sci Food Saf.* 16:96-123.
- Perše M. 2013. Oxidative stress in the pathogenesis of colorectal cancer: Cause or consequence? *BioMed res Int.* 725710.
- Prior RL, Wu X, Schaich K. 2005. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements *J Agric Food Chem.* 53:4290-4302.
- Pulido R, Bravo L, Saura-Calixto F. 2000. Antioxidant activity of dietary polyphenols as determined by a modified ferric reducing/antioxidant power assay. *J Agric Food Chem.* 48:3396–3402.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med.* 26:1231-1237.

- Rice-Evans CA, Miller NJ, Paganga G. 1996. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med.* 20:933–956.
- Rohlík BA, Pipek P, Pánek J. 2013. The effect of natural antioxidants on the colour and lipid stability of paprika salami. *Czech J Food Sci.* 31:307-312.
- Sasso A, Latella G. 2018. Dietary components that counteract the increased risk of colorectal cancer related to red meat consumption. *Int J Food Sci Nutr.* 69:536-548.
- Sies H, Stahl W, Sevanian A. 2005. Nutritional, dietary and postprandial oxidative stress. *J Nutr.* 135:969-972.
- Sirota R, Gorelik S, Harris R, Kohen R, Kanner J. 2013. Coffee polyphenols protect human plasma from postprandial carbonyl modifications. *Mol Nutr Food Res.* 57:916-919.
- Sreejayan N, von Ritter C. 1998. Effect of bile acids on lipid peroxidation: The role of iron. *Free Radic Biol Med.* 25:50-56.
- Tagliazucchi D, Verzelloni E, Conte A. 2010. Effect of dietary melanoidins on lipid peroxidation during simulated gastric digestion: Their possible role in the prevention of oxidative damage. *J Agric Food Chem.* 58:2513-2519.
- Tagliazucchi D, Verzelloni E, Conte A. 2012. The first tract of alimentary canal as an extractor. Release of phytochemicals from solid food matrices during simulated digestion. *J Food Biochem.* 36:555-568.
- Tirosh O, Shpaizer A, Kanner J. 2015. Lipid peroxidation in a stomach medium is affected by dietary oils (Olive/Fish) and antioxidants: The Mediterranean versus Western diet. *J. Agric. Food Chem.* 63:7016-7023.
- Van Hecke T, Ho PL, Goethals S, De Smet S. 2017. The potential of herbs and spices to reduce lipid oxidation during heating and gastrointestinal digestion of a beef product. *Food Res Int.* 102:785-792.
- Wang H, Gan D, Zhang X, Pan Y. 2010. Antioxidant capacity of the extracts from pulp of *Osmanthus fragans* and its components. *LWT Food Sci Technol.* 43:319-325.

Zamora R, Hidalgo FJ. 2016. The triple defensive barrier of phenolic compounds against lipid oxidation-induced damage in food products. *Trends Food Sci Technol.* 54:165-174.

Figure captions

Figure 1. Turkey breast meat lipid peroxidation as affected by Mediterranean Diet salad after *in vitro* gastro-intestinal digestion. A portion of Mediterranean Diet salad contained 200 g of tomato, 25 g of pink onion, 25 g of black olives, 10 g of extra-virgin olive oil and 0.5 g of fresh basil. The above quantities were intended as a salad dish consumed with 100 g of cooked turkey meat. Lipid hydroperoxides were expressed as nmol H₂O₂/g of meat. Different letters indicate that the values are significantly different ($P < 0.05$). n.d. means not detected.

Figure 2. Effect of the Mediterranean Diet salad ingredients and their phenolic-rich fractions on the amount of lipid hydroperoxides measured at the end of the gastro-intestinal digestion of turkey breast meat. Black column represents meat alone. Light grey columns represent the co-digestion of meat with the different food ingredients. Dark grey columns represent the co-digestion of meat with the phenolic fractions extracted from the different food ingredients. The tested ingredients were tomato (200g/100g of meat), onion (25 g/100 g meat), black olives (25 g /100 g of meat), EVOO (10 g /100 g of meat) and fresh basil (0.5 g/100 g of meat). Lipid hydroperoxides were expressed as nmol H₂O₂/g of meat. EVOO: extra-virgin olive oil. Different letters indicate that the values are significantly different ($P < 0.05$). n.d. means not detected.

Figure 3. Occurrence of phenolic classes in the tested vegetables and EVOO. (A) Incidence of phenolic classes in tomato, onion, fresh basil and black olives. (B) Incidence of individual tyrosol-derivatives in black olives and EVOO. EVOO: extra-virgin olive oil.

Figure 4. Principal component analysis of vegetable and EVOO phenolic-rich fractions activities, phenolic classes and lipid peroxidation inhibitory activity. Graph of the biplot of PC1 versus PC2. The symbol ♦ identifies the phenolic classes and the biochemical properties, whereas the symbol ▲ represents the food ingredients. FRAP: ferric reducing power; ABTS: ABTS radical scavenging activity; HO•: hydroxyl radical scavenging activity; O₂•⁻: superoxide anion radical scavenging activity; EVOO: extra-virgin olive oil.

Table 1. Mass spectral characteristics of phenolic compounds identified in the studied vegetables

	<i>Compound</i>	<i>R_t</i> (min)	<i>[M-H]⁻</i> (m/z)	<i>MS² ion fragments (m/z)</i>
1	Hydroxybenzoic acid- <i>O</i> -hexoside isomer	8.7	299	137 (100%)
2	Hydroxybenzoic acid-dihexoside	9.9	461	137 (100%), 299 (62%)
3	Hydroxytyrosol-di- <i>O</i> -hexoside	10.7	477	153 (100%), 315 (53%), 123 (10%)
4	Hydroxytyrosol isomer*	11.2	153	123 (100%)
5	Hydroxytyrosol- <i>O</i> -hexoside isomer	11.3	315	153 (100%), 123 (17%)
6	Caffeoylquinic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside isomer	11.6	677	515 (100%), 353 (23%), 191 (5%)
7	Vanillic acid-4- <i>O</i> -hexoside	11.8	329	167 (100%), 152 (9%)
8	Hydroxytyrosol- <i>O</i> -hexoside isomer	12.0	315	153 (100%), 123 (25%)
9	Hydroxytyrosol isomer	12.1	153	123 (100%)
10	Protocatechuic acid- <i>O</i> -hexoside	12.8	315	153 (100%)
11	Caffeic acid- <i>O</i> -hexoside- <i>O</i> -pentoside isomer	13.7	473	341 (100%), 179 (48%), 135 (7%)
12	Syringic acid-4- <i>O</i> -hexoside	14.1	359	197 (100%), 182 (8%), 167 (5%)
13	Calceolarioside	14.1	477	323 (100%), 315 (90%), 161 (16%)
14	Caffeoylquinic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside isomer	14.1	677	515 (100%), 341 (21%)
15	Caffeoylquinic acid- <i>O</i> -hexoside isomer	14.9	515	341 (100%), 323 (64%), 179 (57%), 353 (34%)
16	Caffeic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside	15.5	503	341 (100%), 179 (17%)
17	3- <i>O</i> -Caffeoylquinic acid	15.6	353	191 (100%), 179 (24%), 135 (19%)
18	Caffeoylquinic acid- <i>O</i> -hexoside isomer	15.6	515	353 (100%), 191 (82%), 179 (12%) 323 (8%)
19	Caffeoyl-coumaroylquinic acid	15.7	499	337 (100%), 173 (36%), 191 (34%)
20	Rosmarinic acid	15.8	359	197 (100%), 161 (6%), 153 (8%)
21	Gallic acid	15.9	169	125 (100%)
22	Protocatechuic acid- <i>O</i> -hexoside- <i>O</i> -pentoside	16.7	447	315 (100%), 271 (43%), 153 (15%)
23	Coumaric acid*	16.7	163	119 (100%)
24	Coumaric acid- <i>O</i> -hexoside isomer	16.9	325	163 (100%), 119 (24%)
25	Caffeic acid- <i>O</i> -hexoside isomer	17.0	341	179 (100%), 135 (36%)
26	Caffeic acid*	17.1	179	135 (100%)
27	Hydroxybenzoic acid- <i>O</i> -hexoside isomer	17.8	299	137 (100%)
28	Di-hydro-coumaric acid- <i>O</i> -hexoside	18.0	327	165 (100%), 121 (4%)
29	Di-hydro-caffeic acid- <i>O</i> -hexoside isomer	18.0	343	181 (100%), 137 (33%)
30	Syringic acid	18.2	197	153 (100%)

31	Caffeic acid- <i>O</i> -hexoside- <i>O</i> -pentoside isomer	18.2	473	341 (100%), 179 (6%), 135 (4%)
32	Feruloylquinic acid- <i>O</i> -hexoside isomer	18.6	529	367 (100%), 191 (17%)
33	Caffeoyl-6 β -hexose	18.7	341	281 (100%), 179 (83%), 251 (42%), 135 (14%), 323 (12%), 221 (9%)
34	Apigenin- <i>O</i> -hexoside	18.8	431	269 (100%)
35	Protocatechuic acid- <i>O</i> -pentoside	19.3	285	153 (100%), 109 (20%)
36	Syringic acid-4- <i>O</i> -pentoside	19.3	329	197 (100%), 182 (6%), 153 (2%)
37	Di-hydro-caffeic acid- <i>O</i> -hexoside isomer	19.5	343	181 (100%), 137 (9%)
38	Caffeoylquinic acid- <i>O</i> -hexoside isomer	19.8	515	323 (100%), 353 (18%), 191 (14%), 341 (5%)
39	4- <i>O</i> -Caffeoylquinic acid <i>cis</i>	19.9	353	173 (100%), 191 (38%)
40	Ferulic acid*	20.4	193	149 (100%), 134 (82%), 178 (36%)
41	Ferulic acid-4- <i>O</i> -hexoside	20.5	355	193 (100%)
42	Quercetin-tri- <i>O</i> -hexoside	20.6	787	625 (100%), 463 (52%)
43	Syringic acid-4- <i>O</i> -acetylhexoside	21.0	401	197 (100%)
44	Syringic acid-dihexoside	21.0	521	197 (100%), 167 (6%), 183 (2%)
45	Coumaric acid- <i>O</i> -hexoside isomer	21.0	325	163 (100%), 119 (24%)
46	Caffeoyl-6 α -hexose	21.0	341	179 (100%), 135 (41%), 281 (21%), 221 (11%), 323 (8%), 251 (5%)
47	Di-hydro-ferulic acid- <i>O</i> -hexoside	21.2	357	195 (100%), 177 (8%), 151 (8%), 136 (6%), 119 (2%)
48	Caffeoylquinic acid- <i>O</i> -hexoside isomer	21.4	515	353 (100%), 341 (74%), 191 (64%), 179 (19%)
49	Di-hydro-caffeic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside	22.0	505	343 (100%), 181 (9%)
50	Medioresinol	22.3	387	207 (100%), 369 (53%), 163 (35%)
51	5- <i>O</i> -Caffeoylquinic acid <i>trans</i>	22.9	353	191 (100%)
52	Sinapic acid-4- <i>O</i> -hexoside	22.9	385	223 (100%), 208 (6%)
53	Cyanidin-3- <i>O</i> -glucoside*	23.0	449 [#]	287 (100%)
54	Cyanidin-di- <i>O</i> -hexoside	23.2	611 [#]	449 (100%), 287 (21%)
55	Apigenin- <i>O</i> -pentoside	23.2	401	269 (100%)
56	4- <i>O</i> -Caffeoylquinic acid <i>trans</i>	23.4	353	173 (100%)
57	Quercetin-3- <i>O</i> -rutinoside- <i>O</i> -hexoside- <i>O</i> -pentoside	23.5	903	741 (100%), 609 (5%), 301 (2%)
58	Feruloylquinic acid- <i>O</i> -hexoside isomer	24.1	529	367 (100%), 191 (60%)
59	Peonidin-3- <i>O</i> -hexoside	24.4	463 [#]	301 (100%)
60	Feruloyl-hexose	24.6	355	193 (100%), 235 (30%), 295 (4%)
61	Caffeic acid- <i>O</i> -hexoside isomer	24.8	341	179 (100%), 135 (36%)
62	Quercetin-3- <i>O</i> -hexoside-7- <i>O</i> -hexoside	24.9	625	463 (100%), 301 (12%), 271 (7%)
63	Taxifolin- <i>O</i> -hexoside	24.9	465	303 (100%)

64	Cyanidin-3- <i>O</i> -malonylhexoside	25.2	535 [#]	287 (100%), 449 (5%)
65	Peonidin-3- <i>O</i> -malonylhexoside	25.2	549 [#]	301 (100%), 463 (6%)
66	Myricetin-di- <i>O</i> -hexoside	25.4	641	479 (100%), 317 (21%)
67	Quercetin-3- <i>O</i> -rutinoside-7- <i>O</i> -hexoside	25.4	771	609 (100%)
68	Sinapoyl-hexose	25.8	385	223 (100%), 208 (2%), 265 (6%), 325 (1%)
69	5- <i>O</i> -Caffeoylquinic acid <i>cis</i>	26.5	353	191 (100%)
70	Oleuropein aglycone isomer	26.5	377	197 (100%), 153 (61%)
71	Caffeic acid- <i>O</i> -malonylhexoside	27.5	457	341 (100%), 179 (14%)
72	Amentoflavone	27.5	537	375 (100%), 179 (14%)
73	Ferulic acid-4- <i>O</i> -pentoside isomer	28.0	325	193 (100%), 149 (36%), 134 (3%)
74	Ferulic acid-4- <i>O</i> -pentoside isomer	28.6	325	193 (100%), 149 (36%), 134 (3%)
75	4- <i>O</i> -Cumarylquinic acid	28.9	337	173 (100%), 163 (17%)
76	Apigenin-6,8-di- <i>C</i> -hexoside	29.1	593	473 (100%), 353 (49%), 383 (33%)
77	5- <i>O</i> -Cumarylquinic acid	29.3	337	191 (100%), 173 (5%) 163 (3%)
78	Isorhamnetin-di- <i>O</i> -hexoside isomer	29.5	639	477 (100%), 315 (5%)
79	Kaempferol-3- <i>O</i> -acetylhexoside	30.0	489	285 (100%), 255 (7%)
80	Quercetin-7- <i>O</i> -hexoside-4'- <i>O</i> -hexoside	31.3	625	463 (100%), 301 (22%)
81	5- <i>O</i> -Feruloylquinic acid	32.2	367	191 (100%), 173 (4%)
82	Quercetin-3- <i>O</i> -hexoside-4'- <i>O</i> -hexoside	33.6	625	463 (100%), 301 (31%), 179 (4%)
83	Kaempferol-3- <i>O</i> -hexoside-7- <i>O</i> -hexoside	33.8	609	285 (100%), 447 (73%), 255 (7%)
84	Isorhamnetin-di- <i>O</i> -hexoside isomer	34.1	639	477 (100%)
85	Myricetin-7- <i>O</i> -hexoside	34.4	479	317 (100%), 289 (65%)
86	Secoisolariciresinol- <i>O</i> -hexoside	35.0	523	361 (100%)
87	Naringenin- <i>C</i> -hexoside	35.1	433	313 (100%)
88	Isorhamnetin-3- <i>O</i> -hexoside-4'- <i>O</i> -hexoside	35.4	639	315 (100%), 477 (63%), 301 (17%), 271 (6%)
89	Quercetin-3- <i>O</i> -rutinoside-7- <i>O</i> -pentoside	36.5	741	609 (100%), 300 (80%)
90	Lariciresinol- <i>O</i> -hexoside	37.3	521	329 (100%), 359 (15%)
91	Apigenin- <i>C</i> -hexoside- <i>O</i> -rhamnoside	37.9	577	341 (100%), 413 (50%), 311 (15%)
92	Eriodictiol- <i>O</i> -hexoside	38.5	449	287 (100%), 151 (42%)
93	Quercetin-3- <i>O</i> -hexoside isomer	39.6	463	301 (100%), 151 (5%), 179 (3%)
94	Di-hydro-quercetin	39.8	303	285 (100%), 267 (54%), 257 (41%)
95	Quercetin-3- <i>O</i> -rutinoside*	39.9	609	301 (100%), 343 (46%), 273 (28%), 243 (13%)
96	Kaempferol-3- <i>O</i> -rutinoside-7- <i>O</i> -pentoside	40.1	725	593 (100%), 285 (30%), 255 (7%), 257 (3%)
97	Luteolin- <i>O</i> -rutinoside isomer	40.1	593	285 (100%), 447 (2%)

98	Phloretin-di- <i>C</i> -hexoside	40.1	597	357 (100%), 387 (91%), 477 (81%)
99	Quercetin-3- <i>O</i> -glucoside*	41.0	463	301 (100%), 151 (23%), 179 (2%)
100	Luteolin- <i>O</i> -hexoside	41.5	447	285 (100%)
101	Naringenin-di- <i>O</i> -hexoside	41.6	595	271 (100%)
102	Luteolin- <i>O</i> -rutinoside isomer	41.7	593	285 (100%), 447 (67%)
103	Pinoresinol- <i>O</i> -hexoside	42.0	519	359 (100%), 151 (2%)
104	Nuzhenide	42.4	685	523 (100%), 453 (93%), 421 (32%), 299 (3%)
105	Phloretin- <i>C</i> -hexoside	42.9	435	315 (100%), 345 (5%)
106	Syringaresinol- <i>O</i> -hexoside	43.3	579	417 (100%), 181 (8%)
107	Verbascoside	43.5	623	461 (100%), 315 (2%)
108	Kaempferol-3- <i>O</i> -rutinoside	44.4	593	285 (100%)
109	4,5-diCaffeoylquinic acid	44.5	515	353 (100%), 179 (18%), 335 (15%), 191 (14%)
110	Quercetin-3- <i>O</i> -acetylhexoside	44.9	505	301 (100%), 463 (67%), 179 (35%)
111	Liquiritigenin-7- <i>O</i> -hexoside	45.0	417	255 (100%)
112	Apigenin- <i>O</i> -hexoside- <i>O</i> -rhamnoside	45.1	577	269 (100%)
113	Apigenin- <i>O</i> -hexoside- <i>O</i> -pentoside	45.3	563	269 (100%), 431 (23%)
114	Kaempferol-3- <i>O</i> -hexoside	45.9	447	284 (100%), 255 (70%), 285 (51%)
115	3,5-diCaffeoylquinic acid	46.0	515	353 (100%), 191 (4%)
116	Isorhamnetin-3- <i>O</i> -hexoside	46.8	477	315 (100%), 300 (12%)
117	Naringenin- <i>O</i> -hexoside isomer	46.8	433	271 (100%)
118	Oleuropein aglycone isomer	47.2	377	307 (100%), 333 (65%), 275 (55%), 139 (12%), 345 (7%)
119	Quercetin-4'- <i>O</i> -hexoside	47.6	463	301 (100%), 179 (12%), 151 (4%)
120	Naringenin- <i>O</i> -hexoside- <i>O</i> -pentoside	47.6	565	271 (100%), 403 (11%)
121	Kaempferol-7- <i>O</i> -hexoside	48.8	447	285 (100%), 257 (11%)
122	Hydroxy-decarboxymethyl-oleuropein aglycone	49.1	335	199 (100%), 181 (29%)
123	Decarboxymethyl-oleuropein aglycone	49.6	319	195 (100%), 165 (18%)
124	Oleuropein	49.8	539	377 (100%), 307 (66%), 275 (32%), 345 (14%)
125	Isorhamnetin-4'- <i>O</i> -hexoside	50.6	477	315 (100%), 299 (15%), 300 (12%)
126	Di-hydroxy-ligstroside aglycone	50.6	393	361 (100%), 257 (79%), 323 (27%), 195 (19%), 151 (16%)
127	β -methoxyverbascoside	51.1	653	491 (100%), 635 (93%)
128	Coumaroyl-caffeoylquinic acid	52.2	499	353 (100%), 191 (10%), 173 (7%)
129	Naringenin- <i>O</i> -hexoside isomer	52.9	433	271 (100%)
130	Ligstroside	56.8	523	361 (100%), 259 (19%)
131	Quercetin	60.8	301	151 (100%), 179 (71%)
132	Ligstroside aglycone	64.0	361	291 (100%), 259 (31%), 223 (4%)

*identified by comparison with authentic standards

#Indicates [M+H]⁺ rather than [M-H]⁻

Table 2. Quantitative results (mg/100 g fresh food) for phenolic compounds identified in the vegetables. Values represent means \pm standard deviation of triplicate determination (n.d. means not detected).

	<i>Compound</i>	<i>Tomato</i>	<i>Onion</i>	<i>Black olives</i>	<i>EVOO</i>	<i>Basil</i>
<i>Hydroxybenzoic acids</i>						
21	Gallic acid	n.d.	< LOQ	n.d.	n.d.	0.08 \pm 0.05
30	Syringic acid	n.d.	< LOQ	n.d.	n.d.	0.56 \pm 0.19
35	Protocatechuic acid- <i>O</i> -pentoside	0.37 \pm 0.07	n.d.	n.d.	n.d.	n.d.
1	Hydroxybenzoic acid- <i>O</i> -hexoside isomer	0.17 \pm 0.04	n.d.	n.d.	n.d.	n.d.
27	Hydroxybenzoic acid- <i>O</i> -hexoside isomer	n.d.	n.d.	n.d.	n.d.	0.19 \pm 0.08
10	Protocatechuic acid- <i>O</i> -hexoside	0.20 \pm 0.02	n.d.	n.d.	n.d.	0.08 \pm 0.02
7	Vanillic acid-4- <i>O</i> -hexoside	0.13 \pm 0.02	n.d.	n.d.	n.d.	0.13 \pm 0.04
36	Syringic acid-4- <i>O</i> -pentoside	n.d.	n.d.	n.d.	n.d.	10.31 \pm 0.16
12	Syringic acid-4- <i>O</i> -hexoside	n.d.	n.d.	n.d.	n.d.	0.14 \pm 0.01
43	Syringic acid-4- <i>O</i> -acetylhexoside	n.d.	n.d.	n.d.	n.d.	0.42 \pm 0.11
22	Protocatechuic acid- <i>O</i> -hexoside- <i>O</i> -pentoside	n.d.	n.d.	n.d.	n.d.	7.17 \pm 0.27
2	Hydroxybenzoic acid-dihexoside	0.32 \pm 0.07	n.d.	n.d.	n.d.	n.d.
44	Syringic acid-dihexoside	n.d.	n.d.	n.d.	n.d.	2.25 \pm 0.31
<i>Total hydroxybenzoic acids</i>		<i>1.19 \pm 0.11</i> <i>(2.4%)</i>	<i>< LOQ</i>	<i>n.d.</i>	<i>n.d.</i>	<i>21.33 \pm 0.49</i> <i>(63.3%)</i>
<i>Hydroxycinnamic acids</i>						
23	Coumaric acid	0.13 \pm 0.01	n.d.	0.04 \pm 0.01	n.d.	n.d.
26	Caffeic acid	0.28 \pm 0.02	n.d.	0.10 \pm 0.01	n.d.	n.d.
40	Ferulic acid	0.73 \pm 0.18	n.d.	n.d.	n.d.	n.d.
24	Coumaric acid- <i>O</i> -hexoside isomer	0.12 \pm 0.03	n.d.	n.d.	n.d.	n.d.
45	Coumaric acid- <i>O</i> -hexoside isomer	0.81 \pm 0.02	n.d.	n.d.	n.d.	n.d.
73	Ferulic acid-4- <i>O</i> -pentoside isomer	n.d.	n.d.	n.d.	n.d.	1.73 \pm 0.13
74	Ferulic acid-4- <i>O</i> -pentoside isomer	n.d.	n.d.	n.d.	n.d.	5.35 \pm 0.49
28	Dihydro-coumaric acid- <i>O</i> -hexoside	0.81 \pm 0.02	n.d.	n.d.	n.d.	n.d.
75	4- <i>O</i> -Cumaroylquinic acid	0.18 \pm 0.08	n.d.	n.d.	n.d.	n.d.
77	5- <i>O</i> -Cumaroylquinic acid	0.53 \pm 0.06	n.d.	n.d.	n.d.	n.d.
25	Caffeic acid- <i>O</i> -hexoside isomer	5.72 \pm 0.56	n.d.	0.67 \pm 0.01	n.d.	0.05 \pm 0.02
33	Caffeoyl-6 β -hexose	1.13 \pm 0.04	n.d.	n.d.	n.d.	n.d.

46	Caffeoyl-6 α -hexose	2.59 ± 0.16	n.d.	0.39 ± 0.04	n.d.	n.d.
61	Caffeic acid- <i>O</i> -hexoside isomer	0.17 ± 0.05	n.d.	n.d.	n.d.	n.d.
29	Dihydro-caffeic acid- <i>O</i> -hexoside isomer	0.55 ± 0.05	n.d.	n.d.	n.d.	n.d.
37	Dihydro-caffeic acid- <i>O</i> -hexoside isomer	0.69 ± 0.12	n.d.	n.d.	n.d.	n.d.
17	3- <i>O</i> -Caffeoylquinic acid	0.18 ± 0.01	n.d.	n.d.	n.d.	n.d.
39	4- <i>O</i> -Caffeoylquinic acid <i>cis</i>	0.38 ± 0.17	n.d.	n.d.	n.d.	n.d.
51	5- <i>O</i> -Caffeoylquinic acid <i>trans</i>	4.57 ± 0.03	n.d.	n.d.	n.d.	n.d.
56	4- <i>O</i> -Caffeoylquinic acid <i>trans</i>	4.61 ± 0.11	n.d.	n.d.	n.d.	n.d.
69	5- <i>O</i> -Caffeoylquinic acid <i>cis</i>	0.90 ± 0.01	n.d.	n.d.	n.d.	n.d.
41	Ferulic acid-4- <i>O</i> -hexoside	1.56 ± 0.46	n.d.	n.d.	n.d.	n.d.
60	Feruloyl-hexose	2.95 ± 0.85	n.d.	n.d.	n.d.	0.05 ± 0.01
47	Dihydro-ferulic acid- <i>O</i> -hexoside	6.65 ± 0.63	n.d.	n.d.	n.d.	n.d.
20	Rosmarinic acid	n.d.	n.d.	n.d.	n.d.	0.08 ± 0.01
81	5- <i>O</i> -Feruloylquinic acid	1.80 ± 0.23	n.d.	n.d.	n.d.	n.d.
52	Sinapic acid-4- <i>O</i> -hexoside	2.21 ± 0.19	1.58 ± 0.19	n.d.	n.d.	2.34 ± 0.19
68	Sinapoyl-hexose	n.d.	5.75 ± 0.33	n.d.	n.d.	n.d.
71	Caffeic acid- <i>O</i> -malonylhexoside	0.23 ± 0.01	n.d.	n.d.	n.d.	n.d.
11	Caffeic acid- <i>O</i> -hexoside- <i>O</i> -pentoside isomer	n.d.	n.d.	n.d.	n.d.	0.16 ± 0.05
31	Caffeic acid- <i>O</i> -hexoside- <i>O</i> -pentoside isomer	n.d.	n.d.	n.d.	n.d.	0.09 ± 0.01
13	Calceolarioside	n.d.	n.d.	0.32 ± 0.09	n.d.	n.d.
19	Caffeoyl-coumaroylquinic acid	0.12 ± 0.01	n.d.	n.d.	n.d.	n.d.
128	Coumaroyl-caffeoylquinic acid	0.10 ± 0.01	n.d.	n.d.	n.d.	n.d.
16	Caffeic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside	0.14 ± 0.01	n.d.	n.d.	n.d.	n.d.
49	Dihydro-caffeic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside	0.19 ± 0.02	n.d.	n.d.	n.d.	n.d.
15	Caffeoylquinic acid- <i>O</i> -hexoside isomer	0.06 ± 0.01	n.d.	n.d.	n.d.	n.d.
18	Caffeoylquinic acid- <i>O</i> -hexoside isomer	0.52 ± 0.09	n.d.	n.d.	n.d.	n.d.
38	Caffeoylquinic acid- <i>O</i> -hexoside isomer	0.38 ± 0.17	n.d.	n.d.	n.d.	n.d.
48	Caffeoylquinic acid- <i>O</i> -hexoside isomer	0.11 ± 0.03	n.d.	n.d.	n.d.	n.d.
109	4,5-diCaffeoylquinic acid	1.12 ± 0.28	n.d.	n.d.	n.d.	n.d.
115	3,5-diCaffeoylquinic acid	0.73 ± 0.03	n.d.	n.d.	n.d.	0.35 ± 0.08
32	Feruloylquinic acid- <i>O</i> -hexoside isomer	1.43 ± 0.01	n.d.	n.d.	n.d.	n.d.
58	Feruloylquinic acid- <i>O</i> -hexoside isomer	1.70 ± 0.42	n.d.	n.d.	n.d.	n.d.

6	Caffeoylquinic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside isomer	0.33 ± 0.06	n.d.	n.d.	n.d.	n.d.
14	Caffeoylquinic acid-3- <i>O</i> -hexoside-4- <i>O</i> -hexoside isomer	0.06 ± 0.01	n.d.	n.d.	n.d.	n.d.
Total hydroxycinnamic acids		47.48 ± 1.46 (94.4%)	7.32 ± 0.39 (19.0%)	1.52 ± 0.09 (3.5%)	n.d.	10.20 ± 0.55 (30.3%)
Flavonols						
131	Quercetin	n.d.	0.04 ± 0.01	n.d.	n.d.	n.d.
114	Kaempferol-3- <i>O</i> -hexoside	n.d.	0.09 ± 0.01	n.d.	n.d.	n.d.
121	Kaempferol-7- <i>O</i> -hexoside	< LOQ	0.19 ± 0.02	n.d.	n.d.	n.d.
93	Quercetin-3- <i>O</i> -hexoside isomer	n.d.	< LOQ	n.d.	n.d.	n.d.
99	Quercetin-3- <i>O</i> -glucoside	0.01 ± 0.01	0.43 ± 0.05	0.03 ± 0.01	n.d.	n.d.
119	Quercetin-4'- <i>O</i> -hexoside	n.d.	7.29 ± 0.40	n.d.	n.d.	n.d.
116	Isorhamnetin-3- <i>O</i> -hexoside	n.d.	0.04 ± 0.01	n.d.	n.d.	n.d.
125	Isorhamnetin-4'- <i>O</i> -hexoside	n.d.	2.35 ± 0.07	< LOQ	n.d.	n.d.
85	Myricetin-7- <i>O</i> -hexoside	n.d.	0.01 ± 0.01	n.d.	n.d.	n.d.
79	Kaempferol-3- <i>O</i> -acetylhexoside	n.d.	< LOQ	n.d.	n.d.	n.d.
110	Quercetin-3- <i>O</i> -acetylhexoside	n.d.	< LOQ	n.d.	n.d.	< LOQ
108	Kaempferol-3- <i>O</i> -rutinoside	< LOQ	n.d.	n.d.	n.d.	n.d.
95	Quercetin-3- <i>O</i> -rutinoside	0.37 ± 0.01	n.d.	n.d.	n.d.	n.d.
83	Kaempferol-3- <i>O</i> -hexoside-7- <i>O</i> -hexoside	n.d.	0.21 ± 0.01	n.d.	n.d.	n.d.
62	Quercetin-3- <i>O</i> -hexoside-7- <i>O</i> -hexoside	n.d.	0.03 ± 0.01	n.d.	n.d.	n.d.
80	Quercetin-7- <i>O</i> -hexoside-4'- <i>O</i> -hexoside	n.d.	0.15 ± 0.01	n.d.	n.d.	n.d.
82	Quercetin-3- <i>O</i> -hexoside-4'- <i>O</i> -hexoside	< LOQ	11.13 ± 0.18	0.03 ± 0.01	n.d.	< LOQ
78	Isorhamnetin-di- <i>O</i> -hexoside isomer	n.d.	< LOQ	n.d.	n.d.	n.d.
84	Isorhamnetin-di- <i>O</i> -hexoside isomer	n.d.	< LOQ	n.d.	n.d.	n.d.
88	Isorhamnetin-3- <i>O</i> -hexoside-4'- <i>O</i> -hexoside	n.d.	0.38 ± 0.01	n.d.	n.d.	n.d.
66	Myricetin-di- <i>O</i> -hexoside	n.d.	0.01 ± 0.01	n.d.	n.d.	n.d.
96	Kaempferol-3- <i>O</i> -rutinoside-7- <i>O</i> -pentoside	< LOQ	< LOQ	n.d.	n.d.	n.d.
89	Quercetin-3- <i>O</i> -rutinoside-7- <i>O</i> -pentoside	0.17 ± 0.01	< LOQ	n.d.	n.d.	n.d.
67	Quercetin-3- <i>O</i> -rutinoside-7- <i>O</i> -hexoside	0.01 ± 0.01	< LOQ	n.d.	n.d.	n.d.
42	Quercetin-tri- <i>O</i> -hexoside	n.d.	0.08 ± 0.01	n.d.	n.d.	n.d.
57	Quercetin-3- <i>O</i> -rutinoside- <i>O</i> -hexoside- <i>O</i> -pentoside	0.01 ± 0.01	< LOQ	n.d.	n.d.	n.d.

<i>Total flavonols</i>		<i>0.56 ± 0.01</i> (1.1%)	<i>22.44 ± 0.45</i> (58.3%)	<i>0.06 ± 0.01</i> (0.1%)	<i>n.d.</i>	<i>< LOQ</i>
<hr/>						
<i>Anthocyanins</i>						
53	Cyanidin-3- <i>O</i> -glucoside	n.d.	1.59 ± 0.02	n.d.	n.d.	n.d.
59	Peonidin-3- <i>O</i> -hexoside	n.d.	0.45 ± 0.01	n.d.	n.d.	n.d.
64	Cyanidin-3- <i>O</i> -malonylhexoside	n.d.	4.29 ± 0.24	n.d.	n.d.	n.d.
65	Peonidin-3- <i>O</i> -malonylhexoside	n.d.	0.84 ± 0.04	n.d.	n.d.	n.d.
54	Cyanidin-di- <i>O</i> -hexoside	n.d.	1.25 ± 0.04	n.d.	n.d.	n.d.
<i>Total anthocyanins</i>		<i>n.d.</i>	<i>8.42 ± 0.24</i> (21.9%)	<i>n.d.</i>	<i>n.d.</i>	<i>n.d.</i>
<hr/>						
<i>Lignans</i>						
50	Medioresinol	0.04 ± 0.01	n.d.	n.d.	n.d.	1.66 ± 0.01
103	Pinoresinol- <i>O</i> -hexoside	n.d.	n.d.	n.d.	n.d.	0.07 ± 0.01
90	Lariciresinol- <i>O</i> -hexoside	n.d.	n.d.	n.d.	n.d.	0.40 ± 0.04
86	Secoisolariciresinol- <i>O</i> -hexoside	n.d.	0.03 ± 0.01	n.d.	n.d.	n.d.
106	Syringaresinol- <i>O</i> -hexoside	n.d.	0.15 ± 0.01	0.04 ± 0.01	n.d.	n.d.
<i>Total lignans</i>		<i>0.04 ± 0.01</i> (0.1%)	<i>0.18 ± 0.15</i> (0.5%)	<i>0.04 ± 0.01</i> (0.1%)	<i>n.d.</i>	<i>2.14 ± 0.23</i> (6.4%)
<hr/>						
<i>Flavones</i>						
55	Apigenin- <i>O</i> -pentoside	0.03 ± 0.01	n.d.	n.d.	n.d.	n.d.
34	Apigenin- <i>O</i> -hexoside	0.07 ± 0.02	n.d.	n.d.	n.d.	n.d.
100	Luteolin- <i>O</i> -hexoside	n.d.	n.d.	0.27 ± 0.02	n.d.	n.d.
72	Amentoflavone	n.d.	n.d.	0.02 ± 0.01	n.d.	< LOQ
113	Apigenin- <i>O</i> -hexoside- <i>O</i> -pentoside	n.d.	n.d.	< LOQ	n.d.	n.d.
91	Apigenin- <i>C</i> -hexoside- <i>O</i> -rhamnoside	0.06 ± 0.01	n.d.	n.d.	n.d.	n.d.
112	Apigenin- <i>O</i> -hexoside- <i>O</i> -rhamnoside	n.d.	n.d.	0.02 ± 0.01	n.d.	n.d.
76	Apigenin-6,8-di- <i>C</i> -hexoside	n.d.	n.d.	0.01 ± 0.01	n.d.	0.02 ± 0.01
97	Luteolin- <i>O</i> -rutinoside isomer	n.d.	n.d.	0.02 ± 0.01	n.d.	n.d.
102	Luteolin- <i>O</i> -rutinoside isomer	n.d.	n.d.	0.04 ± 0.01	n.d.	n.d.
<i>Total flavones</i>		<i>0.16 ± 0.02</i> (0.3%)	<i>n.d.</i>	<i>0.38 ± 0.01</i> (0.9%)	<i>n.d.</i>	<i>0.02 ± 0.01</i> (0.1%)
<hr/>						
<i>Flavanones</i>						
111	Liquiritigenin-7- <i>O</i> -hexoside	0.02 ± 0.01	n.d.	n.d.	n.d.	n.d.
87	Naringenin- <i>C</i> -hexoside	0.01 ± 0.01	n.d.	n.d.	n.d.	n.d.
117	Naringenin- <i>O</i> -hexoside isomer	0.01 ± 0.01	n.d.	n.d.	n.d.	n.d.
129	Naringenin- <i>O</i> -hexoside isomer	0.01 ± 0.01	n.d.	n.d.	n.d.	n.d.
92	Eriodictiol- <i>O</i> -hexoside	0.02 ± 0.01	n.d.	n.d.	n.d.	n.d.


120	Naringenin- <i>O</i> -hexoside- <i>O</i> -pentoside	0.01 ± 0.01	n.d.	n.d.	n.d.	n.d.
101	Naringenin-di- <i>O</i> -hexoside	< LOQ	n.d.	n.d.	n.d.	n.d.
Total flavonones		0.09 ± 0.01 (0.2%)	n.d.	n.d.	n.d.	n.d.
Tyrosol derivatives						
4	Hydroxytyrosol isomer	n.d.	n.d.	0.63 ± 0.14	n.d.	n.d.
9	Hydroxytyrosol isomer	n.d.	n.d.	0.65 ± 0.07	1.50 ± 0.10	n.d.
1	Hydroxytyrosol- <i>O</i> -hexoside isomer	n.d.	n.d.	15.24 ± 0.48	n.d.	n.d.
5	Hydroxytyrosol- <i>O</i> -hexoside isomer	n.d.	n.d.	2.41 ± 1.00	n.d.	n.d.
123	Decarboxymethyl-oleuropein aglycone	n.d.	n.d.	n.d.	0.77 ± 0.01	n.d.
122	Hydroxy-decarboxymethyl-oleuropein aglycone	n.d.	n.d.	n.d.	11.31 ± 0.27	n.d.
132	Ligstroside aglycone	n.d.	n.d.	n.d.	13.66 ± 0.54	n.d.
70	Oleuropein aglycone isomer	n.d.	n.d.	20.98 ± 0.67	n.d.	n.d.
118	Oleuropein aglycone isomer	n.d.	n.d.	n.d.	46.98 ± 2.79	n.d.
126	Di-hydroxy-ligstroside aglycone	n.d.	n.d.	n.d.	0.07 ± 0.03	n.d.
2	Hydroxytyrosol-di- <i>O</i> -hexoside	n.d.	n.d.	0.96 ± 0.11	n.d.	n.d.
130	Ligstroside	n.d.	n.d.	0.11 ± 0.02	n.d.	n.d.
124	Oleuropein	n.d.	n.d.	0.24 ± 0.03	n.d.	n.d.
107	Verbascoside	n.d.	n.d.	0.34 ± 0.01	n.d.	n.d.
127	β-methoxylverbascoside	n.d.	n.d.	0.15 ± 0.01	n.d.	n.d.
104	Nuzhenide	n.d.	n.d.	0.21 ± 0.06	n.d.	n.d.
Total tyrosol derivatives		n.d.	n.d.	41.93 ± 1.13 (95.4%)	74.30 ± 2.85 (100.0%)	n.d.
Dihydroflavonols						
94	Dihydro-quercetin	n.d.	<LOQ	n.d.	n.d.	n.d.
63	Taxifolin- <i>O</i> -hexoside	n.d.	0.02 ± 0.01	n.d.	n.d.	n.d.
Total dihydroflavonols		n.d.	0.02 ± 0.01 (0.1%)	n.d.	n.d.	n.d.
Dihydrochalcones						
105	Phloretin- <i>C</i> -hexoside	n.d.	0.01 ± 0.01	n.d.	n.d.	n.d.
98	Phloretin-di- <i>C</i> -hexoside	0.76 ± 0.01	0.07 ± 0.01	n.d.	n.d.	n.d.
Total dihydrochalcones		0.76 ± 0.01 (1.5%)	0.08 ± 0.01 (0.2%)	n.d.	n.d.	n.d.
Total phenolic compounds		50.27 ± 1.47	38.46 ± 0.66	43.93 ± 0.10	74.30 ± 2.85	33.69 ± 0.77

Table 3. Radical scavenging properties, ferrous ions chelating ability and ferric ions reducing properties of phenolic fractions from vegetable foods and extra-virgin olive oil.


	<i>ABTS radical scavenging</i>	<i>Hydroxyl radical scavenging</i>	<i>Superoxide anion scavenging</i>	<i>Fe³⁺ reducing properties</i>	<i>Fe²⁺ chelating ability</i>
	<i>μmol ascorbic acid equivalent/mg of total phenolic compounds^a</i>				<i>% chelation^b</i>
Tomato	1.60 ± 0.05 ^c	0.89 ± 0.05 ^c	2.77 ± 0.45 ^a	2.04 ± 0.07 ^b	4.65 ± 1.14 ^c
Fresh basil	1.69 ± 0.01 ^c	0.89 ± 0.02 ^c	0.11 ± 0.01 ^c	1.49 ± 0.08 ^b	54.72 ± 6.29 ^a
Onion	2.97 ± 0.20 ^a	1.56 ± 0.07 ^b	1.04 ± 0.05 ^b	1.03 ± 0.02 ^c	6.90 ± 2.98 ^c
Black olives	2.70 ± 0.04 ^b	1.78 ± 0.08 ^a	0.14 ± 0.04 ^c	1.47 ± 0.04 ^b	43.53 ± 1.86 ^b
EVOO	1.61 ± 0.07 ^c	0.87 ± 0.07 ^c	0.27 ± 0.01 ^c	0.63 ± 0.02 ^d	7.54 ± 1.61 ^c

^adata expressed as μmol ascorbic acid equivalent normalized for the total phenolic content as determined by mass spectrometry experiments

^b% of chelated Fe²⁺ by 100 μg of phenolic compounds




Mediterranean diet salad (g/100 g meat)




A


Tomato, 50.3 ± 1.5 mg/100g fresh food




Onion, 38.5 ± 0.7 mg/100g fresh food



Black Olives, 43.9 ± 0.1 mg/100g fresh food





Basil, 33.7 ± 0.8 mg/100g fresh food



Tyrosol derivatives composition

Black Olives, 41.9 ± 1.1 mg/100g fresh food

EVOO, 74.3 ± 2.8 mg/100g fresh food



Biplot PC1 vs PC2

