

This is the peer reviewed version of the following article:

Tocilizumab in COVID-19: finding the optimal route and dose – Authors' reply / Guaraldi, G.; Milic, J.; Cozzi-Lepri, A.; Pea, F.; Mussini, C.. - In: THE LANCET. RHEUMATOLOGY. - ISSN 2665-9913. - 2:12(2020), pp. e739-e740. [10.1016/S2665-9913(20)30333-7]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

13/01/2026 02:11

The Lancet Rheumatology
Tocilizumab for severe COVID-19 pneumonia - Authors' reply 2
--Manuscript Draft--

Manuscript Number:	
Article Type:	Correspondence (Author's Reply)
Keywords:	
Corresponding Author:	Giovanni Guaraldi University of Modena and Reggio Emilia Modena, ITALY
First Author:	Giovanni Guaraldi
Order of Authors:	Giovanni Guaraldi
	Jovana Milic
	Alessandro Cozzi-Lepri
	Federico Pea
	Cristina Mussini
Manuscript Region of Origin:	ITALY

Tocilizumab for severe COVID-19 pneumonia - Authors' reply 2

Dr Jain et al underline that TESEO cohort found no difference in efficacy of subcutaneous (SC) tocilizumab (TCZ) formulation, in comparison to intravenous (IV) route and advocate for its use due to an approximatively 6-fold cost reduction. However, we argue that IV administration has other advantages, i.e. the pharmacokinetic profile that is more linear and predictable, while proteolytic degradation of the SC formulation could be variable. Additionally, elevated levels of IL-6 may downregulate the hepatic cytochromes [1]. This may promote enhanced drug exposure as it has been recently postulated for darunavir [2]. Consistently, we believe that prospective pharmacokinetic studies comparing different administration routes are needed to address both appropriate dose-finding and safety. A formal cost-effectiveness analysis should also be considered.

We agree that the optimal time for TCZ administration in COVID-19 is critical. While a beneficial effect of TCZ on mortality was shown in observational studies, a recent randomized trial did not confirm these results. Besides unmeasured confounding, the case-mix of the target population, number of doses, timing of the intervention are other possible reasons for the conflicting results between observational and randomized studies. Assuming that a causal link could be established, the question when is best to start TCZ should be addressed in a randomized study. The emulation of such a trial in the observational setting requires sophisticated methodology beyond those used in our parental paper and a collaborative study with a much larger sample size. A simple correlation analysis is unlikely to produce the answer that we need.

Regarding the need for monitoring patients with severe renal impairment, chronic kidney disease was found in 14 participants at hospital admission and 7 of these (50%) received TCZ. The primary endpoint was observed in 4/7 (57%) in TCZ in addition to standard of care (SoC) and in 3/7 (43%) in SoC group ($p=1.0$). All but one (who was treated with TCZ) of the 7 participants who experienced the endpoint have died. Therefore, our data, although limited, suggest that TCZ use was not harmful in this subgroup.

To conclude, the challenge of appropriate TCZ use lays on the prediction of the progression of respiratory failure in people who develop a cytokine storm. This is typically accompanied by "respiratory crush" is unlikely to be captured by chest radiology. A recent study demonstrated little benefit with this regard [3]. We can rely on a machine learning algorithms which provides a trustworthy 48 hours prediction of severe respiratory failure, with satisfactory accuracy [4].

Giovanni Guaraldi^{1,2}

Jovana Milic^{2,3}

Alessandro Cozzi-Lepri⁴

Federico Pea⁵

Cristina Mussini^{1,2}

¹ Department of Infectious Diseases, Azienda Ospedaliero-Universitaria Policlinico of Modena, Modena 41124, Italy

² Department of Surgical, Medical, Dental and Morphological Sciences, University of Modena and Reggio Emilia, Modena, Italy

³ Clinical and Experimental Medicine PhD Program, University of Modena and Reggio Emilia, Modena, Italy

⁴ Centre for Clinical Research, Epidemiology, Modelling and Evaluation, Institute for Global Health, University College London, London, UK

⁵ Institute of Clinical Pharmacology, Azienda Ospedaliero-Universitaria Santa Maria Della Misericordia, University of Udine, Udine, Italy

References

1. Morgan ET. Impact of infectious and inflammatory disease on cytochrome P450-mediated drug metabolism and pharmacokinetics. *Clin Pharmacol Ther.* **2009**; 85(4):434–438.
2. Cojutti PG, Londero A, Siega P Della, et al. Comparative Population Pharmacokinetics of Darunavir in SARS-CoV-2 Patients vs. HIV Patients: The Role of Interleukin-6. *Clin Pharmacokinet* [Internet]. **2020**; . Available from: <https://doi.org/10.1007/s40262-020-00933-8>
3. Colombi D, Bodini FC, Petrini M, et al. Well-aerated Lung on Admitting Chest CT to Predict Adverse Outcome in COVID-19 Pneumonia. *Radiology.* **2020**; 296(2):E86–E96.
4. Ferrari D, Milic J, Tonelli R, et al. Machine learning in predicting respiratory failure in patients with COVID-19 pneumonia - challenges, strengths, and opportunities in a global health emergency. *medRxiv* [Internet]. **2020**; :2020.05.30.20107888. Available from: <http://medrxiv.org/content/early/2020/06/03/2020.05.30.20107888.abstract>