UV solar exposure of outdoor workers in Mediterranean area

Chiara Burattini

*Dept. of Astronautical Electric and
Energy Engineering (DIAEE)
Sapienza University
Rome, Italy
chiara.burattini@uniroma1.it*

Laura Pompei

*Dept. of Astronautical Electric and
Energy Engineering (DIAEE)
Sapienza University
Rome, Italy
laura.pompei@uniroma1.it*

Alberto Modenese

*Dept. of Biomedical, Metabolic and
Neural Sciences
University of Modena & Reggio Emilia
Modena, Italy
alberto.modenese@unimore.it*

Giacomo Salvadori

*Dept. of Energy System Territory and
Construction Engineering
University of Pisa
Pisa, Italy
giacomo.salvadori@unipi.it*

Andrea Militello

*Dept. of Occupational and
Environmental Medicine, Epidemiology
and Hygiene, Italian Workers'
Compensation Authority (INAIL-
DiMEILA)
Monte Porzio Catone (Rome), Italy
a.militello@inail.it*

Francesco Leccese

*Dept. of Energy System Territory and
Construction Engineering
University of Pisa
Pisa, Italy
francesco.leccese@unipi.it*

Massimo Borra

*Dept. of Occupational and
Environmental
Medicine, Epidemiology and Hygiene,
Italian Workers' Compensation
Authority
(INAIL-DiMEILA)
Monte Porzio Catone (Rome), Italy
m.borra@inail.it*

Fabriziomaria Gobba

*Dept. of Biomedical, Metabolic and
Neural Sciences
University of Modena & Reggio Emilia
Modena, Italy
fabriziomaria.gobba@unimore.it*

Fabio Bisegna

*Dept. of Astronautical Electric and
Energy Engineering (DIAEE)
Sapienza University
Rome, Italy
fabio.bisegna@uniroma1.it*

Abstract—The exposure to UV solar radiation can have several adverse health effects. During summer in Mediterranean area high UV irradiation levels occur and outdoor workers may incur in overexposure risk. This paper reviews the measurement campaigns of UV solar exposure carried out in Mediterranean area.

Keywords—solar radiation, occupational, erythemat, measurement.

I. INTRODUCTION

The Mediterranean area is extended between the latitude 30° and 47° N and for this it has a temperate climate; nevertheless, in Summer season high UV irradiation levels are recorded. Figure 1 shows the daily erythemat UV Index (UVI) [1] in Europe in a Summer day (21st July) provided from the TEMIS service [2]: in the Mediterranean area UVI values between 9 and 11 have been recorded.

Outdoor workers in Mediterranean area have high risk to incur in adverse health effects consequent to UV solar radiation overexposure. Health diseases related to excessive exposure to UV radiation involve skin and eyes and may be acute or long-term effects of exposure: they include erythema, non-melanoma and melanoma skin cancer, photokeratitis, photoconjunctivitis and cataract.

The assessment of worker's exposure to UV radiation is important under the aspects of prevention, protection and diagnosis. This paper presents a review of the scientific measurement campaigns carried out in Mediterranean area in the recent years with the aim of comparing the experiences.

The European Directive 2006/25/CE established a threshold limit of exposure to the artificial optical radiation of $H_{\text{eff}} = 30 \text{ J/m}^2$ over an interval of 8 hours: it corresponds to

This work has been carried out within the BRIC project financed by INAIL.

a SED of about 100 J/m^2 .

Figure 1. UVI in Europe in a summer day (TEMIS)

II. METHOD

A. Articles selection

The articles search database "Scopus" has been queried with the following keywords: solar radiation + outdoor workers; solar exposure + occupational; measure erythemat exposure.

Results were firstly filtered basing on title and abstract: 31 articles were extracted and analyzed. They treated measurement campaigns of UV solar exposure all over the World: among these, 10 articles regarded the measure of workers' exposure in Mediterranean area and were considered for the analysis. Another article, published in a national Journal of the sector was added, for a total of 11 papers.

The articles, ordered for publication data are: Milon et al. (2007) [3], Serrano et al. (2009) [4], Sisto et al. (2009) [5], Serrano et al. (2013) [6], Miligi et al. (2013) [7], Nardini et al. (2014) [8], Modenese et al. (2019) [9], Wittlich et al. (2020) [10], Kovačić et al. (2020) [11] Modenese et al. (2020c) [12], and Modenese et al. (2020d) [13].

B. Analysis

The 11 selected articles were read and analyzed in the following aspects:

- Work typology;
- Number of participants;
- Period of measurement campaign: season/month, number of days, hours;
- Environmental conditions: whether conditions, setting typology;
- Instruments typology (dosimeters, radiometers) and their position;
- Measured dimensions: UV irradiance, effective radiant exposure H_{eff} , SED, UVI;
- Body districts considered.

For the paper Wittlich et al. [10] only the data referred to the 2 measurement campaigns carried out in the Mediterranean area were considered.

III. RESULTS

The 11 selected articles were published from 2007 to 2020: the temporal distribution is showed in the graph of figure 2.

Figure 2. Year of publication of the papers

The papers can be grouped in three periods with intervals of some years between one and the other: the number of papers published in the 2019-2020 indicates a renewed interest for the problem in the last years.

The measurement campaigns regarded a wide typology of works and some papers investigated more than one work [4][7][13]; the number of workers that participated to the measurement campaign was generally less than 10 people (see Table 1); gender and age of participants were generally not specified.

TABLE I. WORK AND WORKERS INFORMATION

Paper	Work typology	Number of workers
Kovacic et al. (2020)	construction	4
Miligi et al. (2013)	agriculture	From 13 to 22
	fishing	
Milon et al. (2007)	construction	20
Modenese et al. (2020a)	construction	3
Modenese et al. (2020b)	longshore	10
	traffic coordination	4
Modenese et al. (2019)	fishing	7
Nardini et al. (2014)	agriculture	6
Serrano et al. (2009)	gardeners	4
	lifeguards	5
Serrano et al. (2013)	costruction	8
Sisto et al. (2009)	-	From 13 to 22
Wittlich et al. (2020)	construction	4+4

The environmental conditions indicated a small geographic distribution: the campaigns were located in three countries (most of them were carried out in Italy) and often in recurring places of measurements; table 2 indicates also that different environments were investigated.

TABLE II. ENVIRONMENT INFORMATION

Paper	country	location	setting	altitude (m)
Kovacic et al. (2020)	Croatia	Zagreb	-	128
Miligi et al. (2013)	Italy	Siena	rural	300
		41°-43° N	sea	0
Milon et al. (2007)	Switzerland	Valais	mountain	500-600 1400-1500 2000-2500
Modenese et al. (2020a)	Italy	Siena	-	-
Modenese et al. (2020b)	Italy	Trieste	port	0
Modenese et al. (2019)	Italy	North-West Adriatic Sea	sea	0
Nardini et al. (2014)	Italy	Marche	farm	195
Serrano et al. (2009)	Spain	Valencia	university campus	0
		Malvarrosa	beach	0
Serrano et al. (2013)	Spain	Valencia	university campus	0
Sisto et al. (2009)	Italy	Siena	rural	300
Wittlich et al. (2020)	Italy	Siena	-	-
	Croatia	Zagreb	-	-

Table 3 shows that the campaigns of measure were carried out principally in Summer and some of them in Spring, when the risk of exposure is higher; nevertheless, in many cases only few days were recorded and only few studies investigated a long period. Many measurements were performed in cloudless sky conditions, while the information is not available when the campaign lasts many days: probably, in these cases, many sky conditions occurred.

TABLE III. TIMING AND WEATHER INFORMATION

Paper	months	duration	hours	sky
Kovacic et al. (2020)	Jun-Oct	-	07:00-18:00	-
Miligi et al. (2013)	Apr, Jul, Oct	3 d	-	variable
	Jul	-	-	clear
Milon et al. (2007)	Jul-Sep	2-4 h	10:00-16:00	cloudless
Modenes e et al. (2020a)	May-Jun	23 d	09:00-17:00	-
Modenes e et al. (2020b)	Jul	1 d	Full working day	-
Modenes e et al. (2019)	May	2 d	07:00-14:00	cloudless
Nardini et al. (2014)	Apr, May	19 d	10:00-12:00 14:00-16:00	-
	Jun	8 d		
	Jul	8 d		
	Ago	5 d		
	Sep	16 d		
Serrano et al. (2009)	Jun	4 d	06:00-13:00	cloudless
	Jul	4 d	10:00-19:00	cloudless
Serrano et al. (2013)	Jul	5 d	08:00-14:00 15:00-19:00	-
Sisto et al. (2009)	Apr, Jul, Oct	-	-	variable
Wittlich et al. (2020)	May-Sep	5 days per week	07:00-17:00	-
	Jun-Oct			

In many articles the effective radiant exposure or dose not was the only dimension measured, but often the UV irradiance on the horizontal plan was recorded and sometime the UVI or the SED was calculated. Moreover, many studies measured the exposure of several body districts. The following table 4 shows the information regarding the measurement conditions.

TABLE IV. MESUREMENT INFORMATION

Paper	Instrument	weighting	Dimensions	position
Kovacic et al. (2020)	dosimeters	erythematic	UV-A and UVB/C irradiances, SED	Left upper arm
	Meteorological station	-	UV daily dose	-
	TEMIS	Erythematic, vitamin D, DNA-damage	daily dose	
Miligi et	Polysulp	-	-	Arm, back

Paper	Instrument	weighting	Dimensions	position
al. (2013)	hone dosimeters			
	Film dosimeters	-	-	Back, heat, arms
Milon et al. (2007)	Film dosimeters	CIE	Effective daily exposure, SED, percent of ambient exposure	Neck, shoulder, low back, forehead
	dosimeter	-	Static UV irradiance	horizontal
	Baseline Surface Radiation Network	-	Daily irradiation cycles	-
Modenese et al. (2020a)	Electronic dosimeters	erythematic	UV-A and UVB/C irradiances	Left upper arm
	ESA database	erythematic	Total environmental UV dose	-
Modenese et al. (2020b)	Electronic dosimeters	-	Personal UVR exposure, ocular UVA dose	Chest, back, sunglass
	Spectroradiometer	-	Environment UV radiant exposure	-
	ESA	erythematic	Effective radiant exposure	Horizontal plan
Modenese et al. (2019)	Electronic dosimeters	Unweighted, CIE erythematic, ICNIRP erythematic actinic	Solar UV exposure	Back, chest, nape, sunglass
	Spectroradiometer	-	Environment UV exposure	Horizontal plan
Nardini et al. (2014)	Spectroradiometer	-	irradiance	Horizontal plan
	Personal dosimeters	CIE erythematic	UVR exposure	Cheek, nape, forehead, forearm
Serrano et al. (2009)	Spore-film dosimeters	CIE erythematic	Solar erythematic exposure, MED (type II), SED	Shoulder, wrist
	PMA detector		Ambient UVR	roof
Serrano et al. (2013)	Spore-film dosimeters	CIE erythematic	SED	Shoulder, chest
	UVB radiometer	erythematic	Ambient UVR	roof
Sisto et al. (2009)	Polysulp hone film dosimeters	CIE erythematic	UVA and UVB solar irradiance, effective dose	Arm, back
	-	-	Skin reflectivity, external cutaneous temperature	Cheek, inner arm

Paper	Instrument	weighting	Dimensions	position
Wittlich et al. (2020)	Electronic dosimeters	-	UVR exposure	Left upper arm

IV. DISCUSSION

Results of articles selection evidenced that only 11 papers regarding measurement campaigns of the UV solar radiation exposure of outdoor workers in Mediterranean area have been published in 14 years (less than 1 per year); moreover, the measurement campaigns were carried out in only four European countries (Italy, Spain, Croatia and Switzerland), while no articles regard the Asiatic and North African countries. This could be due to the use of a single search database, but it indicates that the problem of workers' overexposure to UV solar radiation and the related health risks are poorly or not considered in many Mediterranean countries, particularly in the South regions where the health risks are greater due to the higher UV irradiation levels.

On the contrary, where the problem of outdoor workers' protection against UV solar radiation has considered, significant measurement campaigns have been carried out. Results of analysis indicated that the UV solar exposure of a large variety of outdoor workers has been measured in several environmental conditions: this demonstrates that this theme has a large extension and involves a big number of workers. At the same time, the collection of many data consents to compare the different exposures and it is a first step through the systematization.

Summer and Spring were the periods mostly investigated: even in these two seasons exposure risks are greater, in the Mediterranean area the erythema exposure should be monitored in all seasons, since high irradiance levels can be recorded also in Winter [14].

Regarding the method of measure, the articles analysis demonstrated a substantial uniformity: about all studies used personal dosimeters erythemally weighted (generally with the CIE curve) and the measure of effective radiant exposure was often combined with the environmental UV radiant exposure measured on the horizontal plan with spectroradiometers or derived from satellite data. The dosimeters were worn in different positions, and this allows to obtain exposure information on many body districts.

V. CONCLUSION

The analysis performed on the selected article indicated that in many regions of the Mediterranean area the health risks connected to overexposure to UV solar radiation of outdoor workers seems underestimated for the absence of publications regarding measurement campaign on this theme.

Instead, the article selected contains significant measured data, of several work typologies, environmental settings and body districts. Results of these measurement campaigns could compose the first step of a database collecting and systematizing the erythema exposure of outdoor workers.

REFERENCES

- [1] World Health Organization, World Meteorological Organization, United Nations Environment Programme, International Commission of Non-Ionizing Radiation Protection, "Global Solar UV Index: A Practical Guide", Geneva, Switzerland, 2002.
- [2] <https://www.temis.nl/uvradiation/UVarchive>
- [3] A. Milon, P-E. Sottas, J-L. Bulliard, D. Vernez, "Effective exposure to solar UV in building workers: influence of local and individual factors", *Journal of Exposure Science and Environmental Epidemiology*, vol.17, pp. 58-68, 2007.
- [4] M.A. Serrano, J. Cañada, J.C. Moreno, "Erythema Ultraviolet Exposure in Two Groups of Outdoor Workers in Valencia, Spain", *Photochemistry and Photobiology*, vol. 85, pp. 1468-1473, 2009.
- [5] R. Sisto, M. Borra, G.R. Casale, A. Militello, A.M. Siani, "Quantitative evaluation of personal exposure to UV radiation of workers and general public", *Radiation Protection Dosimetry*, vol. 137, n. 3-4, pp. 193-196, 2009.
- [6] M.A. Serrano, J. Cañada, J.C. Moreno, "Solar UV exposure in construction workers in Valencia, Spain", *Journal of Exposure Science and Environmental Epidemiology*, vol. 23, pp. 525-530, 2013.
- [7] L. Miligi, A. Benvenuti, P. Legittimo, A.M. Badiali, V. Cacciarini, A. Chiarugi et al., "Solar ultraviolet radiation risk in outdoor workers: a specific project of Tuscany Region (Italy)", *Epidemiologia e Prevenzione*, vol. 37, Gen-Feb. 2013.
- [8] G. Nardini, D. Neri, M. Paroncini, "Measured Anatomical Distribution of Solar UVR on Strawberry Production Workers in Italy", *Journal of Agricultural Safety and Health*, vol. 20, pp. 67-78, 2014.
- [9] A. Modenese, F.P. Ruggieri, F. Bisegna, M. Borra, C. Burattini, E. Della Vecchia, C. Grandi, A. Grasso, L. Gugliemetti, M. Manini, A. Militello, F. Gobba, "Occupational Exposure to Solar UV Radiation of a Group of Fishermen Working in the Italian North Adriatic Sea", *International Journal of Environmental Research and Public Health*, vol. 16, pp. 3001, 2019.
- [10] M. Wittlich, S.M. John, G.S. Tiplica, C.M. Sălăvăstru, A.I. Butacu, A. Modenese, V. Paolucci, D. D'Hauw, F. Gobba, P. Sartorelli, J. Macan, J. Kovačić, K. Grandahl, H. Moldovan, "Personal solar ultraviolet radiation dosimetry in an occupational setting across Europe", *Journal of European Academy of Dermatology and Venereology*, vol. 34, pp. 1835-1841, 2020.
- [11] J. Kovačić, M. Wittlich, S. Malte John, J. Macan, "Personal ultraviolet radiation dosimetry and its relationship with environmental data: A longitudinal pilot study in Croatian construction workers", *Journal of Photochemistry and Photobiology, B: Biology*, vol. 207, pp. 111866, 2020.
- [12] A. Modenese, F. Gobba, V. Paolucci, S. Malte John, P. Sartorelli, M. Wittlich, "Occupational solar UV exposure in construction workers in Italy: results of one-month of monitoring with personal dosimeters", *Proceedings of the 2020 IEEE International Conference on Environment and Electrical Engineering and 2020 IEEE Industrial and Commercial Power Systems Europe*, Madrid, Spain, 9-12 June 2020.
- [13] A. Modenese, F. Bisegna, M. Borra, C. Burattini, L. Gugliemetti, F. Larese Filon, A. Militello, P. Toffanin, F. Gobba, "Occupational Exposure to Solar UV Radiation in a Group of Dock-workers in North-East Italy", *Proceedings of the 2020 IEEE International Conference on Environment and Electrical Engineering and 2020 IEEE Industrial and Commercial Power Systems Europe*, Madrid, Spain, 9-12 June 2020.
- [14] M. Borra, C. Grandi, A. Militello, C. Burattini, L. Gugliemetti, A. Mangione, F. Bisegna, "Developing an algorithm to assess the UV erythema dose for outdoor workers", *Proceedings - 2018 IEEE International Conference on Environment and Electrical Engineering and 2018 IEEE Industrial and Commercial Power Systems Europe, IEEEIC/I and CPS Europe 2018*. DOI: 10.1109/IEEEIC.2018.8494540