

The Interplay Between Problematic Online Pornography Use, Psychological Stress, Emotion Dysregulation and Insomnia Symptoms During the COVID-19 Pandemic: A Mediation Analysis

Alessandro Musetti^{1,*}, Alessio Gori^{2,*}, Alessia Alessandra³, Eleonora Topino⁴, Grazia Terrone⁵, Giuseppe Plazzi^{6,7}, Marco Cacioppo⁴, Christian Franceschini³

¹Department of Humanities, Social Sciences and Cultural Industries, University of Parma, Parma, Italy; ²Department of Health Sciences, University of Florence, Florence, Italy; ³Department of Medicine and Surgery, University of Parma, Parma, Italy; ⁴Department of Human Sciences, Lumsa University of Rome, Rome, Italy; ⁵Department of History, Cultural Heritage, Education and Society, University of Rome Tor Vergata, Rome, Italy; ⁶Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy; ⁷IRCCS Institute of Neurological Sciences of Bologna (ISNB), Bologna, Italy

Correspondence: Alessandro Musetti, Department of Humanities, Social Sciences and Cultural Industries, University of Parma, Borgo Carissimi 10, Parma, 43121, Italy, Tel +39 0521 034820, Email alessandro.musetti@unipr.it

*These authors contributed equally to this work

Purpose: Although a link between problematic online pornographic use (POPU) and insomnia symptoms has been established, psychological and psychopathological mechanisms underlying this relationship are still not clear. Psychological stress and emotion dysregulation have been pointed out as relevant in the development and maintenance of insomnia. This study aims to explore the associations between POPU, psychological stress, emotion dysregulation and insomnia symptoms and to understand the mediating role of psychological stress and emotion dysregulation in the relationship between POPU and insomnia symptoms.

Participants and Methods: A sample of 776 Italian adults aged 19–48 years (51.4% female; M age = 28.49; SD = 7.33) completed questionnaires regarding demographics, COVID-19-related variables, POPU, psychological stress, emotion dysregulation, and insomnia symptoms.

Results: After controlling for demographic covariates and COVID-19-related variables, multiple mediation model showed that higher psychological stress and emotion dysregulation fully mediated the link between POPU and insomnia.

Conclusion: The findings underscore the significance of the negative consequences of POPU and underline the importance of working on this and its effects on psychological stress and emotion dysregulation to limit insomnia.

Keywords: addictive behaviors, online pornography, problematic online pornographic use, stress, emotion dysregulation, clinical psychology, insomnia

Introduction

Insomnia is a condition characterized by difficulty in initiating or maintaining sleep associated with psychiatric and physical comorbidities which constitutes a significant public health problem.¹ It impairs an individual daily life and social functioning by increasing fatigue, daytime sleepiness and cognitive impairments.² While insomnia disorder has an estimated prevalence of 5–15%,³ insomnia symptoms have been observed to occur with estimates ranging 30–50% among the adult population.⁴

Given that insomnia is a heterogeneous clinical condition, several models have been developed to explain its multifaceted pathophysiology. According to the well-established Neurocognitive Model, insomnia is highly dependent on individuals' sleep system response to external stressors and dysfunctional coping strategies.^{5–7} More specifically,

individual proneness represents a predisposing factor that, with the occurrence of stressful life events, could precipitate acute insomnia episodes which may, in turn, develop into chronic insomnia through maladaptive behavioral coping strategies as a result of conditioned arousal.⁷

In this regard, with the diffusion of the Internet and digital technologies, online activities become additional ways to cope with stressful situations.⁸ Although involvement in such activities typically does not imply negative consequences, for a minority of individuals it can lead to reduced social interactions and impaired daily activities.⁹ Notably, recent research provided evidence for the association between different forms of problematic Internet use,¹⁰ and the onset and maintaining of sleep disorders. In fact, excessive exposure to artificial blue-spectrum light, usually for internet browsing, can lead to melatonin suppression.¹¹ Furthermore, users' inability to stop Internet usage (eg, for playing video games or accessing pornography) can negatively affect their sleep-wake schedule.⁸ In this regard, problematic online pornographic use (POPU) can be considered a specific type of problematic Internet use^{12,13} that includes any use of online pornography leading to significant negative interpersonal or personal consequences for the user.¹⁴ In fact, POPU has been associated with higher relational problems^{15,16} and lower mental health.¹⁷ Notably, pornography, for its characteristics, may play a specific role in worsening users' sleep quality¹⁸ because it includes "any material designed to cause or enhance sexual arousal or sexual excitement in the viewer",¹⁹ and excessive pornography use is associated with increased stress and anxiety,²⁰ social impairment and isolation,²¹ sexual dysfunctions²² and sleep disturbances.¹¹ Indeed, Cacioppo et al¹⁶ found that POPU was linked to psychological distress and emotional difficulties, which, in turn, were found to be associated with sleep disturbance.^{23–25} In the same line, effective emotion regulation abilities were found to be associated with sexual satisfaction and lower levels of POPU by Pepping et al.²⁶

Dysfunctional ways of dealing with stressful situations have been seen to affect the sleep-wake schedule. Alterations in cognitive arousal systems are traditionally considered a risk factor of insomnia symptoms⁵ and the ability of an individual to regulate emotion plays a role in decreasing the detrimental effects of emotional stress on sleep physiology.²⁷ Balleisio et al²⁸ observed that higher pre-sleep intrusive thoughts affected sleep continuity. Similarly, Li et al²⁹ and Du et al³⁰ showed that pre-sleep rumination, as a dysfunctional emotion regulation strategy, could exacerbate anxiety resulting in disrupted sleep, and that resilience, as an effective self-adjustment to stressful situations, would moderate stress influence on sleep. A dysregulated stress response has been shown to disrupt prefrontal control involved in emotion regulation, thus leading to one's inability to properly adapt to situational demands³¹ which lead, in turn, to worsening of symptoms such as sleep disturbances. Moreover, maladaptive self-regulation strategies, such as problematic internet use and POPU, are associated with an increase in psychological stress.³²

In sum, the available literature reports consistent associations between POPU, psychological stress, difficulties in regulating emotions and insomnia symptoms.^{27,33} However, to date, no study has investigated the associations between these variables in a single model, thus limiting our ability to comprehend how these related factors operate as a whole. Therefore, this study aims to further understand the link between POPU and insomnia symptoms taking into account the mediation role played in this relationship by psychological stress and emotion dysregulation. In addition, given the negative impact of COVID-19 pandemic on insomnia symptoms,^{34,35} we also controlled for COVID-19 related variables.

The current study has two aims: (1) to explore the multiple mediating role of emotion dysregulation and psychological stress between POPU and insomnia symptoms; (2) to examine the sequential relationship between stress and emotion dysregulation. Specifically, we hypothesized that:

H1. High POPU would be associated with increased insomnia symptoms;

H2. Psychological stress would mediate the association between POPU and insomnia symptoms;

H3. Emotion dysregulation would mediate the association between POPU and insomnia symptoms;

H4. Psychological stress and emotion dysregulation would sequentially mediate the association between POPU and insomnia symptoms.

Materials and Methods

Procedure

Data were collected from general population following a snowball sampling method.

Inclusion criteria were a) age of 18 years or older, b) Italian native language, and c) living in Italy during the second wave of COVID-19. Participants were requested to fill out a 20-min anonymous web survey on the Microsoft Azure platform during the second wave of the COVID-19 pandemic in Italy (18 December 2020–18 January 2021). The web-survey was promoted via university platforms and on different social media and virtual learning environments. Furthermore, respondents were asked to spread the survey among acquaintances. Participants signed the informed electronic consent form before accessing the survey. The study was approved by the local ethics committee (CERIP – Comitato Etico del Centro di Ricerca e di Intervento Psicologico – University of Messina, prot. n. 17758). All procedures were conducted in accordance with the Declaration of Helsinki and its later advancements.

Measures

Socio-Demographic Information

In this section information about gender, age, nationality, ethnicity, education, family (ie, single, married or in a stable relationship, divorced or separated, and widow) and occupational statuses (ie, student, working student, employed, freelance, unemployed, and retired) was collected.

COVID-19-Related Variables

Three ad hoc items (yes/no) were used to assess the participant direct or indirect experience related to COVID-19 (ie, personal or someone close infection of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and quarantine history in the last 3 months).

Problematic Online Pornography Use

The Cyber Pornography Addiction Test (CYPAT)¹⁶ was used for assessing the risk of POPU. It is a self-report scale consisting of 11 items (eg, “I told myself to stop using online pornography but I didn’t succeed”) scored on a 5-point Likert scale (1 = never, 5 = always). The total score ranged from 11 to 55, with higher scores indicating greater POPU. No cut-off score has been developed for this scale. Following Jacobs et al,³⁶ we used a total CYPAT score range of 23 to 55, to identify users with high levels of POPU. Previous studies showed good psychometric properties and validity of this instrument.^{16,37} In the present study, Cronbach’s α was 0.93.

Emotion Dysregulation

The Italian version of the Difficulties in Emotion Regulation Scale (DERS;³⁸ original version by Gratz and Roemer)³⁹ was used to assess participants’ emotion dysregulation. It is a self-report scale containing 36 items (eg, “When I’m upset, I acknowledge my emotions”). Participants were asked to rate their responses on a 5-point Likert scale, ranging from 1 (Almost never) to 5 (Almost always). Higher total scores indicate greater emotion dysregulation. The DERS is a widely used tool that demonstrates good psychometric properties in different settings and cultures.^{40,41} In the present study, Cronbach’s α was 0.93.

Psychological Stress

The Stress subscale of the Italian version of the Depression Anxiety Stress Scales-21 (DASS-21-S;⁴² original version by Lovibond and Lovibond 1995)⁴³ was adopted to evaluate participants psychological stress. The DASS-21-S subscale includes 7 items (eg, “I felt that I was using a lot of nervous energy”) rated on a 4-points Likert scale ranging from 0 (Did not apply to me at all) to 3 (Applied to me very much, or most of the time). Subscale scores range from 0 to 21, where higher scores indicate greater psychological stress. The DASS-21 is an internationally recognized screening tool for psychological stress that showed good psychometric properties.⁴⁴ In the present study, Cronbach’s α was 0.90.

Insomnia Symptoms

The Insomnia Severity Index (ISI)⁴⁵ was used to assess participants’ insomnia symptoms that occurred within the

previous 2 weeks. It is a self-report instrument that includes seven items (eg, “How worried/distressed are you about your current sleep problem?”) rated on a 5-point Likert scale from 0 (Not at all) to 4 (Very much). The total score ranges from 0 to 28, where higher scores indicate higher levels of insomnia symptoms. The ISI has been shown to be a reliable and valid screening tool for insomnia symptoms.⁴⁶ In the present study, Cronbach’s α was 0.85.

Data Analytic Strategy

All statistical analyses were performed using SPSS 21.0 for Mac. Descriptive statistics were computed for all study variables. Univariate normality was examined through skewness and kurtosis.

Correlations between variables were computed using point-biserial correlations between dichotomous and continuous variables and Pearson’s correlation for continuous variables.

The hypothetical mediating role of psychological stress and emotion dysregulation in the relationship between POPU and insomnia was tested using Hayes’ SPSS macro PROCESS.⁴⁷ Specifically, Model 6 was performed to assess the significance of all the hypothesized relation between the study variables, controlling for both demographic covariates (ie, age and gender) and COVID-19 related variables (ie, someone close infected with SARS-CoV-2, having been in quarantine, being infected-with SARS-CoV-2). The 95% bias-corrected confidence interval (CI) was examined on 5000 bootstrap samples. Bootstrapping is a random resampling method that “makes no assumption about the shape of the distributions of the variables or the sampling distribution of the statistic” (p. 722).⁴⁸ The indirect effect of POPU on insomnia through psychological stress and emotion dysregulation is considered statistically significant if CI does not contain zero.⁴⁷ All the results with an α value lower than 0.05 were considered significant.

Results

Sample Socio-Demographics

Analyses were performed on a total of 776 participants, aged 19–48 years ($M = 28.49$, $SD = 7.33$), that voluntarily took part in our survey. Of the total participants, 399 (51.42%) were male, and 756 (97.42%) of Italian origin. As for their ethnicity, 765 (98.58%) were European, 10 (1.29%) were African, and 1 Latino-American (0.13%). Most of the participants completed secondary higher education ($n = 501$, 64.56%) and were single ($n = 432$, 55.67%). With regard to occupational status, 309 (39.82%) participants were employees. There was no significant association between participants’ gender and age ($t_{(774)} = -1.21$, $p = 0.23$).

Preliminary Analysis

Descriptive statistics for the scales used in this study are reported in Table 1. As expected for a nonclinical sample, the mean score of the CYPAT was in the nonclinical range. Specifically, 45 participants (11.3%) showed a high level of POPU. In addition, 385 (49.61%) participants reported to have had a close one infected with SARS-CoV-2, 149 (19.20%), had a quarantine history and 57 (7.35%) had been infected with SARS-CoV-2 in the past 3 months.

As shown in Table 2, correlation analysis revealed a pattern of significant and positive associations between all main investigated variables. In addition, being male was significantly and positively associated with higher levels of POPU, as well as significantly and negatively related with insomnia symptoms and stress. Younger age was significantly and

Table 1 Descriptive Statistics for the Used Scales (N = 776)

	Ranges	Overall Mean	Std. Deviation	Skewness	Kurtosis
CYPAT	1–53	13.61	5.88	2.81	9.01
DASS-21-S	0–21	8.83	5.06	0.31	–0.44
DERS	40–160	83.91	21.80	0.52	–0.40
ISI	0–27	5.54	4.60	1.10	1.27

Abbreviations: ISI, Insomnia Severity Index; CYPAT, Cyber Pornography Addiction Test; DERS, Difficulties in Emotion Regulation Scale; DASS-21-S, Depression Anxiety Stress Scales-21-Stress.

Table 2 Intercorrelations Between Variables of Interest

	1	2	3	4	5	6	7	8	9
1. Age	1	0.04	0.00	0.08*	0.02	-0.04	-0.03	-0.21**	-0.22**
2. Gender		1	-0.03	-0.09*	-0.07*	-0.08*	0.20**	-0.03	-0.08*
3. COVID1			1	0.29**	0.16**	-0.06	0.07	-0.02	-0.05
4. COVID2				1	0.44**	-0.02	-0.05	-0.04	-0.01
5. COVID3					1	-0.02	-0.03	-0.04	-0.13
6. ISI						1	0.10**	0.34**	0.46**
7. CYPAT							1	0.30**	0.14**
8. DERS								1	0.54**
9. DASS-21-S									1

Notes: Bold values indicate significant p-values. *Correlation is significant at the 0.05 level (2-tailed). **Correlation is significant at the 0.01 level (2-tailed).

Abbreviations: COVID1, someone close infected with SARS-CoV-2; COVID2, having been in quarantine; COVID3, being infected with SARS-CoV-2; ISI, Insomnia Severity Index; CYPAT, Cyber Pornography Addiction Test; DERS, Difficulties in Emotion Regulation Scale; DASS-21-S, Depression Anxiety Stress Scales-21-Stress.

positively associated with higher psychological stress and emotion dysregulation. No significant associations were found between COVID-19-related variables and main investigated variables.

The Mediation Model

In order to test the mediating effect of stress and emotion dysregulation on the relationship between POPU use and insomnia, a multiple mediational analysis has been conducted. Results supported the hypothesized model. Controlling for age ($B = 0.05$, $t = 2.35$, $p = 0.019$), gender (females coded as 1, males coded as 2; $B = -0.46$, $t = -1.52$, $p = 0.13$), and COVID-19 related variables (someone close infected with SARS-CoV-2 coded as 1, someone close not infected with SARS-CoV-2 coded as 2, $B = -0.34$, $t = -1.11$, $p = 0.27$; having been in quarantine coded as 1, having not been in quarantine coded as 2, $B = -0.11$, $t = -0.25$, $p = 0.80$; being infected with SARS-CoV-2 coded as 1, having not been infected with SARS-CoV-2 coded as 2, $B = -0.12$, $t = -0.21$, $p = 0.83$), POPU positively predicted psychological stress ($B = 0.14$, $t = 4.63$, $p < 0.001$) and emotion dysregulation ($B = 0.87$, $t = 4.81$, $p < 0.001$). Psychological stress positively predicted emotion dysregulation ($B = 2.06$, $t = 15.78$, $p < 0.001$). Both psychological stress and emotion dysregulation positively predicted insomnia ($B = 0.36$, $t = 10.36$, $p < 0.001$ and $B = 0.03$, $t = 3.62$, $p < 0.001$, respectively). Direct effect ($B = 0.01$, $t = 0.53$, $p = 0.60$), and total effect ($B = 0.10$, $t = 3.50$, $p < 0.001$) of POPU on insomnia were not significant and significant, respectively.

Analysis of the bias-corrected confidence intervals of the indirect effects in the bootstrapped samples further revealed that all indirect effects of the model were significant, further corroborating the sequential mediation of psychological stress and emotion dysregulation (with scores of psychological stress predicting emotion dysregulation) in the relationship between POPU and insomnia. Thus, higher psychological stress and emotion dysregulation fully mediated the link between POPU and insomnia. The total model was significant, $F(8767) = 29.45$, $p < 0.001$ and accounted for a significant amount of variance in adolescent problematic SNSs use ($R^2 = 0.235$), thus supporting the hypothesized path in the relationships between the investigated variables.

Discussion

This study aims to investigate the interplay between POPU psychological stress, emotion dysregulation and insomnia symptoms during the second wave of the COVID-19 in Italy. We hypothesized that psychological stress and emotion dysregulation would play a mediating role in the relationship between POPU and insomnia symptoms. In particular, we expected that both psychological stress and emotion dysregulation would be associated to higher levels of insomnia symptoms mediating its association with POPU. In addition, given the documented negative implications of the COVID-

19 on several aspects of sleep quality,⁴⁹ we also examined the role of COVID-19 related variables. Surprisingly, no significant associations were found between COVID-19-related variables and insomnia symptoms, as well as, POPU, psychological stress, and emotion dysregulation. This result must be seen in light of the characteristics of the sample. In fact, the participants of this study were mostly indirectly affected by the COVID-19 pandemic. More specifically, about half of the participants reported having had a close one infected with SARS-CoV-2, while only a small minority was personally infected. It is possible that, due to the snowball sampling method used, people with more negative experiences with COVID-19 pandemic decided not to take part in the survey. Therefore, further research is needed on more representative samples.

As for the relationship between investigated variables and age, in line with the literature, we found that younger age was significantly and positively associated with higher psychological stress⁴² and emotion dysregulation.⁵⁰ Indeed, the pandemic had a significant influence on mental health^{51,52} with particularly marked consequences for some subcategories of the population, including young adults.⁵³ Among young adults, in fact, an increase in both internalizing (eg, withdrawal, somatic complaints, anxiety, and depression) and externalizing (eg, aggressive and rule-breaking behaviors) problems were found.⁵⁴ Consistent with previous study we also found that being female was significantly and positively associated with higher levels insomnia symptoms⁵⁵ and psychological stress.⁵¹ Indeed, females were found to exhibit higher anxiety and perception of danger to the COVID-19 virus than males⁵⁶ and this could impact the sleep quality.⁵⁷ Furthermore, a significant and positive association between being male and POPU was shown, in line with previous research.^{16,58}

Concerning our hypotheses, these have all been confirmed. As a result of the performed correlation analysis, a significant positive association was found between POPU and insomnia symptoms. This echoes previous evidence,¹⁸ and is also in line with research into the broader dimension of problematic Internet use.⁵⁹ However, mediation analyses also highlighted that the relationship between POPU and insomnia symptoms occurs mainly through an indirect way, with the total mediation by psychological stress and emotion dysregulation.

Indeed, POPU was found to be associated with psychological stress and this corroborates previous studies that pointed out that not so much of the average daily use of pornography itself, but rather the levels of compulsiveness and perceived uncontrolled use are significantly associated with emotional distress,^{14,60} with difficulty in controlling the use of pornography related to severe negative consequences.⁶¹ Consistently, other research also showed that the problematic use of online sexual activities was associated with an increase in symptoms of anxiety, depression, and dissociation (see Hermand et al⁶² for a review).

Furthermore, both POPU and psychological stress had a significant positive effect on emotion dysregulation. Individuals with POPU have in fact been described as having poor coping strategies and ineffective emotion regulation skills.^{60,63} In this regard, although the literature highlights the etiological role of emotion dysregulation for both substance related^{64,65} and behavioral addictions,^{64,66,67} including the sexual one,⁶⁸ the data presented suggest that this can be further fueled by addictive behavior. The online pornography use can become problematic if used as a dysfunctional coping strategy for the regulation of affective states: this does not favor the development of more adaptive strategies in the subject, but on the contrary, stimulates a compulsive attitude towards cyberpornography feeding the state of dysregulation in the long term, in line with when the subject with sexual addiction is found.^{68,69}

Finally, data showed that both psychological stress and emotion dysregulation were associated with greater insomnia symptoms, further supporting and enriching the associations highlighted in scientific literature concerning the impact of distress and arousal (see Bos and Macedo⁷⁰ for a review). Indeed, Tsypes et al⁷¹ found that individuals with generalized anxiety disorder showed greater difficulties in emotion regulation, with a significant influence on a wide range of sleep problems. Parallely, previous research also highlighted that negative effects influenced sleep quality, both directly and indirectly through emotion dysregulation.⁷²

Summing up, this study highlights a significant total effect of the POPU on insomnia symptoms which is mainly explained through an indirect path that involves the effects of psychological stress and emotion dysregulation. This appears particularly significant, especially in light of the effects that POPU (see Mauer-Vakil and Bahji⁷³ for a review), psychological stress,⁷⁴ emotion dysregulation,⁷⁵ and insomnia symptoms⁷⁶ may have on mental health. Furthermore,

these data appear particularly consistent and relevant also with respect to the current evidence related to the pandemic, which highlights high levels of POPU, psychological stress, emotion dysregulation, and insomnia symptoms.^{58,77–79}

However, these findings have some limitations that should be noted. First, the cross-sectional nature of the study makes it impossible to draw inferences about causality in the associations between the variables. In fact, the mediation model was evoked based on theory, and it is not possible to rule out a reverse causation model in which increased insomnia symptoms generate psychological stress, emotion dysregulation and POPU. Therefore, we used the term “mediation” only in the statistical sense. Future research should use longitudinal designs to determine the direction of these associations. Furthermore, the use of a convenience snowball sample does not allow to calculate the response rate and the online diffusion of the survey limits our knowledge about participants' motivation to take part in the research. Thus, the generalizability of the findings is necessarily limited. Further studies with random or nationwide representative samples are warranted. Another limitation of this study includes using a sample with low levels of POPU and insomnia symptoms. However, our results are consistent with previous studies on POPU¹⁶ and insomnia symptoms⁸⁰ conducted in the normal population. Further studies with clinical samples are needed to confirm and expand our findings.

Finally, the use of self-report measure exposes to the possibility of well-known biases (eg, the desirability one). The use of a multi-method technique for assessment and measurement (eg, integrating with interviews) could be an important goal for future research to overcome this issue.

Conclusion

This study offered an in-depth exploration of the relationship between POPU and insomnia symptoms, showing the significant mediation role of psychological stress and emotion dysregulation. These data may have important practical implication and provided useful indication to elaborate effective clinical interventions. Indeed, these results enrich the framework of the negative consequences of POPU and underline the importance of working on this and its effects on psychological stress and emotion dysregulation to limit insomnia symptoms. In fact, our results suggest that taking into account negative consequences of POPU on users' psychological functioning could be more informative than solely screening for POPU (eg, by the use of criteria derived from substance use disorders). This process-based approach to clinical practice (eg, Kinderman et al⁸¹) may allow to avoid the overpathologization of a common behavior, such as online pornography use, and it may help clinicians to recognize psychological mechanisms underlying symptom presentation.

Ethics Statement

All participants provided written informed consent and were informed about the purpose of the study. Data were collected and used complying with the Italian ethical standards and with the Declaration of Helsinki. The study was approved by the local ethics committee (CERIP – Comitato Etico del Centro di Ricerca e di Intervento Psicologico – University of Messina, prot. n. 17758) as the University of Parma's Research Ethics Board was not established yet at the time of project evaluation.

Disclosure

Alessandro Musetti and Alessio Gori are co-first authors for this study. GP participated in advisory board for UCB Pharma, Jazz Pharmaceuticals, Bioproject, Takeda and Idorsia. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. Bjorvatn B, Jernelöv S, Pallesen S. Insomnia – a heterogenic disorder often comorbid with psychological and somatic disorders and diseases: a narrative review with focus on diagnostic and treatment challenges. *Front Psychol.* 2021;12. doi:10.3389/fpsyg.2021.639198.
2. Edinger JD, Arnedt JT, Bertisch SM, et al. Behavioral and psychological treatments for chronic insomnia disorder in adults: an American Academy of Sleep Medicine systematic review, meta-analysis, and GRADE assessment. *J Clin Sleep Med.* 2021;17(2):263–298. doi:10.5664/JCSM.8988
3. Ohayon MM. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Med Rev.* 2002;6(2):97–111. doi:10.1053/smr.2002.0186

4. Buysse DJ, Angst J, Gamma A, Ajdacic V, Eich D, Rössler W. Prevalence, course, and comorbidity of insomnia and depression in young adults. *Sleep*. 2008;31(4):473–480. doi:10.1093/sleep/31.4.473
5. Harvey AG. A cognitive model of insomnia. *Behav Res Ther*. 2002;40(8):869–893. doi:10.1016/S0005-7967(01)00061-4
6. Kalmbach DA, Anderson JR, Drake CL. The impact of stress on sleep: pathogenic sleep reactivity as a vulnerability to insomnia and circadian disorders. *J Sleep Res*. 2018;27(6):e12710. doi:10.1111/jsr.12710
7. Perlis ML, Giles DE, Mendelson WB, Bootzin RR, Wyatt JK. Psychophysiological insomnia: the behavioural model and a neurocognitive perspective. *J Sleep Res*. 1997;6(3):179–188. doi:10.1046/j.1365-2869.1997.00045.x
8. Kardefelt-Winther D. A conceptual and methodological critique of internet addiction research: towards a model of compensatory internet use. *Comput Human Behav*. 2014;31(1):351–354. doi:10.1016/j.chb.2013.10.059
9. Billieux J, Schimmenti A, Khazaal Y, Maurage P, Heeren A. Are we overpathologizing everyday life? A tenable blueprint for behavioral addiction research. *J Behav Addict*. 2015;4(3):119–123. doi:10.1556/2006.4.2015.009
10. Alimoradi Z, Lin CY, Broström A, et al. Internet addiction and sleep problems: a systematic review and meta-analysis. *Sleep Med Rev*. 2019;47:51–61. doi:10.1016/j.smrv.2019.06.004
11. Small GW, Lee J, Kaufman A, et al. Brain health consequences of digital technology use. *Dialogues Clin Neurosci*. 2020;22(2):179–187. doi:10.31887/DCNS.2020.22.2/gsmall
12. Brand M, Young KS, Laier C, Wöfling K, Potenza MN. Integrating psychological and neurobiological considerations regarding the development and maintenance of specific Internet-use disorders: an Interaction of Person-Affect-Cognition-Execution (I-PACE) model. *Neurosci Biobehav Rev*. 2016;71:252–266. doi:10.1016/j.neubiorev.2016.08.033
13. Laier C, Brand M. Empirical evidence and theoretical considerations on factors contributing to cybersex addiction from a cognitive-behavioral view. *Sex Addict Compulsivity*. 2014;21(4):305–321. doi:10.1080/10720162.2014.970722
14. Grubbs JB, Volk F, Exline JJ, Pargament KI. Internet pornography use: perceived addiction, psychological distress, and the validation of a brief measure. *J Sex Marital Ther*. 2015;41(1):83–106. doi:10.1080/0092623X.2013.842192
15. Manning JC. The impact of internet pornography on marriage and the family: a review of the research. *Sex Addict Compulsivity*. 2006;13(2–3):131–165. doi:10.1080/10720160600870711
16. Cacioppo M, Gori A, Schimmenti A, Baiocco R, Laghi F, Caretti V. Development of a new screening tool for cyber pornography: psychometric properties of the cyber pornography addiction test (CYPAT). *Clin Neuropsychiatry*. 2018;15(1):60–65.
17. Nelson LJ, Padilla-Walker LM, Carroll JS. “I believe it is wrong but I still do it”: a comparison of religious young men who do versus do not use pornography. *Psycholog Relig Spiritual*. 2010;2(3):136–147. doi:10.1037/a0019127
18. Shirk SD, Saxena A, Park D, Kraus SW. Predicting problematic pornography use among male returning US veterans. *Addict Behav*. 2021;112:106647. doi:10.1016/j.addbeh.2020.106647
19. Kraus S, Rosenberg H. The pornography craving questionnaire: psychometric properties. *Arch Sex Behav*. 2014;43(3):451–462. doi:10.1007/s10508-013-0229-3
20. Odaci H, Çikrikci Ö. An exploration of the associations among internet use, depression, anxiety and stress among youths. *Mediterr J Clin Psychol*. 2017;5(3):1–16. doi:10.6092/2282-1619/2017.5.1635
21. Camilleri C, Perry JT, Sammut S. Compulsive Internet pornography use and mental health: a cross-sectional study in a sample of university students in the United States. *Front Psychol*. 2021;11:1–24. doi:10.3389/fpsyg.2020.613244
22. Whelan G, Brown J. Pornography addiction: an exploration of the association between use, perceived addiction, erectile dysfunction, premature (early) ejaculation, and sexual satisfaction in males aged 18–44 years. *J Sex Med*. 2021;18(9):1582–1591. doi:10.1016/j.jsxm.2021.06.014
23. Lin CY, Broström A, Griffiths MD, Pakpour AH. Investigating mediated effects of fear of COVID-19 and COVID-19 misunderstanding in the association between problematic social media use, psychological distress, and insomnia. *Internet Interv*. 2020;21:100345. doi:10.1016/j.invent.2020.100345
24. Galbiati A, Sforza M, Fasiello E, et al. The association between emotional dysregulation and REM sleep features in insomnia disorder. *Brain Cogn*. 2020;146:105642. doi:10.1016/j.bandc.2020.105642
25. Kalmbach DA, Cuamatzi-Castelan AS, Tonnu CV, et al. Hyperarousal and sleep reactivity in insomnia: current insights. *Nat Sci Sleep*. 2018;10:193–201. doi:10.2147/NSS.S138823
26. Pepping CA, Cronin TJ, Lyons A, Caldwell JG. The effects of mindfulness on sexual outcomes: the role of emotion regulation. *Arch Sex Behav*. 2018;47(6):1601–1612. doi:10.1007/s10508-017-1127-x
27. Vandekerckhove M, Wang YL. Emotion, emotion regulation and sleep: an intimate relationship. *AIMS Neurosci*. 2018;5(1):1–17. doi:10.3934/Neuroscience.2018.1.1
28. Ballesio A, Ghezzi V, Vacca M, Ottaviani C, Lombardo C. Effects of presleep cognitive intrusions on subjective sleep and next-day cognitive performance in insomnia. *Behav Ther*. 2020;51(5):688–699. doi:10.1016/j.beth.2019.09.003
29. Li Y, Gu S, Wang Z, et al. Relationship between stressful life events and sleep quality: rumination as a mediator and resilience as a moderator. *Front Psychiatry*. 2019;10:1–9. doi:10.3389/fpsyg.2019.00348
30. Du C, Zan MCH, Cho MJ, et al. Increased resilience weakens the relationship between perceived stress and anxiety on sleep quality: a moderated mediation analysis of higher education students from 7 countries. *Clocks Sleep*. 2020;2(3):334–353. doi:10.3390/clocksleep2030025
31. Roos LE, Knight EL, Beauchamp KG, et al. Acute stress impairs inhibitory control based on individual differences in parasympathetic nervous system activity. *Biol Psychol*. 2017;125:58–63. doi:10.1016/j.biopsycho.2017.03.004
32. Deatherage S, Servaty-Seib HL, Aksoz I. Stress, coping, and internet use of college students. *J Am Coll Health*. 2014;62(1):40–46. doi:10.1080/07448481.2013.843536
33. Pettorrosso M, Valle S, Cavic E, Martinotti G, Di Giannantonio M, Grant JE. Problematic Internet Use (PIU), personality profiles and emotion dysregulation in a cohort of young adults: trajectories from risky behaviors to addiction. *Psychiatry Res*. 2020;289:113036. doi:10.1016/j.psychres.2020.113036
34. Liu C, Pan W, Li L, Li B, Ren Y, Ma X. Prevalence of depression, anxiety, and insomnia symptoms among patients with COVID-19: a meta-analysis of quality effects model. *J Psychosom Res*. 2021;147:5. doi:10.1016/j.jpsychores.2021.110516
35. Jahrami H, BaHammam AS, Bragazzi NL, Saif Z, Faris M, Vitiello MV. Sleep problems during the COVID-19 pandemic by population: a systematic review and meta-analysis. *J Clin Sleep Med*. 2021;17(2):299–313. doi:10.5664/jcsm.8930

36. Jacobs T, Geysmans B, Van HG, et al. Associations between online pornography consumption and sexual dysfunction in young men: multivariate analysis based on an international web-based survey. *JMIR Public Heal Surveill.* 2021;7(10):e32542. doi:10.2196/32542
37. Fernandez DP, Griffiths MD. Psychometric instruments for problematic pornography use: a systematic review. *Eval Heal Prof.* 2021;44(2):111–141. doi:10.1177/0163278719861688
38. Sighinolfi C, Pala AN, Chiri LR, Marchetti I, Sica C. Difficulties in Emotion Regulation Scale (DERS): traduzione e adattamento italiano. [Difficulties in Emotion Regulation Scale (DERS): the Italian translation and adaptation.]. *Psicoter Cogn e comport.* 2010;16(2):141–170.
39. Gratz KL, Roemer M. Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of the Difficulties In Emotion Regulation Scale. *J Psychopathol Behav Assess.* 2004;26(1):41–54. doi:10.1023/B:JOBA.0000007455.08539.94
40. Hallion LS, Steinman SA, Tolin DF, Diefenbach GJ. Psychometric properties of the Difficulties in Emotion Regulation Scale (DERS) and its short forms in adults with emotional disorders. *Front Psychol.* 2018;9:1–12. doi:10.3389/fpsyg.2018.00539
41. Sörman K, Garke M, Isacson NH, et al. Measures of emotion regulation: convergence and psychometric properties of the difficulties in emotion regulation scale and emotion regulation questionnaire. *J Clin Psychol.* 2021;1–17. doi:10.1002/jclp.23206.
42. Bottesi G, Ghisi M, Altoè G, Conforti E, Melli G, Sica C. The Italian version of the depression anxiety stress scales-21: factor structure and psychometric properties on community and clinical samples. *Compr Psychiatry.* 2015;60:170–181. doi:10.1016/j.comppsy.2015.04.005
43. Lovibond SH, Lovibond PF. *Manual for the Depression Anxiety Stress Scales.* Psychology Foundation of Australia; 1995.
44. Gomez R, Summers M, Summers A, Wolf A, Summers J. Depression anxiety stress scales-21: measurement and structural invariance across ratings of men and women. *Assessment.* 2014;21(4):418–426. doi:10.1177/1073191113514106
45. Bastien CH, Vallières A, Morin CM. Validation of the insomnia severity index as an outcome measure for insomnia research. *Sleep Med.* 2001;2(4):297–307. doi:10.1016/S1389-9457(00)00065-4
46. Manzar MD, Jahrami HA, Bahammam AS. Structural validity of the insomnia severity index: a systematic review and meta-analysis. *Sleep Med Rev.* 2021;60:101531. doi:10.1016/j.smrv.2021.101531
47. Hayes AF. *Introduction to Mediation, Moderation, and Conditional Process Analysis.* Guilford Press; 2018.
48. Preacher KJ, Hayes AF. SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behav Res Methods.* 2004;36(4):717–731. doi:10.3758/BF03206553
49. Franceschini C, Musetti A, Zenesini C, et al. Poor sleep quality and its consequences on mental health during the COVID-19 lockdown in Italy. *Front Psychol.* 2020;11:1–15. doi:10.3389/fpsyg.2020.574475
50. Giromini L, Ales F, de Campora G, Zennaro A, Pignolo C. Developing age and gender adjusted normative reference values for the Difficulties in Emotion Regulation Scale (DERS). *J Psychopathol Behav Assess.* 2017;39(4):705–714. doi:10.1007/s10862-017-9611-0
51. Gori A, Topino E. Across the covid-19 waves; assessing temporal fluctuations in perceived stress, post-traumatic symptoms, worry, anxiety and civic moral disengagement over one year of pandemic. *Int J Environ Res Public Health.* 2021;18(11):5651. doi:10.3390/ijerph18115651
52. Gori A, Topino E, Caretti V. The impact of COVID-19 lockdown on perceived stress: the role of defence mechanisms and coping strategies. *J Contingencies Crisis Manag.* 2021:1–12. doi:10.1111/1468-5973.12380
53. Cao W, Fang Z, Hou G, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res.* 2020;287:112934. doi:10.1016/j.psychres.2020.112934
54. Rossi R, Succi V, Talevi D, et al. COVID-19 pandemic and lockdown measures impact on mental health among the general population in Italy. *Front Psychiatry.* 2020;11:7–12. doi:10.3389/fpsyg.2020.00790
55. Gualano MR, Lo Moro G, Voglino G, Bert F, Siliquini R. Effects of COVID-19 lockdown on mental health and sleep disturbances in Italy. *Int J Environ Res Public Health.* 2020;17(13):1–13. doi:10.3390/ijerph17134779
56. Rodriguez-Besteiro S, Tornero-Aguilera JF, Fernández-Lucas J, Clemente-Suárez VJ. Gender differences in the covid-19 pandemic risk perception, psychology and behaviors of Spanish university students. *Int J Environ Res Public Health.* 2021;18(8):3908. doi:10.3390/ijerph18083908
57. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res.* 2020;288:112954. doi:10.1016/j.psychres.2020.112954
58. Sallie SN, Ritou VJE, Bowden-Jones H, Voon V. Assessing online gaming and pornography consumption patterns during COVID-19 isolation using an online survey: highlighting distinct avenues of problematic internet behavior. *Addict Behav.* 2021;123:107044. doi:10.1016/j.addbeh.2021.107044
59. Younes F, Halawi G, Jabbar H, et al. Internet addiction and relationships with insomnia, anxiety, depression, stress and self-esteem in university students: a cross-sectional designed study. *PLoS One.* 2016;11(9):e0161126. doi:10.1371/journal.pone.0161126
60. Vaillancourt-Morel MP, Blais-Lecours S, Labadie C, Bergeron S, Sabourin S, Godbout N. Profiles of cyberpornography use and sexual well-being in adults. *J Sex Med.* 2017;14(1):78–85. doi:10.1016/j.jsxm.2016.10.016
61. Spenhoff M, Kruger THC, Hartmann U, Kobs J. Hypersexual behavior in an online sample of males: associations with personal distress and functional impairment. *J Sex Med.* 2013;10(12):2996–3005. doi:10.1111/jsm.12160
62. Hermand M, Benyamina A, Donnadieu-Rigole H, et al. Addictive use of online sexual activities and its comorbidities: a systematic review. *Curr Addict Rep.* 2020;7(2):194–209. doi:10.1007/s40429-020-00301-3
63. Wéry A, Billieux J. Problematic cybersex: conceptualization, assessment, and treatment. *Addict Behav.* 2017;64:238–246. doi:10.1016/j.addbeh.2015.11.007
64. Caretti V, Gori A, Craparo G, Giannini M, Iraci-Sareri G, Schimmenti A. A new measure for assessing substance-related and addictive disorders: the addictive behavior questionnaire (ABQ). *J Clin Med.* 2018;7(8):194. doi:10.3390/jcm7080194
65. Zdankiewicz-ścigala E, Ścigala DK. Attachment style, early childhood trauma, alexithymia, and dissociation among persons addicted to alcohol: structural equation model of dependencies. *Front Psychol.* 2020;10:1–13. doi:10.3389/fpsyg.2019.02957
66. Topino E, Gori A, Cacioppo M. Alexithymia, dissociation, and family functioning in a sample of online gamblers: a moderated mediation study. *Int J Environ Res Public Health.* 2021;18(24):13291. doi:10.3390/ijerph182413291
67. Gori A, Topino E, Pucci C, Griffiths MD. The relationship between alexithymia, dysmorphic concern, and exercise addiction: the moderating effect of self-esteem. *J Pers Med.* 2021;11(11):1111. doi:10.3390/jpm11111111
68. Walton MT, Cantor JM, Lykins AD. An online assessment of personality, psychological, and sexuality trait variables associated with self-reported hypersexual behavior. *Arch Sex Behav.* 2017;46(3):721–733. doi:10.1007/s10508-015-0606-1

69. Adams KD, Robinson DW. Shame reduction, affect regulation, and sexual boundary development: essential building blocks of sexual addiction treatment. *Sex Addict Compulsivity*. 2001;8(1):23–44. doi:10.1080/10720160127559
70. Bos SC, Macedo AF. Literature review on Insomnia (2010–2016). *Biol Rhythm Res*. 2019;50(1):94–163. doi:10.1080/09291016.2017.1413766
71. Tsybes A, Aldao A, Mennin DS. Emotion dysregulation and sleep difficulties in generalized anxiety disorder. *J Anxiety Disord*. 2013;27(2):197–203. doi:10.1016/j.janxdis.2013.01.008
72. Hoag JR, Tennen H, Stevens RG, Coman E, Wu H. Affect, emotion dysregulation, and sleep quality among low-income women. *Sleep Health*. 2016;2(4):283–288. doi:10.1016/j.sleh.2016.08.006
73. Mauer-Vakil D, Bahji A. The addictive nature of compulsive sexual behaviours and problematic online pornography consumption: a review. *Can J Addict*. 2020;11(3):42–51. doi:10.1097/CXA.000000000000091
74. Teh HC, Archer JA, Chang W, Chen SHA. Mental well-being mediates the relationship between perceived stress and perceived health. *Stress Health*. 2015;31(1):71–77. doi:10.1002/smi.2510
75. Pellerone M, Cascio MI, Costanzo G, Gori A, Pace U, Craparo G. Alexithymia and psychological symptomatology: research conducted on a non-clinical group of Italian adolescents. *Int J Cult Ment Health*. 2017;10(3):300–309. doi:10.1080/17542863.2017.1307434
76. Biddle DJ, Hermens DF, Lallukka T, Aji M, Glozier N. Insomnia symptoms and short sleep duration predict trajectory of mental health symptoms. *Sleep Med*. 2019;54:53–61. doi:10.1016/j.sleep.2018.10.008
77. Di Giunta L, Lunetti C, Fiasconaro I, et al. COVID-19 impact on parental emotion socialization and youth socioemotional adjustment in Italy. *J Res Adolesc*. 2021;31(3):657–677. doi:10.1111/jora.12669
78. Wang C, Pan R, Wan X, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun*. 2020;87:40–48. doi:10.1016/j.bbi.2020.04.028
79. Morin CM, Vézina-Im L-A, Ivers H, et al. Prevalent, incident, and persistent insomnia in a population-based cohort tested before (2018) and during the first-wave of COVID-19 pandemic (2020). *Sleep*. 2021:1–6. doi:10.1093/sleep/zsab258.
80. Gerber M, Lang C, Lemola S, et al. Validation of the German version of the insomnia severity index in adolescents, young adults and adult workers: results from three cross-sectional studies. *BMC Psychiatry*. 2016;16(1):1–14. doi:10.1186/s12888-016-0876-8
81. Kinderman P, Schwannauer M, Pontin E, Tai S. Psychological processes mediate the impact of familial risk, social circumstances and life events on mental health. *PLoS One*. 2013;8(10):e76564. doi:10.1371/journal.pone.0076564

Nature and Science of Sleep

Dovepress

Publish your work in this journal

Nature and Science of Sleep is an international, peer-reviewed, open access journal covering all aspects of sleep science and sleep medicine, including the neurophysiology and functions of sleep, the genetics of sleep, sleep and society, biological rhythms, dreaming, sleep disorders and therapy, and strategies to optimize healthy sleep. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/nature-and-science-of-sleep-journal>