

Study of five penicillinase producing *Neisseria gonorrhoeae* isolated in Italy

Carla Sabia, Moreno Bondi, Patrizia Messi, Simona De Niederhäusern, Giuliano Manicardi

Department of Biomedical Sciences, University of Modena and Reggio Emilia, Modena, Italy

SUMMARY

Five penicillinase producing *Neisseria gonorrhoeae* (PPNG) were isolated from urethral specimens of men admitted to the "Santa Chiara" Hospital (Trento, Italy). All strains proved to be resistant to penicillin and ampicillin, and sensitive to cefuroxime, erythromycin, tetracycline, spectinomycin, nalidixic acid and ciprofloxacin. PPNG plasmid profiles showed that four of the isolates carried the 3.2 MDa "Africa" plasmid and one the 4.5 MDa "Asia" plasmid, the two well-known phenotypes reported in the USA and Europe as well as in Asian and African countries. Membrane matings were performed using *N. gonorrhoeae* carrying the 24.5 MDa conjugative plasmid as donors and *E. coli* K12 J 53 as recipient. The transfer of β -lactamic antibiotic resistance was supported by the presence of 4.5 or 3.2 MDa plasmid bands and by β -lactamase production in the transconjugants. Restriction analysis of Asian and African plasmids is reported.

KEY WORDS: PPNG, *Neisseria gonorrhoeae*, plasmids, conjugation

Received April 5, 2005

Accepted April 26, 2005

INTRODUCTION

Gonorrhoea is a common transmissible bacterial infection, and WHO (1999) estimates place the yearly number of new gonococcal infections worldwide at nearly 60 million. Gonorrhoea begins as an infection of the superficial mucous membranes (Brown, 1999), such as those of the endocervix, urethra, pharynx or rectum (LiMaye, 2003). Approximately 10-20% of infected women develop salpingitis, with resultant tubal scarring and either infertility or ectopic pregnancy (Westrom and Eschenbach, 1999; Zhu *et al.*, 2004). Many cases are asymptomatic and therefore remain undiagnosed. Untreated or inade-

quately treated gonococcal infections may have repercussions such as blood dissemination (LiMaye, 2003) and spread to other people.

Neisseria gonorrhoeae has developed chromosomal and plasmid resistance to several antibiotics, and the number of penicillinase-producing *N. gonorrhoeae* (PPNG) increased rapidly from 1987 to 1991 (Fox *et al.*, 1997), slightly falling off when penicillin therapy against gonorrhoea was discontinued (Lind, 1997). It has been suggested that the decrease of PPNGs is due to the therapeutic use of active drugs other than penicillins and to the use of fluoroquinolones, which can attack β -lactamase plasmids (Kam *et al.*, 1995).

Penicillinase-producing *N. gonorrhoeae* carrying the Asian plasmid (4.4-4.7 MDa) were first isolated in the USA in 1976 (Ashford *et al.*, 1976) from United States military servicemen who had relocated from Southeast Asia, and these agents are named according to their epidemiological origin in the Far East. The first strains carrying the African plasmid phenotype (3.2-3.4 MDa) were instead isolated in the UK in 1976 (Phillips, 1976).

Corresponding author

Dr. Moreno Bondi
Dipartimento di Scienze Biomediche
Università degli Studi di Modena
e Reggio Emilia
Via Campi, 287
41100 Modena, Italy
e-mail: bondi.moreno@unimore.it

Two other deletion derivatives were detected in 1984 by separate research groups and designated Toronto (3.05 MDa) (Yeung *et al.*, 1986) and Rio plasmids (2.9 MDa) (van Embden *et al.*, 1985). Other plasmids have been reported, including Nimes (4.0 MDa) (Gouby *et al.*, 1986), New Zealand (6.0 MDa) (Brett, 1989) and an unnamed variant (3.9 MDa) isolated in the Philippines (Knapp *et al.*, 1997), but global dissemination of these plasmids has not as yet followed their emergence (Pagotto *et al.*, 2000, Palmer *et al.*, 2000). Studies, based on restriction endonuclease, hybridisation and heteroduplex analysis by electron microscopy have shown that the African, Toronto, Nimes, Rio, and New Zealand-type plasmids are deletion or insertion derivatives of the prototype Asian plasmid (Yeung *et al.*, 1986; Dillon *et al.*, 1989). In particular, the African plasmid presents a 2.1 Kb deletion as compared to the Asian plasmid.

Penicillinase production in these plasmids is mediated by a TEM1-type β -lactamase encoded by the Tn2 transposon carried on several related plasmids (Palmer *et al.*, 2000) that is cut and includes 84% of *tnpR* noncoding sequences, the entire *bla* gene, and the right inverted repeat (IR-R) (Fayet *et al.*, 1982; Pagotto *et al.*, 2000).

In spite of a worldwide prevalence (Lee *et al.*, 2000), PPNG strains have rarely been reported in Italy, probably due to the limited use of penicillin in gonococcal urethritis therapy. The isolation of *N. gonorrhoeae* less sensitive to penicillin was first reported up to 1980 (Coppini *et al.*, 1980). Subsequently, we isolated two PPNG

strains carrying the Asian and African plasmids (Bondi *et al.*, 1984; Manicardi *et al.*, 1992). Considering the lack of new reports on PPNG in Italy, the present work aims to study the plasmid phenotypes of five penicillinase-producing *Neisseria gonorrhoeae* isolated at the "Santa Chiara" Hospital of Trento.

MATERIALS AND METHODS

Bacterial strains

The penicillinase producing *Neisseria gonorrhoeae* (PPNG) used in this study are listed in Table 1. The strains were isolated from urethral specimen of men admitted to the "Santa Chiara" Hospital (Trento, Italy) and identified by standard biochemical methods (API NH, bioMérieux, France). β -lactamase production was assayed using the chromogenic cephalosporin test (Nitrocefim, Oxoid, Milano, Italy) (O'Callaghan *et al.*, 1972). All *N. gonorrhoeae* were grown on GC agar base (Oxoid) supplemented with 1% IsoVitalax (Oxoid) and ampicillin (10 mg/ml, Sigma Chemical Co. St. Louis, MO, USA), and incubated for up to 24h at 37°C in a humid atmosphere supplemented with 5% CO₂.

Determination of minimum inhibitory concentration

The minimum inhibitory concentration (MIC) was determined in GC agar base supplemented with 1% IsoVitalax by breakpoint criteria, as defined by the National Committee for Clinical

TABLE 1 - MIC of the five PPNG strains

	MIC (μ g/ml)							
	P	Ap	Cf	E	Sp	T	Na	Cip
<i>N. gonorrhoeae</i> 57	>128	>128	0,004	0,008	2	1	1	0.25
<i>N. gonorrhoeae</i> 98	>128	>128	0,004	0,016	8	0.06	0.5	0.03
<i>N. gonorrhoeae</i> 157	>128	>128	0,008	0,016	4	1	0.06	0.015
<i>N. gonorrhoeae</i> 237	>128	>128	0,016	0,008	16	0.06	0.5	0.125
<i>N. gonorrhoeae</i> 243	>128	>128	0,008	0,008	1	0.06	0.25	0.03

P = penicillin; Ap = ampicillin; Cf = cefuroxime; E = erythromycin; T = tetracycline; Sp = spectinomycin; Na = nalidixic acid; Cip = ciprofloxacin.

Laboratory Standards (NCCLS, 2000). Twofold serial dilutions of the following antibiotic powders were tested: ampicillin (Ap), penicillin (P), cefuroxim (Cf), erythromycin (E), tetracycline (T), spectinomycin (St), nalidixic acid (Na), ciprofloxacin (Cip) (all from Sigma).

PPNG plasmid profiles

Plasmid DNA was isolated by the lyses method of Birnboim and Doly (1979). Electrophoresis was conducted on 0.7% agarose, horizontal slab gels, in TRIS acetate buffer at pH 8.0, using a steady voltage of 75 V for 120 min. Purified plasmids of *E. coli* V517 (Macrina *et al.*, 1978) were used as size reference plasmid for molecular weight determinations.

Conjugation experiments

Matings were performed by a modified Roberts and Falkow (1977) method, using the PPNGs carrying the 24.5 MDa conjugative plasmid as donors and *E. coli* K12 J 53 Na^R, Ap^S as recipient. For each conjugation, equal volumes of donor and recipient suspensions were inoculated in 10 ml of GC broth (Oxoid) with 2 mM CaCl₂ and 2 mM MgCl₂ added. After overnight incubation at 37°C, matings were performed on a membrane filter (0.45 µm-pore-size filter; Millipore Corp., Bedford, Mass.) then placed in a GC-agar base containing Kellogg's supplement (Difco, Detroit, MI). After 6 h of further incubation the cells were resuspended in 2 ml GC broth by vigorous shaking and spread on McConkey agar (Oxoid) supplemented with 10 µg/ml ampicillin and 50 µg/ml nalidixic acid. The developed colonies were tested for 4.5 and 3.2 MDa plasmid bands and for β-lactamase activity.

Asian and African plasmids restriction analysis

Asian and African plasmids were recovered from agarose gel by the Maniatis technique (Maniatis, 1982). Plasmids were digested with *Bam*HI, *Hind*III, *Pst*I according to the manufacturer's instructions (Promega, Madison, WI). In all analysis lambda phage DNA (Roche Applied Science, Indianapolis, IN) digested with *Bam*HI, *Hind*III and *Eco*RI (Boehringer, Mannheim, Germany) was used as molecular size marker. Digests were separated by electrophoresis on 1% (w/v) agarose gel in TRIS acetate buffer at pH 8 at 30V for 16h.

RESULTS

Minimum inhibitory concentration

MIC values of PPNGs are reported in Table 1. All strains were resistant to penicillin and ampicillin (MIC >128 mg/ml), whereas they were sensitive to other antibiotics tested at the following MIC ranges: cefuroxime 0.004-0.016 mg/ml, erythromycin 0.008-0.016 mg/ml, tetracycline 0.06-1 mg/ml, spectinomycin 1-16 mg/ml, nalidixic acid 0.06 -1 mg/ml, ciprofloxacin 0.0015-0.25 mg/ml.

PPNG and *E. coli* transconjugant plasmid profiles

N. gonorrhoeae 343, 237, 157 and 98 presented the 3.2 MDa African plasmid and *N. gonorrhoeae*




FIGURE 1 - PPNG strains and *E. coli* NGC Asian transconjugant plasmid profiles. Lane 1: molecular size markers prepared from *E. coli* V517 (35.8, 3.7, 3.4, 2.6, 2.0, 1.8 MDa); lane 2: *E. coli* NGC 57 Asian transconjugant; lane 3: *N. gonorrhoeae* 57 with Asian profile; lanes 4, 5, 6, 7: *N. gonorrhoeae* 237, *N. gonorrhoeae* 157, *N. gonorrhoeae* 343, *N. gonorrhoeae* 98, all with African profile.

57 showed the "Asia" configuration, characterized by the 4.5 MDa plasmid. *N. gonorrhoeae* 157, 98 and 57 harboured the 24.5 MDa conjugative plasmid. All strains had the 2.6 MDa cryptic plasmid.

African and Asian PPNG strains carrying the 24.5 MDa conjugative plasmid (*N. gonorrhoeae* 157,

98 and 57) transferred β -lactamic resistance to *E. coli* K12 J 53. This transfer was supported by 4.5 or 3.2 MDa plasmid bands observed in the transconjugants and confirmed by β -lactamase production. Plasmid profiles of PPNG strains and *E. coli* K12 NGC Asia transconjugant are shown in Figure 1.




FIGURE 2 - Restriction analysis of *N. gonorrhoeae* 57 Asian plasmid and *N. gonorrhoeae* 237 African plasmid. Lanes 1, 2, 3: Asian plasmid digest with *Bam*HI and *Pst*I (lane 1), with *Bam*HI and *Hind*III (lane 2), with *Bam*HI (lane 3); lanes 4, 5: African plasmid digested with *Bam*HI and *Pst*I (lane 4) and with *Bam*HI (lane 5).




FIGURE 3 (a, b) - Restriction endonuclease map of the *N. gonorrhoeae* 57 Asian plasmid (a) and *N. gonorrhoeae* 237 African plasmid (b). In both plasmids the *Pst*I site is located in the common region containing the portion of the *Tn2* transposon between the two *Bam*HI sites, and this explains why two bands with equal molecular weight are formed. The Asian plasmid, which shares a region containing a *Hind* III site was also cleaved with *Bam*HI and *Hind*III into three fragments.

Asian and African plasmids restriction analysis

Figure 2 illustrates the restriction analysis of 4.5 and 3.2 MDa plasmids. *Bam*HI cleaves the plasmids into two characteristic bands (3.16, 1.34 MDa for Asian plasmid and 1.86, 1.34 MDa for African plasmid), as reported by McNicol *et al.*, (1983). Using both *Bam*HI and *Pst*I enzymes, each plasmid is cleaved into three fragments (1.86, 0.86, 0.47 MDa for African plasmid and 3.16, 0.86, 0.47 for Asian plasmid). In both plasmids, the *Pst*I site is located in the common region containing the portion of the Tn2 transposon between the two *Bam*HI sites, and this explains why two bands with equal molecular weight are formed. The Asian plasmid, which shares a region containing a *Hind*III site with another R factor carried by *Haemophilus ducrey* (Brunton *et al.*, 1981), was also cleaved with *Bam*HI and *Hind*III into three fragments (2.16, 1.34 and 1.00 MDa) (McNicol *et al.*, 1983). The plasmidial maps are shown in Figure 3 (a, b).

DISCUSSION

Gonorrhoea remains a major cause of bacterial sexually transmitted disease (STD) worldwide. In Africa, for example, *N. gonorrhoeae* is the most important cause of male urethritis, and the prevalence of gonococcal infection among women is high, in particular among prostitutes and workers in the sex industry (Germain *et al.*, 1997). The high prevalence of *N. gonorrhoeae* in developing countries is largely related to poor accessibility to diagnostic tests and appropriate treatments (Alary *et al.*, 1996). The control of gonococcal infection is important considering the high incidence of complications. Moreover, gonorrhoea may play a role in facilitating HIV acquisition and transmission, due to mucosal inflammation and greater release of the virus in semen (Wasserheit, 1992; Bhuiyan *et al.*, 1999).

Antimicrobial resistance among *N. gonorrhoeae* isolates in many developing countries has increased remarkably (Nissinen *et al.*, 1997). Surveys by the WHO Western Pacific Region in 1997 (WHO, 1998) showed that PPNG was widely distributed throughout the region. PPNG is particularly prevalent in the Philippines (81.8%), Korea (79.3%), Singapore (61%), Vietnam (64%)

and Malaysia (41%). Interestingly, rates are very low in other countries, such as Japan (2.3%), Australia (6.4%) and New Zealand (7.4%). The literature reports high levels of penicillin resistance that may be a consequence of therapy for unrelated illnesses or self-medication with penicillin or penicillin-related drugs. For this reason the Centres for Disease Control have recommended a first-line therapeutic regimen based on fluoroquinolones and cephalosporin (CDC, 1993). Although in some parts of Europe penicillins are still used effectively to treat gonorrhoeae, resistance is frequent in imported cases (Tapsall, 2001). Even in our work the gonococci studied had high MIC values for penicillin and ampicillin, and considering the lack of PPNG reports in Italy, these strains were probably imported. Resistance was not detected against cefuroxime, erythromycin, tetracycline or nalidixic acid. Our isolates were also susceptible to spectinomycin and ciprofloxacin, which remain valuable alternatives for treating gonorrhoea.

In our isolates, plasmid analysis showed that four strains carried the 3.2 MDa "Africa" and only one the 4.5 MDa "Asia" plasmids, which are responsible for the two well-known phenotypes reported in the USA and Europe as well as in Asian and African countries. The presence of the 24.5 MDa conjugative plasmid in both phenotypes suggests the possibility of a horizontal spread of plasmid-mediated penicillin resistance. For example, the host range of an Asia-type plasmid was shown to include gonococci, members of the family *Enterobacteriaceae* and *Haemophilus influenzae* (Pagotto and Dillon, 2001). This occurrence is supported by the transferability of β -lactamic resistance from PPNG to bacteria belonging to different genera, as observed for *E. coli* in our study.

REFERENCES

- ALARY, M., LAGA, M., VUYLSTEKE, B., NZILA, N., AND PIOT, P. (1996). Signs and symptoms of prevalent and incident cases of gonorrhoea and genital chlamydial infection among female prostitutes in Kinshasa, Zaire. *Clin Infect Dis*, **22**, 477-484.
- ASHFORD, W.A., GOLASH, R.G., AND HEMMING, V.G. (1976). Penicillase-producing *Neisseria Gonorrhoeae*. *Lancet*, **2**, 657-658.
- BIRNBOIM, H.C., AND DOLY, J. (1979). A rapid alkaline

- extraction procedure for screening recombinant plasmid DNA. *Nucleic Res*, **7**, 1513-1523.
- BHUIYAN, B.U., RAHMAN, M., MIAH, M.R., NAHAR, S., ISLAM, N., AHMED, M., RAHMAN, K.M., AND ALBERT, M.J. (1999). Antimicrobial susceptibilities and plasmid contents of *Neisseria gonorrhoeae* isolates from commercial sex workers in Dhaka, Bangladesh: emergence of high-level resistance to ciprofloxacin. *J Clin Microbiol*, **37**, 1130-1136.
- BONDI, M., MANICARDI, G., PICCININI, L., CASOLARI, C., AND FABIO, U. (1984). Emergence in Italy of a PPNG strain carrying "Asia" plasmid. *Microbiologica*, **7**, 257-261.
- BRETT, M. (1989). A novel gonococcal β -lactamase plasmid. *J Antimicrob Chemother*, **23**, 653-654.
- BROWN, T.J., YEN-MOORE, A., AND TYRING, S.K. (1999). Disseminated gonococcal infection. *J Acad Dermatol*, **41**, 517-518.
- BRUNTON, J., BENNET, P., AND GRINSTED, J. (1981). Molecular nature of a plasmid specifying beta-lactamase production in *Haemophilus ducreyi*. *J Bacteriol*, **148**, 788-795.
- CENTERS FOR DISEASE CONTROL. (1993). 1993 sexually transmitted diseases treatment guidelines. *Morbidity Mortal Weekly Rep*, **42** (RR-14), 4-5.
- COPPINI, M., BONDI, M., CASOLARI, C., PICCININI, L., AND MANICARDI, G. (1980). Considerazioni microbiologiche ed epidemiologiche su ceppi di *Neisseria gonorrhoeae* isolati nel modenese. *Giornale Italiano di Dermatologia e Venereologia*, **115**, 393-395.
- DILLON, J.A., AND YENG, K.H. (1989). Beta-lactamase plasmids and chromosomally mediated antibiotic resistance in pathogenic *Neisseria* species. *Clin Microbiol Rev*, **2**, Suppl, S125-133.
- FAYET, O., FROMET, Y., AND PIFFARETTI, J.C. (1982). β -Lactamase specifying plasmids isolated from *Neisseria gonorrhoeae* have retained an intact right part of Tn3-like transposon. *J Bacteriol*, **149**, 136-144.
- FOX, K.K., KNAPP, J.S., HOLMES, K.K., HOOK, E.W. 3RD, JUDSON, F.N., THOMPSON, S.E., WASHINGTON, J.A., AND WHITTINGTON, W.L. (1997). Antimicrobial resistance in *Neisseria gonorrhoeae* in the United States, 1998-1994: the emergence of decreased susceptibility to the fluoroquinolones. *J Infect Dis*, **175**, 1396-1403.
- GERMAIN, M., ALARY, M., GUÈDÈMÈ, A., PADONOU, F., DAVO, N., ADJOVI, C., VAN DYCK, E., JOLY, J.R., AND MAHONY, J.B. (1997). Evaluation of a screening algorithm for the diagnosis of genital infections with *Neisseria gonorrhoeae* and *Chlamydia trachomatis* among female sex workers in Bénin. *Sex Transm Dis*, **24**, 109-115.
- GOUBY, A., BOURG, G., AND RAMUZ, M. (1986). Previously undescribed 6.6-kilobase R plasmid in penicillinase-producing *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother*, **29**, 1095-1097.
- KAM, K.M., LO, K.K., HO, N.K., AND CHEUNG, M.M. (1995). Rapid decline in penicillase-producing *Neisseria gonorrhoeae* in Hong Kong associated with emerging 4-fluoroquinolone resistance. *Genitourin Med*, **71**, 141-144.
- KANPP, J.S., MESOLA, V.P., NEAL, S.W., WI, T.E., TUAZOM, C., AND MANALASTAS, R. (1997). Molecular epidemiology, in 1994, of *Neisseria gonorrhoeae* in Manila and Cebu City, Republic of the Philippines. *Sex Transm Dis*, **24**, 2-7.
- YEUNG, K.H., DILLON, J.R., PAUZE, M., AND WALLACE, E. (1986). A novel 4.9-kilobase plasmid associated with an outbreak of penicillinase-producing *Neisseria gonorrhoeae*. *J Infect Dis*, **153**, 1162-1165.
- LEE, K., SHIN, J.W., LIM, J.B., KIM, Y.A., YONG, D., OH, H.B., AND CHONG, Y. (2000). Emerging antimicrobial resistance, plasmid profile and pulsed-field gel electrophoresis pattern of the endonuclease-digested genomic DNA of *Neisseria gonorrhoeae*. *Yonsei Med J*, **41**, 381-386.
- LIMAYE, A.R. (2003). Disseminated gonococcal infection in women. *Primary Care Update for OB/GYNs*, **10**, 186-190.
- LIND, I. (1997). Antimicrobial resistance in *Neisseria gonorrhoeae*. *Clin Infect Dis*, **24** Suppl, 93-97.
- MACRINA, F.L., KOPECKO, D.J., JONES, K.R., AYERS, D.J., AND M.C., COWEN, S.M. (1978). A multiple plasmid-containing *Escherichia coli* strain: convenient source of size reference plasmid molecules. *Plasmid*, **1**, 417-420.
- MANIATIS, T., FRITSCH, E.F., AND SAMBROOK, J. (1982). Molecular Cloning. A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- MANICARDI, G., ALOISI, P., NEGLIA, R., AND BONDI, M. (1992). Plasmid analysis in PPNG strains isolated in Italy. *Microbiologica*, **15**, 1-6.
- MCNICOL, P.J., ALBRITTON, W.L., AND RONALD, A.R. (1983). Characterization of ampicillin resistance plasmid of *Haemophilus ducreyi* and *Neisseria gonorrhoeae* with regard to location of origin of transfer and mobilization by a conjugative plasmid of *Haemophilus ducreyi*. *J Bacteriol*, **156**, 437-440.
- NATIONAL COMMITTEE FOR CLINICAL LABORATORY STANDARDS. (2000). Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically. Approved Standard M7-A5. *National Committee for Clinical Laboratory Standards: Wayne, PA*.
- NISSINEN, A., JÄRVINEN, H., LIIMATAINEN, O., JÄHKOLA, M., HUOVINEN, P. (1997). Antimicrobial resistance in *Neisseria gonorrhoeae* in Finland, 1976-1995. The Finnish Study Group for Antimicrobial Resistance. *Sex Transm Dis*, **24**, 576-581.
- O'CALLAGHAN, C.H., MORRIS, A., KIRBY, S.M., AND SHINGLER, A.H. (1972). Novel method for detection of beta-lactamase by using a chromogenic cephalosporin substrate. *Antimicrob Agent Chemother*, **1**, 283-288.
- PAGOTTO, F., AMAN, A.T., LAI-KING, N.G., YEUNG, K.H.,

- BRETT, M., AND DILLON, J.A. (2000). Sequence Analysis of the Family of Penicillinase-Producing Plasmids of *Neisseria gonorrhoeae*. *Plasmid*, **43**, 24-34.
- PAGOTTO, F., AND DILLON, J.A. (2001). Multiple origins and replication proteins influence biological properties of beta-lactamase-producing plasmids from *Neisseria gonorrhoeae*. *J Bacteriol*, **183**, 5472-5481.
- PALMER, H.M., LEEMING, J.P., AND TURNER, A. (2000). A multiplex polymerase chain reaction to differentiate beta-lactamase plasmids of *Neisseria gonorrhoeae*. *J Antimicrob Chemother*, **45**, 777-782.
- PHILLIPS, I. (1976). Beta-lactamase-producing, penicillin-resistant gonococcus. *Lancet*, **2**, 656-657.
- ROBERTS, M., AND FALKOW, S. (1977). Conjugal transfer of R-plasmids in *Neisseria gonorrhoeae*. *Nature*, **226**, 630-631.
- TAPSALL, J. (2001). Antimicrobial resistance in *Neisseria gonorrhoeae*. Geneva, World Health Organisation, WHO/CDS/CSR/DRS/2001, **3**, 1-65.
- VAN EMBDEN, J. D., DESSES-KROON, M., AND VAN KLINGEREN, B. (1985). New β -lactamase plasmid in *Neisseria gonorrhoeae*. *J Antimicrob Chemother*, **15**, 247-50.
- WASSERHEIT, J.N. (1992). Epidemiological synergy: interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Transm Dis*, **19**, 61-77.
- WESTROM, L., AND ESCHENBACH, D. (1999). Pelvic inflammatory disease. In: Holmes KK, Mardh PA, Sparling PF, et al., editors, *Sex Transm Dis*, New York: McGraw-Hill, 783-809.
- World Health Organisation, WHO Western Pacific REGION GONOCOCCAL ANTIMICROBIAL SURVEILLANCE PROGRAMME. (1998). Resistance in gonococci isolated in the WHO Western Pacific Region to various antimicrobials used in the treatment of gonorrhoea, 1997. *Commun Dis. Intell*, **13**, 288-291.
- WHO. Gonorrhoea estimates, 1999. (2001). Global prevalence and incidence of selected curable sexually transmitted infections. Geneva: World Health Organization. Available via <http://www.emro.who.int/ASD/backgrounddocuments/uae03/surv/stdoverview.pdf>
- ZHU, W., THOMAS, C.E., AND SPARLING, P.F. (2004). DNA Immunization of mice with a plasmid encoding *Neisseria gonorrhoeae* PorB protein by intramuscular injection and epidermal particle bombardment. *Vaccine*, **22**, 661-670.

