

This is a pre print version of the following article:

Karyotype variations in Italian populations of the peach-potato aphid *Myzus persicae* (Hemiptera: Aphididae) / Rivi, Marco; Monti, Valentina; E., Mazzoni; Cassanelli, Stefano; M., Panini; D., Bizzaro; Mandrioli, Mauro; Manicardi, Gian Carlo. - In: BULLETIN OF ENTOMOLOGICAL RESEARCH. - ISSN 0007-4853. - STAMPA. - 102:6(2012), pp. 663-671. [10.1017/S0007485312000247]

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

13/01/2026 02:01

## Proof Delivery Form

## Bulletin of Entomological Research

Date of delivery:

Journal and vol/article ref:

ber

0\_0/1200024

Number of pages (not including this page): 9

This proof is sent to you on behalf of Cambridge University Press. Please print out the file and check the proofs carefully. Make any corrections necessary on a hardcopy and answer queries on each page of the proofs.

Please return the **marked proof** within

2

days of receipt to:

Laura Bates, at the following email address: batesla10@yahoo.co.uk

OR post to her at:

Laura Bates, Westcroft, West End, Crail, Fife KY10 3RH, Scotland.

**Authors are strongly advised to read these proofs thoroughly because any errors missed may appear in the final published paper. This will be your ONLY chance to correct your proof. Once published, either online or in print, no further changes can be made.**

To avoid delay from overseas, please send the proof by airmail or courier.

If you have **no corrections** to make, please email **batesla10@yahoo.co.uk** to save having to return your paper proof. If corrections are light, you can also send them by email, quoting both page and line number.

- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal. Only **one** set of corrections are permitted.
- Corrections which do NOT follow journal style will not be accepted.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required.

• If you have problems with the file please contact

**nmarshall@cambridge.org**

Please note that this pdf is for proof checking purposes only. It should not be distributed to third parties and may not represent the final published version.

**Important:** you must return any forms included with your proof. We cannot publish your article if you have not returned your signed copyright form

NOTE - for further information about **Journals Production** please consult our **FAQs** at [http://journals.cambridge.org/production\\_faqs](http://journals.cambridge.org/production_faqs)

---

**Author queries:**

- Q1** Please check that all names have been spelled correctly and appear in the correct order. Please also check that all initials are present. Please check that the author surnames (family name) have been correctly identified by a pink background. If this is incorrect, please identify the full surname of the relevant authors. Occasionally, the distinction between surnames and forenames can be ambiguous, and this is to ensure that the authors' full surnames and forenames are tagged correctly, for accurate indexing online. Please also check all author affiliations..

# Offprint order form



CAMBRIDGE  
UNIVERSITY PRESS

PLEASE COMPLETE AND RETURN THIS FORM. WE WILL BE UNABLE TO SEND OFFPRINTS UNLESS A RETURN ADDRESS AND ARTICLE DETAILS ARE PROVIDED.

VAT REG NO. GB 823 8476 09

## Bulletin of Entomological Research (BER)

Volume:

no:

### Offprints

To order offprints, please complete this form and send it to **the publisher (address below)**. Please give the address to which your offprints should be sent. They will be despatched by surface mail within one month of publication. For an article by **more than one author this form is sent to you as the first named author**.

Number of offprints required: \_\_\_\_\_

Email: \_\_\_\_\_

Offprints to be sent to (print in BLOCK CAPITALS): \_\_\_\_\_

Post/Zip Code: \_\_\_\_\_

Telephone: \_\_\_\_\_ Date (dd/mm/yy): \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_

Author(s): \_\_\_\_\_

Article Title: \_\_\_\_\_

All enquiries about offprints should be addressed to **the Publisher: Journals Production Department, Cambridge University Press, University Printing House, Shaftesbury Road, Cambridge CB2 8BS, UK.**

**Charges for offprints (excluding VAT)** Please circle the appropriate charge:

Number of copies	25	50	100	150	200	per 50 extra
1-4 pages	£68	£109	£174	£239	£309	£68
5-8 pages	£109	£163	£239	£321	£399	£109
9-16 pages	£120	£181	£285	£381	£494	£120
17-24 pages	£131	£201	£331	£451	£599	£131
Each Additional 1-8 pages	£20	£31	£50	£70	£104	£20

### Methods of payment

If you live in Belgium, France, Germany, Ireland, Italy, Portugal, Spain or Sweden and are not registered for VAT we are required to charge VAT at the rate applicable in your country of residence. If you live in any other country in the EU and are not registered for VAT you will be charged VAT at the UK rate.

If registered, please quote your VAT number, or the VAT number of any agency paying on your behalf if it is registered.

VAT Number: \_\_\_\_\_

Payment **must** be included with your order, please tick which method you are using:

- ☐ Cheques should be made out to Cambridge University Press.
- ☐ Payment by someone else. Please enclose the official order when returning this form and ensure that when the order is sent it mentions the name of the journal and the article title.
- ☐ Payment may be made by any credit card bearing the Interbank Symbol.

Card Number:

Expiry Date (mm/yy): \_\_\_\_\_ / \_\_\_\_\_ Card Verification Number:

The card verification number is a 3 digit number printed on the **back** of your **Visa** or **Master card**, it appears after and to the right of your card number. For **American Express** the verification number is 4 digits, and printed on the **front** of your card, after and to the right of your card number.

Signature of card holder: \_\_\_\_\_ Amount (Including VAT if appropriate): £ \_\_\_\_\_

Please advise if address registered with card company is different from above

# Karyotype variations in Italian populations of the peach-potato aphid *Myzus persicae* (Hemiptera: Aphididae)

M. Rivi<sup>1</sup>, V. Monti<sup>1,2</sup>, E. Mazzoni<sup>3</sup>, S. Cassanelli<sup>1</sup>,  
M. Panini<sup>3</sup>, D. Bizzaro<sup>4</sup>, M. Mandrioli<sup>2</sup>  
and G.C. Manicardi<sup>1\*</sup>

<sup>1</sup>Dipartimento di Scienze Agrarie e degli Alimenti, Università di Modena e Reggio Emilia, Reggio Emilia, Italy; <sup>2</sup>Dipartimento di Biologia, Università di Modena e Reggio Emilia, Modena, Italy; <sup>3</sup>Istituto di Entomologia e Patologia vegetale, Università Cattolica del Sacro Cuore, Piacenza, Italy; <sup>4</sup>Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, Ancona, Italy

## Abstract

In this study, we present cytogenetic data regarding 66 *Myzus persicae* strains collected in different regions of Italy. Together with the most common  $2n=12$  karyotype, the results showed different chromosomal rearrangements:  $2n=12$  with A1–3 reciprocal translocation,  $2n=13$  with A1–3 reciprocal translocation and A3 fission,  $2n=13$  with A3 fission,  $2n=13$  with A4 fission,  $2n=14$  with X and A3 fissions. A  $2n=12-13$  chromosomal mosaicism has also been observed. Chromosomal aberrations (and in particular all strains showing A1–3 reciprocal translocation) are especially frequent in strains collected on tobacco plants, and we suggest that a clastogenic effect of nicotine. Further benefited by the holocentric nature of aphid chromosomes, could be at the basis of the observed phenomenon.

**Keywords:** karyotype variations, chromosomal rearrangements, holocentric chromosomes, nicotine, clastogenic effect, *Myzus persicae*, Aphididae

(Accepted 27 March 2012)

## Introduction

Classical and molecular cytogenetics provide an integrated approach for structural, functional and evolutionary analyses of chromosomes. This ranges from karyotype analyses to molecular mapping of chromosomes.

To date, studies concerning chromatin structure and organization have been mainly focused on eukaryotes having monocentric chromosomes, whereas species possessing holocentric/holokinetic chromosomes have been rather

neglected. Chromosomes with diffused centromeric activity have been found in Protista, as well as in plant and animal species (Wrench *et al.*, 1994). The chromosomes of aphids, like those of other hemipteran insects, have diffuse centromeres so that kinetic activity is dispersed along the entire length of each chromatid at least in mitotic divisions, thus influencing chromosome behaviour (White, 1973). In organisms possessing this kind of chromatin organization, chromosome fusions and fissions can occur without any duplication or loss of centromeres. This has consequences for the survival of the *de novo* chromosomal changes through mitosis and meiosis, and hence for karyotype evolution. Autosomal fusions and fissions, particularly the latter, seemed to play a pivotal role in aphid karyotype evolution (Blackman, 1980), although this view is at present somewhat speculative due

\*Author for correspondence

Fax: +54 11 4576-3354

E-mail: giancarlo.manicardi@unimore.it

Table 1. List of the Italian populations of *M. persicae* analyzed.

Northern Italy				Central-southern Italy			
Population	Chromosome number	Color	Host plant	Population	Chromosome number	Color	Host plant
Torino 1	12	G	Peach	Pisa 1	12	G	Peach
Torino 2	12	G	Peach	Pisa 2	12	G	Peach
Torino 3	12	G	Peach	Pisa 3	12	G	Peach
Torino 4	12	G	Peach	Ascoli 1	12	G	Peach
Cuneo 1	12	G	Peach	Pescara 1	12	G	Tobacco
Como 1	12	G	Peach	Pescara 2	12t + frm3	R	Tobacco
Lodi 1	12	G	Peach	Chieti 1	12t	R	Tobacco
Padova 1	12	G	Peach	Chieti 2	12t	R	Tobacco
Padova 2	12	G	Peach	Chieti 3	12t	R	Tobacco
Piacenza 1	12	G	Tomato	Chieti 4	12t	R	Tobacco
Piacenza 2	12	G	Tomato	Salerno 1	12t	R	Tobacco
Piacenza 3	12	G	Tomato	Salerno 2	12/13 frm3	R	Tobacco
Piacenza 4	12	G	Tomato	Salerno 3	13 frm3	R	Tobacco
Piacenza 5	12	G	Tomato	Benevento 1	12t	R	Tobacco
Piacenza 6	12	G	Peach	Cosenza 1	13 frm 4	G	Peach
Piacenza 7	12	G	Peach	Cosenza 2	14 frm X+3	G	Peach
Piacenza 8	12	G	Peach	Catanzaro 1	12	G	Potato
Piacenza 9	12	G	Aubergine	Catanzaro 2	12	G	Peach
Piacenza 10	13 frm 4	G	Aubergine	Cagliari	12	G	Peach
Piacenza 11	12	G	Peach				
Bologna 1	12	G	Peach				
Bologna 2	12	G	Peach				
Bologna 3	12	G	Peach				
Bologna 4	12	G	Peach				
Bologna 5	12	G	Peach				
Bologna 6	12	G	Peach				
Ferrara 1	12	G	Peach				
Ferrara 2	12	G	Peach				
Ferrara 3	12	G	Peach				
Ferrara 4	12	G	Peach				
Ferrara 5	12	G	Peach				
Ferrara 6	12	R	Peach				
Ferrara 7	12	G	Peach				
Ravenna 1	12	G	Peach				
Ravenna 2	12	G	Peach				
Ravenna 3	12	G	Peach				
Ravenna 4	12	G	Peach				
Ravenna 5	12	G	Peach				
Ravenna 6	13 frm 4	G	Peach				
Ravenna 7	12	G	Peach				
Ravenna 8	12	G	Peach				
Ravenna 9	12	G	Peach				
Ravenna 10	12	R	Peach				
Ravenna 11	12	G	Peach				
Ravenna 12	12	G	Peach				
Forlì 1	13 frm 3	G	Peach				
Forlì 2	12	G	Peach				

52 to a lack of knowledge concerning the mechanisms involved in  
53 rearrangements of the holocentric chromosomes (Spence &  
54 Blackman, 2000).  
55 A recurrent chromosomal rearrangement found in the  
56 peach-potato aphid *Myzus persicae* (Sulzer) (Hemiptera:  
57 Aphididae) populations collected worldwide involves a  
58 A1–3 reciprocal translocation associated with increased levels  
59 of resistance to organophosphate and carbamate insecticides  
60 (Blackman *et al.*, 1978; Spence & Blackman, 1998).  
61 The standard female karyotype of this species is  $2n = 12$ , but  
62 specimens with a chromosome complement of either  $2n = 13$  or  
63 14 have also been reported (Blackman, 1980; Lauritzen, 1982).  
64 On the basis of relative chromosome lengths, Blackman (1971)  
65 concluded that the  $2n = 13$  karyotype raised from a break in

one autosome of the pair A3, whereas a break in one  
66 chromosome of either the A2 and A3 pairs led to a  $2n = 14$   
67 karyotype. Rare cases of strain possessing  $2n = 11$  and  $3n = 18$   
68 have also been reported (Blackman, 1980; Yang & Zhang,  
69 2000). Very recently, the analysis of mitotic metaphase  
70 chromosomes of a *M. persicae* laboratory strain revealed  
71 different chromosome numbers, ranging from 12 to 17, within  
72 each embryo (intraclonal genetic variation *sensu* Loxdale &  
73 Lushai (2003)). Chromosome length measurements revealed  
74 that the observed chromosomal mosaicism is due to recurrent  
75 fragmentations of chromosomes X, 1 and 3 (Monti *et al.*, 2012).  
76 The present study shows cytogenetic data regarding 66  
77 *M. persicae* strains collected in different Italian regions  
78 showing several chromosomal rearrangements, the most  
79



Fig. 1. Geographic distribution of the sampling sites.

common being the A1–3 reciprocal translocation, which we here reported for the first time in Italy. We have also looked for the presence of a relationship between karyotype variations and the host plants.

#### Material and methods

*Myzus persicae* populations were collected mainly from peach (*Prunus persicae* L.) orchards (48), but also from

herbaceous hosts like tobacco (10), tomato (5), potato (1) and aubergine (2) at various locations in different areas of Italy (see table 1, fig. 1) and maintained as parthenogenetic female colonies on pea-seedlings (*Pisum sativum* cv 'Meraviglia d'Italia') under constant environmental conditions: 21°C, 16 h light:8 h dark photoperiod.

For chromosome spreads, adult females were dissected in Ringer saline solution and embryos were kept in a 1% hypotonic solution of sodium citrate for 30 min. The embryos





Fig. 2. Metaphase plate of the *M. persicae* strain Ferrara 03 stained with (a) Giemsa and (b) relative karyotype. Arrows indicate X chromosomes. Bar corresponds to 10  $\mu$ m.

were then transferred to minitubes and centrifuged at 350 g for 3 min. Methanol-acetic acid 3:1 was added to the pellet, which was made to flow up and down for 1 min through a needle of a 1 ml hypodermic syringe to obtain disaggregation of the material followed by a further centrifugation at 1000  $\times$  g for 3 min. This step was repeated with fresh fixative. Finally, the pellet was resuspended in new fixative, and 20  $\mu$ l of cellular suspension was dropped onto clean slides and stained with 5% Giemsa solution in Soerensen buffer, pH 6.8 for 10 min. Silver staining of nucleolar organizing regions (NORs) was achieved following Howell & Black (1980). Slides were examined using a Nikon Eclipse 80i fluorescence microscope with UV filters, and photographs were taken using Nikon digital sight DS-U1. Morphometric analyses of mitotic plates were carried out on 30 metaphases using the software MicroMeasure, freely available at the Biology Department at Colorado State University website (<http://rydberg.biology.colostate.edu/MicroMeasure>). Male induction for Salerno 03, Pescara 02, Cosenza 02 and Pisa 01 strains was evaluated by exposing parthenogenetic female aphids to short photoperiods (8 h light:16 h dark) according to Crema (1979).

## Results

The analysis of mitotic cells of embryos, obtained from parthenogenetic females, confirmed that  $2n=12$  is the standard chromosome number in *M. persicae* (fig. 2), but 14 out of 66 strains analysed showed intraspecific karyotype variants due to both structural and numerical variations in chromosome complements (table 1, figs 3–6).

The most frequent chromosomal rearrangement found in Italian populations is related to the A1–3 reciprocal translocation, which was found either alone (fig. 3) or together with an A3 fission (in one strain; fig. 6a, b). Other chromosome fissions involved A3 (found in two cases; fig. 4) and A4 (found in three cases; fig. 5), whereas a strain possessing 14 chromosomes as a consequence of both X and A3 fissions was also found (fig. 6c, d). Lastly, we identified a strain showing an intra-individual chromosome mosaicism due to the presence of mitotic plates with 12 (24% of the observed plates) and 13 (76%) chromosomes as a consequence of an A3 fission (fig. 4b).

NOR staining (figs 3a, c, g, h and 6c) revealed the presence of heteromorphism in the size of rDNA genes in strains Salerno 3 (fig. 4c) and Cosenza 2 (fig. 6c) and evidenced that the fission of the X chromosomes observed in Cosenza 2 always occurred in the X chromosome bearing the smallest

NOR-positive telomere and involved the X telomere opposite to the rDNA-bearing one (fig. 6c).

Considering the geographical distribution, it is evident that almost all karyotype variations (11 out of 14) were present in central and southern Italian regions, whereas only three were found in northern locations. Furthermore, all but one of the strains collected on tobacco showed chromosomal rearrangements; and, in particular, all the strains possessing the A1–3 reciprocal translocation were found on this plant and were red in colour.

Male induction revealed that the *M. persicae* strains Salerno 03, Pescara 02 and Cosenza 02, all possessing different kinds of karyotype variations, are anholocyclic since it was not possible to induce the sexual generation differently from that obtained under the same experimental conditions with the *M. persicae* strain Pisa 1, which showed a normal karyotype.

## Discussion

The typical aphid karyotype consists of pairs of rod-like chromosomes, whose number is typically stable within a genus, as shown in the large genus *Aphis*, where the typical chromosome number is eight with the exception of *A. farinosa* with  $2n=6$  (Blackman, 1980; Hales *et al.*, 1997). Nevertheless, exceptions have been published as revealed in the genus *Amphorophora*, where the chromosome number varies from  $2n=4$  to  $2n=72$  (Blackman, 1980).

Rearrangements most commonly involved autosomes, as shown in *M. persicae*, where, despite a standard chromosome number of  $2n=12$ , several strains possessing karyotypes consisting of 11–14 chromosomes have previously been reported (Blackman, 1980). On the contrary, Hales (1989) and Monti *et al.* (2012) demonstrated a complex pattern of associations and fissions occurring on both autosomes and X chromosomes in *Schoutedenia lutea* (van der Goot) (Hemiptera: Aphididae) and *M. persicae*, respectively, suggesting different scenarios for understanding aphid karyotype evolution.

The most common chromosomal variant described in *M. persicae* complement is a reciprocal translocation between the first and the third autosome pairs, leading to females with  $2n=12$  karyotype showing a marked structural heterozygosity (Blackman, 1980).

The empirical data, as presented in this paper, reveal for the first time that this chromosomal aberration also occurs in Italy since seven out of the 14 strains showed karyotype variations due to the A1–3 reciprocal translocation. In view of the absence of any primary constriction, which is typical of the holocentric chromosomes, together with the lack of specific banding patterns after conventional banding procedures, we combined procedures of standard chromosome staining (such as Giemsa and silver staining) with chromosome length evaluation. In particular, we used silver staining to confirm the exclusive localization of NORs regions on X chromosome telomeres in *M. persicae* and analyzed the involvement of sex chromosomes in the translocation event (Manicardi *et al.*, 2002). Afterwards, in the absence of any other cytogenetic markers, the morphometric analysis was employed to identify autosomes A1 and A3 as the chromosomes engaged in the rearrangement.

According to the literature, a link exists between the A1–3 chromosomal reciprocal translocation and resistance to organophosphate and carbamate insecticides due to E4 gene amplification (Blackman *et al.*, 1995), perhaps involving the removal of a repressor gene away from the structural genes in






Fig. 3. *M. persicae* chromosome complements showing A1-3 reciprocal translocation. (a) Benevento 01 is silver stained, (b) Salerno 01, (c) Chieti 02 and (e) Chieti 03 are stained with Giemsa, whereas (d, g) Chieti 1 and (f, h) Chieti 4 are both Giemsa and silver stained. The (i) karyotype is derived from (c) Chieti 02. Arrows indicate X chromosomes. Asterisks indicate A1-3 translocated chromosomes. Bar corresponds to 10 μm.

controls (Blackman *et al.*, 1978). Preliminary data involving PCR and southern blot analysis revealed that, in one of the Italian populations with this chromosomal aberration (Chieti 1), the FE4 gene (electrophoretically fast variant (allele) of the normal expressed carboxylesterase 4 (E4) enzyme) only was present (Rivi *et al.*, 2009). This strain showed a moderate increase in esterase activity and was considered an S/R1 (susceptible/first resistance level) strain *sensu* Devonshire *et al.* (1992). The aforementioned data allows us to suggest that this is the first *M. persicae* strain possessing the A1-3 chromosomal reciprocal translocation linked to an FE4 and not directly related to a high level of esterase-based insecticide resistance. Experiments currently in progress are aimed to extend this experimental procedure to all Italian strains possessing A1-3 reciprocal translocations, in order to better clarify the relationships between this chromosomal rearrangement and the insecticide resistance in *M. persicae* populations.

Other fissions relatively frequent in the studied Italian *M. persicae* populations occurred at autosomes 3 and 4,

whereas in one case only the fission involved the X chromosome. Different autosome fragmentations have been repeatedly described in *M. persicae* populations collected worldwide, whereas the X fragmentation has been observed only in a *M. persicae* laboratory strain characterised by an extensive chromosomal mosaicism (Monti *et al.*, 2012). In this connection, it must be emphasized that in both such cases, the X fission occurs in X chromosomes possessing a low number of rDNA genes and in the telomeric region opposite to the NORs-bearing one. The recurrent fission of the same chromosomes in the same region argues that the *M. persicae* genome possesses some fragile/labile sites that could be the basis for the observed changes in the chromosome number.

For many years, chromosome evolution has been generally explained by considering the random-breakage model (Becker & Lenhard, 2007). On the contrary, a number of comparative cytogenetic studies evidences a relationship between chromosomal rearrangements and specific chromosomal architecture and suggests a role of the repetitive DNAs in chromosome




Fig. 4. *M. persicae* chromosome complements showing A3 fission. (a) Forlì 01 and (b) Salerno 02 are stained with Giemsa, whereas (c) Salerno 03 and (d) its relative karyotype are silver stained. Arrow heads indicate chromosomes involved in the fission. Bar corresponds to 10  $\mu$ m.




Fig. 5. Giemsa staining of *M. persicae* chromosome complements showing A4 fission: (a) Cosenza 01, (b) Ravenna 06 and (c) Piacenza 10. The (d) karyotype is derived from (b) Ravenna 06. Arrow heads indicate chromosomes involved in the fission. Bar corresponds to 10  $\mu$ m.

240 rearrangements. The nature of the repetitive DNA within  
241 chromosomal breakpoint regions varies significantly, from  
242 clusters of rRNA and tRNA genes to simple di- and  
tri-nucleotide expansions (Caceres *et al.*, 1999; Carlton *et al.*, 243  
2002; Coghlan & Wolfe, 2002; Kellis *et al.*, 2003; Renciuk *et al.*, 244  
2011). The data reported in this paper confirmed recent 245




Fig. 6. (a) Pescara 02 complement stained with  $\text{AgNO}_3$  and (b) relative karyotype. (c) Cosenza 02 complement silver stained with (d) relative karyotype. Arrow heads indicate chromosomes involved in the fissions. Asterisk indicates A1–3 translocated chromosomes. Bar corresponds to 10  $\mu\text{m}$ .

observations regarding the recurrent fission of the same chromosomes in the same region (Monti *et al.*, 2012), allowing us to further support the hypothesis concerning the presence of fragile/labile sites in the *M. persicae* holocentric chromosomes.

Chromosomal rearrangements in aphids have been hypothesized to affect some complex phenotypic traits, such as the host plant choice (Blackman, 1987; French-Constant *et al.*, 1988). For example, karyotypic variants observed in the corn leaf aphid *Rhopalosiphum maidis* (Fitch) have been associated with changes in the host choice. Similarly, an association of chromosome number with host plant has been described within the *Sitobion* genus, which shows  $2n=12$  on ferns and  $2n=18$  on grasses (Brown & Blackman, 1988; Hales *et al.*, 1997).

A peculiar example of host adaptation concerns *M. persicae* strains feeding on tobacco. Morphometric analyses of specific taxonomic markers revealed that they are distinguishable from those living on other host plant so that the tobacco-feeding form was elevated to the status of a separate species by Blackman (1987). Further molecular evidences failed to confirm the genetic isolation of the population living on tobacco (Field *et al.*, 1994; Clements *et al.*, 2000), although other data, as well as behavioural/pheromonal evidence, suggests that the two forms undergone some significant degree of ecological-evolutionary divergence (Kephalogianni *et al.*, 2002; Margaritopoulos *et al.*, 2003; Blackman *et al.*, 2007).

Our data put in evidence that all but one of the strains collected on tobacco plants showed karyotype variations, whereas only four of the 56 population collected on other hosts (corresponding to about 7% of the total) displayed chromosomal rearrangements. A suggestive explanation for the observed relationships between chromosomal rearrangements

and tobacco plants could rely in the clastogenic effect of nicotine.

Nicotine is a naturally occurring alkaloid found primarily in members of the solanaceous plant family, including *Nicotiana tabacum*. Several reports showed that nicotine, as a consequence of DNA replication fork stress (Richards, 2001; Freudenreich, 2005), produces genotoxic effects on Chinese hamster ovarian (CHO) cells (Trivedi *et al.*, 1990, 1993) and sister chromatid exchanges and chromosome aberrations in bone marrow cells of mice (Sen *et al.*, 1991). Extensive chromosomal rearrangements have also been described in a mice population known as 'tobacco mice' since they live close to kiln for drying tobacco (Fraguedakis-Tsolis *et al.*, 1997). In addition, DNA fragmentation by nicotine has been demonstrated both in peripheral lymphocytes (Sassen *et al.*, 2005) and in human spermatozoa (Arabi, 2004). Nicotine, together with ultraviolet exposure, has also been considered an exogenous factor which can contribute to the generation of mutations which could be at the basis of chromosomal mosaicism (De, 2011), a very rare phenomenon we have observed in Salerno 02, one of the strains collected on tobacco plants.

Even if there are no literature data analyzing nicotine effects on organisms possessing holocentric chromosomes, the previously reported data allows us to propose at least that chromosome architecture, rather than random breakages, has a pivotal role in aphid chromosome evolution and rearrangements.

The high telomerase expression, previously reported in *M. persicae* (Monti *et al.*, 2011), that stabilized chromosomes involved in fragmentations, coupled to reproduction by obligate apomictic parthenogenesis, could be at the basis of the stabilization of the observed chromosome instability on *M. persicae* strains collected on tobacco plants favouring the inheritance of the variant karyotypes.

## References

- Arabi, M. (2004) Nicotinic infertility: assessing DNA and plasma membrane integrity of human spermatozoa. *Andrologia* **36**, 305–310.
- Becker, T.S. & Lenhard, B. (2007) The random versus fragile breakage models of chromosome evolution: a matter of resolution. *Molecular Genetics and Genomics* **278**, 487–491.
- Blackman, R.L. (1971) Variation in the photoperiodic response within natural populations of *Myzus persicae* (Sulz.). *Bulletin of Entomological Research* **60**, 533.
- Blackman, R.L. (1980) Chromosome numbers in the Aphididae and their taxonomic significance. *Systematic Entomology* **5**, 7–25.
- Blackman, R.L. (1987) Morphological discrimination of a tobacco-feeding form from *Myzus persicae* (Sulzer) (Hemiptera: Aphididae), and a key to new world *Myzus* (*Nectarosiphon*) species. *Bulletin of Entomological Research* **77**, 713–730.
- Blackman, R.L., Takada, H. & Kawakami, K. (1978) Chromosomal rearrangement involved in insecticide resistance of *Myzus persicae*. *Nature* **271**, 450–452.
- Blackman, R.L., Spence, J.M., Field, L.M. & Devonshire, A.L. (1995) Chromosomal location of the amplified esterase genes conferring resistance to insecticides in *Myzus persicae* (Homoptera: Aphididae). *Heredity* **75**, 297–302.
- Blackman, R.L., Malarky, G. & Margaritopoulos, J.T. (2007) Distribution of common genotypes of *Myzus persicae* (Hemiptera: Aphididae) in Greece, in relation to life cycle and host plant. *Bulletin of Entomological Research* **97**, 253–263.
- Brown, P.A. & Blackman, R.L. (1988) Karyotype variation in the corn leaf aphid, *Rhopalosiphum maidis* (Fitch), species complex (Hemiptera: Aphididae) in relation to host-plant and morphology. *Bulletin of Entomological Research* **78**, 351–363.
- Caceres, M., Ranz, J.M., Barbadilla, A., Long, M. & Ruiz, A. (1999) Generation of a widespread *Drosophila* inversion by a transposable element. *Science* **285**, 415–418.
- Carlton, J.M., Angiuoli, S.V., Suh, B.B., Kooij, T.W., Perte, M., Silva, J.C., Ermolaeva, M.D., Allen, J.E., Selengut, J.D., Koo, H.L., Peterson, J.D., Pop, M., Kosack, D.S., Shumway, M.F., Bidwell, S.L., Shallom, S.J., van Aken, S. E., Riedmuller, S.B., Feldblyum, T.V., Cho, J.K., Quackenbush, J., Sedegah, M., Shoaibi, A., Cummings, L. M., Florens, L., Yates, J.R., Raine, J.D., Sinden, R.E., Harris, M.A., Cunningham, D.A., Preiser, P.R., Bergman, L. W., Vaidya, A.B., van Lin, L.H., Janse, C.J., Waters, A.P., Smith, H.O., White, O.R., Salzberg, S.L., Venter, J.C., Fraser, C.M., Hoffman, S.L., Gardner, M.J. & Carucci, D.J. (2002) Genome sequence and comparative analysis of the model rodent malaria parasite *Plasmodium yoelii yoelii*. *Nature* **419**, 512–519.
- Clements, K.M., Sorenson, C.E., Wiegmann, B.M., Neese, P.A. & Roe, R.M. (2000) Genetic, biochemical, and behavioural uniformity among populations of *Myzus nicotianae* and *Myzus persicae*. *Entomologia Experimentalis et Applicata* **95**, 269–281.
- Coghlan, A. & Wolfe, K.H. (2002) Fourfold faster rate of genome rearrangement in nematodes than in *Drosophila*. *Genome Research* **12**, 857–867.
- Crema, R. (1979) Egg viability and sex determination in *Megoura viciae* (Homoptera: Aphididae). *Entomologia Experimentalis et Applicata* **26**, 152–156.
- De, S. (2011) Somatic mosaicism in healthy human tissues. *Trends in Genetics* **27**, 217–223.
- Devonshire, A.L., Devine, G.J. & Moores, G.D. (1992) Comparison of microplate esterase assays and immunoassay for identifying insecticide resistant variants of *Myzus persicae* (Homoptera: Aphididae). *Bulletin of Entomological Research* **82**, 459–463.
- French-Constant, R.H., Byrne, F.J., Stribley, M.F. & Devonshire, A.L. (1988) Rapid identification of the recently recognised *Myzus antirrhinii* (Macchiati) (Hemiptera: Aphididae) by polyacrylamide gel electrophoresis. *Entomologist* **107**, 20–23.
- Field, L.M., Javed, N., Stribley, M.F. & Devonshire, A.L. (1994) The peach-potato aphid *Myzus persicae* and the tobacco aphid *Myzus nicotianae* have the same esterase-based mechanisms of insecticide resistance. *Insect Molecular Biology* **3**, 143–148.
- Fraguedakis-Tsolis, S., Hauffe, H.C. & Searle, J.B. (1997) Genetic distinctiveness of a village population of house mice: Relevance to speciation and chromosomal evolution. *Proceedings of the Royal Society of London, Series B: Biological Science* **264**, 355–360.
- Freudenreich, C.H. (2005) Molecular mechanisms of chromosome fragility. *ChemTracks-Biochemistry and Molecular Biology* **18**, 141–152.
- Hales, D.F. (1989) The chromosomes of *Schoutedenia lutea* (Homoptera, Aphidoidea, Greenideinae), with an account of meiosis in the male. *Chromosoma* **98**, 295–300.
- Hales, D.F., Tomiuk, J., Wohrmann, K. & Sunnucks, P. (1997) Evolutionary and genetic aspects of aphid biology: A review. *European Journal of Entomology* **94**, 1–55.
- Howell, W.M. & Black, D.A. (1980) Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. *Experientia* **36**, 1014–1015.
- Kellis, M., Patterson, N., Endrizzi, M., Birren, B. & Lander, E.S. (2003) Sequencing and comparison of yeast species to identify genes and regulatory elements. *Nature* **423**, 241–254.
- Lauritzen, M. (1982) Q-Band and G-Band Identification of 2 chromosomal rearrangements in peach-potato aphids, *Myzus persicae* (Sulzer), resistant to insecticides. *Hereditas* **97**, 95–102.
- Manicardi, G.C., Mandrioli, M., Bizzaro, D. & Bianchi, U. (2002) Cytogenetic and molecular analysis of heterochromatic areas in the holocentric chromosomes of different aphid species. pp. 47–56 in Sobti, R.C., Obe, G. & Athwal, R.S. (Eds), *Some Aspects of Chromosome Structure and Functions*. New Delhi, India, Narosa Publishing House.
- Kephalogianni, T.E., Tsitsipis, J.A. & Margaritopoulos, J.T. (2002) Variation in the life cycle and morphology of the tobacco host-race of *Myzus persicae* (Hemiptera: Aphididae) in relation to its geographical distribution. *Bulletin of Entomological Research* **92**, 301–307.
- Loxdale, H.D. & Lushai, G. (2003) Rapid changes in clonal lines: the death of a 'sacred cow'. *Biological Journal of the Linnean Society* **79**, 3–16.
- Margaritopoulos, J.T., Blackman, R.L. & Tsitsipis, J.A. (2003) Co-existence of different host-adapted forms of the *Myzus persicae* group (Hemiptera: Aphididae) in southern Italy. *Bulletin of Entomological Research* **93**, 131–135.
- Monti, V., Giusti, M., Bizzaro, D., Manicardi, G.C. & Mandrioli, M. (2011) Presence of a functional (TTAGG)<sub>n</sub> telomere-telomerase system in aphids. *Chromosome Research* **19**, 625–633.
- Monti, V., Mandrioli, M., Rivi, M. & Manicardi, G.C. (2012) The vanishing clone: karyotypic evidence for extensive intra-clonal genetic variation in the peach potato aphid, *Myzus persicae* (Hemiptera: Aphididae). *Biological Journal of the Linnean Society* **105**, 350–358.

- 440 **Renciuk, D., Kypr, J. & Vorlickova, M.** (2011) CGG Repeats  
441 associated with fragile X chromosome form left-handed  
442 Z-DNA structure. *Biopolymers* **95**, 174–181.
- 443 **Richards, R.** (2001) Fragile and unstable chromosomes in  
444 cancer: causes and consequences. *Trends in Genetics* **17**,  
445 339–345.
- 446 **Rivi, M., Mazzoni, E., Criniti, A., Cassanelli, S., Bizzaro, D. &**  
447 **Manicardi, G.C.** (2009) Relationship between chromosomal  
448 translocation and FE4 gene amplification in an Italian  
449 population of the peach-potato aphid *Myzus persicae*  
450 (Hemiptera: Aphididae). *Redia* **92**, 229–231.
- 451 **Sassen, A., Richter, E., Semmler, M., Harreus, U., Gamarra, F. &**  
452 **Kleinsasser, N.** (2005) Genotoxicity of nicotine in mini-organ  
453 cultures of human upper aerodigestive tract epithelia RID  
454 A-3601-2008. *Toxicological Sciences* **88**, 134–141.
- 455 **Sen, S., Sharma, A. & Talukder, G.** (1991) Inhibition of clastogenic  
456 effects of nicotine by chlorophyllin in mice bone-marrow cells  
457 *in vivo*. *Phytotherapy Research* **5**, 130–133.
- 458 **Spence, J.M. & Blackman, R.L.** (1998) Chromosomal rearrange-  
459 ments in the *Myzus persicae* group and their evolutionary  
460 significance. pp. 113–118 in Nieto Nafria, J.M. & Dixon, A.F.  
461 G. (Eds) *Chromosomal Rearrangements in the Myzus persicae*  
Group and their Evolutionary Significance. León, Spain, 462  
Universidad De León Secretariado de Publicacions. 463
- Spence, J.M. & Blackman, R.L.** (2000) Inheritance and meiotic  
464 behaviour of a de novo chromosome fusion in the aphid  
465 *Myzus persicae* (Sulzer). *Chromosoma* **109**, 490–497. 466
- Trivedi, A.H., Dave, B.J. & Adhvaryu, S.G.** (1990) Assessment of  
467 genotoxicity of nicotine employing *in vitro* mammalian test  
468 system. *Cancer Letters* **54**, 89–94. 469
- Trivedi, A.H., Dave, B.J. & Adhvaryu, S.G.** (1993) Genotoxic  
470 effects of tobacco extract on Chinese hamster ovary cells.  
471 *Cancer Letters* **70**, 107–112. 472
- White, M.J.D.** (1973) *Animal Cytology and Evolution*. Cambridge,  
473 UK, Cambridge University Press. 474
- Wensch, D.L., Ketheley, J.B. & Norton, R.A.** (1994) Cytogenetic  
475 of holokinetic chromosomes and inverted meiosis: keys to the  
476 evolutionary success of mites with generalization on eu-  
477 karyotes. pp. 282–343 in Houck, M.A. (Ed.), *Mites: Ecological*  
478 *and Evolutionary Analysis of Life-History Patterns*. New York,  
479 USA, Chapman & Hall. 480
- Yang, X.-W. & Zhang, X.** (2000) Karyotype polymorphism in  
481 different geographic populations of green peach aphid  
482 *Myzus persicae* (Sulzer) in China. *Entomologia Sinica* **7**, 29–35. 483  
484