

This is the peer reviewed version of the following article:

Joint Complete Records occurrence over FGM sequences / Khorrami Chokami, Amir; Montagna, Silvia. - (2024). (Intervento presentato al convegno The 52nd Scientific Meeting of the Italian Statistical Society tenutosi a Università degli Studi di Bari Aldo Moro, Bari nel 17-20 Giugno 2024).

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

15/10/2024 10:16

(Article begins on next page)

Joint Complete Records occurrence over FGM sequences

Amir Khorrami Chokami¹ and Silvia Montagna²

¹ ESOMAS Department, Università di Torino and Collegio Carlo Alberto, Torino, Italy

`amir.khorramichokami@unito.it`,

² Dipartimento di Economia “Marco Biagi”, Università degli Studi di Modena e Reggio Emilia, via Berengario 51, Modena 41121, Italy

Abstract. A record over a sequence of iid absolutely continuous random variables is defined as a random variable surpassing all preceding ones. The extension to the multivariate case is not obvious. In this project, we focus on the so-called complete records. We examine the probability of appearance of two complete records at two different times, whenever the within sequences of random vectors where the dependence among components follows the Farlie-Gumbel-Morgenstern copula.

Keywords: complete records, time dependence, standard max stable distribution, FGM copula.

1 Introduction

In recent times, there has been a growing interest on the occurrence of extreme events. The study of maxima in a multivariate context presents the challenge of tackling the dependence among extreme events, in connection with the parent copula of the underlying sequence of random vectors (see [9], [8] among others). Let us consider now a sequence of iid random variables X_1, X_2, \dots . The rv X_m is called a *record* if $X_m > \max(X_1, \dots, X_{m-1})$, $m \geq 2$ (obviously, X_1 is the first record because it is the first variable appearing in the sequence). In this simple case, various results concerning the time of appearance of a record, the asymptotic distribution of the m -th record as $m \rightarrow \infty$, increments between records, are available (see [1], [4, Sections 6.2 and 6.3], [2] and [3], among others). In particular, if we call $I_m^R = \mathbb{1}(X_m > \max(X_1, \dots, X_{m-1}))$, $m \in \mathbb{N}$, then

$$\mathbb{P}(I_j^R = 1, I_k^R = 1) = \mathbb{P}(I_j^R = 1) \mathbb{P}(I_k^R = 1) = j^{-1}k^{-1}, \quad j < k, \quad (1)$$

indicating independence between the appearance of records at two different times.

The extension to the multivariate case is not an easy task. The first issue we have to face, regards the non-unique definition of a multivariate record. In this work, we consider the so-called complete records, defined as follows. Let $\mathbf{X}_1, \mathbf{X}_2, \dots$, be iid random vectors (rv) in \mathbb{R}^d . Put for integers $j < k$

$$\mathbf{M}_j := \max_{1 \leq i \leq j} \mathbf{X}_i, \quad \mathbf{M}_j^k := \max_{j \leq i \leq k} \mathbf{X}_i,$$

where the maximum is taken componentwise. a rv \mathbf{X}_m is a *complete record* if each component is a univariate record itself, i.e., $\mathbf{X}_m > \mathbf{M}_{m-1}$. The event indicating the appearance of a complete record at time m is then $I_m^{\text{CR}} := \mathbf{1}(\mathbf{X}_m > \mathbf{M}_{m-1})$. Some findings regarding complete records across sequences of random vectors with independent components and with Marshall-Olkin copula are available in [2] and [7], respectively.

This paper studies the left-hand side of Equation (1) in a bivariate context, precisely when the copula of the underlying sequence of random vectors is of the Farlie-Gumbel-Morgenstern (FGM) type:

$$F(x, y) = e^{(x+y)} (1 + \lambda (1 - e^x) (1 - e^y)), \quad x, y \leq 0 \text{ and } |\lambda| \leq 1. \quad (2)$$

with bivariate density given by

$$f(x, y) = e^{x+y} (1 + \lambda (2e^x - 1) (2e^y - 1)). \quad (3)$$

The challenge in studying records arises from the immediate complexity of computations required when transitioning from independent cases to those where components are dependent. The FGM copula, albeit with difficulties, enables the derivation of exact formulas for complete record occurrences. This copula is valuable due to its simple structure and straightforward interpretation of the parameter λ , which captures dependence characteristics. These qualities have made the FGM distribution a popular choice for various applications, including capital allocation and the so-called collective risk models. For a thorough understanding of this copula in an extreme-value context, refer to [5]. We highlight our findings are independent of the marginal distribution function F_{X_i} , $i = 1, 2$, provided that they are continuous. Moreover, we consider a sequence $\boldsymbol{\eta}_1, \boldsymbol{\eta}_2, \dots \in \mathbb{R}^2$ of *standard max-stable* rv (i.e. with Negative-Exp(1) margins and such that $\mathbf{M}_n \stackrel{d}{=} \boldsymbol{\eta}_1/n$) and FGM copula. This is also the framework of [6], which deals with single complete records.

2 Complete Records Occurrences

Theorem 1. *Let $\boldsymbol{\eta}_1, \boldsymbol{\eta}_2, \dots$ be a sequence of bivariate standard max-stable rv with FGM copula. Then, the probability of appearance of a CR is*

$$\mathbb{P}(I_j^R = 1, I_k^R = 1) = \sum_{i=1}^{16} p_i \quad (4)$$

for $j, k \in \mathbb{N}, j < k$. The p_i 's are given in the proof for readability.

Proof. For what concerns the probability of the joint events we have

$$\begin{aligned}
 \mathbb{P}(I_j^{\text{CR}} = 1, I_k^{\text{CR}} = 1) &= \mathbb{P}(\boldsymbol{\eta}_j > \mathbf{M}_{j-1}, \boldsymbol{\eta}_k > \mathbf{M}_{k-1}) \\
 &= \mathbb{P}\left(\boldsymbol{\eta}_j > \frac{\boldsymbol{\eta}_1}{j-1}, \boldsymbol{\eta}_k > \max\left(\boldsymbol{\eta}_j, \frac{\boldsymbol{\eta}_2}{k-j-1}\right)\right) \\
 &= \int_{(-\infty, 0]^2} \int_{(-\infty, \mathbf{y}]} \mathbb{P}(\boldsymbol{\eta}_1 < (j-1)\mathbf{x}, \boldsymbol{\eta}_2 < (k-j-1)\mathbf{y}) \, d\mathbb{P}_{\boldsymbol{\eta}}(\mathbf{x}) \, d\mathbb{P}_{\boldsymbol{\eta}}(\mathbf{y}) \\
 &= \int_{(-\infty, 0]^2} \mathbb{P}(\boldsymbol{\eta}_2 < (k-j-1)\mathbf{y}) \int_{(-\infty, \mathbf{y}]} \mathbb{P}(\boldsymbol{\eta}_1 < (j-1)\mathbf{x}) \, d\mathbb{P}_{\boldsymbol{\eta}}(\mathbf{x}) \, d\mathbb{P}_{\boldsymbol{\eta}}(\mathbf{y}).
 \end{aligned}$$

Let us compute the internal integral, which will yield a function of \mathbf{y} .

$$\begin{aligned}
 \int_{(-\infty, \mathbf{y}]} \mathbb{P}(\boldsymbol{\eta}_1 < (j-1)\mathbf{x}) \, d\mathbb{P}_{\boldsymbol{\eta}}(\mathbf{x}) &= \int_{-\infty}^{y_2} \int_{-\infty}^{y_1} \mathbb{P}(\eta_1^{(1)} < (j-1)x_1, \eta_1^{(2)} < (j-1)x_2) \, d\mathbb{P}_{\boldsymbol{\eta}}(\mathbf{x}) = \\
 &= \int_{-\infty}^{y_2} \int_{-\infty}^{y_1} e^{(j-1)(x_1+x_2)} \left(1 + \lambda \left(1 - e^{(j-1)x_1}\right) \left(1 - e^{(j-1)x_2}\right)\right) \cdot \\
 &\quad \cdot e^{x_1+x_2} \left(1 + \lambda(2e^{x_1} - 1)(2e^{x_2} - 1)\right) \, dx_1 \, dx_2 = \\
 &= \int_{-\infty}^{y_2} \int_{-\infty}^{y_1} e^{j(x_1+x_2)} \, dx_1 \, dx_2 \\
 &\quad + \int_{-\infty}^{y_2} \int_{-\infty}^{y_1} e^{j(x_1+x_2)} \lambda \left(1 - e^{(j-1)x_1}\right) \left(1 - e^{(j-1)x_2}\right) \, dx_1 \, dx_2 \\
 &\quad + \int_{-\infty}^{y_2} \int_{-\infty}^{y_1} e^{j(x_1+x_2)} \lambda (2e^{x_1} - 1)(2e^{x_2} - 1) \, dx_1 \, dx_2 \\
 &\quad + \int_{-\infty}^{y_2} \int_{-\infty}^{y_1} e^{j(x_1+x_2)} \lambda^2 \left(1 - e^{(j-1)x_1}\right) \left(1 - e^{(j-1)x_2}\right) (2e^{x_1} - 1)(2e^{x_2} - 1) \, dx_1 \, dx_2 \\
 &=: I_1 + I_2 + I_3 + I_4.
 \end{aligned}$$

It can be shown that

$$\begin{aligned}
 I_1 &= \frac{1}{j^2} e^{j(y_1+y_2)} \\
 I_2 &= \lambda \prod_{i=1}^2 \left(\frac{e^{jy_i}}{j} - \frac{e^{(2j-1)y_i}}{2j-1} \right) \\
 I_3 &= \lambda \prod_{i=1}^2 \left(2 \frac{e^{(j+1)y_i}}{j+1} - \frac{e^{jy_i}}{j} \right) \\
 I_4 &= \lambda^2 \prod_{i=1}^2 \left(\frac{2}{j+1} e^{(j+1)y_i} - \frac{e^{jy_i}}{j} - \frac{e^{2jy_i}}{j} + \frac{e^{(2j-1)y_i}}{2j-1} \right).
 \end{aligned}$$

Thus, we have

$$\begin{aligned}
& \int_{(-\infty, 0]^2} \mathbf{P}(\boldsymbol{\eta}_2 < (k-j-1)\mathbf{y}) \int_{(-\infty, \mathbf{y}] } \mathbf{P}(\boldsymbol{\eta}_1 < (j-1)\mathbf{x}) d\mathbf{P}_{\boldsymbol{\eta}}(\mathbf{x}) d\mathbf{P}_{\boldsymbol{\eta}}(\mathbf{y}) \\
&= \int_{-\infty}^0 \int_{-\infty}^0 \left[\sum_{i=1}^4 I_i \right] \cdot F((k-j-1)y_1, (k-j-1)y_2) f(y_1, y_2) dy_1 dy_2 \\
&== \sum_{i=1}^4 \int_{-\infty}^0 \int_{-\infty}^0 I_i \cdot F((k-j-1)y_1, (k-j-1)y_2) f(y_1, y_2) dy_1 dy_2,
\end{aligned}$$

where F and f are as in Equation (2) and Equation (3), respectively. Solving the bivariate integral above, requires the computation of the following 16 integrals, each one relative to a different p_i in Equation (4). This is because $F((k-j-1)y_1, (k-j-1)y_2) f(y_1, y_2)$ is a sum of four different contributes, each of them being multiplied with each one of the 4 integrals.

For $F \cdot f \cdot I_1$: The corresponding double integral is the sum of the following

$$\begin{aligned}
p_1 &= \frac{1}{j^2 k^2} \\
p_2 &= \frac{\lambda}{j^2} \left(\frac{1}{k} - \frac{1}{2k-j-1} \right)^2 \\
p_3 &= \frac{\lambda}{j^2} \left(\frac{2}{k+1} - \frac{1}{k} \right)^2 \\
p_4 &= \frac{\lambda^2}{j^2} \left(\frac{2}{k+1} - \frac{1}{k} - \frac{2}{2k-j} + \frac{1}{2k-j-1} \right)^2
\end{aligned}$$

For $F \cdot f \cdot I_2$: The corresponding double integral is the sum of the following

$$\begin{aligned}
p_5 &= \lambda \left(\frac{1}{jk} - \frac{1}{(2j-1)(k+j-1)} \right)^2 \\
p_6 &= \lambda \left(\frac{1}{jk} - \frac{1}{(2j-1)(k+j-1)} - \frac{1}{j(2k-j-1)} - \frac{1}{(2j-1)(2k-3)} \right)^2 \\
p_7 &= \lambda \left(\frac{2}{j(k+1)} - \frac{2}{(2j-1)(k+j)} - \frac{1}{jk} - \frac{1}{(2j-1)(k+j-1)} \right)^2 \\
p_8 &= \lambda^2 \left(\frac{2}{j(k+1)} - \frac{2}{(2j-1)(k+j)} - \frac{1}{jk} + \frac{1}{(2j-1)(k+j-1)} - \frac{2}{j(2k-j)} + \frac{2}{(2j-1)(2k-1)} \right. \\
&\quad \left. + \frac{1}{j(2k-j-1)} - \frac{2}{(2j-1)(2k-3)} \right)^2
\end{aligned}$$

For $F \cdot f \cdot I_3$: The corresponding double integral is the sum of the following

$$\begin{aligned}
 p_9 &= \lambda \left(\frac{2}{(j+1)(k+1)} - \frac{1}{jk} \right)^2 \\
 p_{10} &= \lambda \left(\frac{2}{(j+1)(k+1)} - \frac{1}{jk} - \frac{2}{(j+1)(2k-j)} + \frac{1}{j(2k-j-1)} \right)^2 \\
 p_{11} &= \lambda \left(\frac{4}{(j+1)(k+2)} - \frac{2}{j(k+1)} - \frac{2}{(j+1)(k+1)} + \frac{1}{jk} \right)^2 \\
 p_{12} &= \lambda^2 \left(\frac{4}{(j+1)(k+2)} - \frac{2}{j(k+1)} - \frac{2}{(j+1)(k+1)} + \frac{1}{jk} - \frac{k}{(j+1)(2k-j+1)} + \frac{2}{j(2k-j)} \right. \\
 &\quad \left. + \frac{2}{(j+1)(2k-j)} - \frac{1}{j(2k-j-1)} \right)^2
 \end{aligned}$$

For $F \cdot f \cdot I_4$: The corresponding double integral is the sum of the following

$$\begin{aligned}
 p_{13} &= \lambda^2 \left(\frac{2}{(j+1)(k+1)} - \frac{1}{jk} - \frac{1}{j(k+j)} + \frac{1}{(2j-1)(k+j-1)} \right)^2 \\
 p_{14} &= \lambda \left(\frac{2}{(j+1)(k+1)} - \frac{1}{jk} - \frac{1}{j(k+j)} + \frac{1}{(2j-1)(k+j-1)} \right. \\
 &\quad \left. - \frac{2}{(j+1)(2k-j)} + \frac{1}{j(2k-j-1)} + \frac{1}{j(2k-1)} - \frac{1}{(2j-1)(2k-2)} \right)^2 \\
 p_{15} &= \lambda^2 \left(\frac{4}{(j+1)(k+1)} - \frac{2}{j(k+1)} - \frac{2}{j(k+j+1)} + \frac{2}{(2j-1)(k+j)} \right. \\
 &\quad \left. - \frac{2}{(j+1)(k+1)} + \frac{1}{jk} + \frac{1}{j(k+j)} - \frac{1}{(2j-1)(k+j-1)} \right)^2 \\
 p_{16} &= \lambda^2 \left(\frac{4}{(j+1)(k+2)} - \frac{2}{j(k+1)} - \frac{2}{j(k+j+1)} + \frac{2}{(2j-1)(k+j)} \right. \\
 &\quad - \frac{2}{(j+1)(k+1)} + \frac{1}{jk} + \frac{1}{j(k+j)} - \frac{1}{(2j-1)(k+j-1)} + \\
 &\quad - \frac{4}{(j+1)(k+1)} + \frac{2}{j(2k-j)} + \frac{1}{jk} - \frac{2}{(2j-1)(2k-1)} + \\
 &\quad \left. + \frac{2}{(j+1)(2k-j)} - \frac{1}{j(2k-j-1)} - \frac{1}{j(2k-1)} + \frac{1}{(2j-1)(2k-2)} \right)^2.
 \end{aligned}$$

This concludes the proof.

3 Conclusion

In conclusion, this work addresses the challenge of studying the probability of occurrence of two complete records over independent and identically distributed

bivariate sequences of standard max-stable random variables with the FGM copula. The authors suggest the use of mathematical softwares to simplify the expressions given in the proof of Theorem 1. Ongoing projects regard the extension the case of complete records to sequences characterized by a general copula, and the study of alternative definitions of multivariate records.

References

1. Arnold, B.C., Balakrishnan, N., Nagaraja, H.N.: Records. Wiley Series in Probability and Statistics. Wiley, New York (1998)
2. Falk, M., Khorrami Chokami, A., Padoan, S.: Some results on joint record events. *Statistics & Probability Letters* **135**, 11–19 (2018). DOI <https://doi.org/10.1016/j.spl.2017.11.011>. URL <https://www.sciencedirect.com/science/article/pii/S0167715217303656>
3. Falk, M., Khorrami Chokami, A., Padoan, S.A.: Records for time-dependent stationary gaussian sequences. *Journal of Applied Probability* **57**, 78–96 (2020). DOI [10.1017/jpr.2019.77](https://doi.org/10.1017/jpr.2019.77)
4. Galambos, J.: The Asymptotic Theory of Extreme Order Statistics, 2 edn. Krieger, Malabar (1987)
5. Hashorva, E., Hüsler, J.: Extreme values in FGM random sequences. *Journal of Multivariate Analysis* **68**(2), 212–225 (1999). DOI <https://doi.org/10.1006/jmva.1998.1795>. URL <https://www.sciencedirect.com/science/article/pii/S0047259X9891795X>
6. Khorrami Chokami, A.: Complete records over independent FGM sequences. *Book of Short Papers CLADAG 2023* (2023)
7. Khorrami Chokami, A., Padoan, S.A.: On multivariate records over sequences of random vectors with marshall-olkin dependence of components. *Proceedings of the LI Meeting of the Italian Statistical Society*. (2023)
8. Leonetti, P., Khorrami Chokami, A.: The maximum domain of attraction of multivariate extreme value distributions is small. *Electronic Communications in Probability* **27**, 1 – 8 (2022). DOI [10.1214/22-ECP501](https://doi.org/10.1214/22-ECP501). URL <https://doi.org/10.1214/22-ECP501>
9. Resnick, S.I.: Extreme Values, Regular Variation, and Point Processes, *Applied Probability*, vol. 4. Springer, New York (1987). DOI [10.1007/978-0-387-75953-1](https://doi.org/10.1007/978-0-387-75953-1)