

This is a pre print version of the following article:

Monitoring Caffeine Intake: The Relevance of Adequate Assessment in the Population / Mattioli, Anna Vittoria; Manenti, Antonio; Farinetti, Alberto. - In: JOURNAL OF THE AMERICAN NUTRITION ASSOCIATION. - ISSN 2769-7061. - 43:1(2024), pp. 110-112. [10.1080/27697061.2023.2225095]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

04/11/2024 03:45

(Article begins on next page)

Monitoring Caffeine Intake: The Relevance of Adequate Assessment in the Population

Anna Vittoria Mattioli, Antonio Manenti and Alberto Farinetti

QUERY SHEET

This page lists questions we have about your paper. The numbers displayed at left are hyperlinked to the location of the query in your paper.

The title and author names are listed on this sheet as they will be published, both on your paper and on the Table of Contents. Please review and ensure the information is correct and advise us if any changes need to be made. In addition, please review your paper as a whole for typographical and essential corrections.

Your PDF proof has been enabled so that you can comment on the proof directly using Adobe Acrobat. For further information on marking corrections using Acrobat, please visit <https://authorservices.taylorandfrancis.com/how-to-correct-proofs-with-adobe/>

The CrossRef database (www.crossref.org/) has been used to validate the references.


AUTHOR QUERIES

- Q1** Please check and confirm whether the author affiliations and corresponding details have been set correctly.
- Q2** This journal requires a structured abstract.
- Q3** The disclosure statement has been inserted. Please correct if this is inaccurate.
- Q4** Please provide missing volume number for the "(Ref. 1)." references list entry.
- Q5** Please note that the ORCID section has been created from information supplied with your manuscript submission/ CATS. Please correct if this is inaccurate.



ARTICLE COMMENTARY

Monitoring Caffeine Intake: The Relevance of Adequate Assessment in the Population

Anna Vittoria Mattioli , Antonio Manenti and Alberto Farinetti

Department of Medical and Surgical Sciences for Children and Adults, University of Modena and Reggio Emilia, Modena, Italy

ABSTRACT

The present letter to editor comments on the manuscript “Bulczak EM, Chmurzyńska AU. Caffeine Consumption in Polish Adults: Development and Validation of a Polish Questionnaire for Assessing Caffeine Intake. *J Am Nutr Assoc.* 2023 Feb 1:1–7. doi:10.1080/27697061.2023.2172749. Epub ahead of print. PMID: 36725370.” regarding adequate monitoring of the consumption of caffeine.

ARTICLE HISTORY

Received 6 February 2023

Revised 24 May 2023

Accepted 9 June 2023

KEYWORDS

Monitoring; caffeine; pandemic; stress

Dear Editor,

We have read with great interest the article “Caffeine Consumption in Polish Adults: Development and Validation of a Polish Questionnaire for Assessing Caffeine Intake” by Bulczak and Chmurzyńska, and we found it of relevance with a view to caffeinated beverages consumption (1). This article aims to design and validate an online food frequency questionnaire (FFQ) on caffeine intake and to use it to estimate caffeine consumption in Polish adults.

The authors concluded that the average caffeine consumption among Polish adults slightly exceeds the safe consumption dose established by the European Food Safety Authority.

We find this manuscript of great interest and would like to contribute to the discussion.

The creation of validated questionnaires for a specific population is essential, especially in the evaluation of substances such as caffeine and which are present in various drinks and foods. Coffee is the most common source of caffeine among adults, while young people prefer energy drinks.

The habits of preparing and drinking coffee vary greatly in different countries, and in the European nations, different varieties of preparation are distinguished, for example, espresso coffee, mocha, and the coffee pot, which provides different filtration of the various substances contained in coffee in addition to caffeine (2–4). Furthermore, habits change in different countries, and in the Mediterranean countries, coffee is taken in relation to the meal. To this we must add the habit of integrating the drink with milk and sugar, all confounding factors when one wants to carry out a precise analysis of caffeine intake (5).

Validation of a caffeine intake questionnaire is crucial to ensure that it accurately reflects the individual's consumption and provides reliable data. Furthermore, the questionnaire can also be compared with other methods of assessing caffeine intake, such as biologic markers, to further validate its accuracy.

Chlorogenic acid (CGA) is the major polyphenolic constituents of coffee; it increases the activity of the factor nuclear factor erythroid 2-related factor 2 (Nrf2), inhibits the action of NFκB (directly and indirectly) and activates sirtuin-1 (SIRT-1). Moreover, chlorogenic acid modulates a number of important biochemical pathways involved in inflammation (6).

To date, there are no specific biomarkers that evaluate the activity of coffee on health. However, based on large studies, it seems that coffee may exert some direct effects related to the reduction of inflammatory biomarkers, a phenomenon confirmed in most of the systematic reviews and meta-analyses available. It also seems that one of the main mechanisms of the anti-inflammatory effect of coffee is mediated by the increase in the concentration of adiponectin (7).

Interesting perspectives may be further elucidated from the analysis of coffee metabolites in the microbiota. These metabolites may be used as biomarkers of efficacy and absorption (8, 9).

It is important to note that validated questionnaires should be culturally and linguistically appropriate as well as easily understandable by the intended population.

The study by Bulczak and Chmurzyńska (1) was conducted between 2019 and 2020 and represents the reality in the Polish population just prior to and in the early stages of the COVID-19 pandemic. However, the data collected during the pandemic have highlighted profound changes in eating habits worldwide. This effect has been reported in several studies in which changes in diet and intake of single foods have been largely attributed to a response to stress (10–14). Several studies show that the COVID-19 pandemic has had a disproportionate economic impact for women and lower-income groups; women have suffered greatly from the economic crisis induced by the closure of productive activities and also because they are very active in tourism and in the manufacturing industry (15, 16).

In our studies performed during the pandemic phase, we found an increase in caffeine consumption attributable in young people to an increase in energy drinks and in adults to an increase in coffee consumption (10, 17, 18).

In a previous manuscript, we analyzed the effects of the pandemic on coffee consumption and caffeine intake in women, noting that there was an increase in coffee consumption that was associated with an increase in smoking (19).

The association between the consumption of coffee and use of cigarettes has been known for some time and the increase of these two substances easily occurs simultaneously (20).

In a very interesting experiment, Papke and coworkers evaluated the influence of coffee on receptor response to nicotine. The experimental model used oocytes that were surgically removed from mature female *Xenopus laevis* frogs (20). Smoking populations with high-sensitivity (HS) forms of $\alpha 4\beta 2$ nAChR receptors are limited in response to high concentrations of nicotine. Upon waking, after the previous day's nicotine has been extensively metabolized, the HS receptors will be primed for the day's first dose of the drug. The delivery of 1-methylpyridinium (n-MP) in breakfast coffee may then tune that response to the first cigarettes of the day, decreasing the response of upregulated HS receptors and perhaps increasing the activity of low-sensitivity receptors that will respond to the full dose of nicotine. These effects on the receptors exert an action on the brain; however, it is not known whether they can also influence the effects of the beneficial substances contained in coffee, in particular the antioxidants such as phenolic compounds (chlorogenic acids, cafestol, kahweol).

The relationship between coffee and smoking is complex and is strongly influenced by the type of coffee that is consumed (as previously mentioned, the method of preparation modifies the composition of the ingredients that filter into the drink) and by the quantity of cigarettes smoked.

More research is needed to understand whether the positive effects associated with coffee consumption are mitigated with cigarette smoking. Several studies have correlated the beneficial effects of coffee consumption on cardiovascular diseases and cardiovascular risk factors (6). A meta-analysis including more than 12,000,000 participants concluded that there was a nonlinear protective association between long-term coffee consumption and cardiovascular events. Specifically, compared to non-habitual coffee drinkers, the relative risk of cardiovascular disease is 0.85 (95% confidence interval, 0.80–0.90) for a median of 3.5 cups consumed per day (21). Similar data have been recently reported in the US population (22)

Cicero and coworkers performed a subanalysis of the Brisighella Heart Study (Italy) and compared central and peripheral blood pressure (BP) values in a subcohort of 720 men (47.9%) and 783 women (52.1%) reporting the drinking of different amounts of coffee each day (21). They found that regular coffee drinking is associated with lower systolic BP, peripheral pulse pressure (PP), aortic BP, and aortic PP, but with similar arterial stiffness (23). It is quite difficult to compare these results with other European cohorts due to

the different dietary pattern in Northern Italy compared to that in other countries, including a different method of coffee preparation and consumption and a different amount and biovariability of bioactive peptides and polyphenols included in daily consumed foods (3, 23, 24).

Coffee and caffeine are important dietary components with a strong influence on health. Increasing knowledge about these foods is important for understanding the mechanisms of action and giving more precise indications in personalized prevention. In this view, the article by Bulczak and Chmurzyńska (1) is relevant to increase our knowledge of the habits of different countries to proceed with targeted social awareness campaigns.

Disclosure statement

No potential conflict of interest was reported by the authors.

ORCID

Anna Vittoria Mattioli  <http://orcid.org/0000-0003-1487-9530>

References

- Bulczak EM, Chmurzyńska AU. Caffeine consumption in polish adults: development and validation of a polish questionnaire for assessing caffeine intake. *J Am Nutr Assoc.* 2023;1–7. Epub ahead of print. PMID: 36725370 doi:10.1080/27697061.2023.2172749.
- Mattioli AV. Effects of caffeine and coffee consumption on cardiovascular disease and risk factors. *Future Cardiol.* 2007;3(2):203–12. doi:10.2217/14796678.3.2.203.
- Rothschild JA, Kilding AE, Plews DJ. Pre-exercise nutrition habits and beliefs of endurance athletes vary by sex, competitive level, and diet. *J Am Coll Nutr.* 2021;40(6):517–28. Epub 2020 Sep 14. PMID: 32926647. doi:10.1080/07315724.2020.1795950.
- Mattioli AV, Pennella S, Farinetti A, Manenti A. Energy drinks and atrial fibrillation in young adults. *Clin Nutr.* 2018;37(3):1073–4. doi:10.1016/j.clnu.2017.05.002.
- Román GC, Jackson RE, Gadhia R, Román AN, Reis J. Mediterranean diet: the role of long-chain ω -3 fatty acids in fish; polyphenols in fruits, vegetables, cereals, coffee, tea, cacao and wine; probiotics and vitamins in prevention of stroke, age-related cognitive decline, and Alzheimer disease. *Rev Neurol (Paris).* 2019;175(10):724–41. Epub 2019 Sep 11. doi:10.1016/j.neuro.2019.08.005.
- Surma S, Sahebkar A, Banach M. Coffee or tea: anti-inflammatory properties in the context of atherosclerotic cardiovascular disease prevention. *Pharmacol Res.* 2023;187:106596. doi:10.1016/j.phrs.2022.106596.
- Linden-Torres E, Zambrano-Galván G, Sahebkar A, Ríos-Mier M, Simental-Mendía LE. Coffee consumption has no effect on circulating markers of liver function but increases adiponectin concentrations: a systematic review and meta-analysis of randomized controlled trials. *Nutr Res.* 2022;106:24–34. doi:10.1016/j.nutres.2022.07.010.
- Hegde S, Shi DW, Johnson JC, Geesala R, Zhang K, Lin YM, Shi XZ. Mechanistic study of coffee effects on gut microbiota and motility in rats. *Nutrients.* 2022;14(22):4877. doi:10.3390/nu14224877.
- Lawrence G, Midtved I, Samuelsen SO, Kristoffersen AK, Enersen M, Håheim LL. The blood microbiome and its association to cardiovascular disease mortality: case-cohort study. *BMC Cardiovasc Disord.* 2022;22(1):344. doi:10.1186/s12872-022-02791-7.

- 227 10. Coppi F, Nasi M, Farinetti A, Manenti A, Gallina S, Mattioli AV. Physical activity, sedentary behaviour, and diet in menopausal
228 women: comparison between COVID19 “first wave” and “second
229 wave” of pandemic in Italy. *Progress in Nutrition*. 2021;23(2):11755.
230 doi:10.23751/pn.v23i2.11755.
- 231 11. Beato AF, da Costa LP, Nogueira R. “Everything is gonna be al-
232 right with me”: the role of self-compassion, affect, and coping in
233 negative emotional symptoms during coronavirus quarantine.
234 *IJERPH*. 2021;18(4):2017. doi:10.3390/ijerph18042017.
- 235 12. Al-Musharaf S, Aljuraiban G, Bogis R, Alnafisah R, Aldhwayan
236 M, Tahrani A. Lifestyle changes associated with COVID-19 quar-
237 antine among young Saudi women: a prospective study. *PLoS*
238 *One*. 2021;16(4):e0250625. doi:10.1371/journal.pone.0250625.
- 239 13. Coppi F, Farinetti A, Stefanelli C, Mattioli AV. Changes in food
240 during the COVID-19 pandemic: the different roles of stress and
241 depression in women and men. *Nutrition*. 2023;108:111981.
242 doi:10.1016/j.nut.2023.111981.
- 243 14. Giannini DT, Tavares CM, Takey M, Aloise MLR, da Costa AJ, de
244 Carvalho DS, da Silva SC, Pontes MHP, Monteiro CB. Adolescents
245 emotional state and behavioral and dietary habit changes during
246 isolation due to the COVID-19 pandemic. *J Am Nutr Assoc*.
247 2022;41(4):415–23. doi:10.1080/07315724.2021.1897899.
- 248 15. Mattioli AV, Coppi F, Nasi M, Gallina S. Stress and cardiovascu-
249 lar risk burden after the pandemic: current status and future
250 prospects. *Expert Rev Cardiovasc Ther*. 2022;20(7):507–13. doi:10
251 .1080/14779072.2022.2092097.
- 252 16. Vogel B, Acevedo M, Appelman Y, Bairey Merz CN, Chieffo A,
253 Figtree GA, Guerrero M, Kunadian V, Lam CSP, Maas AHEM,
254 et al. The lancet women and cardiovascular disease commission:
255 reducing the global burden by 2030. *Lancet*. 2021;397(10292):2385–
256 438. doi:10.1016/S0140-6736(21)00684-.
- 257 17. Mattioli AV, Sabatini S. Changes in energy drink consumption
258 during the COVID-19 quarantine. *Clin Nutr ESPEN*. 2021;45:516–
259 7. doi:10.1016/j.clnesp.2021.06.034.
- 260 18. Madhyamapurush W. Analysis on factors affecting tourist in-
261 volvement in coffee tourism after the COVID-19 pandemic in
262 Thailand. *F1000Res*. 2022;11:1269. doi:10.12688/f1000re-
263 search.123759.2.
- 264 19. Coppi F, Migaldi M, Stefanelli C, Farinetti A, Mattioli AV. Changes
265 in coffee and caffeine intake during the pandemic in
266 women smokers and non-smokers: a future challenge for cardio-
267 vascular prevention. *Acta Biomed*. 2023;94(2):e2023114.
268 doi:10.23750/abm.v94i2.14118.
- 269 20. Papke RL, Karaffa M, Horenstein NA, Stokes C. Coffee and cig-
270 arettes: modulation of high and low sensitivity α 4 β 2 nicotinic
271 acetylcholine receptors by n-MP, a biomarker of coffee consump-
272 tion. *Neuropharmacology*. 2022;216:109173. doi:10.1016/j.neuro-
273 pharm.2022.109173.
- 274 21. Ding M, Bhupathiraju SN, Satija A, van Dam RM, Hu FB. Long-term
275 coffee consumption and risk of cardiovascular disease:
276 a systematic review and a dose-response meta-analysis of pro-
277 spective cohort studies. *Circulation*. 2014;129(6):643–59.
278 doi:10.1161/CIRCULATIONAHA.113.005925.
- 279 22. Di Maso M, Boffetta P, Negri E, La Vecchia C, Bravi F. Caffeinated
280 coffee consumption and health outcomes in the us population: a
281 dose-response meta-analysis and estimation of disease cases and
282 deaths avoided. *Adv Nutr*. 2021;12(4):1160–76. doi:10.1093/ad-
283 vances/nmaa177.
- 284 23. Cicero AFG, Fogacci F, D’Addato S, Grandi E, Rizzoli E, Borghi
285 C ; on behalf of the Brisighella Heart Study. Self-reported coffee
286 consumption and central and peripheral blood pressure in the
287 cohort of the brisighella heart study. *Nutrients*. 2023;15(2):312.
288 doi:10.3390/nu15020312.
- 289 24. Mattioli AV, Migaldi M, Farinetti A. Coffee in hypertensive wom-
290 en with asymptomatic peripheral arterial disease: a potential nu-
291 traceutical effect. *J Cardiovasc Med (Hagerstown)*. 2018;19(4):183–
292 5. doi:10.2459/JCM.0000000000000626.
- 293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344