

# Real-world safety and effectiveness of zanubrutinib vs ibrutinib in CLL: the CLL-ZANU2024 Italian cohort

Enrica Antonia Martino,<sup>1</sup> Annalisa Pitino,<sup>2</sup> Ernesto Vigna,<sup>1</sup> Raffaella Pasquale,<sup>3</sup> Isacco Ferrarini,<sup>4</sup> Riccardo Moia,<sup>5</sup> Andrea Visentin,<sup>6</sup> Alessandro Sanna,<sup>7</sup> Marina Motta,<sup>8</sup> Massimo Moratti,<sup>3</sup> Paolo Sportoletti,<sup>9</sup> Annalisa Chiarenza,<sup>10</sup> Alessandro Maggi,<sup>11</sup> Valentina Zammit,<sup>12</sup> Michele Merli,<sup>13</sup> Idanna Innocenti,<sup>14</sup> Claudia Giordano,<sup>15</sup> Laura Nocilli,<sup>16</sup> Massimiliano Postorino,<sup>17</sup> Caterina Stelitano,<sup>18</sup> Andrea Ferrario,<sup>19</sup> Anna Maria Frustaci,<sup>20</sup> Marcello Riva,<sup>21</sup> Sara Pepe,<sup>22</sup> Adalberto Ibatici,<sup>23</sup> Stefania Scardino,<sup>24</sup> Paola Anticoli Borza,<sup>25</sup> Laura Ballotta,<sup>26</sup> Salvatrice Mancuso,<sup>27</sup> Francesco Malaspina,<sup>28</sup> Anna Mele,<sup>29</sup> Sara Galimberti,<sup>30</sup> Gioacchino Catania,<sup>31</sup> Annamaria Giordano,<sup>32</sup> Ilaria Angeletti,<sup>33</sup> Luana Schiattone,<sup>34</sup> Elsa Pennese,<sup>34</sup> Rosanna Miccolis,<sup>35</sup> Angelo Fama,<sup>36</sup> Giulio Giordano,<sup>37</sup> Catello Califano,<sup>38</sup> Antonella Bruzzese,<sup>1</sup> Santino Caserta,<sup>1</sup> Giuliana Farina,<sup>39</sup> Pietro Bulian,<sup>40</sup> Giacomo Loseto,<sup>41</sup> Barbara Pocali,<sup>42</sup> Vanessa Innao,<sup>43</sup> Piero Galieni,<sup>44</sup> Vincenzo Fraticelli,<sup>45</sup> Candida Vitale,<sup>46,47</sup> Azzurra Romeo,<sup>48</sup> Marco Rossi,<sup>49,50</sup> Ilaria Scortechini,<sup>51</sup> Federico Vozella,<sup>52</sup> Luigi Malandrucolo,<sup>53</sup> Marzia Varettoni,<sup>54</sup> Lucia Morello,<sup>55</sup> Giuseppe Pietrantonio,<sup>56</sup> Esmeralda Conte,<sup>57</sup> Martina Cantelli,<sup>58</sup> Roberta Murru,<sup>59</sup> Daniele Caracciolo,<sup>50</sup> Enrico Derenzini,<sup>60</sup> Valentina Di Martina,<sup>61</sup> Roberto Marasca,<sup>62</sup> Maria Ilaria Del Principe,<sup>17</sup> Amalia Figuera,<sup>10</sup> Francesco Angotzi,<sup>6</sup> Marta Coscia,<sup>19</sup> Nicola Di Renzo,<sup>24</sup> Luca Laurenti,<sup>14</sup> Nicola Amodio,<sup>50</sup> Pellegrino Musto,<sup>32,63</sup> Francesco Di Raimondo,<sup>10</sup> Arcangelo Liso,<sup>33</sup> Alessandra Tedeschi,<sup>20</sup> Livio Trentin,<sup>6</sup> Gianluca Gaidano,<sup>5</sup> Francesca Romana Mauro,<sup>22</sup> Giovanni Tripepi,<sup>64</sup> Fortunato Morabito,<sup>65</sup> Valter Gattel,<sup>40,\*</sup> and Massimo Gentile<sup>1,66,\*</sup>

<sup>1</sup>Hematology Unit, Department of Onco-hematology, Azienda Ospedaliera Annunziata, Cosenza, Italy; <sup>2</sup>Institute of Clinical Physiology, National Research Council, Rome, Italy; <sup>3</sup>Azienda Sanitaria Universitaria Friuli Centrale, SOC Clinica Ematologia, Udine, Italy; <sup>4</sup>Department of Engineering for Innovation Medicine, University of Verona, Verona, Italy; <sup>5</sup>Division of Hematology, Department of Translational Medicine, Università del Piemonte Orientale, Novara, Italy; <sup>6</sup>Hematology Unit, Department of Medicine, University of Padova, Padova, Italy; <sup>7</sup>Hematology Unit, Department of Oncology, Azienda Ospedaliero Universitaria Careggi, Florence, Italy; <sup>8</sup>Department of Hematology, Azienda Socio-Sanitaria Territoriale Spedali Civili di Brescia, Lombardy, Italy; <sup>9</sup>Centro di Ricerca Emato-Oncologica, Dipartimento di Medicina e Chirurgia Università di Perugia, Perugia, Italy; <sup>10</sup>Division of Hematology, Azienda Policlinico-S. Marco, University of Catania, Catania, Italy; <sup>11</sup>Haematology, Ospedale G. Moscati, Taranto, Italy; <sup>12</sup>Oncoematology and Trapianto di Midollo Osseo Unit, Oncology Department Ospedale La Maddalena Palermo, Palermo, Italy; <sup>13</sup>Hematology, Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Ca' Granda, Ospedale Maggiore Policlinico, Milan, Italy; <sup>14</sup>Dipartimento di Scienze di Laboratorio ed Ematologiche, Area Ematologia, Fondazione Policlinico Universitario A. Gemelli Istituto di Ricovero e Cura a Carattere Scientifico, Università Cattolica del Sacro Cuore, Rome, Italy; <sup>15</sup>Department of Clinical Medicine and Surgery, Federico II University Medical School, Naples, Italy; <sup>16</sup>Department of Hematology, Azienda Ospedaliera Papardo, Messina, Italy; <sup>17</sup>Hematology Department of Biomedicine and Prevention, Tor Vergata University of Rome, Rome, Italy; <sup>18</sup>Department of Hematology, Azienda Ospedaliera Bianchi Melacrino Morelli, Reggio Calabria, Italy; <sup>19</sup>Department of Medicine and Surgery, University of Insubria and Department of Oncology, Azienda Socio-Sanitaria Territoriale Sette Laghi, Ospedale di Circolo, Varese, Italy; <sup>20</sup>Department of Haematology, Niguarda Cancer Center, Azienda Socio-Sanitaria Territoriale Grande Ospedale Metropolitano Niguarda, Milan, Italy; <sup>21</sup>Hematology Department, Ospedale San Bortolo di Vicenza, Vicenza, Italy; <sup>22</sup>Hematology, Department of Translational and Precision Medicine, Sapienza University, Rome, Italy; <sup>23</sup>Hematology and Cellular Therapy Unit, Istituto di Ricovero e Cura a Carattere Scientifico Ospedale Policlinico San Martino, Genoa, Italy; <sup>24</sup>Hematology and Stem Cell Transplant Unit, "Vito Fazzi" Hospital, Lecce, Italy; <sup>25</sup>Unità Operativa Complessa Ematologia, Azienda Ospedaliera San Giovanni-Addolorata, Rome, Italy; <sup>26</sup>Dipartimento Clinico di Scienze Mediche, Chirurgiche e della Salute, Università degli Studi di Trieste, Trieste, Italy; <sup>27</sup>Department of Health Promotion, Mother and Child Care, Internal Medicine and Medical Specialties, Hematology Unit, University of Palermo, Palermo, Italy; <sup>28</sup>Hematology Unit, Istituto di Ricovero e Cura a Carattere Scientifico, IRST Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori "Dino Amadori," Meldola, Italy; <sup>29</sup>Haematology, Ospedale Cardinale Panico, Lecce, Italy; <sup>30</sup>Division of Hematology, University Hospital of Pisa, Pisa, Italy; <sup>31</sup>Struttura Complessa a Direzione Universitaria Ematologia-Azienda Ospedaliero-Universitaria di Sassari Antonio e Biagio e C. Arrigo, Alessandria, Italy; <sup>32</sup>Unit of Hematology and Stem Cell Transplantation, Azienda Ospedaliero-Universitaria Careggi Policlinico, Bari, Italy; <sup>33</sup>Department of Medicine and Surgery, University of Perugia and Hematology/Oncology Unit, Terni Hospital, Terni, Italy; <sup>34</sup>Dipartimento di Ematologia Clinica Ospedale civile S. Spirito Pescara, Pescara, Italy; <sup>35</sup>Department of Hematology, Azienda Sanitaria Locale-Barletta Andria Trani, Barletta, Italy; <sup>36</sup>Hematology Unit, Department of Internal Medicine, Ospedale G. Mazzini, Teramo, Italy; <sup>37</sup>Unità Operativa Complessa, Medicina Servizio e Ambulatorio di Ematologia Ospedale di Riferimento Regionale Antonio Cardarelli, Campobasso, Italy; <sup>38</sup>Onco-Hematology Unit, "A. Tortora" Hospital, Pagani, Italy; <sup>39</sup>Hematology, Hospital "Sant'Anna e San Sebastiano," Caserta, Italy; <sup>40</sup>Clinical and Experimental Onco-Hematology Unit, Centro di Riferimento Oncologico di Aviano, Istituto di Ricovero e Cura a Carattere Scientifico, Aviano, Italy; <sup>41</sup>Hematology and Cell Therapy Unit, Istituto di Ricovero e Cura a Carattere Scientifico Istituto Tumori "Giovanni Paolo II," Bari, Italy; <sup>42</sup>Division of Hematology and Stem Cell Transplantation Unit, Cardarelli Hospital, Naples, Italy; <sup>43</sup>Hematology Unit, Azienda Ospedaliera di Rilievo Nazionale e di Alta Specializzazione Garibaldi, Catania, Italy; <sup>44</sup>Department of Haematology and Stem Cell Transplantation Unit, C. e G. Mazzoni Hospital, Ascoli Piceno, Italy; <sup>45</sup>Department of Hematology, Gemelli Molise, Campobasso, Italy; <sup>46</sup>Division of Hematology, Azienda Ospedaliero Universitaria Città della Salute e della Scienza di Torino, Turin, Italy; <sup>47</sup>Department of Molecular Biotechnology and Health Sciences, University of Turin, Turin, Italy; <sup>48</sup>Hematology Unit, Ospedale Santa Maria Goretti, Latina, Italy; <sup>49</sup>Department of Hematology-Oncology, Azienda Universitaria Ospedaliera Renato Dulbecco, Catanzaro, Italy; <sup>50</sup>Department of Experimental and Clinical Medicine Magna Grecia University, Catanzaro, Italy; <sup>51</sup>Azienda Ospedaliero Universitaria delle Marche, Ancona, Italy; <sup>52</sup>Hematology, Stem Cell Transplantation, Fondazione Policlinico Universitario Campus Bio medico di Roma, Rome, Italy; <sup>53</sup>Hematology Unit Ospedale "Fabrizio Spaziani," Frosinone, Italy; <sup>54</sup>Division of Hematology, Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo di Pavia, Pavia, Italy; <sup>55</sup>Unità Operativa di Ematologia, Ospedale Guglielmo da Saliceto, Piacenza, Italy; <sup>56</sup>Istituto di Ricovero e Cura a Carattere Scientifico Oncology Referral Cancer Center of Basilicata, Rionero in Vulture, Italy; <sup>57</sup>Haematology Unit,

Submitted 15 October 2025; accepted 18 November 2025; prepublished online on *Blood Advances* First Edition 7 January 2026; final version published online 27 February 2026. <https://doi.org/10.1182/bloodadvances.20250118757>.

\*V.G. and M.G. contributed equally as senior authors.

The data that support the findings of this study are available from the corresponding author, Massimo Gentile ([massimo.gentile@unicat.it](mailto:massimo.gentile@unicat.it)), on request. The data are not publicly available because of privacy or ethical restrictions.

The full-text version of this article contains a data supplement.

© 2026 American Society of Hematology. Published by Elsevier Inc. Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

## Key Points

- In a large real-world Italian cohort, zanubrutinib showed significantly lower treatment discontinuation rates compared with ibrutinib.
- Zanubrutinib-treated patients experienced fewer off-target AEs supporting its preferential use than ibrutinib in CLL with comorbidities.

Bruton tyrosine kinase inhibitors (BTKis) have dramatically changed the therapeutic landscape of chronic lymphocytic leukemia (CLL), with ibrutinib, first-in-class, demonstrating durable efficacy even in high-risk patients. However, off-target adverse events (AEs) have raised concerns, prompting the development of more selective second-generation BTKis, such as zanubrutinib, designed to improve tolerability while maintaining efficacy. Despite encouraging results from clinical trials, real-world data comparing zanubrutinib with ibrutinib remain limited. In this multicenter, retrospective study, we analyzed 934 patients with CLL treated outside clinical trials, including 393 receiving zanubrutinib and 541 receiving ibrutinib. We evaluated time to treatment discontinuation (TTD) and time to next treatment or death (TTNTD) in both the overall cohort and a propensity score-matched population. Patients who were treated with zanubrutinib experienced lower 12-month discontinuation rates (overall: 12.6% vs 21.4%; matched: 12.4% vs 20.2%) and higher 12-month TTNTD rates (overall: 91.9% vs 83.0%; matched: 93.2% vs 83.4%). Multivariable analyses confirmed zanubrutinib as an independent predictor of longer TTD and TTNTD, whereas high-risk features, including age, relapsed/refractory disease, Binet stage C, *TP53* disruption, Eastern Cooperative Oncology Group 2 to 3, and congestive heart failure, were consistently associated with poorer outcomes. AEs leading to discontinuation, particularly atrial fibrillation, bleeding, and infections, were less frequent with zanubrutinib, reflecting its favorable safety profile. These findings provide real-world evidence that zanubrutinib offers more durable disease control and improved persistence compared with ibrutinib, reinforcing its clinical value as a preferred second-generation BTKi. Nevertheless, the relatively short follow-up for zanubrutinib warrants cautious interpretation of long-term outcomes, and underscores the need for ongoing observation to fully characterize its durability and safety.

## Introduction

Bruton tyrosine kinase inhibitors (BTKis) have significantly improved outcomes of patients with chronic lymphocytic leukemia (CLL) and small lymphocytic leukemia (SLL).<sup>1</sup> Ibrutinib, the first-in-class BTKi to receive regulatory approval, represented a major advance in the treatment landscape of CLL/SLL.<sup>2</sup> Its efficacy has been demonstrated in seminal studies both as monotherapy and in combination with anti-CD20 monoclonal antibodies, in treatment-naïve (TN) and relapsed/refractory (R/R) settings.<sup>2-9</sup> Notably, ibrutinib has shown durable efficacy even in patients with high-risk cytogenetic abnormalities, such as del17p and TP53 mutations, subgroups typically resistant to conventional chemoimmunotherapy.<sup>10,11</sup> However, long-term follow-up data from major clinical trials and real-world cohorts have consistently reported high rates of treatment discontinuation, primarily due to off-target toxicities.<sup>9</sup>

Second-generation BTKis have been developed with greater selectivity for BTK, aiming to reduce off-target adverse events (AEs).<sup>12</sup>

Among these, zanubrutinib, which is a highly selective BTKi, was approved by the European Medicines Agency in November 2022 for the treatment of CLL,<sup>13</sup> and subsequently received reimbursement approval by the Agenzia Italiana del Farmaco in October 2023 for use as monotherapy.<sup>14</sup> These approvals were based on the results of 2 pivotal phase 3 trials: SEQUOIA (NCT03336333)<sup>15</sup> and ALPINE (NCT03734016).<sup>16</sup> In addition to ibrutinib and zanubrutinib, acalabrutinib is also an approved BTKi for CLL/SLL.<sup>17,18</sup>

The SEQUOIA trial evaluated zanubrutinib in patients with CLL/SLL who were TN. After a median follow-up of 61.2 months, zanubrutinib demonstrated a significant improvement in progression-free survival (PFS) compared with chemoimmunotherapy (median PFS not reached vs 44.1 months), irrespective of immunoglobulin heavy chain variable (*IGHV*) mutational status. Similarly, favorable outcome was observed in the cohort of patients with del(17p), with both median PFS and overall survival not reached.<sup>15</sup>

The ALPINE trial, a randomized phase 3 study in R/R CLL/SLL, compared zanubrutinib with ibrutinib. Zanubrutinib showed a

**Table 1. Baseline characteristics of the patients with CLL treated with zanubrutinib or ibrutinib**

	Zanubrutinib (N = 393)	Ibrutinib (N = 541)	P value	SMD
Median age (range), y	76 (44-94)	70 (37-88)	<b>&lt;.001</b>	0.53
<b>Sex, n (%)</b>				
Male	237 (60.3)	353 (65)		
Female	156 (39.7)	188 (35)	.122	0.10
<b>Line of therapy, n (%)</b>				
TN	239 (60.7)	221 (41)		
R/R	154 (39.3)	320 (59)	<b>&lt;.001</b>	-0.41
<b>Binet stage, n (%)</b>				
A	41 (10.4)	65 (12)		
B	151 (38.4)	237 (44)		
C	201 (51.1)	239 (44)	.109	0.13
<b>IGHV mutational status, n (%)</b>				
Mutated	103 (26.2)	181 (33)		
Germ line	242 (61.6)	360 (67)		
Unknown	48 (12.2)	-	.26	0.08
<b>TP53 disruption, n (%)*</b>				
Absent	254 (72.5)	269 (49.7)	<b>&lt;.001</b>	-0.32
Present	98 (18.3)	206 (38.1)		
Unknown	36 (9.2)	66 (12.2)		
<b>ECOG-PS, n (%)</b>				
0-1	290 (74)	399 (74)		
2-3	103 (26)	142 (26)	1	0.001
<b>Cardiovascular risk factor, n (%)</b>				
Hypertension	155 (39.4)	213 (39)	.98	0.001
Atrial fibrillation	51 (13)	70 (13)	.98	0.001
Hypercholesterolemia	60 (15.3)	83 (15.3)	.97	-0.002
Congestive heart failure	24 (6.1)	33 (6)	.99	0.06
Peripheral arterial disease	12 (3.1)	17 (3)	.93	0.03
Cerebrovascular disease	13 (3.3)	18 (3.1)	.98	0.03
Diabetes	67 (17)	92 (17.1)	.98	0.001
Myocardial infarction	48 (12.2)	66 (12)	.98	0.0004

Boldface values indicate statistically significant results.

SMD, standardized mean difference.

\*TP53 disruption was determined by the documentation of 17p deletion and/or TP53 mutation by fluorescence in situ hybridization and sequencing, respectively; present, del17p positive and/or TP53 mutated; absent, del17p negative and TP53 negative.

higher overall response rate (85.6% vs 75.4%) and superior PFS, with a 32% relative reduction in the risk of progression or death. In high-risk patients harboring del(17p) and/or TP53 mutations, zanubrutinib was associated with a 49% improvement in PFS compared with ibrutinib.<sup>16</sup> Both trials reported a favorable safety profile for zanubrutinib, including lower rates of treatment discontinuation due to AEs compared with control arms.<sup>15,16</sup>

Despite these results, real-world evidence on the use of zanubrutinib outside of clinical trials remains limited, particularly in comparison with first-generation BTKis, such as ibrutinib.<sup>19-21</sup>


To address this gap, we conducted a retrospective, multicenter analysis of patients with CLL treated with zanubrutinib in routine clinical practice. Additionally, we performed a comparative evaluation of outcomes in patients treated with zanubrutinib vs ibrutinib,

focusing on time to treatment discontinuation (TTD) and time to next treatment or death (TTNTD), both in the overall cohort and in a propensity score-matched population.

## Materials and methods

### Patients

We retrospectively analyzed CLL databases from 52 Italian centers, including all consecutive patients with TN or R/R CLL who received zanubrutinib or ibrutinib outside of a clinical trial. Data sets contained comprehensive clinical and laboratory information, including age, sex, date of diagnosis, Rai and Binet stage, laboratory parameters, biological markers (interphase fluorescence in situ hybridization analyses for the short arm of chromosome 17 -del(17p)-, mutational status of the *IGHV*, and *TP53* mutations),



**Figure 1. Kaplan-Meier curves of TTD for all 934 patients with CLL treated with zanubrutinib or ibrutinib.** (A) Kaplan-Meier curve of TTD according to treatment arm in the original cohorts. (B) Kaplan-Meier curve of TTD according to treatment arm in the propensity score-matched cohorts.

Eastern Cooperative Oncology Group-performance status (ECOG-PS), comorbidities, treatment history, and date of last follow-up or death. All data were abstracted from clinical records at the time of treatment initiation and updated prospectively. The assessment of *IGHV* status and *TP53* abnormalities (including deletion and/or mutation) was performed locally in certified laboratories adhering to European Research Initiative on CLL, as previously described.<sup>22,23</sup> Treatment-emergent AEs were collected and graded according to the Common Terminology Criteria for Adverse Events, version 5.0.

This research is part of the CLL-ZANU2024 study, an Italian, observational, multicenter, longitudinal secondary data analysis evaluating the safety and effectiveness of zanubrutinib in the real-world post-marketing setting. This observational comparative study used registry data, with retrospective data collection for ibrutinib (from June 2013 to June 2019; these data have been partially published<sup>22-32</sup>) and ambispective (retrospective/prospective) data collection for zanubrutinib (from October 2023 onward).

**Table 2. Univariate and multivariate analyses for TTD**

	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P value	HR (95% CI)	P value
Age*	1.01 (0.99-1.02)	.280	1.01 (0.99-1.03)	.250
<b>Sex</b>				
Male vs female	0.96 (0.72-1.28)	.771	0.92 (0.66-1.26)	.596
<b>Therapy</b>				
Zanubrutinib vs ibrutinib	0.49 (0.36-0.69)	<b>.000</b>	0.45 (0.31-0.65)	<b>.000</b>
<b>Line of therapy</b>				
R/R vs TN	1.56 (1.17-2.07)	<b>.002</b>	1.33 (0.98-1.81)	.071
<b>Binet stage</b>				
B vs A	1.41 (0.82-2.41)	.215	1.31 (0.76-2.28)	.334
C vs A	1.85 (1.1-3.13)	.022	1.85 (1.08-3.16)	<b>.024</b>
<b>IGHV mutational status</b>				
Germ line vs mutated	1.04 (0.77-1.4)	.816	–	
<b>TP53 disruption†</b>				
Present vs absent	1.67 (1.24-2.24)	<b>.001</b>	1.57 (1.16-2.12)	<b>.003</b>
<b>ECOG-PS</b>				
2-3 vs 0-1	1.56 (1.17-2.08)	<b>.003</b>	1.52 (1.11-2.07)	<b>.009</b>
<b>Cardiovascular risk factor</b>				
Hypertension	1.19 (0.9-1.57)	.214	–	
Atrial fibrillation	1.03 (0.68-1.56)	.884	–	
Hypercholesterolemia	0.93 (0.62-1.38)	.717	–	
Congestive heart failure	1.82 (1.16-2.87)	<b>.009</b>	1.7 (1.06-2.72)	<b>.028</b>
Peripheral arterial disease	0.86 (0.38-1.93)	.710	–	
Cerebrovascular disease	0.91 (0.41-2.06)	.828	–	
Diabetes	0.91 (0.62-1.33)	.629	–	
Myocardial infarction	0.89 (0.57-1.37)	.582	–	

Boldface values indicate statistically significant results.

\*The effect corresponds to 1 year of increase in age.

†TP53 disruption was determined by the documentation of 17p deletion and/or TP53 mutation by fluorescence in situ hybridization and sequencing, respectively; present, del17p positive and/or TP53 mutated; absent, del17p negative and TP53 negative.

## Statistical analysis

Continuous variables were summarized by medians and ranges, whereas categorical variables were reported as absolute frequencies and percentages, as appropriate.


The main end point was TTD, defined as the interval from treatment initiation to permanent discontinuation. Treatment discontinuation was defined as the permanent cessation of the BTKi for any reason, including AEs, disease progression, transformation to Richter syndrome, patient decision, or death. A discontinuation event required a treatment gap of >30 consecutive days without BTKi reinitiation. Temporary interruptions of treatment (eg, for intercurrent illness, planned procedures, or toxicity) lasting ≤30 days were not classified as discontinuation. Dose reductions or physician-directed treatment holds were considered treatment modifications rather than discontinuations, provided that therapy was resumed within 30 days. Discontinuation events were not centrally adjudicated; however, participating centers applied standardized criteria aligned with real-world clinical practice and International Workshop on Chronic Lymphocytic Leukemia recommendations. All AEs resulting in definitive treatment discontinuation were classified as grade >3 and irreversible.

TTNTD was defined as the time from treatment initiation to the earlier of 2 events: the start of a subsequent therapy or death. Patients not receiving a new line of therapy were censored at their last known clinical activity. All subsequent therapies were given according to International Workshop on Chronic Lymphocytic Leukemia criteria.<sup>33</sup> Reasons for treatment discontinuation were documented from clinical charts for a subset of patients.

Follow-up was truncated at 24 months to minimize differences in observation time between treatment groups, as patients treated with ibrutinib had longer follow-up periods than those treated with zanubrutinib. Patients lost to follow-up were not excluded, but treated as censored observations in the survival analysis.

Univariable Cox regression models were used to assess the association between covariates and TTD or TTNTD. Covariates included treatment type (zanubrutinib vs ibrutinib), age, gender, Binet stage, *IGHV* mutational status, *TP53* disruption [del(17p) and/or *TP53* mutation], line of therapy, β2-microglobulin, lactate dehydrogenase, ECOG-PS, and cardiovascular risk factors.

To reduce treatment-selection bias (or confounding by indication), a propensity score model was developed using multivariable



**Figure 2. Kaplan-Meier curves of TTNTD for all 934 patients with CLL treated with zanubrutinib or ibrutinib.** (A) Kaplan-Meier curve of TTNTD according to treatment arm in the original cohorts. (B) Kaplan-Meier curve of TTNTD according to treatment arm in the propensity score-matched cohorts.

logistic regression, incorporating covariates with a standardized mean difference  $\geq 0.1$  between treatment groups. This method is considered more robust than relying on  $P$  value thresholds, which can be overly sensitive to sample size. The model also included an interaction term between age and the number of previous lines of therapy. The dependent variable was the treatment received (zanubrutinib vs ibrutinib), and the resulting predicted probabilities were used as propensity scores. Matching was performed using a caliper width of 0.02 standard deviations of the logit of the propensity score, in a 1:1 ratio (1 ibrutinib-treated control for each

zanubrutinib-treated patient). No weighting was applied as matching was 1-to-1; all patients in the matched cohort received a weight of 1.

Univariable and multivariable Cox regression models were fitted on the matched cohort to identify factors associated with TTD and TTNTD. In all multivariable models, we always forced age and sex. Variables significantly associated with the outcomes on univariable analyses were included in multivariable models. Hazard ratios (HRs), 95% confidence intervals (CIs), and  $P$ -values were

**Table 3. Univariate and multivariate analyses for TTNTD**

	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P value	HR (95% CI)	P value
Age*	1.01 (0.99-1.02)	.364	1.01 (0.99-1.03)	.189
<b>Sex</b>				
Male vs female	0.96 (0.7-1.33)	.806	0.89 (0.62-1.27)	.509
<b>Therapy</b>				
Zanubrutinib vs ibrutinib	0.42 (0.29-0.62)	<b>.000</b>	0.35 (0.22-0.56)	<b>.000</b>
<b>Line of therapy</b>				
R/R vs TN	1.76 (1.28-2.43)	<b>.001</b>	1.46 (1.03-2.06)	<b>.033</b>
<b>Binet stage</b>				
B vs A	1.33 (0.73-2.43)	.347	1.28 (0.69-2.38)	.426
C vs A	1.91 (1.07-3.42)	<b>.029</b>	2.03 (1.12-3.66)	<b>.019</b>
<b>IGHV mutational status</b>				
Germ line vs mutated	1 (0.72-1.4)	.985	–	
<b>TP53 disruption†</b>				
Present vs absent	1.87 (1.35-2.59)	<b>.000</b>	1.75 (1.25-2.44)	<b>.001</b>
<b>ECOG-PS</b>				
2-3 vs 0-1	1.39 (1-1.93)	.047	1.35 (0.95-1.92)	.094
<b>Cardiovascular risk factor</b>				
Hypertension	1.19 (0.87-1.61)	.282	–	
Atrial fibrillation	1.03 (0.65-1.63)	.892	–	
Hypercholesterolemia	0.95 (0.61-1.48)	.824	–	
Congestive heart failure	2.14 (1.34-3.42)	.001	1.91 (1.17-3.11)	<b>.009</b>
Peripheral arterial disease	0.87 (0.36-2.11)	.750	–	
Cerebrovascular disease	1.16 (0.51-2.63)	.717	–	
Diabetes	0.92 (0.6-1.41)	.704	–	
Myocardial infarction	0.74 (0.44-1.25)	.260	–	

Boldface values indicate statistically significant results.

\*The effect corresponds to 1 year of increase in age.

†TP53 disruption was determined by the documentation of 17p deletion and/or TP53 mutation by fluorescence in situ hybridization and sequencing, respectively; present, del17p positive and/or TP53 mutated; absent, del17p negative and TP53 negative.

reported for all Cox models. All statistical analyses were performed using R version 4.5.0 (Copyright [C] 2025 The R Foundation for Statistical Computing: package survival, survminer, TableOne, cobalt, MatchIt). Additional analyses were conducted using IBM SPSS Statistics, version 29.

All participating centers obtained ethics approval, and the study was conducted in accordance with the Declaration of Helsinki. Written informed consent has been obtained from all patients.

## Results

### Study population and baseline characteristics

A total of 934 patients with CLL who initiated treatment with either zanubrutinib or ibrutinib met the inclusion criteria and were included in this analysis. Of these, 541 (57.9%) received ibrutinib, and 393 (42.1%) received zanubrutinib (Table 1). The median age was 76 years (range, 44-94) in patients of the zanubrutinib cohort, and 70 years (range, 37-88) in those of the ibrutinib cohort (Table 1). The 2 cohorts were generally balanced in terms of sex,

IGHV mutational status, ECOG-PS, and cardiovascular risk factors. Beyond age, they also differed as for line of therapy, Binet stage, and TP53 disruption status. Specifically, the ibrutinib group had higher rates of patients who were R/R (59% vs 39.3%) and of TP53 disruption (38.1% vs 18.3%) than the zanubrutinib group.

### TTD

Over a median follow-up of 13 months for the zanubrutinib cohort and 18 months for the ibrutinib cohort, the median TTD was not reached in both groups (Figure 1A). At 12 months, the discontinuation rate was significantly lower for zanubrutinib (12.6%) than for ibrutinib (21.4%;  $P < .001$ ; Figure 1A).

On univariable analysis, the following variables were significantly associated with shorter TTD: >1 previous line of therapy (HR, 1.56; 95% CI, 1.17-2.07;  $P = .002$ ), Binet stage C (HR, 1.85; 95% CI, 1.10-3.13;  $P = .02$ ), TP53 disruption (HR, 1.67; 95% CI, 1.24-2.24;  $P = .001$ ), ECOG-PS 2 to 3 (HR, 1.56; 95% CI, 1.17-2.08;  $P = .003$ ), and congestive heart failure (HR, 1.82; 95% CI, 1.16-2.87;  $P = .009$ ; Table 2).

**Table 4. Reasons for zanubrutinib or ibrutinib discontinuation in patients with CLL**

	Zanubrutinib	Ibrutinib
Total patients, N	393	541
Patients who discontinued treatment, n (%)	49 (12.5)	155 (28.6)
Median follow-up, mo	13	18
<b>Reasons for discontinuation</b>		
Toxic effect of therapy	14 (3.6)	71 (13.1)
Arthralgia/myalgia/arthritis	0	21 (3.9)
Atrial fibrillation	1 (0.3)	11 (2.0)
Bleeding episodes	5 (1.3)	16 (2.9)
Rash	1 (0.3)	11 (2.0)
Infection	6 (1.6)	4 (0.7)
Diarrhea	0	8 (1.5)
Hepatitis B virus reactivation	1 (0.3)	0
Disease progression		
CLL	16 (4.1)	63 (11.6)
Richter syndrome	1 (0.3)	24 (4.4)
Second neoplasia	3 (0.8)	4 (0.7)
Patient request	1 (0.3)	0
Other	1 (0.3)	1 (0.2)
Non-CLL-related death	6 (1.5)	4 (0.7)
Death from unknown causes	7 (1.8)	12 (2.2)

On multivariable analysis, treatment with zanubrutinib was independently associated with significantly longer TTD (HR, 0.45; 95% CI, 0.31-0.65;  $P < .001$ ; Table 2). In the same model, Binet stage C (HR, 1.85; 95% CI, 1.08-3.16;  $P = .02$ ), TP53 disruption (HR, 1.57; 95% CI, 1.16-2.12;  $P = .003$ ), ECOG-PS 2 to 3 (HR, 1.52; 95% CI, 1.11-2.07;  $P = .009$ ), and congestive heart failure (HR, 1.70; 95% CI, 1.06-2.72;  $P = .028$ ) were independently associated with shorter TTD.

### TTNTD

The median TTNTD was not reached in either cohort (Figure 2A). At 12 months, TTNTD estimates were significantly improved in the zanubrutinib cohort (91.9%) compared with the ibrutinib cohort (83.0%;  $P < .001$ ; Figure 2A). Univariable analysis identified the following variables as significantly associated with shorter TTNTD:  $>1$  previous line of therapy (HR, 1.76; 95% CI, 1.28-2.43;  $P = .001$ ), Binet stage C (HR, 1.91; 95% CI, 1.07-3.42;  $P = .029$ ), TP53 disruption (HR, 1.87; 95% CI, 1.35-2.59;  $P < .001$ ), ECOG-PS 2 to 3 (HR, 1.39; 95% CI, 1.00-1.93;  $P = .047$ ), and congestive heart failure (HR, 2.14; 95% CI, 1.34-3.42;  $P = .001$ ; Table 3). Conversely, zanubrutinib treatment was significantly associated with a reduced risk of TTNTD (HR, 0.42; 95% CI, 0.29-0.62;  $P < .001$ ). In multivariable analysis,  $>1$  previous line of therapy (HR, 1.46; 95% CI, 1.03-2.06;  $P = .033$ ), Binet stage C (HR, 2.03; 95% CI, 1.12-3.66;  $P = .019$ ), TP53 disruption (HR, 1.75; 95% CI, 1.25-2.44;  $P = .001$ ), and congestive heart failure (HR, 1.91; 95% CI, 1.17-3.11;  $P = .009$ ) were independently associated with lower TTNTD, whereas zanubrutinib therapy was associated with a significantly improved TTNTD (HR, 0.35; 95% CI, 0.22-0.56;  $P < .001$ ; Table 3).

**Table 5. Baseline characteristics of the patients with CLL treated with zanubrutinib or ibrutinib after propensity score matching**

	Zanubrutinib (N = 270)	Ibrutinib (N = 270)	SMD
Mean age (SD), y	72.1 (9.81)	71.2 (9.16)	0.095
<b>Sex, n (%)</b>			
Male	173 (64.1)	168 (62.2)	
Female	97 (35.9)	102 (37.8)	-0.018
<b>Line of therapy, n (%)</b>			
1L	146 (54.1)	145 (53.7)	
2L+	124 (45.9)	125 (46.3)	-0.0037
<b>Binet stage, n (%)</b>			
A	30 (10.4)	37 (12)	-0.0259
B	106 (38.4)	113 (44)	-0.0259
C	134 (51.1)	120 (44)	0.0519
<b>IGHV mutational status, n (%)</b>			
Mutated	80 (29.6)	88 (32.6)	
Germ line	190 (70.4)	182 (67.4)	0.0296
<b>TP53 disruption, n (%)*</b>			
Absent	183 (67.8)	183 (67.8)	
Present	87 (32.2)	87 (32.2)	0.0000
<b>ECOG-PS, n (%)</b>			
0-1	212 (78.5)	197 (73)	
2-3	58 (21.5)	73 (27)	-0.0556
<b>Cardiovascular risk factor, n (%)</b>			
Hypertension	106 (39.3)	109 (40.4)	-0.0111
Atrial fibrillation	35 (13)	35 (13)	0.0000
Hypercholesterolemia	36 (13.3)	47 (17.4)	-0.0407
Congestive heart failure	13 (4.8)	17 (6.3)	-0.015
Peripheral arterial disease	10 (3.7)	10 (3.7)	0.00001
Cerebrovascular disease	8 (3)	9 (3.3)	-0.0037
Diabetes	40 (14.8)	46 (17)	-0.0222
Myocardial infarction	34 (12.6)	33 (12.2)	0.0037

\*TP53 disruption was determined by the documentation of 17p deletion and/or TP53 mutation by FISH and sequencing, respectively; present=del17p positive and/or TP53 mutated; absent=del17p negative and TP53 negative. 1L, first line; 2L+, 2 or more prior lines of therapy; SMD, standardized mean difference.

### Reasons for discontinuation

Overall, 12.5% of patients (n = 49) in the zanubrutinib cohort and 28.6% of patients (n = 155) in the ibrutinib cohort discontinued treatment during the study period (Table 4).

In both cohorts, toxicity (14/49 patients [28.6%] for zanubrutinib; 71/155 patients [46.3%] for ibrutinib) and disease progression (17/49 patients [34.7%] for zanubrutinib; 63/155 patients [41%] for ibrutinib) were the most frequently reported reasons for discontinuation. Discontinuation due to disease progression was more common in the ibrutinib group (Table 4). Supplemental Table 1 reports the most common grade 3 to 4 treatment-emergent AEs observed in both cohorts. The incidence of AEs was generally comparable between the 2 treatment groups.

**Table 6. Univariate and multivariate analyses for TTD after propensity score matching**

	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P value	HR (95% CI)	P value
Age*	1.03 (1.01-1.06)	<b>.008</b>	1.02 (1-1.05)	<b>.028</b>
<b>Sex</b>				
Male vs female	0.98 (0.65-1.48)	.942	0.99 (0.65-1.51)	.947
<b>Therapy</b>				
Zanubrutinib vs ibrutinib	0.49 (0.33-0.74)	<b>.001</b>	0.51 (0.35-0.76)	<b>.001</b>
<b>Line of therapy</b>				
2L+ vs 1L	1.51 (1.03-2.22)	<b>.035</b>	1.44 (0.96-2.15)	.079
<b>Binet stage</b>				
B vs A	1.27 (0.64-2.52)	.499		
C vs A	1.69 (0.87-3.28)	.121		
<b>IGHV mutational status</b>				
Germ line vs mutated	1.22 (0.79-1.89)	.362		
<b>TP53 disruption†</b>				
Present vs absent	1.44 (0.98-2.11)	.063		
<b>ECOG-PS</b>				
2-3 vs 0-1	2.05 (1.39-3.04)	<b>.000</b>	1.74 (1.15-2.63)	<b>.009</b>
<b>Cardiovascular risk factor</b>				
Hypertension	1.38 (0.94-2.03)	.100	–	
Atrial fibrillation	1.19 (0.69-2.06)	.523	–	
Hypercholesterolemia	0.76 (0.4-1.42)	.383	–	
Congestive heart failure	3.4 (1.96-5.88)	<b>.000</b>	2.46 (1.3-4.64)	<b>.006</b>
Peripheral arterial disease	1.24 (0.47-3.23)	.666	–	
Cerebrovascular disease	1.43 (0.49-4.13)	.514	–	
Diabetes	1.02 (0.6-1.72)	.947	–	
Myocardial infarction	1.07 (0.62-1.85)	.800	–	

Boldface values indicate statistically significant results.

1L, first line; 2L+, 2 or more prior lines of therapy.

\*The effect corresponds to 1 year of increase in age.

†TP53 disruption was determined by the documentation of 17p deletion and/or TP53 mutation by fluorescence in situ hybridization and sequencing, respectively; present, del17p positive and/or TP53 mutated; absent, del17p negative and TP53 negative.

However, arthralgia, diarrhea, and atrial fibrillation occurred significantly more frequently in patients receiving ibrutinib than in those treated with zanubrutinib.

### Propensity score-matching analysis

After 1:1 propensity score matching, 2 well-balanced cohorts of 270 patients each were selected (Table 5). With a median follow-up of 13 months for zanubrutinib and 16 months for ibrutinib, the 12-month discontinuation rate was significantly lower in the zanubrutinib group (12.4%) than in the ibrutinib group (20.2%;  $P < .001$ ; Figure 1B).

On univariable analysis, age (HR, 1.03; 95% CI, 1.01-1.06;  $P = .008$ ), >1 previous line of therapy (HR, 1.51; 95% CI, 1.03-2.22;  $P = .035$ ), ECOG-PS 2 to 3 (HR, 2.0; 95% CI, 1.39-3.04;  $P < .001$ ), and congestive heart failure (HR, 2.46; 95% CI, 1.30-4.64;  $P = .006$ ) were independently associated with an increased risk of discontinuing therapy. Conversely, zanubrutinib treatment was significantly associated with a reduced risk of discontinuation (HR, 0.49; 95% CI, 0.33-0.74;  $P < .001$ ; Table 6).

On multivariable analysis, age (HR, 1.02; 95% CI, 1.01-1.05;  $P = .028$ ), ECOG-PS 2 to 3 (HR, 1.74; 95% CI, 1.15-2.63;  $P = .009$ ), and congestive heart failure (HR, 2.46; 95% CI, 1.30-4.64;  $P = .006$ ) were independently associated with an increased risk of discontinuing therapy. Conversely, zanubrutinib treatment was significantly associated with a reduced risk of discontinuation (HR, 0.51; 95% CI, 0.35-0.76;  $P = .001$ ; Table 6).

As for TTNTD, the 12-month estimates were again improved for the zanubrutinib (93.2%) vs the ibrutinib (83.4%) cohorts ( $P < .001$ ; Figure 2B).

In univariable analysis, age (HR, 1.03; 95% CI, 1.01-1.06;  $P = .039$ ), >1 previous line of therapy (HR, 1.78; 95% CI, 1.12-2.82;  $P = .015$ ), TP53 disruption (HR, 1.63; 95% CI, 1.06-2.51;  $P = .026$ ), ECOG PS (HR, 1.74; 95% CI, 1.12-2.70;  $P = .014$ ), and congestive heart failure (HR, 3.53; 95% CI, 1.97-6.33;  $P < .001$ ) were independently associated with inferior TTNTD, whereas zanubrutinib therapy was independently associated with significantly improved TTNTD (HR, 0.37; 95% CI, 0.22-0.60;  $P < .001$ ; Table 7).



and congestive heart failure as negative prognostic factors for TTNTD, whereas zanubrutinib remained independently associated with improved outcomes. These benefits were observed in both first-line and R/R settings, although the shorter follow-up in the zanubrutinib cohort warrants cautious interpretation.

Given the higher prevalence of unfavorable features in the ibrutinib group, propensity score matching was used to mitigate imbalance. Although residual confounding cannot be excluded, age and congestive heart failure resulted in consistently poorer TTD and TTNTD, and zanubrutinib retained its favorable association across all models. ECOG-PS 2 to 3 negatively affected TTD, whereas *TP53* disruption and R/R disease affected TTNTD. Notably, in the ibrutinib setting, *TP53* disruption influenced outcomes primarily when mutation and 17p deletion co-occurred;<sup>24,38</sup> whether this applies to second-generation BTKis warrants further study. Other biomarkers, including CD49d expression, unavailable in our cohort but prognostically relevant for ibrutinib and acalabrutinib,<sup>39-41</sup> could also be explored for patients who were treated with zanubrutinib.

Two smaller Chinese retrospective studies reported favorable outcomes with zanubrutinib, but lacked comparators and had limited follow-up.<sup>20,21</sup>

A recent US abstract by Hou et al<sup>19</sup> described >3000 patients treated with ibrutinib, acalabrutinib, or zanubrutinib, and showed that zanubrutinib had the most favorable cardiovascular safety profile, and highest 6- and 12-month treatment persistence. Median TTNTD was not reached for zanubrutinib, compared with 30.2 and 35.8 months for ibrutinib and acalabrutinib, respectively, consistent with our findings.

Complementary real-world evidence from the Flatiron Health database compared acalabrutinib and ibrutinib.<sup>42</sup> Acalabrutinib showed longer therapy duration, primarily due to improved tolerability, in line with the ELEVATE-RR trial.

Direct comparisons between acalabrutinib and zanubrutinib remain lacking, but early observational data suggest potential advantages for zanubrutinib in cardiovascular safety and treatment continuity. Overall, real-world data, including ours, support the clinical transition toward second-generation BTKis, particularly for patients at high risk of treatment-related toxicity.

Several limitations of our study should be acknowledged. The follow-up for the zanubrutinib cohort was necessarily shorter due to later regulatory approval, potentially limiting long-term outcome assessment, although early divergence of survival curves supports robustness. Given the retrospective nature of data collection, reliable grading of AEs according to Common Terminology Criteria for Adverse Events was feasible mainly for laboratory-based parameters, whereas subjective or clinically reported events may have been underestimated. Treatment allocation was non-randomized, and may have been influenced by physician or institutional preferences. Temporal differences existed between cohorts (ibrutinib, June 2013–June 2019; zanubrutinib, October 2023 onward), potentially introducing residual confounding related to the treatment era. To reduce bias from unequal observation periods, follow-up was truncated at 24 months. Although this harmonization mitigates differences, residual effects cannot be entirely excluded. Despite these limitations, propensity score

matching, multivariable adjustment, and the multicenter design strengthen the generalizability of our findings.

In conclusion, this study provides the first real-world comparative effectiveness analysis of zanubrutinib vs ibrutinib in CLL, confirming superior tolerability and treatment durability of zanubrutinib. Patients treated with zanubrutinib experienced fewer off-target AEs, particularly atrial fibrillation, diarrhea, and arthralgia, supporting its preferential use in patients with CLL with comorbidities or a higher risk of treatment-related toxicity. When contextualized with emerging real-world data, including acalabrutinib, our findings reinforce the clinical transition toward second-generation BTKis in routine practice, zanubrutinib appears particularly well suited for patients with a high comorbidity burden or increased risk of AEs. Direct head-to-head comparisons with acalabrutinib will be critical to further refine BTKi selection and optimize outcomes.

## Acknowledgment

This study was supported by the Piano Nazionale di Ripresa e Resilienza (project code MAD-2022-12375673).

## Authorship

Contribution: All authors contributed equally to this study.

Conflict-of-interest disclosure: R. Moia reports participation in advisory boards and speaker bureaus for AbbVie, Johnson & Johnson, BeOne, and AstraZeneca. G. Gaidano reports participation in advisory boards for, and speaker's bureau honoraria from AbbVie, AstraZeneca, BeOne, Hikma, Incyte, Johnson & Johnson, and Lilly. The remaining authors declare no competing financial interests.

ORCID profiles: E.A.M., 0000-0002-8504-3231; A.P., 0000-0002-7768-3925; R.P., 0000-0003-2253-8883; I.F., 0000-0001-9867-8335; A.V., 0000-0003-0271-7200; A.M.F., 0000-0003-2587-7901; A.I., 0009-0001-7363-8684; F. Malaspina, 0000-0003-3041-9606; S.G., 0000-0001-7899-5966; G.C., 0000-0002-1743-9525; E.P., 0009-0004-8041-6424; A. Fama, 0000-0003-1214-0193; G. Giordano, 0000-0002-6932-5028; S.C., 0000-0002-0698-1781; C.V., 0000-0002-2592-8724; M.R., 0000-0002-7258-475X; G.P., 0000-0002-1717-8337; E.C., 0000-0003-4957-2045; R. Murru, 0000-0001-9792-9542; E.D., 0000-0002-7154-8140; R. Marasca, 0000-0002-6431-6878; M.I.D.P., 0000-0002-3958-0669; A. Figuera, 0000-0002-3863-9884; F.A., 0000-0001-7835-2573; M.C., 0000-0003-2123-7675; L.L., 0000-0002-8327-1396; P.M., 0000-0003-3277-6594; L.T., 0000-0003-1222-6149; G. Gaidano, 0000-0002-4681-0151; G.T., 0000-0002-3580-4602; F. Morabito, 0000-0002-2585-7073; M.G., 0000-0002-5256-0726.

Correspondence: Massimo Gentile, Hematology Unit, AO of Cosenza, viale della Repubblica snc, 87100 Cosenza, Italy; email: [massimo.gentile@unical.it](mailto:massimo.gentile@unical.it); Ernesto Vigna, Hematology Unit, AO of Cosenza, viale della Repubblica snc, 87100 Cosenza, Italy; email: [ernesto.vigna@aocs.it](mailto:ernesto.vigna@aocs.it); and Valter Gattei, Clinical and Experimental Onco-Hematology Unit, Centro di Riferimento Oncologico di Aviano, Istituto di Ricovero e Cura a Carattere Scientifico, Via Franco Gallini 2, 33081 Aviano, Italy; email: [vgattei@cro.it](mailto:vgattei@cro.it).

## References

1. Eichhorst B, Robak T, Montserrat E, et al. Chronic lymphocytic leukaemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2021;32(1):23-33.
2. Byrd JC, Furman RR, Coutre SE, et al. Targeting BTK with ibrutinib in relapsed chronic lymphocytic leukemia. *N Engl J Med*. 2013;369(1):32-42.
3. Byrd JC, Brown JR, O'Brien S, et al; RESONATE Investigators. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med*. 2014;371(3):213-223.
4. Burger JA, Tedeschi A, Barr PM, et al; RESONATE-2 Investigators. Ibrutinib as initial therapy for patients with chronic lymphocytic leukemia. *N Engl J Med*. 2015;373(25):2425-2437.
5. Shanafelt TD, Wang XV, Kay NE, et al. Ibrutinib-rituximab or chemoimmunotherapy for chronic lymphocytic leukemia. *N Engl J Med*. 2019;381(5):432-443.
6. Woyach JA, Ruppert AS, Heerema NA, et al. Ibrutinib regimens versus chemoimmunotherapy in older patients with untreated CLL. *N Engl J Med*. 2018;379(26):2517-2528.
7. Moreno C, Greil R, Demirkan F, et al. Ibrutinib plus obinutuzumab versus chlorambucil plus obinutuzumab in first-line treatment of CLL (iLLUMINATE): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol*. 2019;20(1):43-56.
8. Munir T, Brown JR, O'Brien S, et al. Final analysis from RESONATE: up to six years of follow-up on ibrutinib in previously treated CLL or SLL. *Am J Hematol*. 2019;94(12):1353-1363.
9. Barr PM, Owen C, Robak T, et al. Up to 8-year follow-up from RESONATE-2: first-line ibrutinib treatment for patients with CLL. *Blood Adv*. 2022;6(11):3440-3450.
10. O'Brien S, Furman RR, Coutre S, et al. Single-agent ibrutinib in treatment-naïve and relapsed/refractory CLL: a 5-year experience. *Blood*. 2018;131(17):1910-1919.
11. Strati P, Keating MJ, O'Brien SM, et al. Outcomes of first-line treatment for CLL with 17p deletion. *Haematologica*. 2014;99(8):1350-1355.
12. Tam C, Thompson PA. BTK inhibitors in CLL: second-generation drugs and beyond. *Blood Adv*. 2024;8(9):2300-2309.
13. European Medicines Agency. Brukinsa (zanubrutinib) EPAR. Accessed 25 March 2025. <https://www.ema.europa.eu/en/medicines/human/EPAR/bruksina>
14. Agenzia Italiana del Farmaco. Determina 640-2023 Brukinsa. Accessed 25 March 2025. [https://www.aifa.gov.it/documents/20142/961234/Determina\\_640-2023\\_Bruksina.pdf](https://www.aifa.gov.it/documents/20142/961234/Determina_640-2023_Bruksina.pdf)
15. Shadman M, Munir T, Robak T, et al. Zanubrutinib versus bendamustine and rituximab in treatment-naïve CLL/SLL: median 5-year follow-up of SEQUOIA. *J Clin Oncol*. 2025;43(7):780-787.
16. Brown JR, Eichhorst B, Lamanna N, et al. Sustained benefit of zanubrutinib vs ibrutinib in patients with R/R CLL/SLL: final comparative analysis of ALPINE. *Blood*. 2024;144(26):2706-2717.
17. Byrd JC, Wierda WG, Schuh A, et al. Acalabrutinib monotherapy in patients with relapsed/refractory chronic lymphocytic leukemia: updated phase 2 results. *Blood*. 2020;135(15):1204-1213.
18. Martino EA, Bruzzese A, Vigna E, et al. Acalabrutinib in chronic lymphocytic leukemia. *Expert Opin Pharmacother*. 2023;24(5):545-549.
19. Hou J, Blanc S, Maglente GA, et al. Real-world BTKi treatment patterns and outcomes among patients with CLL/SLL in US community oncology practices. *J Clin Oncol*. 2024;42(16 suppl):e19020.
20. Zhu H, Xia Y, Miao Y, et al. A real-world study of zanubrutinib treatment in CLL/SLL. *Blood*. 2023;142(suppl 1):6554.
21. Luo J, Zhang J, Liu L, et al. Real-world outcomes of zanubrutinib monotherapy in CLL: a multicenter retrospective study. *Blood*. 2023;142(suppl 1):6540.
22. Bomben R, Zucchetto A, Laureana R, et al. CD49d expression is included in a revised 4-factor model predicting outcome in CLL patients treated with ibrutinib: a multicenter real-world experience. *Hemasphere*. 2024;8(7):e128.
23. Martino EA, Mauro FR, Reda G, et al. Ibrutinib as first-line therapy in CLL patients over 80 years old: a retrospective real-life multicenter Italian cohort. *Hematol Oncol*. 2024;42(1):e3249.
24. Bomben R, Rossi FM, Vit F, et al. Clinical impact of TP53 disruption in CLL patients treated with ibrutinib: a campus CLL study. *Leukemia*. 2023;37(4):914-918.
25. Visentin A, Mauro FR, Catania G, et al. Obinutuzumab plus chlorambucil versus ibrutinib in previously untreated CLL patients without TP53 disruptions: a real-life CLL campus study. *Front Oncol*. 2022;12:1033413.
26. Levi S, Bronstein Y, Goldschmidt N, et al. Efficacy of front-line ibrutinib versus FCR in CLL: a retrospective multicenter real-world study. *Am J Hematol*. 2023;98(2):E24-E27.
27. Reda G, Mattiello V, Frustaci AM, et al. Ibrutinib in patients over 80 with CLL: a multicenter Italian cohort. *Blood Adv*. 2023;7(4):525-528.
28. Visentin A, Mauro FR, Cibien F, et al. Continuous treatment with ibrutinib in 100 untreated patients with TP53-disrupted CLL: a real-life campus CLL study. *Am J Hematol*. 2022;97(3):E95-E99.

29. Morabito F, Tripepi G, Del Poeta G, et al. Effectiveness of ibrutinib as first-line therapy for CLL patients and indirect comparison with rituximab-bendamustine outside clinical trials. *Am J Hematol*. 2021;96(8):E269-E272.
30. Morabito F, Tripepi G, Del Poeta G, et al. Assessment of the 4-factor score: retrospective analysis of 586 CLL patients receiving ibrutinib. *Am J Hematol*. 2021;96(5):E168-E171.
31. Morabito F, Tripepi G, Del Poeta G, et al. Comparison of ibrutinib and idelalisib plus rituximab in real-life relapsed/resistant CLL cases. *Eur J Haematol*. 2021;106(4):493-499.
32. Gentile M, Morabito F, Del Poeta G, et al. Survival risk score for real-life relapsed/refractory CLL patients receiving ibrutinib: a campus CLL study. *Leukemia*. 2021;35(1):235-238.
33. Hallek M, Cheson BD, Catovsky D, et al. iwCLL guidelines for diagnosis, indications for treatment, response assessment, and supportive management of CLL. *Blood*. 2018;131(25):2745-2760.
34. Mato AR, Nabhan C, Thompson MC, et al. Toxicities and outcomes of 616 ibrutinib-treated patients in the United States: a real-world analysis. *Haematologica*. 2018;103(5):874-879.
35. Hardy-Abeloos C, Pinotti R, Gabrilove J. Ibrutinib dose modifications in the management of CLL. *J Hematol Oncol*. 2020;13(1):66.
36. Ysebaert L, Aurran-Schleinitz T, Dartigeas C, et al. Real-world results of ibrutinib in relapsed/refractory CLL in France: early results on a large series of 428 patients. *Am J Hematol*. 2017;92(8):E166-E168.
37. UK CLL Forum. Ibrutinib for relapsed/refractory CLL: a UK and Ireland analysis of outcomes in 315 patients. *Haematologica*. 2016;101(12):1563-1572.
38. Morabito F, Del Poeta G, Mauro FR, et al. TP53 disruption as a risk factor in the era of targeted therapies: a multicenter retrospective study of 525 CLL cases. *Am J Hematol*. 2021;96(8):E306-E310.
39. Alsadhan A, Chen J, Gaglione EM, et al. CD49d expression identifies a biologically distinct subtype of CLL with inferior progression-free survival on BTK inhibitor therapy. *Clin Cancer Res*. 2023;29(18):3612-3621.
40. Tissino E, Pozzo F, Benedetti D, et al. CD49d promotes disease progression in CLL: new insights from CD49d bimodal expression. *Blood*. 2020;135(15):1244-1254.
41. Tissino E, Benedetti D, Herman SEM, et al. Functional and clinical relevance of VLA-4 (CD49d/CD29) in ibrutinib-treated CLL. *J Exp Med*. 2018;215(2):681-697.
42. Roeker LE, DerSarkissian M, Ryan K, et al. Real-world comparative effectiveness of acalabrutinib and ibrutinib in patients with CLL. *Blood Adv*. 2023;7(16):4291-4301.