

Clinical and experimental efficacy of gemtuzumab ozogamicin in core binding factor acute myeloid leukemia

Michele Gottardi,¹ Federico Mosna,¹ Sergio de Angeli,² Cristina Papayannidis,³ Anna Candoni,⁴ Marino Clavio,⁵ Cristina Tecchio,⁶ Andrea Piccin,⁷ Marta Campo dell'Orto,⁸ Fabio Benedetti,⁶ Giovanni Martinelli,³ Filippo Gherlinzoni¹

¹Department of Hematology, General Hospital, Treviso; ²Laboratory of Cell Cryopreservation, General Hospital, Treviso; ³Department of Hematology, Institute "L.A. Seragnoli," University of Bologna; ⁴Department of Hematology, University of Udine; ⁵Department of Hematology, University of Genoa; ⁶Department of Hematology, University of Verona; ⁷Department of Hematology, General Hospital, Bolzano; ⁸Department of Pathology, General Hospital, Treviso, Italy

Abstract

Leukemia-initiating cells of core binding factor (CBF) acute myeloid leukemia (AML) likely derive from early committed hematopoietic precursors expressing CD33. As such, targeting CD33 could ameliorate the chance of cure of CBF AML patients. We compared 12 CBF AML patients treated with Fludarabine, Cytarabine, Idarubicin and Gemtuzumab Ozogamicin (FLAI-GO regimen) with 25 CBF AML patients treated with the same schedule, but without GO. With the limit of small numbers, we observed a consistent trend toward better overall survival, disease free survival and event free survival in the FLAI-GO group. We also demonstrated the ability of GO to induce the disappearance *in vitro* of the AML1-ETO molecular transcript in a polymerase chain reaction-positive graft without decreasing the clonogenic potential of CD34+/CD38- cells. This represents the proof of principle for using GO in a purging strategy before autologous stem cell transplantation. Therefore, our data argue in favor of the reinstatement of GO in the therapy of CBF AML.

Brief Report

Core Binding Factor (CBF) Acute Myeloid Leukemia (AML) patients experience treatment failure in the order of 40-50%.¹ As such, efforts have been made to increase the dose intensity of first line treatment over the standard Daunorubicin/Cytarabine 3+7 induction, under the assumption that better survival could be achieved by obtaining a deeper early clearance of blasts before clonal evolution of the disease. Among other attempts, the addition of a third chemotherapeutic drug, such as Fludarabine, has been tested in various clinical trials,^{2,3} with conflicting results regarding Overall Survival (OS), but, in most cases, with an increase in the rate of complete remission (CR) and Disease-Free Survival (DFS).^{2,3} Moreover, novel agents with compatible safety profile have been tested in addition to chemotherapy; among these, the immunoconjugate Gemtuzumab Ozogamicin (GO), which combines in a single agent a monoclonal antibody targeting CD33 with the DNA damaging toxin calicheamicin. Following initial studies, GO has been combined to chemotherapy to improve efficacy, particularly in the case of CBF AML, in which blasts highly express the target antigen.^{1,3} Three-drug Fludarabine-based regimens combined with GO proved successful mainly in the setting of cytogenetically favorable, such as CBF AML, or intermediate-risk AML,^{3,4} whereas results have somehow been disappointing in adverse risk patients.^{3,4} Despite this, the clinical development of GO has suffered from concerns raised by an increased incidence of hepatotoxicity and veno-occlusive disease when GO was used at the dose of 9 mg/m² twice during induction,⁵ and from the early results of a randomized phase-3 trial that showed no advantage in OS and a significant increase of Treatment-Related Mortality (TRM) (5% vs 1%) in the GO-treated group.⁶ This ultimately led to the withdrawal of the drug from the market in 2010. Later studies^{3,7,8} showed, on the opposite, an unequivocal survival benefit by GO, even if mainly restricted to CBF AML, at the lower schedule of 3 to 6 mg/m², without neither increased hepatotoxicity nor higher TRM.^{3,7,8} This led to the reevaluation of GO, which, unfortunately, has not yet resulted in the reinstatement of the drug to clinical practice.⁵

In order to address the role of GO in the treatment of CBF AML, we retrospectively reviewed 12 CBF AML patients [t(8;21) n=8; inv(16) n=4] treated from 2006 to 2009 with the FLAI-GO regimen (Fludarabine 30

Correspondence: Michele Gottardi, Hematology, Department of Specialty Medicine, General Hospital "Santa Maria di Ca' Foncello", piazza Ospedale 1, 31100 Treviso, Italy.
Tel.: +39.0422.322221 - Fax: +39.0422.322542
E-mail: migottardi@ulss.tv.it

Key words: Core binding factor; acute myeloid leukemia; gemtuzumab ozogamicin; immunotherapy; autologous stem cell transplantation.

Acknowledgments: the authors are grateful to Leukemia-Lymphoma-Myeloma ONLUS Association (AIL), Section of Treviso (Italy), for the financial support.

Contributions: MG and FM designed and performed the research, analyzed data and wrote the manuscript; CP, AC, MC, CT, AP and FB collected data and performed research; SDA performed the clonogenic assays; MCD performed the molecular tests; FM performed the statistical analysis; GM and FG critically revised the manuscript; all authors approved the final manuscript.

Conflict of interest: the authors declare no potential conflict of interest.

Received for publication: 4 January 2017.
Revision received: 23 April 2017.
Accepted for publication: 27 April 2017.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright M. Gottardi et al., 2017
Licensee PAGEPress, Italy
Hematology Reports 2017; 9:7028
doi:10.4081/hr.2017.7028

mg/m² on days 1-5; Cytarabine 2 gr/m² on days 1-5; Idarubicin 10 mg/m² on days 1,3,5; GO 3 mg/m² on day 6),⁴ and consolidated with two high-dose cytarabine (HiDAC)-based cycles (overall dose 24 gr/m²/cycle). We applied a regimen that was previously described in an independent series of AML patients.⁴ Patients with c-KIT tyrosine kinase domain mutation at codon 816 (TKD⁸¹⁶) at diagnosis (n=2) or with the persistence of molecular transcript, assessed by sequential polymerase chain reaction (PCR), at the end of consolidation (n=5), *i.e.* Minimal Residual Disease (MRD)-positivity, were then consolidated with either allogeneic or autologous hematopoietic stem cell transplantation (HSCT) based on the availability of a donor. We decided to further intensify the treatment of KIT mutated patients early on, based on initial studies showing an adverse prognosis of these patients as compared to KIT wild-type CBF

AML.⁹ Conversely, the persistence of the molecular transcript at the end of consolidation was considered a predictor of adverse prognosis based on previous experiences,^{10,11} as well as our own unpublished data.

We compared this group with 25 CBF AML patients [t(8;21) n=13; inv(16) n=12] treated according to the same criteria and with the same schedule, but without GO, in the years 2003-2006 and 2010-2013. The two groups were comparable in all clinical and laboratory features (Table 1). In the latter group, autologous HSCT was performed in 5 patients because of MRD-positivity, and allogeneic HSCT was

performed in 2 patients because of the lack of cytogenetic response after induction therapy.

Patients in the two groups reached a comparable rate of CR after induction (n=12/12 vs 22/25, P=0.540); no death in induction was observed, and all patients completed their therapeutic schedule. At median follow-up of 69.2 months, 3 patients in the FLAI-GO group relapsed at 8, 14 and 42 months after the achievement of CR; of these, two out of three achieved second CR following rescue therapy and underwent allogeneic HSCT. Conversely, at median DFS 14.7 months, 11 patients of the FLAI group relapsed; among these, 7 out of 10 achieved second CR and were consolidated with allogeneic HSCT.

As shown in Figure 1, we observed a consistent, yet non-statistical trend towards better OS, DFS and, most importantly in evaluating the efficacy of first line therapy, Event Free Survival (EFS) in the FLAI-GO group (5-yrs OS 69.4% vs 48.6%, P=0.202; 5-yrs DFS 54.7% vs 42.4%, P=0.327; 5-yrs EFS 54.7% vs 36.9%, P=0.136; Figure 1). Besides, patients tended to relapse later when treated with FLAI-GO (median DFS: unreached vs 14.7 months; Figure 1). We believe these differences did not reach the statistical significance mainly due to small

numbers. The achievement of MRD-negativity (FLAI-GO=100% of the 4 patients analyzed; FLAI=63% of the 13 patient analyzed), assessed by PCR, was of pivotal prognostic importance (P<0.001 for either OS, DFS and EFS).

The potential of GO in the treatment of CBF AML is based on strong biological bases: first of all, most t(8;21) AML blasts do not express the transporter P-glycoprotein (Pgp),¹² i.e. the Multi-Drug Resistance 1 (MDR1) gene product, and this seems related to a selective repression of the promoter of MDR1 by AML1-ETO.¹³ Several studies have pointed out how GO extrusion by Pgp affects clinical response.¹⁴ Moreover, CBF fusion transcripts promote leukemogenesis by inducing the initial expansion of a preleukemic myeloid cell compartment predisposed to secondary mutations and characterized by CD33+ early committed myeloid precursors incorporating the AML1-ETO or CBFβ-MYH11 transcripts.¹⁵ According to this model, founding Leukemia-Initiating Cells of CBF AML, differently from other types of AML, would arise from these early committed myeloid precursors rather than the more immature Hematopoietic Stem Cells,¹⁵ and therefore be more sensitive to GO therapy because of their markedly higher expression




Figure 1. Survival by treatment with Gemtuzumab Ozogamicin (GO). Patients receiving FLAI-GO induction therapy showed a consistent, yet nonstatistical, trend towards better Overall Survival (OS), Disease-Free Survival (DFS) and, most importantly in determining the effect of first line treatment, Event-Free Survival (EFS), as compared to a group of patients treated with FLAI. Numbers of the two groups are showed as n; only patients achieving complete remission after induction therapy were considered in determining DFS. Inclusion criteria and treatment schedule was the same in the two groups apart from the addition of GO at 3 mg/m² to induction therapy at day 6 of the FLAI-GO regimen.

Table 1. Baseline patient characteristics.

	FLAI5	My-FLAI5	P
Median age	41.3 (18-66)	46.3 (29-67)	0.2602
Sex	12 M + 13 F	6 M + 6 F	0.909
Secondary acute myeloid leukemia	0	0	NA
Hepatomegaly	4	2	1.0
Splenomegaly	3	2	1.0
Sarcoma	0	1	1.0
Hemoglobin gr/dL	8.4 (4.2-11)	8.5 (5-13.6)	0.9100
White blood cells ×10 ³ /L	19.0 (1.6-95)	18.7 (4.5-45.5)	0.9706
N ×10 ³ /L	1.86 (0.33-6.35)	2.58 (0.45-11.36)	0.3680
Mo ×10 ³ /L	0.95 (0.01-2.56)	0.58 (0.01-2.68)	0.1330
Ly ×10 ³ /L	2.67 (0.50-6.80)	2.86 (0.60-7.73)	0.8016
Blasts ×10 ³ /L	10.9 (0.01-65.55)	11.6 (0.22-32.0)	0.9074
Platelets ×10 ³ /L	60.66 (8-255)	73.72 (6-531)	0.7150
Elevated LDH	18	10	0.638
DIC	2	2	0.305
Acute renal failure	0	1	0.314
t(8;21)/inv(16)	13/12	8/4	0.491
FLT3-ITD	3	2	1.0
NPM1 mutated	0	0	NA
KIT TKD ⁸¹⁶ mutated	1	1	1.0
Packed BM (>80%)	12	5	0.717
Additional cytogenetic abnormalities	None: 14 pts; 1: 7 pts; 2: 3 pts; 3: 1 pts	None: 4; 1: 5 pts; 2: 1 pts; 3: 2 pts	0.387

of CD33. The distinctive chemosensitivity shown by CBF AML¹ could also be explained by these biological differences.¹⁵

As such, autologous HSCT could be used to increase the dose intensity of first line therapy in selected patients, improving OS, as shown by some studies.¹⁶ Results are best when MRD-negativity in the bone marrow and in the products of leukapheresis is obtained before transplantation.¹⁷ Recently, a 5-year EFS of 93% was reported in a small series of CBF AML patients undergoing autologous HSCT with a PCR-negative graft, with the disappearance of CBF transcripts after HSCT in 8 out of 10 previously MRD-positive patients.¹⁸ Nonetheless, in a previous series we observed that clinical outcome of patients could be improved also when autologous HSCT had been performed with CD34⁺ grafts with persistent molecular transcripts.¹⁶ We explain this finding by the higher overall dose-intensity achieved by first line treatment by including autologous HSCT.

The achievement of MRD-negativity has

been related to a better prognosis by many trials.¹⁹ Autologous HSCT, performed at the end of first line treatment, might be beneficial for selected CBF AML patients to achieve the disappearance of MRD, especially when performed using MRD-negative Peripheral Blood Stem Cells (PBSC).¹⁷ Monoclonal antibodies are thus being tested to provide *in vivo* purging of PBSC. This approach resembles the use of Rituximab in the therapy of patients affected by CD20⁺ lymphomas and undergoing autologous HSCT. Concerns are there, though, that treatment with GO might affect hematopoietic reconstitution by reducing the number of hematopoietic long-term repopulating cells collected by leukapheresis.

We therefore evaluated the clonogenic growth of PBSC collected at the end of consolidation therapy from five patients with CBF AML. Briefly, mononuclear cells were cultured in RPMI medium in the presence or absence of GO at a concentration of 5 µg/mL for two hours, a previously *in vitro* dose

proved able to induce the near-complete saturation of the CD33 antigen sites.²⁰ Cells were then collected, GO removed by centrifugation, and cells reseeded in Petri dishes containing semisolid MethoCult GF-H4434 medium. The number of Colony-Forming Units (CFU) was determined after 14 days incubation.

Using unsorted cells, we observed a significant decrease in the number of CFU-GEMM (clonal efficiency 0.000697 vs 0.01037, P=0.016), CFU-GM (0.002606 vs 0.004071, P=0.038) and BFU-E colonies (0.003213 vs 0.004697, P=0.031) in the cells exposed to GO. We then performed the same experiments on samples from the same PBSC units after the enrichment in CD34⁺/CD38⁻ cells by immunomagnetic sorting. The efficacy of this sorting was proved by either immunophenotypic analysis and functional assays, which resulted in significantly more numerous CFU-GEMM, CFU-GM, BFU-E (*data not shown*). This time, though, we did not observe a significant decrease in the clonogenic potential of CD34⁺/CD38⁻ cells by the exposure to GO (0.01518 vs 0.02631, P=0.351), thus confirming the preservation of more immature hematopoietic precursors. Moreover, in one MRD-positive patient we could observe the disappearance of the AML1-ETO molecular transcript following *in vitro* incubation with GO at a concentration of 5 µg/mL for two hours (Figure 2).

Therefore, with the limit of small numbers, the results that we report suggest the possibility of using GO in a purging strategy that would possibly act on residual CBF AML cells without affecting the repopulating ability of PBSC. In order to avoid the limitation of *in vitro* purging, GO could be used *in vivo* before CD34⁺ cell collection in MRD-positive patients.




Figure 2. Disappearance of AML1-ETO molecular transcript following *in vitro* purging of PBSC with Gemtuzumab Ozogamicin. Samples from PBSC collected from 5 patients affected by CBF AML at the end of consolidation were cultured in the presence or absence of Gemtuzumab Ozogamicin at a concentration of 5 µg/mL for two hours. In one patient (*i.e.* TG) affected by t(8;21) AML with persistent AML1-ETO transcript at the end of consolidation and in the PBSC, incubation with GO obtained the disappearance of cells expressing AML1-ETO, as tested by either direct and nested PCR. Abelson (ABL) amplification was used as internal control. TG: initials of the patient's name; Ctrl-/Ctrl+: negative and positive controls.

References

1. Paschka P. Core binding factor acute myeloid leukemia. *Semin Oncol* 2008; 35:410-7.
2. Borthakur G, Kantarjian H, Wang X, et al. Treatment of core-binding-factor acute myelogenous leukemia with fludarabine, cytarabine and granulocyte colony-stimulating factor results in improved event-free survival. *Cancer* 2008;113:3181-5.
3. Burnett AK, Hills RK, Milligan D, et al. Identification of patients with acute myeloblastic leukemia who benefit from the addition of gemtuzumab ozogamicin: results of the MRC AML 15

- trial. *J Clin Oncol* 2011;29:369-77.
4. Candoni A, Martinelli G, Toffoletti E, et al. Gemtuzumab-ozogamicin in combination with fludarabine, cytarabine, idarubicin (FLAI-GO) as induction therapy in CD33-positive AML patients younger than 65 years. *Leuk Res* 2008;32:1800-8.
 5. Rowe JM, Lowenberg B. Gemtuzumab Ozogamycin in acute myeloid leukemia: a remarkable saga about an active drug. *Blood* 2013;121:4838-41.
 6. Petersdorf SH, Kopecky KJ, Slovak ML, et al. A phase III study of gemtuzumab ozogamicin during induction and post-consolidation therapy in younger patients with acute myeloid leukemia. *Blood* 2013;121:4854-60.
 7. Castaigne S, Pautas C, Terré C, et al. Effect of gemtuzumab ozogamicin on survival of adult patients with de-novo acute myeloid leukemia (ALFA-0701): a randomised, open label, phase 3 study. *Lancet* 2012;379:1508-16.
 8. Burnett AK, Russell NH, Hills RK, et al. Addition of gemtuzumab ozogamicin to induction chemotherapy improves survival in older patients with acute myeloid leukemia. *J Clin Oncol* 2012; 30:3924-31.
 9. Pashka P, Marcucci G, Ruppert AS, et al. Adverse prognostic significance of KIT mutations in adult acute myeloid leukemia with inv(16) and t(8;21): a Cancer and Leukemia Group B Study. *J Clin Oncol* 2006;24:3904-11.
 10. Tobal K, Newton J, Macheta M, et al. Molecular quantitation of minimal residual disease in acute myeloid leukemia with t(8;21) can identify patients in durable remission and predict clinical relapse. *Blood* 2000;95:815-9.
 11. Martin G, Barragan E, Bolufer P, et al. Relevance of presenting white blood cell count and kinetics of molecular remission in the prognosis of acute myeloid leukemia with CBFbeta/MYH11 rearrangement. *Haematologica* 2000;85:699-703.
 12. Del Poeta G, Venditti A, Aronica G, et al. P-glycoprotein expression in de novo acute myeloid leukemia. *Leuk Lymphoma* 1997;27:257-74.
 13. Lutterbach B, Sun D, Schuetz J, Hiebert SW. The MYND motif is required for repression of basal transcription from the multidrug resistance 1 promoter by the t(8;21) fusion protein. *Mol Cell Biol* 1998;18:3604-11.
 14. Linenberger ML, Hong T, Flowers D, et al. Multidrug-resistance phenotype and clinical responses to gemtuzumab ozogamicin. *Blood* 2001;98:988-94.
 15. Mosna F, Gottardi M. Stem cell modeling of core binding factor acute myeloid leukemia. *Stem Cell Int* 2016;2016:7625827.
 16. Mosna F, Papayannidis C, Martinelli G, et al. Complex karyotype, older age, and reduced first-line dose intensity determine poor survival in core binding factor acute myeloid leukemia patients with long-term follow-up. *Am J Hematol* 2015;90:515-23.
 17. Gorin NC, Giebel S, Labopin M, et al. Autologous stem cell transplantation for adult acute leukemia in 2015: time to rethink? Present status and future prospects. *Bone Marrow Transplant* 2015;50:1495-502.
 18. Nakasone H, Izutsu K, Wakita S, et al. Autologous stem cell transplantation with PCR-negative graft would be associated with a favourable outcome in core-binding factor acute myeloid leukemia. *Biol Blood Marrow Transplant* 2008;14:1262-9.
 19. Grimwade D, Freeman SD. Defining minimal residual disease in acute myeloid leukemia: which platforms are ready for "prime time"? *Hematol Am Soc Hematol Educ Program* 2014; 2014:222-33.
 20. van Der Velden VH, te Marvelde JG, Hoogeveen PG, et al. Targeting of the CD33-calicheamicin immunoconjugate Mylotarg (CMA-676) in acute myeloid leukemia: in vivo and in vitro saturation and internalization by leukemic and normal myeloid cells. *Blood* 200;97: 3197-204.