This is the peer reviewd version of the followng article:

Allele specific CRISPR/Cas9 editing of dominant Epidermolysis Bullosa Simplex in human epidermal stem cells / Cattaneo, C; Enzo, E; De Rosa, L; Sercia, L; Consiglio, F; Forcato, M; Bicciato, S; Paiardini, A; Basso, G; Tagliafico, E; Paganelli, A; Fiorentini, C; Magnoni, C; Latella, M C; De Luca, M. - In: MOLECULAR THERAPY. - ISSN 1525-0016. - 32:2(2023), pp. 372-383. [10.1016/j.ymthe.2023.11.027]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

08/05/2024 23:57

Allele specific CRISPR/Cas9 editing of dominant Epidermolysis Bullosa Simplex in human epidermal stem cells.

C. Cattaneo, E. Enzo, L. De Rosa, L. Sercia, F. Consiglio, M. Forcato, S. Bicciato, A. Paiardini, G. Basso, E. Tagliafico, A. Paganelli, C. Fiorentini, C. Magnoni, M.C. Latella, M. De Luca

PII: S1525-0016(23)00660-3

DOI: https://doi.org/10.1016/j.ymthe.2023.11.027

Reference: YMTHE 6251

To appear in: *Molecular Therapy*

Received Date: 22 May 2023

Accepted Date: 30 November 2023

Please cite this article as: Cattaneo C, Enzo E, De Rosa L, Sercia L, Consiglio F, Forcato M, Bicciato S, Paiardini A, Basso G, Tagliafico E, Paganelli A, Fiorentini C, Magnoni C, Latella MC, De Luca M, Allele specific CRISPR/Cas9 editing of dominant Epidermolysis Bullosa Simplex in human epidermal stem cells., *Molecular Therapy* (2024), doi: https://doi.org/10.1016/j.ymthe.2023.11.027.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Author(s).





1	Allele specific CRISPR/Cas9 editing of dominant Epidermolysis Bullosa Simplex in human
2	epidermal stem cells.
3	Cattaneo C. ¹ , Enzo E. ¹ , De Rosa L. ¹ , Sercia L. ¹ , Consiglio F. ² , Forcato M. ³ , Bicciato S. ³ ,
4	Paiardini A. ⁴ , Basso G. ⁵ , Tagliafico E. ⁶ , Paganelli A. ⁷ , Fiorentini C. ⁷ , Magnoni C. ⁷ , Latella M.C.* ² ,
5	De Luca M.* ¹ .
6	* These authors contributed equally to this work.
7	Affiliations
8	• ¹ Centre for Regenerative Medicine "Stefano Ferrari", Department of Life Sciences,
9	University of Modena and Reggio Emilia, 41125, Modena, Italy.
10	• ² Holostem Terapie Avanzate, s.r.l., 41125, Modena, Italy.
11	• ³ Department of Life Sciences, University of Modena and Reggio Emilia, 41125, Modena,
12	Italy.
13	• ⁴ Department of Biochemical Sciences 'A. Rossi Fanelli', Sapienza Università di Roma,
14	Rome, 00185, Italy.
15	• ⁵ Genomic Units, IRCCS Humanitas Research Hospital, 20089, Rozzano, Milan, Italy
16	• ⁶ Department of Laboratory Medicine and Pathology, Diagnostic hematology and Clinical,
17	Genomics Unit, Modena University Hospital, 41124, Modena, Italy.
18	• ⁷ Regenerative and Oncological Dermatological Surgery Unit, Modena University Hospital,
19	41124 Modena, Italy.
20	Correspondence should be addressed to M.D.L.: <u>michele.deluca@unimore.it</u> ; tel. +39 059
21	2058057; Fax. +39 059 2058115
22	

24 Abstract

25 Epidermolysis Bullosa Simplex (EBS) is a rare skin disease inherited mostly in an autosomal dominant manner. Patients display a skin fragility that leads to blisters and erosions caused by minor 26 27 mechanical trauma. EBS phenotypic and genotypic variants are caused by genetic defects in 28 intracellular proteins whose function is to provide the attachment of basal keratinocytes to the 29 basement membrane zone and most of EBS cases display mutations in keratin 5 (KRT5) and keratin 30 14 (*KRT14*) genes. Besides palliative treatments, there is still no long-lasting effective cure to correct 31 the mutant gene and abolish dominant negative effect of the pathogenic protein over its wild-type 32 counterpart. Here, we propose a molecular strategy for EBS01 patient's keratinocytes carrying a 33 monoallelic c.475/495del21 mutation in KRT14 exon1. Through the CRISPR/Cas9 system we performed a specific cleavage only on the mutant allele and restore a normal cellular phenotype and 34 a correct intermediate filament network, without affecting the epidermal stem cell, referred to as 35 holoclones, which play a crucial role in epidermal regeneration. 36

37 Introduction

38 Inherited EB is a heterogeneous group of rare, autosomal genetic disorders caused by molecular 39 defects within genes encoding structural proteins forming the epidermal-dermal junction. EB is 40 characterized by recurrent blistering and erosions of the skin (and other stratified epithelia) that arise, 41 spontaneously or upon minimal mechanical stress, within the epidermis in EB simplex (EBS), the 42 lamina lucida in Junctional EB (JEB) and beneath the lamina densa in Dystrophic EB (DEB). EBS is the most common EB form, with a prevalence of 1/30000 - 1/50000.^{1,2} Its clinical manifestations are 43 usually less severe than those of JEB and DEB, which can be devastating and even early lethal. 44 45 However, some EBS forms are marked by a severe phenotype and several clinical variants have been identified based on the mutated gene, site of blister formation, anatomical distribution and mode of 46 inheritance.^{3–5} 47

JEB and DEB are mostly recessively inherited, whilst the vast majority of EBS are inherited in a 48 49 dominant manner. In fact, approximately 75% patients suffering from EBS harbours dominant 50 mutations in *KRT5* and *KRT14*, the genes encoding keratin 5 (K5) and keratin 14 (K14), respectively. 51 K5/K14 pairs form the basal keratinocyte intermediate filaments, which are part of the 52 hemidesmosomal protein complex tethering the epidermal basal layer to the basement membrane and 53 the underlying dermis. Mutant keratins exert a dominant negative effect on the functional keratins 54 encoded by the normal allele, hence perturbing the basal keratinocyte intermediate filament network 55 and leading to intraepidermal blister formation. Thus, whilst JEB and DEB can be tackled by the addition of a corrected copy of the mutated gene in the genome of epidermal stem cells⁶⁻¹⁴, a 56 57 potentially successful combined ex vivo cell and gene therapy of EBS strictly requires editing of the 58 mutated allele.

Here, we outlined an allele specific CRISPR/Cas9 based gene editing approach that is able to disrupt specifically the *KRT14* mutant gene and fully restore functional intermediate filaments in epidermal stem cells, cultivated from an EBS patient carrying a *de novo* monoallelic c.475/495del21 dominant mutation in exon 1 of *KRT14*.

This approach takes advantage of a tailored CRISPR/Cas9 system to induce double strand breaks 63 64 (DSBs) specifically on the mutant allele, leading to non-homologous end joining (NHEJ) repairing process. These rearrangements are likely to generate frameshift mutations resulting in both 65 pathogenic allele expression abolishment and phenotypic and mechanical stress resilience restoration. 66 Besides the remarkable efficacy of this approach, we also demonstrate the correction of the epithelial 67 68 stem cells compartment, which is mandatory for the long-term skin regeneration. This highly effective and safe gene editing strategy would therefore enable a translation to clinical application for the 69 70 treatment of other dominant form of EB.

72 **Results**

73 Novel monoallelic *KRT14* deletion causing a dominant form of Epidermolysis Bullosa Simplex

An 8-year-old EBS patient (referred to as EBS01) suffered from a *de novo* heterozygous dominant mutation (c.475/495del21) within exon 1 of *KRT14*. The patient developed bullous skin lesions few months after birth and currently presents blisters in the palmoplantar region causing postural and ambulation problems. No other cases of EB were known among his relatives. Besides the palliative care and a regular multidisciplinary follow-up, no resolutive treatment is available for this patient (Fig. 1A).

80 The EBS01 variant results in the deletion of seven in-frame amino acids, leading to a shorter K14 protein. The AlphaFold2¹⁵ suite for protein structure prediction and modeling was used to predict the 81 3D-structure of the shorter K14 (Fig. 1B). The c.475/495del21 variant affects the protein structure 82 resulting in the absence of an extended loop, Linker L1, encompassing residues 159–165 (Fig. 1B).¹⁶ 83 84 The L1 structural motif of K14, whose function is still poorly characterized, is predicted to assume a highly flexible, non-helical β -turn, with the pivotal function of connecting coil 1A with 1B (Fig. 85 1C).¹⁷ The importance of the L1 motif is also reflected by its high evolutionary conservation in 86 keratins.¹⁸ Indeed, mutations affecting the linker regions of intermediate filaments have been 87 88 previously observed and related to severe cases of inherited skin blistering diseases, highlighting the unexpected sensitivity of these regions to structural alterations.¹⁷ 89

90

91 Efficient and precise *KRT14* allele-specific editing in EBS primary keratinocytes

The EBS01 genetic variation enables the mutant K14 to exert a dominant negative effect over its wild-type counterpart expressed by the other allele. Thus, a tailored CRISPR/Cas9 system was employed to target the monoallelic *de novo* mutation (*c.475/495del21*) within exon 1 of *KRT14* to induce a deleterious double strand break on the mutant allele and promote NHEJ, inducing specific

disruption of the mutant *KRT14* open reading frame. We designed a sgRNA tailored to specifically
target only the mutant *KRT14* allele and employed the SpCas9 to specifically recognize the "NGG"
PAM present near the deletion site. For the desired specificity, the 19 nucleotides long guide RNA
was sketched straddling both the terminal sides of the 21 base pair deletion and directly flanking the
"AGG" PAM sequence within its 3' end (Fig. 2A).

To first assess the ability of the designed CRISPR/Cas9 system to specifically abolish the expression
of the *KRT14* mutant allele, preliminary experiments employed a lentiviral vector to deliver the gene
editing machinery to EBS01-derived primary keratinocytes.

104 Strikingly, the gene editing machinery was able to disrupt the expression of the mutant *KRT14* allele 105 with an efficiency up to 94%, without affecting the wild-type allele (Fig. S1A). We then performed 106 an *in silico* genome analysis to assess the site specificity of sgRNA del21 using Cas-OFFinder, 107 followed by CCTop and COSMID. We could not detect off-target sites with either none or single 108 mismatch and DNA bulge size=0. However, introducing 2 random mismatches led to the discovery 109 of 2 potential off-targets, both located in intergenic sequences. When 3 random mismatches were 110 introduced, we identified a total of 86 potential off-targets. The majority of these off-target sites were 111 found in intergenic regions (41) and introns (44), with only one potential off-target located in a coding 112 region (3' untranslated region of ZNF641 gene). We focused on six off-targets out of these candidates, 113 based on their potential relevance to gene function in the epidermis.

TIDE analysis outlined the presence of unwanted cleavage in one of the predicted off-target sites (Fig. S1B). These preliminary data suggested that the editing strategy designed to tackle the EBS01 mutation is indeed appropriate.

117 Lentiviral-mediated genome integration of the CRISPR/Cas9 components and their constitutive 118 expression may increase off-target cleavage of similar genomic sequences overtime and trigger 119 unwanted immunological response due to the Cas9 bacterial origin.¹⁹ In fact, stable genome

120 integration of the lentiviral-mediated gene editing cassette is neither needed, nor desirable for clinical 121 application, in that transient expression of both endonuclease and guide RNA is sufficient to attain a 122 stable gene editing.

However, human keratinocytes are hard to transfect and genome access turns out to be a major issue.
Thus, different transfection methods were attempted to deliver the gene editing machinery into
EBS01 keratinocytes.

Plasmids expressing CRISPR/Cas9 components were initially delivered into cells using commercial lipofectamine reagents, which turned out to be highly inefficient. To implement the transfection efficacy and assure an optimal editing efficiency, additional electroporation procedures were investigated, all of which were highly toxic for normal human keratinocytes (data not shown).

To overcome these hurdles, EBS01 primary keratinocytes were directly electroporated with a
ribonucleoprotein complex (RNP) composed of the SpCas9 endonuclease protein and the guide RNA
specific for the mutant allele (sgRNA_del21) (sequence in Table S1).

Following RNP nucleofection, edited EBS01 keratinocytes (eEBS01) underwent further analysis to
 characterize genotypically and phenotypically the editing impact, always comparing them with a not
 transfected EBS01 sample.

Genomic DNA was extracted from sub-confluent eEBS01 and EBS01 cultures and the locus around the 21 base pair deletion was amplified by PCR and used to perform Sanger sequencing. The amplicon sequences were analysed using TIDE analysis (sensitivity >1-5%) and validated with next-generation sequencing (NGS).

As shown in Figure 2B, TIDE analysis outlined a strikingly high allele specificity of our tailored gene editing approach. In eEBS01 cells, the wild-type allele was virtually untouched, whereas the mutant allele displayed a significant amount of InDels abolishing the open reading frame and expression of

pathogenic K14. NGS analysis of three independent RNP nucleofections confirmed the remarkable 143 144 allele specificity (Table S2), with a mutant allele specific gene editing greater than 95% (Fig. 2C). 145 Of note, ddPCR (Fig. S2A) and western blot analysis (Fig. S2B) showed that editing of the mutant 146 allele restored both KRT14 mRNA and K14 expression in eEBS01 keratinocytes, as compared to 147 EBS01 cells. Gene editing specificity was confirmed in an agarose gel analysis, in which PCR products were amplified with oligonucleotides specific for the wild-type (268 bp) and mutant alleles 148 149 (266bp). As shown in Figure 2D, the introduction of several mismatches, after editing and error prone 150 DNA repair, caused the inability of the forward mutant allele specific oligonucleotide 151 (KRT14_seq_del21 primer, see Table S1) to anneal in the edited region, determining the formation 152 of a feebler eEBS01 mutant allele amplification product. Finally, NGS data were employed to calculate the frequency of deletion, insertion and substitution in eEBS01 keratinocytes, outlining 153

deletion as the most frequent modification type (75%), whereas insertion and substitution were detected at a lower frequency (19% and 6% respectively) (Fig. 2E). An independent analysis of the NGS data was performed to gain insights into the prevalence of out-of-frame sequences (80% of the total sequences). The most frequent specific rearrangements are illustrated in Figure 2F.

158

159 Off-Target analysis supports RNP complex-mediated editing specificity and safety

To properly address safety issues, a comprehensive analysis assessing potential off-target sites was performed. Genomic DNA purified from sub-confluent EBS01 keratinocytes electroporated with the RNP complex was used to amplify and sequence off-target genomic regions previously identified after lentiviral transduction. TIDE analysis of eEBS01 shows no undesirable changes in any of the 6 predicted off-target sites, including *PPFIBP1*, thus recovering the unwanted cut chance determined by the genomic lentiviral vector-mediated integration and stable expression of the CRISPR/Cas9 components (Fig. S3).

167 Unbiased genome wide GUIDE-seq analysis was carried-out, the DNA library was sequenced using 168 Illumina Miseq. The subsequent datasets were analysed using GUIDE-seq Bioconductor package 169 software. The analysis confirmed preservation of the KRT14 wild-type allele but outlined 9 "putative" 170 off-target sites with a frequency below 3% but above 1% (Table S3). However, the vast majority of 171 the implicated gene regions are either intergenic or intronic. The unique coding region involved is 172 attributed to the ZNF320 gene, whose expression is anyhow low in basal and superbasal keratinocytes 173 and with a very low GUIDE-seq predicted editing rate. We NGS validated two off-target sites 174 identified in intronic regions (MIPOL and BAIAP) and in the only coding region (ZNF320) detected 175 with GUIDE-seq, for which we obtained appropriate quality PCR products. NGS showed editing 176 efficiencies of 4.1%, 5.3%, and 6.9% for these respective targets (Table S3; NGS raw data published 177 on GSE246345).

Overall, this data indicates the presence of a small number of potential off-target sites, which have a
minimal effect on gene function, underscoring the safety of this non-viral CRISPR/Cas9 approach.

Editing of *KRT14* mutant allele restores functional intermediate filament network in EBS01 keratinocytes

Given the role of K5/K14 pairs in the assembly of keratinocyte intermediate filaments, we investigated whether the unmodified wild-type *KRT14* allele would suffice in restoring normal structural and functional phenotype in edited keratinocytes.

Healthy donor (NHEK), eEBS01 and EBS01 keratinocytes were seeded onto glass coverslips.
Immunostained colonies clearly showed that corrected eEBS01 keratinocytes contained a properly
functioning intermediate filament network, virtually indistinguishable from that of healthy NHEK.
Keratin filaments were well organized and capable of branching throughout the plasma membrane.
In contrast, EBS01 cells displayed a pathologic, roughly fragmentated keratin pattern (Fig. 3A). We
have analysed the intermediate filament assembly in 636 NHEK, 722 EBS01 and 1090 eEBS01 cells.

Such analysis is summarized in Fig. S4 (panel D). The magnified areas shown in Fig. S4 (A-C) are strictly representative of the several images that we have analysed in several independent experiments. This data confirmed the efficacy of this gene editing approach in restoring EBS altered cellular phenotype Of note, the ablation of the mutant *KRT14* allele results in *KRT14* haploinsufficiency, which is comparable to the condition characterizing heathy carriers of recessive EBS mutations, who are completely asymptomatic.^{20–22}

Additional functional assays evaluated the regain of mechanical strength in eEBS01 keratinocytes,
 which is mandatory for a more comprehensive validation of an appropriate functional correction.²³

Heat shock assay is one of the easiest and most reproducible tests demonstrating the instability and thermal sensitivity of mutant keratins in EBS cells. The transient increase in thermal energy of the system results in evident depolymerization and impairment in affected keratinocytes' filament network remodelling, which may render cells vulnerable to cytolysis *in vivo* and support the increased EBS blisters formation in warm environments.²⁴

To this end, EBS01, eEBS01 and NHEK keratinocyte colonies were submitted to thermal shock. At time zero and fifteen minutes after heat shock, EBS01 keratinocyte's intermediate filaments showed an increased disruption of keratin filaments, particularly nearby the nuclear region, which was partially recovered approximately 60 minutes after thermal shock, with aggregates limited to a small portion of the cytoplasm.^{25,26} Such cytoplasmatic aggregates persistence was not observed in NHEK or eEBS01 (Figure 3B).

Since intermediate filaments also play a role in controlling cell mechanical stress, cultured epidermal sheets prepared from NHEK, EBS01 and eEBS01 were detached from the vessel by incubation with Dispase II protease (Fig. 3C). EBS01 cultured sheets disintegrated into small pieces when subjected to high inversion force, whereas eEBS01 sheet showed a structural compactness comparable to that of the healthy donor (NHEK) (Fig. 3D). The greater eEBS01 mechanical strength was also quantitatively evaluated counting sheet-derived fragments and confirmed that specific deletion of the 217 mutated *KRT14* allele confers to eEBS01 cells the capability to reassemble proper and resistant 218 cohesive structures (Fig. 3E).

219

220 Edited eEBS01 cells revert the disease phenotype in skin equivalents

We have generated human skin equivalents containing dermal and epidermal compartments resembling morphological characteristics of human skin. This was achieved by cultivating keratinocytes on a decellularized human dermal matrix.

Figure 4A shows haematoxylin/eosin staining of sections of decellularized dermal matrixes without cells (Dermis) and those overlaid by a fully stratified epidermis (NHEK). Figure 4B (left panels) illustrates the decellularized dermal matrix seeded with EBS01 cells. The regenerated epidermis shows the presence of blisters within the epidermal basal layer (at arrows). In contrast, no blisters were observed in the epidermis generated by gene-edited eEBS01keratinocytes (Figure 4B, right panels).

In summary, these results show that the allelic-specific gene editing of mutant *KRT14* restores proper
expression of K14 and functional intermediate filaments in primary clonogenic EBS keratinocytes.

232

233 CRISPR/Cas9-mediated gene editing via RNP complex electroporation preserves epidermal 234 stem cells

Epidermal regeneration and repair processes rely on long-lived stem cells producing short-lived transient amplifying (TA) progenitors that eventually give rise to terminally differentiated keratinocytes. Keratinocyte stem cells and TA progenitors are located in the basal layer of all stratified epithelia and generate different clonal types, referred to as Holoclones and Meroclones/Paraclones respectively.^{27–31} In view of future clinical applications, the essential feature of any cultured epithelial graft is an adequate number of Holoclones-forming cells, which are mandatory for a stable long-term regeneration of all squamous epithelia.^{10,29,32–34} Clonal analysis of EBS01 clonogenic keratinocytes

confirmed the presence of each clonal type in the culture (Holoclones, Meroclones and Paraclones),
excluding an impact of this *de novo* mutation on the EBS01 derived keratinocyte stem cell
compartment.

245 To first investigate a potential impact of gene editing on the distribution of the different keratinocyte 246 clonal types, we took advantage of single keratinocyte RNA sequencing analysis. To this end, we 247 performed transcriptomic profile analysis of EBS01 and eEBS01 keratinocytes following the same pipeline recently published²⁸, obtaining 5,350 and 7,200 cells, respectively. As reported, both samples 248 249 contained the previously identified 5 keratinocytes clusters (Fig. 5A-C), three of which expressed 250 clonogenic markers (Holoclones, Meroclones and Paraclones clusters), the other two expressing 251 differentiation markers (Terminally differentiated 1 and 2 clusters) (Fig. S5A-B). In particular, the Holoclone (stem cell) cluster displayed a "holoclone signature" able to distinguish it from the other 252 253 clusters (Fig. S5A-B).

To further investigated whether Holoclones were preserved and properly corrected after the gene editing procedure, two clonal analyses of EBS01 and eEBS01 keratinocytes were performed, as described in Material and Methods. The classification of the clonal type confirmed the comparable percentage of Holoclones, Meroclones and Paraclones in both samples (Fig. 5D-E). Genomic DNA, analysed by PCR amplification using primers specific to amplified wild-type and mutant alleles (see Table S1) confirmed that holoclone-forming cells have been edited (Fig. 5F).

As shown in Supplemental Fig. S6, both clonogenicity (A) and percentage of aborted colonies (B) (and growth rate) were comparable in long-term cultures generated by EBS01 and eEBS01 keratinocytes, indicating the absence of clonal selection and/or selective advantages of eEBS01 keratinocytes over the EBS01 cells.

264 These data demonstrate that the gene editing procedure was able to preserve and edit the population

265 of epidermal stem cells crucial to a future *ex vivo* gene therapy aimed at full epidermal restoration.

266

267 **Discussion**

Epidermolysis Bullosa simplex (EBS) is a rare mechanobollous disease inherited mainly in an autosomal dominant fashion and affecting a few thousands of people worldwide. Mutations affecting *KRT14* account for approximately 30% of the reported cases. Mutant *KRT14* exerts a dominant negative effect on the normal allele, perturbing the basal keratinocyte intermediate filament network and leading to intraepidermal blister formation. Hence, a great portion of EBS patients carrying *KRT14* mutations could potentially benefit from the correction of the genetic defect.³⁵

Gene replacement strategies, whereby a functional copy of the defective gene is introduced in the 274 genome of clonogenic keratinocytes, has been successfully exploited in other forms of EB, such as 275 the LAMB3-dependent JEB and RDEB, which are recessively inherited.^{6,8,10-13} However, gene 276 277 replacement is not appropriate for the treatment of dominantly inherited genetic diseases, which 278 instead require either the selective disruption of the mutated allele or the precise editing of the specific mutation.^{36,37} An appropriately designed gene editing machinery would allow to discriminate and 279 280 inactivate, via error prone NHEJ pathway, only the mutant allele, leaving the wild-type counterpart functionally intact. Gene editing has been employed to permanently to repair both dominant^{38–41} and 281 recessive^{42–44} mutations related to EB using knockout and homologous recombination techniques. 282 283 The end-joining pathways, often harnessed for gene knockout, represent the most efficient repair 284 mechanisms for double-strand breaks.⁴⁵ Consequently, gene knockout is the most effective form of 285 gene editing.

This work presents the first evidence of therapeutically relevant allele-specific genetic correction in primary cultures of EBS-derived epidermal stem cells. CRISPR/Cas9 based allele-specific editing platform using a guide RNA specific successfully edited a *de novo* c.475/495del21 mutation within exon 1 of *KRT14* gene in a dominant form of EBS. Our tailored approach was highly effective in disrupting the mutant *KRT14* allele on EBS01 primary keratinocytes (editing efficiency greater than 95%) with no editing on the wild-type allele. To ensure transient expression of editing machinery

suitable for clinical purposes, the sgRNA – Cas9 ribonucleoprotein complex (RNP) has been directly
delivered to EBS01 keratinocytes. Since InDels generation at off-target sites sill poses a risk to the
use of engineered nucleases, we also demonstrated a low occurrence of non-specific CRISPR/Cas9
mediated cleavages through unbiased GUIDE-seq analysis.

Thus, this approach succeeded in abolishing the mutant *KRT14* allele expression and the pathogenic keratin almost entirely, and fully restored a functional intermediate filament network. More importantly, we provide evidence that the long-lived self-renewing stem cells have been targeted and corrected by the gene editing machinery without any cytotoxicity, thus maintaining the ability to regenerate a virtually indistinguishable functional epidermis.

301 The selection of a gene editing tool for allele-specific genetic correction depends on factors such as 302 the specific genetic mutation, the delivery method, the desired level of precision and, perhaps more 303 importantly, on the cell type, mainly when specific somatic stem cells need to be targeted. In the case of EBS01 cells, the deletion of 21 base pairs (c.475/495del21) within exon 1 of KRT14 led us to 304 305 design a sgRNA tailored to specifically target the mutant allele. We opted for a gene editing approach 306 utilizing SpCas9, known for its high cutting efficiency. We have introduced the sgRNA and SpCas9 307 as a ribonucleoprotein (RNP) complex that overcomes many of the challenges associate with mRNA 308 delivery, as the translation steps and the folding of the Cas protein. The RNP complex is immediately 309 active as it is fully developed. Emerging gene editing tools may offer more precise approaches than 310 traditional methods, but often have an efficiency not sufficient to fully tackle a specific population of 311 epidermal stem cells, which represent a small percentage of clonogenic keratinocytes. Promising gene editing tools for allele-specific genetic correction include prime editing^{46,47}, base editing^{48,49}, and 312 Cpf1/Cas12-based editing⁵⁰⁻⁵² which provides versatility by targeting distinct DNA sequences, 313 314 widening the scope of targetable genetic mutations. These tools hold great therapeutic promise due 315 to their precision and reduced off-target effects.

In the context of clinical translation and safety of gene editing strategies, a key aspect is the analysis of off-target effects. As sequencing technologies and data analysis tools continue to advance, there may be, in the near future, more cost-effective and high-throughput options for off-target analysis than whole-genome sequencing.

Our study provides a clear demonstration of the efficacy and potential safety of an allele-specific CRISPR based gene editing approach, which we envision to further translate into a long-lasting decisive clinical treatment for patients suffering from EBS and possibly other related skin-blistering diseases. Although we did not observe abnormal clonal expansion, clinical translation would require additional efforts to determine the optimal RNP dosage to achieve maximal on-target efficiency and minimal off-target impact and thoroughly verify the absence of genotoxicity and genomic instabilities.

327

328 Materials and Methods

329 **Patient, Clinal data and treatments**

EBS patient (EBS01) displayed skin lesions to exclusively the acral regions and no mucosal blisters
 nor erosions were ever reported either by him or his parents. Clinical description and genetic
 counselling with genetic analysis of the family in Supplemental Materials and Methods.

333 Structural Modeling

The structure predictions were performed in a standalone platform of AlphaFold2 and AlphaFold-Multimer ¹⁵ as implemented in ColabFold, which was set up on a local computer with a Linux operating system and accelerated with two NVIDIA GeForce RTX 2080 Ti GPU. The "Template mode" using PDB 3TNU ¹⁶ was used for this purpose. The other parameters were kept at their default values.

339 Primary human cell culture from Epidermolysis Bullosa Simplex patient

340 A skin biopsy has been collected from the EBS01 patient, after obtaining the informed consent. 341 Briefly, skin biopsy was minced and treated with 0,05% trypsin/ 0,01% EDTA for 4 h at 37°C. Every 342 30 min keratinocytes were collected, plated (2,5-3 x 10^4 cells/cm²) on lethally irradiated 3T3-J2-Y 343 cells and grown at 37°C, 5% CO2 in humidified atmosphere in KGM medium: Dulbecco's modified 344 Eagle's (DMEM) and Ham's F12 media (2:1 mixture) containing fetal bovine serum (FBS) (10%), penicillin-streptomycin (50 IU/ml), glutamine (4mM), adenine (0,18mM), insulin (5mg/ml), cholera 345 346 toxin (0,1nM), hydrocortisone (1,1mM), triiodothyronine (Lithyronine Sodium. 2nM), epidermal growth factor (EGF, 10ng/ml). When sub-confluent, cell cultures were serially propagated. 347

348 3T3-J2 cell line

Mouse 3T3-J2 cells were a gift of Professor Howard Green, Harvard Medical School (Boston MA, USA).⁵³ Fibroblasts were cultivated in DMEM supplemented with 10% g-irradiated donor adult bovine serum, penicillin-streptomycin (50IU/ml) and glutamine (4mM). GmbH, (Idar-Oberstein, Germany) produce a GMP clinical grade 3T3-J2 cell bank. That have been authorized for clinical use by national and European regulatory authorities.

354 Ribonucleoprotein (RNP) complex formation and Nucleofection

The synthetic guide RNA was designed straddling both the terminal sides of the c475/495del21 mutation (5'-GCTGAGGTTCAAGACCATTG-3') and directly flanking an "AGG" PAM. It was modified to drive the maximum editing efficiency (Invitrogen, #A35514, Table S1) and was mixed in a 1,1:1 molar ratio with the Cas9 protein (Alt-R S.p. Cas9 Nuclease V3, IDT, #1081058). 5 x 10⁵ keratinocytes were resuspended in 100 μ l primary cell nucleofection solution (P3 Primary Cell 4D-Nucleofector Kit, Lonza, #LOV4XP3024), mixed with the RNP complex solution and 4 μ M Cas9 electroporation enhancer (Alt-R Cas9 Electroporation Enhancer, IDT, #1075915). Cells were

- 362 electroporated using a 4D-Nucleofector (4D-Nucleofector Core Unit, Lonza, #AAF-1001B; 4D-
- 363 Nucleofector X Unit, Lonza, #AAF-1001X) using the program DS-138.

364 Editing analysis by sequence decomposition (TIDE)

eEBS01 and EBS01 keratinocytes genomic DNA was isolated using the QIAamp DNA Mini Kit
(QIAGEN, #51304). A 500 base pair region around eEBS01 and EBS01 genomic target site was
amplified by PCR (primers in Table S1). PCR amplicons were subjected to conventional Sanger
sequencing. The resulting sequence trace files were uploaded on TIDE web tool with the guide RNA
sequence as input.

370 PCR and allele characterization

371 Screening of the allele pattern, in eEBS01 and EBS01 keratinocyte, was done by PCR. The 372 "KRT14_seq_del21" forward primer was specific only for the mutant allele in EBS01 keratinocyte, 373 "KRT14_seq_wt" forward primer was specific only for the wild-type allele. Reverse primer was 374 specific for a sequence common to both alleles (Table S1).

375 **OFF-target analysis**

376 CRISPR/Cas9 online predictors were used to identify the genomic regions which may present the 377 greatest probability of off-target cuttings. Off-target probability was evaluated on the basis of 378 mismatches numbers and the genomic loci of the most probable in silico off-target sites were 379 sequenced and analysed. The resulting sequence trace files were uploaded on TIDE web tool with the 380 guide RNA sequence as input.

381 NGS analysis

The region near the target site was amplified by specific PCR primers with sequence adaptor (Supplementary table 1, KRT14ex1_NGS and KRT14intr1_NGS) and 25 µl of purified amplicon was used to NGS analysis using Illumina sequencing platform. The clipping of reads was performed using

Trimmomatic (v 0.36) and paired-end reads were merged using software FLASH2 (v 2.2.00) to obtain a single, longer read that covers the full target region. The processed reads were mapped, using BMWA MEM (v 0.7.15), to the reference sequence (the wild type *KRT14* exon1 sequence) with default alignment parameters. Only high quality, merged, on target reads were considered for further processing. Finally, the identification and quantification of sequences alleles using CRISPResso (v 1.0.13) occurred. The NGS raw data are available in Gene Expression Omnibus with accession number GSE246345.

392 Clonal analysis

Sub-confluent keratinocytes mass cultures were trypsinized and 0,5-1 cell was plated into each well of a 96-well plate after serial dilution. After 7 days of cultivation, single clones were identified under an inverted microscope and treated with 0,05% trypsin and 0,01% EDTA at 37° for 15-20 minutes. One quarter of the clone was plated onto a 100mm indicator dish, cultivated for 12 days and stained with Rhodamine B for the classification of clonal type. The remaining three quarter was subcultivated on an adequate plastic support and used for further analyses.³⁴

399 Immunofluorescence

Keratinocytes were plated at 2500 cell/well onto glass coverslips. After the formation of small colonies, cells were fixed with ice cold Methanol-Acetone (1:1) at -20°C for 10 minutes. Cells were permeabilized with PBS/Triton 0,5% for 15 minutes. Blocking solution (BSA 5%, 0.3% Triton in 1X PBS) was added for 30 minutes at 37°C and sections were incubated with primary and secondary antibody (Table S1). Cell nuclei were stained with DAPI. Fluorescence signals were monitored under a Zeiss AXIO Imager A.1 Manual Operation Fluorescence Microscope with EC Plan-Neofluar 40X/0,75 objective.

407 **GUIDE-seq analysis**

408 5 x 10^5 primary EBS01 keratinocytes were nucleofected (as described in RNP paragraph) with the 409 RNP complex and 40nM of the annealed dsODN. Treated keratinocytes were then plated (4-6 x 10^3 410 cells/cm2) on lethally irradiated 3T3-J2 cells and cultured until sub-confluence. Keratinocytes' 411 genomic DNA was extracted and 14µg was sent to Creative Biogene for GUIDE-seq library 412 preparation, sequencing in order to identify RGN (CRISPR RNA-guided nucleases)-dependent and -413 independent genomic breakpoint "hotspots". DNA library was sequenced using Illumina Miseq. The 414 subsequent datasets were analysed using the *GUIDEseq* Bioconductor package software.

415 Encapsulation with 10X Genomics chromium system and bioinformatic analysis on single-cell 416 RNA-seq data

Fully confluent keratinocytes were detached and cells were accurately resuspended to obtain a single cell suspension. About 10.000 cells of each eEBS01 and EBS01 samples were loaded into two channels of the Chromium Chip B using the Single Cell reagent kit v3.1 (10X Genomic) for Gel bead Emulsion generation. Following capture and lysis, cDNA was synthesized and amplified. Fifty nanograms of the amplified cDNA were then used for each sample to construct Illumina sequencing libraries. Sequencing was performed on the NextSeq550 Illumina sequencing platform following the 10X Genomics instruction for read generation, reaching at least 50000 reads as mean reads per cell.

424 For the bioinformatic analysis, the Cell Ranger pipeline (version 3.1.0) was used to generate FASTQ 425 files, to align reads to the reference transcriptome (GRCh38) and to calculate UMI counts from the mapped reads. Expression data were imported in R version 3.6.3 and analyzed using Seurat⁵⁴ (version 426 427 3.1.5) R package. Cells were classified using an annotated scRNA-seq dataset of human keratinocytes²⁸ as reference and the FindTransferAnchors and TransferData functions in Seurat with 428 429 default parameters. We assessed the quality of the assigned labels monitoring the expression of known 430 markers. Expression data are available in Gene Expression Omnibus with accession number 431 GSE246345.

432 Heat shock assay

Keratinocytes were plated at 10000 cell/well onto glass coverslip. Cells culture medium was replaced
with KGM medium at 43°C and the well plate was immediately placed in a water bath set at 47°C.
After 15 minutes of heat stress, the medium was immediately replaced with fresh KGM medium at
37°C and the cells were allowed to recover in the incubator at 37°C, 5% CO2 in humidified
atmosphere. Coverslips were removed at 15-minute intervals thereafter and immunostained.

438 Dispase-based keratinocyte dissociation assay (DDA)

4-6 x 10³ cells/cm² keratinocytes were plated, on lethally irradiated 3T3-J2 cells onto a 6 well plate
and cultured. After 20 days, the epidermal sheets were washed with grafting wash and incubated with
Dispase II (Roche, #0494207801), 2,5U/ml diluted in PBS, for 1 hour at 37°C. After detachment, one
sheet was subjected to low force stress with orbital rotation (200 rpm) for 5 minutes at 37°C and the
other monolayer was transferred in a 15ml Falcon tube with 5ml of 1X PBS and exposed to high
mechanical stress by 20-50 inversions. Fragments count was performed with ImageJ.

445 Decellularized dermal matrixes preparation, cryopreservation and sectioning.

Decellularized human dermal matrixes were obtained using human skin samples from surgical waste 446 (abdominoplasty or mammoplasty). Briefly, skin biopsies were sectioned in fragments of 447 448 approximately 1,5 cm², immersed in sterile PBS at 60°C for thirty seconds under constant stirring 449 and then in sterile cold PBS for 1 minute. The epidermis was then mechanically detached from the 450 dermis using forceps. After decellularization, samples were rinsed in KGM for 24 hours. The 451 following day, the decellularized dermal matrixes were seeded with primary human keratinocytes (1 x 10^5 cells per scaffold) onto lethally irradiated 3T3-J2 cells (5 x 10^4 cells per scaffold) in KGM. 452 453 After 10 days in submerged culture, the media was carefully removed, and the samples were gently 454 moved in millicell® cell culture (Merk) and were further cultured for 20-24 days in air-liquid interface (ALI) condition to induce epidermal differentiation. 3D human skin equivalents were dehydrated in 455

a sucrose gradient 0.9M and 2M for thirty minutes respectively at RT, embedded in Killik-OCT
cryostat embedding medium (Bio-Optica) and frozen. 7μm sections of embedded skin equivalents
were obtained with a histological cryomicrotome (Leica CM1850 UV).

459 Haematoxylin and eosin staining.

Hematoxylin and eosin staining was performed on 7µm cryosections of decellularized dermal
matrixes (Harris haematoxylin for 1 min, running tap water for 1 min, eosin Y 50% in ethanol for
thirty seconds, 95% ethanol for 1 min, 100% ethanol for 1 min, two rinses in fresh 100% ethanol for
1 min each) and observed with Zeiss Microscope Axio Imager M2 with an EC Plan-Neofluar 10X/0.3
M27 air objective.

465

466 Acknowledgments

467 This project was supported by the European Research Council (ERC) Advanced Grant HOLO-GT
468 (No. 101019289) and Telethon (Grant number: GGP20088). We thank Le ali di Camilla for providing
469 assistance to patients.

470

471 Author Contributions

C.C. performed experiments, analyzed data, assembled all input data. E.E. defined single cells RNA
seq analyses and revised the manuscript. L.D.R. defined and analyzed functional assay and revised
the manuscript. L.S. performed 3D skin equivalent assays. F.C. performed clonal analysis. M.F. and
S.B. conducted bioinformatics analyses. A.P. performed protein structure prediction and modeling.
G.B. made scRNA-seq library. E.T. performed NGS to identify gene variants. A. Paganelli, C.F. and
C.M. provided EBS01 patient clinical management. M.C.L. and M.D.L. coordinated the study,
defined strategic procedures, administered the experiments, and wrote the manuscript.

480 **Conflict of Interest**

M.D.L. is co-founder and member of the Board of Directors of Holostem Terapie Avanzate (HTA)
s.r.l in liquidation, Modena, Italy, as well as consultants for J-TEC-Japan Tissue Engineering, Ltd.
The other authors state no conflict of interest.

484

485 Keywords

486 Epidermolysis bullosa, genetic disease, gene therapy, stem cells, keratinocytes biology, gene editing

487

488 Data Availability Statement

489 Data availability Sequencing data have been deposited to Gene Expression Omnibus with accession 490 number GSE246345. We declare that the data supporting the findings of this study are available 491 within the paper and its Supplementary Information Files or from the authors upon request.

492

493 **References**

Coulombe, P.A., Kerns, M.L., and Fuchs, E. (2009). Epidermolysis bullosa simplex: a paradigm
 for disorders of tissue fragility. J. Clin. Invest. *119*, 1784–1793.

Chamcheu, J.C., Siddiqui, I.A., Syed, D.N., Adhami, V.M., Liovic, M., and Mukhtar, H. (2011).
 Keratin gene mutations in disorders of human skin and its appendages. Arch. Biochem. Biophys.
 508, 123–137.

499 3. Fine, J.-D., Bruckner-Tuderman, L., Eady, R.A.J., Bauer, E.A., Bauer, J.W., Has, C., Heagerty,
500 A., Hintner, H., Hovnanian, A., Jonkman, M.F., et al. (2014). Inherited epidermolysis bullosa:

501	Updated recommendations on diagnosis and classification. J. Am. Acad. Dermatol. 70, 1103-
502	1126.

4. Fine, J.-D., Eady, R.A.J., Bauer, E.A., Bauer, J.W., Bruckner-Tuderman, L., Heagerty, A.,
Hintner, H., Hovnanian, A., Jonkman, M.F., Leigh, I., et al. (2008). The classification of inherited
epidermolysis bullosa (EB): Report of the Third International Consensus Meeting on Diagnosis
and Classification of EB. J. Am. Acad. Dermatol. *58*, 931–950.

507 5. Sprecher, E. (2010). Epidermolysis Bullosa Simplex. Dermatol. Clin. 28, 23–32.

508 6. Bauer, J.W., Koller, J., Murauer, E.M., De Rosa, L., Enzo, E., Carulli, S., Bondanza, S., Recchia,

A., Muss, W., Diem, A., et al. (2017). Closure of a Large Chronic Wound through Transplantation

510 of Gene-Corrected Epidermal Stem Cells. J. Invest. Dermatol. 137, 778–781.

511 7. De Rosa, L., Enzo, E., Zardi, G., Bodemer, C., Magnoni, C., Schneider, H., and De Luca, M.
512 (2021). Hologene 5: A Phase II/III Clinical Trial of Combined Cell and Gene Therapy of
513 Junctional Epidermolysis Bullosa. Front. Genet. *12*, 705019.

De Rosa, L., Carulli, S., Cocchiarella, F., Quaglino, D., Enzo, E., Franchini, E., Giannetti, A.,
 De Santis, G., Recchia, A., Pellegrini, G., et al. (2014). Long-Term Stability and Safety of
 Transgenic Cultured Epidermal Stem Cells in Gene Therapy of Junctional Epidermolysis Bullosa.
 Stem Cell Rep. 2, 1–8.

Eichstadt, S., Tang, J.Y., Solis, D.C., Siprashvili, Z., Marinkovich, M.P., Whitehead, N., Schu,
 M., Fang, F., Erickson, S.W., Ritchey, M.E., et al. (2019). From Clinical Phenotype to Genotypic
 Modelling: Incidence and Prevalence of Recessive Dystrophic Epidermolysis Bullosa (RDEB).

521 Clin. Cosmet. Investig. Dermatol. *Volume 12*, 933–942.

522	10. Hirsch, T., Rothoeft, T., Teig, N., Bauer, J.W., Pellegrini, G., De Rosa, L., Scaglione, D.,
523	Reichelt, J., Klausegger, A., Kneisz, D., et al. (2017). Regeneration of the entire human epidermis
524	using transgenic stem cells. Nature 551, 327–332.
525	11. Kueckelhaus, M., Rothoeft, T., De Rosa, L., Yeni, B., Ohmann, T., Maier, C., Eitner, L., Metze,
526	D., Losi, L., Secone Seconetti, A., et al. (2021). Transgenic Epidermal Cultures for Junctional
527	Epidermolysis Bullosa — 5-Year Outcomes. N. Engl. J. Med. 385, 2264–2270.
528	12. Mavilio, F., Pellegrini, G., Ferrari, S., Di Nunzio, F., Di Iorio, E., Recchia, A., Maruggi, G.,
529	Ferrari, G., Provasi, E., Bonini, C., et al. (2006). Correction of junctional epidermolysis bullosa
530	by transplantation of genetically modified epidermal stem cells. Nat. Med. 12, 1397–1402.
531	13. Siprashvili, Z., Nguyen, N.T., Gorell, E.S., Loutit, K., Khuu, P., Furukawa, L.K., Lorenz, H.P.,
532	Leung, T.H., Keene, D.R., Rieger, K.E., et al. (2016). Safety and Wound Outcomes Following
533	Genetically Corrected Autologous Epidermal Grafts in Patients With Recessive Dystrophic
534	Epidermolysis Bullosa. JAMA 316, 1808.
535	14. Chakravarti, S., Enzo, E., Rocha Monteiro de Barros, M., Maffezzoni, M.B.R., and Pellegrini, G.
536	(2022). Genetic Disorders of the Extracellular Matrix: From Cell and Gene Therapy to Future

537 Applications in Regenerative Medicine. Annu. Rev. Genomics Hum. Genet. 23, 193–222. 10.

538 15. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool,
539 K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction
540 with AlphaFold. Nature *596*, 583–589.

541 16. Lee, C.-H., Kim, M.-S., Chung, B.M., Leahy, D.J., and Coulombe, P.A. (2012). Structural basis
542 for heteromeric assembly and perinuclear organization of keratin filaments. Nat. Struct. Mol.
543 Biol. *19*, 707–715.

544	17. Rugg, E.L., Morley, S.M., Smith, F.J.D., Boxer, M., Tidman, M.J., Navsaria, H., Leigh, I.M., and
545	Lane, E.B. (1993). Missing links: Weber–Cockayne keratin mutations implicate the L12 linker
546	domain in effective cytoskeleton function. Nat. Genet. 5, 294–300.
547	18. Parry, D.A.D., Marekov, L.N., Steinert, P.M., and Smith, T.A. (2002). A Role for the 1A and L1
548	Rod Domain Segments in Head Domain Organization and Function of Intermediate Filaments:
549	Structural Analysis of Trichocyte Keratin. J. Struct. Biol. 137, 97–108.
550	19. Charlesworth, C.T., Deshpande, P.S., Dever, D.P., Camarena, J., Lemgart, V.T., Cromer, M.K.,
551	Vakulskas, C.A., Collingwood, M.A., Zhang, L., Bode, N.M., et al. (2019). Identification of
552	preexisting adaptive immunity to Cas9 proteins in humans. Nat. Med. 25, 249–254.
553	20. Batta, K., Rugg, E.L., Wilson, N.J., West, N., Goodyear, H., Lane, E.B., Gratian, M., Dopping-
554	Hepenstal, P., Moss, C., and Eady, R.A.J. (2000). A keratin 14 `knockout' mutation in recessive
555	epidermolysis bullosa simplex resulting in less severe disease. Br. J. Dermatol., 7.
556	21. Yiasemides, E., Trisnowati, N., Su, J., Dang, N., Klingberg, S., Marr, P., Melbourne, W., Tran,
557	K., Chow, C.W., Orchard, D., et al. (2008). Clinical heterogeneity in recessive epidermolysis
558	bullosa due to mutations in the keratin 14 gene, KRT14. Clin. Exp. Dermatol. 33, 689–697.
559	22. Sørensen, C.B., Andresen, B.S., Jensen, U.B., Jensen, T.G., Jensen, P.K.A., Gregersen, N., and
560	Bolund, L. (2003). Functional testing of keratin 14 mutant proteins associated with the three
561	major subtypes of epidermolysis bullosa simplex. Exp. Dermatol. 12, 472–479.
562	23. Tan, T.S., Ng, Y.Z., Badowski, C., Dang, T., Common, J.E.A., Lacina, L., Szeverényi, I., and
563	Lane, E.B. (2016). Assays to Study Consequences of Cytoplasmic Intermediate Filament
564	Mutations. In Methods in Enzymology (Elsevier), pp. 219–253.

565	24.	Morley, S.M., Dundas, S.R., James, J.L., Gupta, T., Brown, R.A., Sexton, C.J., Navsaria, H.A.,
566		Leigh, I.M., and Lane, E.B. (1995). Temperature sensitivity of the keratin cytoskeleton and
567		delayed spreading of keratinocyte lines derived from EBS patients. J. Cell Sci. 108, 3463–3471.
568	25.	Coulombe, P.A., Hutton, M.E., Letal, A., Hebert, A., Paller, A.S., and Fuchs, E. (1991). Point
569		mutations in human keratin 14 genes of epidermolysis bullosa simplex patients: Genetic and
570		functional analyses. Cell 66, 1301–1311.
571	26.	Kitajima, Y., Inoue, S., and Yaoita, H. (1989). Abnormal organization of keratin intermediate
572		filaments in cultured keratinocytes of epidermolysis bullosa simplex. Arch. Dermatol. Res. 281,
573		5–10.
574	27.	Barrandon, Y., and Green, H. (1987). Three clonal types of keratinocyte with different capacities
575		for multiplication. Proc. Natl. Acad. Sci. 84, 2302–2306.
576	28.	Enzo, E., Secone Seconetti, A., Forcato, M., Tenedini, E., Polito, M.P., Sala, I., Carulli, S.,
577		Contin, R., Peano, C., Tagliafico, E., et al. (2021). Single-keratinocyte transcriptomic analyses
578		identify different clonal types and proliferative potential mediated by FOXM1 in human
579		epidermal stem cells. Nat. Commun. 12, 2505.
580	29.	Pellegrini, G., Golisano, O., Paterna, P., Lambiase, A., Bonini, S., Rama, P., and De Luca, M.
581		(1999). Location and Clonal Analysis of Stem Cells and Their Differentiated Progeny in the
582		Human Ocular Surface. J. Cell Biol. 145, 769–782.
583	30.	Rochat, A., Kobayashi, K., and Barrandon, Y. (1994). Location of stem cells of human hair
584		follicles by clonal analysis. Cell 76, 1063–1073.
585	31.	Polito, M.P., Marini, G., Palamenghi, M., and Enzo, E. (2023). Decoding the Human Epidermal
586		Complexity at Single-Cell Resolution. Int. J. Mol. Sci. 24, 8544.

587	32.	De Rosa, L., Enzo, E., Palamenghi, M., Sercia, L., and Luca, M.D. (2023). Stairways to Advanced
588		Therapies for Epidermolysis Bullosa. Cold Spring Harb Perspect Biol 15(4).
589	33.	Rama, P., Matuska, S., Paganoni, G., Spinelli, A., De Luca, M., and Pellegrini, G. (2010). Limbal
590		Stem-Cell Therapy and Long-Term Corneal Regeneration. N. Engl. J. Med. 363, 147–155.
591	34.	Enzo, E., Cattaneo, C., Consiglio, F., Polito, M.P., Bondanza, S., and De Luca, M. (2022). Clonal
592		analysis of human clonogenic keratinocytes. In Methods in Cell Biology (Elsevier), pp. 101–116.
593	35.	Bolling, M.C., Lemmink, H.H., Jansen, G.H.L., and Jonkman, M.F. (2011). Mutations in KRT5
594		and KRT14 cause epidermolysis bullosa simplex in 75% of the patients: KRT5 and KRT14
595		mutations in 75% of EBS patients. Br. J. Dermatol., 637–44.
596	36.	György, B., Nist-Lund, C., Pan, B., Asai, Y., Karavitaki, K.D., Kleinstiver, B.P., Garcia, S.P.,
597		Zaborowski, M.P., Solanes, P., Spataro, S., et al. (2019). Allele-specific gene editing prevents
598		deafness in a model of dominant progressive hearing loss. Nat. Med. 25, 1123–1130.
599	37.	Latella, M.C., Di Salvo, M.T., Cocchiarella, F., Benati, D., Grisendi, G., Comitato, A., Marigo,
600		V., and Recchia, A. (2016). In vivo Editing of the Human Mutant Rhodopsin Gene by
601		Electroporation of Plasmid-based CRISPR/Cas9 in the Mouse Retina. Mol. Ther Nucleic Acids
602		5, e389.
603	38.	Kocher, T., Peking, P., Klausegger, A., Murauer, E.M., Hofbauer, J.P., Wally, V., Lettner, T.,
604		Hainzl, S., Ablinger, M., Bauer, J.W., et al. (2017). Cut and Paste: Efficient Homology-Directed
605		Repair of a Dominant Negative KRT14 Mutation via CRISPR/Cas9 Nickases. Mol. Ther. 25,
606		2585–2598.

39. March, O.P., Lettner, T., Klausegger, A., Ablinger, M., Kocher, T., Hainzl, S., Peking, P.,
Lackner, N., Rajan, N., Hofbauer, J.P., et al. (2019). Gene Editing–Mediated Disruption of

- Epidermolytic Ichthyosis–Associated KRT10 Alleles Restores Filament Stability in
 Keratinocytes, J. Invest. Dermatol. *139*, 1699-1710.
- 611 40. Luan, X.-R., Chen, X.-L., Tang, Y.-X., Zhang, J.-Y., Gao, X., Ke, H.-P., Lin, Z.-Y., and Zhang,
- 612 X.-N. (2018). CRISPR/Cas9-Mediated Treatment Ameliorates the Phenotype of the
- 613 Epidermolytic Palmoplantar Keratoderma-like Mouse. Mol. Ther. Nucleic Acids 12, 220–228.
- 41. Aushev, M., Koller, U., Mussolino, C., Cathomen, T., and Reichelt, J. (2017). Traceless Targeting

and Isolation of Gene-Edited Immortalized Keratinocytes from Epidermolysis Bullosa Simplex

616 Patients. Mol. Ther. - Methods Clin. Dev. 6, 112–123.

- 42. Webber, B.R., Osborn, M.J., McElroy, A.N., Twaroski, K., Lonetree, C., DeFeo, A.P., Xia, L.,
- Eide, C., Lees, C.J., McElmurry, R.T., et al. (2016). CRISPR/Cas9-based genetic correction for
- 619 recessive dystrophic epidermolysis bullosa. Npj Regen. Med. 1, 16014.
- 43. Hainzl, S., Peking, P., Kocher, T., Murauer, E.M., Larcher, F., Del Rio, M., Duarte, B., Steiner,
- M., Klausegger, A., Bauer, J.W., et al. (2017). COL7A1 Editing via CRISPR/Cas9 in Recessive
 Dystrophic Epidermolysis Bullosa. Mol. Ther. 25, 2573–2584.
- 623 44. Wu, W., Lu, Z., Li, F., Wang, W., Qian, N., Duan, J., Zhang, Y., Wang, F., and Chen, T. (2017).
- Efficient in vivo gene editing using ribonucleoproteins in skin stem cells of recessive dystrophic
 epidermolysis bullosa mouse model. Proc. Natl. Acad. Sci. *114*, 1660–1665.
- 45. Danner, E., Bashir, S., Yumlu, S., Wurst, W., Wefers, B., and Kühn, R. (2017). Control of gene
 editing by manipulation of DNA repair mechanisms. Mamm. Genome 28, 262–274.
- 628 46. Nelson, J.W., Randolph, P.B., Shen, S.P., Everette, K.A., Chen, P.J., Anzalone, A.V., An, M.,
- 629 Newby, G.A., Chen, J.C., Hsu, A., et al. (2022). Engineered pegRNAs improve prime editing
- 630 efficiency. Nat. Biotechnol. 40, 402–410.

- 631 47. Petri, K., Zhang, W., Ma, J., Schmidts, A., Lee, H., Horng, J.E., Kim, D.Y., Kurt, I.C., Clement,
- 632 K., Hsu, J.Y., et al. (2022). CRISPR prime editing with ribonucleoprotein complexes in zebrafish
- 633 and primary human cells. Nat. Biotechnol. 40, 189–193.
- 48. Huang, T.P., Newby, G.A., and Liu, D.R. (2021). Precision genome editing using cytosine and
 adenine base editors in mammalian cells. Nat. Protoc. *16*, 1089–1128.
- 49. Antoniou, P., Hardouin, G., Martinucci, P., Frati, G., Felix, T., Chalumeau, A., Fontana, L.,
 Martin, J., Masson, C., Brusson, M., et al. (2022). Base-editing-mediated dissection of a γ-globin
 cis-regulatory element for the therapeutic reactivation of fetal hemoglobin expression. Nat.
 Commun. *13*, 6618.
- 50. Kleinstiver, B.P., Sousa, A.A., Walton, R.T., Tak, Y.E., Hsu, J.Y., Clement, K., Welch, M.M.,
- Horng, J.E., Malagon-Lopez, J., Scarfò, I., et al. (2019). Engineered CRISPR–Cas12a variants
 with increased activities and improved targeting ranges for gene, epigenetic and base editing. Nat.
 Biotechnol. *37*, 276–282.
- 51. Strecker, J., Jones, S., Koopal, B., Schmid-Burgk, J., Zetsche, B., Gao, L., Makarova, K.S.,
 Koonin, E.V., and Zhang, F. (2019). Engineering of CRISPR-Cas12b for human genome editing.
 Nat. Commun. *10*, 212.
- 52. McMahon, M.A., Prakash, T.P., Cleveland, D.W., Bennett, C.F., and Rahdar, M. (2018).
 Chemically Modified Cpf1-CRISPR RNAs Mediate Efficient Genome Editing in Mammalian
 Cells. Mol. Ther. 26, 1228–1240.
- 53. Todaro, G.J., Green, H., and Goldberg, B.D. (1964). TRANSFORMATION OF PROPERTIES
 OF AN ESTABLISHED CELL LINE BY SV40 AND POLYOMA VIRUS. Proc. Natl. Acad.
 Sci. 51, 66–73.

54. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data.
Cell *177*, 1888-1902.

656

657 Figures legends

Figure 1: A novel monoallelic *KRT14* deletion causing a dominant form of Epidermolysis bullosa simplex

A) Clinical images of the patient EBS01: bullous lesions are present on the palmoplantar region. 660 Unremitting blister on the heel, dorsal medial part of the left foot and right peri-malleolar region are 661 shown in the panel below. **B**) Amino acid sequence of the wild-type and mutant K14. The invariant 662 663 residues are displayed in black boxes and the red box highlights the mutant region with the 7 amino acids deletion. The secondary structures are represented above the sequences as ribbons (helices) 664 and arrows (strands), respectively; blanks indicate unassigned regions. The Head and Tail domains 665 666 of KRT14 are represented as N-terminal and C-terminal grey boxes, respectively. Coil and Linker regions are represented as yellow and cyan boxes, respectively. C) Predicted three-dimensional 667 668 structures of human wild-type (yellow) and mutant (grey) K14 protein. The lack of the Linker 1 region, which is visible in the c475/495del21 protein, is circled in red. 669

670 Figure 2: Gene-editing of *KRT14* mutated allele

A) Representation of the EBS01 patient's genotype and allele profile of *KRT14* gene on human chromosome 17. The expanded section of the exon 1 illustrates the 21 nucleotides absent in *KRT14* exon 1 mutant allele (light blue). Mutant allele specific guide RNA is depicted in purple, straddling both the terminal sides of the deletion, and the -AGG- PAM sequence in red. **B**) Graphic representation of TIDE analysis. Both the wild-type and mutant allele were detected in the untreated sample (EBS01). In the treated sample (eEBS01) the mutant allele has been deleted by gene editing.

D) Histogram representation of NGS analysis of three technical replicates post gene editing. Edited 677 678 and not edited percentages on both wild-type and mutant alleles are calculated on the total number 679 of reads. D) PCR analysis of three technical replicates (EXP1, EXP2, EXP3). For each experiment, 680 a PCR product was amplified both from the EBS01 and eEBS01 keratinocytes' genome with primers 681 specific to anneal to wild-type (268 bp) and mutant alleles (266bp). E) Graphic representation of 682 editing derived InDels in mutant KRT14 allele (N= 3). F) NGS analysis of PCR products 683 surrounding the Cas9 target sites in the genome of eEBS01 showed a wide variety of Indel mutations 684 mediated by NHEJ at the targeted exon 1. The top sequence is the mutant allele unmodified 685 sequence, the dotted line represents the Cas9 cleavage upstream of the PAM.

686 Fig

Figure 3: Restoration of intermediate filaments

687 A) Immunofluorescent staining of K14 intermediate filaments in healthy donor derived 688 keratinocytes (NHEK), untreated and treated EBS01 keratinocytes (EBS01 and eEBS01 689 respectively). Nuclei are stained in blue by DAPI. Scale bar= 20µm. B) Immunofluorescent staining 690 of K14 filament before and after heat shock assay. At time 0 and after 15 minutes recovery from 691 heat shock, NEHK and eEBS01 do not show any changes in the organization of cytoplasmatic 692 network, whereas EBS01 keratinocytes display disruption of intermediate filament also in the 693 perinuclear region. After 60 minutes recovery, EBS01 intermediate filament aggregates revert to a 694 condition similar to that before the hear shock. Nuclei are stained in blue by DAPI. Scale bar= 695 20µm. C) Dispase-mediated epidermal sheet dissociation assay of NHEK, EBS01 and eEBS01 696 epidermal cultures before (C) and after (D) mechanical stress. No fragmentation was detected in 697 samples subjected to low-force orbital rotation. In contrast, high force stress induced fragmentation 698 of EBS01, but not NHEK and eEBS01 sheets (D). E) Number of fragments derived from each 699 sample has been counted using ImageJ software and unpaired t-test showed a statistically significant 700 P-value between EBS01 and eEBS01 number of fragments (N=3; asterisk indicate P-value<0,05).

701 Figure 4: 3D skin equivalents

(A) Haematoxylin and eosin staining of sections (7-µm thick) of decellularized dermal matrixes
(Dermis) seeded with normal human epidermal keratinocytes (NHEK). (B) The decellularized
dermal matrix was seeded with EBS01 or eEBS01 cells. Black arrowheads show blisters originated
in the epidermal basal layer only in EBS01 3D cultures. Scale bar, 100µm.

706 Figure 5: Genetic correction of epidermal stem cells

707 **A-B)** Uniform manifold approximation and projection (UMAP) of the scRNA-seq experiment. 708 EBS01 and eEBS01 keratinocytes mass cultures were profiled, integrated and classified in all their 709 clonogenic and differentiated clusters (H in red, M light blue, P grey, TD1 light brown, TD2 brown) 710 Distribution and extension of the clusters are comparable between the two samples. Feeder layer 711 derived fibroblast and low-quality keratinocytes are shown in light gray. C) Table shows the 712 percentages of cells in each EBS01 and eEBS01 clusters after scRNA-seq analysis. This quantitative 713 analysis demonstrates a comparable percentage between the treated and not treated samples. D) 714 Graphical representation of clonal analysis assay results. Clonal population percentages represent the 715 average value of Holoclones (H), Meroclones (M) and Paraclones (P) of two independent experiments 716 (EBS01: Holoclones 5%, Meroclones 52%, Paraclones 43%; eEBS01: Holoclones 5%, Meroclones 717 48%, Paraclones 47%). As shown, there are no differences in clonal populations between EBS01 and 718 eEBS01 keratinocytes. E) Representative indicator dishes for EBS01 and eEBS01 Holoclones, 719 Meroclones and Paraclones. F) PCR products of holoclones eEBS01's wild-type and mutant alleles. 720 Wild-type alleles result untouched, whereas all the mutated alleles display editing and lack of primer 721 annealing and amplification. Not treated EBS01 sample was used as control (CP) in which both the 722 wild-type and mutant allele are present and amplified.



B Keratin 14 amino acid sequence C Keratin 14 protein structure

	Head
XRT14_wt	1 10 20 20 10 10 60
KRT14_wt KRT14_c.475/4956ki21	
	Head
XRT14_wt	
KRT14_wt KRT14_0.475/495del21	A COLOROTORIST DI LE LO AGUITO DO LO MALLO COLORADO LA VOLTA VOLTA VOLTA VOLTA VOLTA VOLTA VOLTA VOLTA VOLTA VO
	Coil 1A Linker 1
KRTH_at	
KRT14_vt KRT14_c.475/495del21	NEW WITH WATER TRANSPORTED BY THE PARTY OF T
	Coil 1B
XRT14_mt	
KRT14_wt KRT14_c.475/495dwl21	
	Linker 12
XXT14,mt	210101010101010101010101010101010101010
KRT14_wt KRT14_c.475/495del21	OLKEELAVEN KENELEMIKALNO, VOLOTIVIEN DAAPSVOLOISILKEMIS OVERMAANNA SIREELAVENKEREMIKALNO, VOSOTIVIEN DAAPSVOLSIILKEMIS OVERMAAPSV
	Coil 2
ARTI4_art	Coil 2
XRTH4_wt KRT14_wt KRT14_c47566556621	
XRT14_ed KRT14_vd KRT14_c-475/495del21	Coil 2
ARTIN_M NRTIA_M NRTIA_c475H850H21 ARTIN_M	Coil 2
NTT14_wt NTT14_c4754855421 NTT14_c4754855421 NTT14_c4754855421	Coil 2 Coil 2
88714_e4 N9714_v4 N9714_c4754856421 88714_e4 K9714_c4754856421	Coil 2 Coil 2 Tail
ARTI4_et NRT14_v6 RRT14_c47548558821 ART14_c47548558821 RRT14_v6 RRT14_v6 RRT14_c47548558821 ART14_c47548558821	Coil 2 Coil 2 Tail

KRT14 wild-type RT14 C 475/495del21 mutant Linker 12 Coll 2 Tail Linker 1 Coll A

А





De Luca and colleagues propose an allele specific gene editing strategy for EBS01 patient affected by a rare skin disease inherited in an autosomal manner (EBS). This work aims to restore a normal cellular phenotype and a correct intermediate filament network in epidermal stem cells.

ournal Prendroo