

31. Vanacker J-M, Pettersson K, Gustafsson J-Å, Laudet V 1999 Transcriptional targets shared by estrogen receptor related receptors (ERRs) and estrogen receptor (ER) α but not by ER β . *EMBO J* 18: 4270–4279
32. Heard DJ, Norby PL, Holloway J, Vissing H 2000 Human ERR γ , a third member of the estrogen receptor-related receptor (ERR) subfamily of orphan nuclear receptors: tissue specific isoforms are expressed during development in the adult. *Mol Endocrinol* 14:382–392
33. Solomin L, Johansson CB, Zetterström RH, Bissonnette RP, Heyman RA, Olson L, Lendahl U, Friesen J, Perlmann T 1998 Retinoid-X receptor signalling in the developing spinal cord. *Nature* 395: 398–402
34. de Urquiza AM, Solomin L, Perlmann T 1999 Feedback-inducible nuclear-receptor-driven reporter gene expression in transgenic mice. *Proc Natl Acad Sci USA* 96: 13270–13275
35. Mader S, Leroy P, Chen JY, Chambon P 1993 Multiple parameters control the selectivity of nuclear receptors for their response elements. Selectivity and promiscuity in response element recognition by retinoic acid receptors and retinoid X receptors. *J Biol Chem* 268:591–600
36. McKnight SL, Kingsbury R 1982 Transcriptional control signals of a eukaryotic protein-coding gene. *Science* 217:316–324
37. Luckow B, Schutz G 1987 CAT constructions with multiple unique restriction sites for the functional analysis of eukaryotic promoters and regulatory elements. *Nucleic Acids Res* 15:5490
38. Wang Y, DeMayo FJ, Tsai SY, O'Malley BW 1997 Ligand-inducible and liver-specific target gene expression in transgenic mice. *Nat Biotechnol* 15:239–243
39. McKnight RA, Shamay A, Sankaran L, Wall RJ, Hennighausen L 1992 Matrix-attachment regions can impart position-independent regulation of a tissue-specific gene in transgenic mice. *Proc Natl Acad Sci USA* 89: 6943–6947
40. Sambrook J, Fritsch EF, Maniatis T 1989 *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Plainview, NY
41. Ma ZQ, Spreafico E, Pollio G, Santagati S, Conti E, Cattaneo E, Maggi A 1993 Activated estrogen receptor mediates growth arrest and differentiation of a neuroblastoma cell line. *Proc Natl Acad Sci USA* 90:3740–3744
42. Ciana P, Braliou GG, Demay FG, von Lindern M, Baretino D, Beug H, Stunnenberg HG 1998 Leukemic transformation by the v-ErbA oncoprotein entails constitutive binding to and repression of an erythroid enhancer in vivo. *EMBO J* 17:7382–7394
43. Bradford MM 1976 A rapid and sensitive methods for the quantitation of microgram quantities of proteins utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248–251

Erratum

In the article “Pleiotropic Effects of Substitutions of a Highly Conserved Leucine Residue in Transmembrane Helix III of the Human Lutropin/Choriogonadotropin Receptor with Respect to Constitutive Activation and Hormone Responsiveness” by Hiromitsu Shinozaki, Francesca Fanelli, Xuebo Liu, Julie Jaquette, Kazuto Nakamura, and Deborah L. Segaloff (*Molecular Endocrinology* **15**: 972–984, 2001), The heading for the right column of Table 1 should read “ ^{125}I -hCG Bound (ng/ 10^6 cells)”, not “ K_d (nM)”.