

This is the peer reviewed version of the following article:

Hamiltonian/Stroh formalism for anisotropic media with microstructure / Nobili, Andrea; Radi, Enrico. - In: PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY OF LONDON SERIES A: MATHEMATICAL PHYSICAL AND ENGINEERING SCIENCES. - ISSN 1364-503X. - 380:2231(2022), pp. N/A-N/A.
[10.1098/rsta.2021.0374]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

19/07/2024 02:45

(Article begins on next page)



Subject Areas:

74B05, 37K06

Keywords:

Stroh formalism, couple-stress, canonical system, variational principles, microstructure

¹

Author for correspondence:

Andrea Nobili

e-mail: andrea.nobili@unimore.it

Hamiltonian/Stroh formalism for anisotropic media with microstructure

Andrea Nobili^{1,3} and Enrico Radi^{2,3}

¹Department of Engineering Enzo Ferrari, University of Modena and Reggio Emilia, via Vivarelli 10, 41125 Modena, Italy

²Department of Sciences and Methods of Engineering, University of Modena and Reggio Emilia, via Amendola 2, 42122 Reggio Emilia, Italy

³Centre En&Tech, Tecnopolo, p.le Europa 1, 42124 Reggio Emilia, Italy

Moving from variational principles, we develop the Hamiltonian formalism for generally anisotropic microstructured materials, in an attempt to extend the celebrated Stroh formulation. Microstructure is expressed through the indeterminate (or Mindlin-Tiersten's) theory of couple stress elasticity. The resulting canonical formalism appears in the form of a Differential Algebraic system of Equations (DAE), which is then recast in purely differential form. This structure is due to the internal constraint that relates the micro- to the macro- rotation. The special situations of plain and anti-plane deformations are also developed and they both lead to a 7-dimensional coupled linear system of differential equations. In particular, the antiplane problem shows remarkable similarity with the theory of anisotropic plates, with which it shares the Lagrangian. Yet, unlike for plates, a classical Stroh formulation cannot be obtained, owing to the difference in the constitutive assumptions. Nonetheless, the canonical formalism brings new insight into the problem's structure and highlights important symmetry properties.

1. Introduction

The celebrated Stroh formalism [27] is a reformulation of the equations of elasticity which proves particularly useful for solving problems in plane anisotropic elastostatics [28]. These are reduced to a six-dimensional eigenvalue problem, of which they share all the relevant features. Besides, the method may be readily extended to steady-state elastodynamics [29]. In particular, the formalism is especially suited for discussing travelling wave propagation and it has gained considerable attention since it allows to prove existence of surface waves in generally anisotropic materials, a result that has eluded early researchers dealing with leaky waves [26]. **As an example, in [10] it is illustrated how to derive the Stroh form of the governing equations for localized edge vibration modes in a circular isotropic Kirchhoff-Love shell, and then use the impedance matrix to efficiently compute the real roots of the frequency equation.** It is now established that the secular equation governing surface waves is always real and that, whenever a surface wave exists, it is unique [3].

Only recently, it could be recognized that the essence of the formalism lies in its Hamiltonian nature, thereby a space variable is treated as a time-like coordinate [8]. Despite this knowledge having been already noted in passing [2], the realization of its full potential is a recent progress, which has been put to advantage to systematically generalise the formalism to more complicated situations. For example, it could bring constrained elasticity [5] and laminated plates [7,9] in Stroh form. **Also, it provided a basis to develop asymptotic reduced models for near-resonance disturbances in anisotropic media [11].** Indeed, in the absence of such a guidance, the right first-order formulation may only be developed by trial and error, such as it occurred for plates, see [8] and references therein. Furthermore, to the best of the authors' knowledge, no similar attempts appear in the literature in the direction of applying the Stroh formalism to complex media. As a remarkable exception, we mention the extension of the Stroh formalism to piezoelectricity in the form of a 8-dimensional eigenvalue problem [4], and to piezo-magneto-elastic or magneto-electro-elastic media, as a 10-dimensional eigenproblem [1]. Similarly, anisotropic micropolar elasticity is considered in [15], where a 14-dimensional system is found for generalized plane strain and 6-dimensional for plane strain. It is worth emphasizing that all such papers develop the Stroh formalism through ad-hoc assumptions, in a trial and error approach, and it is not entirely clear how conjugate variable have been selected (that is whether they are the conjugate momenta of the variational principle or a linear combination thereof). In similar fashion, we mention the extension of the Stroh formalism to self similar problems in elastodynamics by the Smirnov-Sobolev method [30]. Although moving from a different perspective, that is directed at the problem's solution rather than at elucidating the underlying variational structure, the paper reveals that a Stroh-like formalism still holds in dynamics.

In this paper, we apply the Hamiltonian formalism to systematically develop the canonical form of the governing equations of elastostatics for a microstructured medium. Microstructure is described in the spirit of the indeterminate (or Mindlin-Tiersten's) couple stress theory, which is a Cosserat theory wherein the couple stress is related to the gradient of the continuum (or macro) rotation [14,17,24]. Introduction of the microstructure has important downfalls on fracture mechanics [18,21] as well as body [12], Rayleigh [20,25], Stoneley [23] and Love [6] wave propagation, with important potential for applications [22]. It is therefore natural to investigate the variational structure of the associated Hamiltonian system. We show that the internal constraint relating the micro to the macro behaviour prevents reaching a classical Stroh formalism. This is especially surprising for antiplane problems, whose variational structure parallels that of anisotropic plates, which are amenable to the Stroh formalism. Still, the canonical system provides new insight into the fundamental structure of the equations.

49 2. Couple stress elasticity

Let us consider a Cartesian co-ordinate system (O, x_1, x_2, x_3) , with the triad of unit vectors $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3)$ directed alongside the relevant axes, attached to an elastic couple stress (CS) material. This is a polar material, for which, alongside the classical Cauchy force stress tensor \mathbf{s} , we define the couple stress tensor $\boldsymbol{\mu}$ (a table of symbols is presented in the Appendix). Across any surface of unit normal \mathbf{n} , an internal reduced couple vector acts, determined by the couple stress tensor as $\mathbf{q}_n^e = \boldsymbol{\mu}\mathbf{n}$. It is expedient to decompose the force stress tensor \mathbf{s} into its symmetric and skew-symmetric parts, respectively

$$\boldsymbol{\sigma} = \text{Sym } \mathbf{s} = \frac{1}{2} (\mathbf{s} + \mathbf{s}^T), \quad \text{and} \quad \boldsymbol{\tau} = \text{Skw } \mathbf{s} = \frac{1}{2} (\mathbf{s} - \mathbf{s}^T),$$

where the superscript T denotes the transposed tensor. Componentwise, we have $s_{ij} = \sigma_{ij} + \tau_{ij}$, with

$$\sigma_{ij} = s_{(ij)} = \frac{1}{2} (s_{ij} + s_{ji}), \quad \tau_{ij} = s_{\langle ij \rangle} = \frac{1}{2} (s_{ij} - s_{ji}).$$

In addition, the couple stress tensor $\boldsymbol{\mu}$ is split into its deviatoric and spherical parts

$$\boldsymbol{\mu} = \boldsymbol{\mu}^D + \boldsymbol{\mu}^S, \quad \boldsymbol{\mu}^S = \frac{1}{3} (\boldsymbol{\mu} \cdot \mathbf{1}) \mathbf{1},$$

50 where $\mathbf{1}$ is the rank 2 identity tensor and \cdot denotes the scalar product, i.e. componentwise,
51 $\mathbf{1}_{ij} = \delta_{ij}$, with δ_{ij} indicating Kronecker's delta symbol, while $\mathbf{A} \cdot \mathbf{B} = A_{ij} B_{ij}$ and Einstein's
52 summation convention over twice repeated subscripts is assumed.

53 In terms of kinematics, we introduce the displacement vector \mathbf{u} and the rotation vector
54 $\boldsymbol{\varphi}$. Unlike Cosserat micro-polar theories, for which displacements and micro-rotations are
55 independent fields, CS theory relates one to the other, through [14, Eqs.(4.9)]

$$\boldsymbol{\varphi} = \frac{1}{2} \text{curl } \mathbf{u}, \quad \Leftrightarrow \quad \varphi_i = \frac{1}{2} e_{ijk} u_{k,j}, \quad (2.1)$$

56 where \mathbf{E} is the rank-3 permutation (Levi-Civita) tensor, whose components are denoted by
57 e_{ijk} , and a subscript comma denotes partial differentiation, e.g. $(\text{grad } u)_{kj} = u_{k,j} = \partial u_k / \partial x_j$. It
58 follows that $\boldsymbol{\varphi}$ is solenoidal

$$\text{div } \boldsymbol{\varphi} = 0, \quad \Leftrightarrow \quad \varphi_{j,j} = 0. \quad (2.2)$$

As in CE, we define the linear strain tensor

$$\boldsymbol{\varepsilon} = \text{Sym grad } \mathbf{u}, \quad \Leftrightarrow \quad \varepsilon_{ij} = u_{(i,j)}.$$

For a linear elastic anisotropic material, we have

$$\boldsymbol{\sigma} = \mathbb{C}\boldsymbol{\varepsilon},$$

where \mathbb{C} is the rank 4 tensor of elastic moduli, i.e. $\mathbb{C}_{ijkl} = c_{ijkl}$, possessing the major and the minor symmetry property, i.e. $c_{ijkl} = c_{klij}$ and $c_{ijkl} = c_{jikl} = c_{ijlk}$, respectively. By $\mathbb{C}\boldsymbol{\varepsilon}$ we mean the rank 2 tensor obtained by double summation over the last pair of indices: $c_{ijkl}\varepsilon_{kl}$. For isotropic materials, we have

$$\mathbb{C} = 2G\mathbb{I} + \lambda\mathbf{1} \otimes \mathbf{1}, \quad \Leftrightarrow \quad c_{ijkl} = 2G\delta_{ik}\delta_{jl} + \lambda\delta_{kl}\delta_{ij},$$

59 where \mathbb{I} is the rank 4 identity tensor and λ and $G > 0$ are Lamé moduli. Here, the dyadic product
60 is introduced for rank 2 tensors such that $(\mathbf{A} \otimes \mathbf{B})\mathbf{C} = (\mathbf{B} \cdot \mathbf{C})\mathbf{A}$, for any triple of rank 2 tensors
61 \mathbf{A} , \mathbf{B} and \mathbf{C} .

62 We introduce the *torsion-flexure (wryness or curvature) tensor* as the gradient of the rotation field

$$\boldsymbol{\chi} = \text{grad } \boldsymbol{\varphi}, \quad \Leftrightarrow \quad \chi_{ij} = \varphi_{i,j}. \quad (2.3)$$

63 In light of the connection (2.2) and recalling that $\text{tr grad} \equiv \text{div}$, it is seen that $\boldsymbol{\chi}$ is purely deviatoric,
64 i.e. $\boldsymbol{\chi} = \boldsymbol{\chi}^D$. Here, the divergence operator on rank 2 tensors operates on the *second* component,
65 i.e. $(\text{div } \boldsymbol{s})_i = s_{ij,j}$.

For a linear elastic CS material, we have the constitutive law

$$\boldsymbol{\mu} = \ell^2 \mathbb{G} \boldsymbol{\chi},$$

66 where $\ell > 0$ is a characteristic length related to the microstructure and \mathbb{G} is the rank 4 tensor
 67 of couple stress moduli possessing the major symmetry property $g_{ijkl} = g_{klij}$. Immediately, it
 68 appears that, to any effect, $\boldsymbol{\mu}$ may be replaced by $\boldsymbol{\mu}^D$ and in fact, the CS theory is named
 69 indeterminate after the observation that the first invariant of the couple stress tensor, i.e. $\text{tr } \boldsymbol{\mu} =$
 70 $\boldsymbol{\mu} \cdot \mathbf{1} = \mu_{11} + \mu_{22} + \mu_{33}$, rests indeterminate. Therefore, we are free to set it equal to zero without
 71 any loss of generality, i.e. $g_{iikl} = g_{kl ii} = 0$. In the following, for the sake of brevity, we shall write
 72 $\boldsymbol{\mu}$ with the understanding that $\boldsymbol{\mu}^D$ is meant. For isotropic materials, we have

$$\boldsymbol{\mu} = 2G\ell^2 (\boldsymbol{\chi} + \eta \boldsymbol{\chi}^T) \Leftrightarrow g_{ijkl} = 2G (\delta_{jl} \delta_{ik} + \eta \delta_{jk} \delta_{il}), \quad (2.4)$$

73 where $-1 < \eta < 1$ is a dimensionless number similar to Poisson's ratio.

The equilibrium equations, in the absence of body forces, read [19, Eq.(2)]

$$\text{div } \boldsymbol{s} = \boldsymbol{o}, \quad (2.5a)$$

$$\text{axial } \boldsymbol{\tau} - \frac{1}{2} \text{div } \boldsymbol{\mu} = \boldsymbol{o}, \quad (2.5b)$$

74 where $\text{axial } \boldsymbol{\tau} = \frac{1}{2} \mathbf{E} \boldsymbol{\tau}$, i.e. $(\text{axial } \boldsymbol{\tau})_i = \frac{1}{2} e_{ijk} \tau_{jk}$, denotes the axial vector attached to a skew-
 75 symmetric tensor. It is observed that, introducing the curl of a tensor as $(\text{curl } \mathbf{W})_{ij} = e_{jkl} W_{il,k}$, it
 76 can be easily proved that

$$2 \text{ axial curl } \mathbf{W} = \text{div} \left[(\text{tr } \mathbf{W}) \mathbf{1} - \mathbf{W}^T \right].$$

77 Consequently, Eqs.(2.5) admit a solution in terms of the Günther stress tensors \mathbf{W} and \mathbf{Z} [13,15]

$$\boldsymbol{s} = \text{curl } \mathbf{W}, \quad \boldsymbol{\mu} = \text{curl } \mathbf{Z} + (\text{tr } \mathbf{W}) \mathbf{1} - \mathbf{W}^T. \quad (2.6)$$

78 However, as pointed out in [15], this representation leads to a formalism that is not closed.
 79 Through the inverse formula

$$\boldsymbol{\tau} = \mathbf{E} \text{ axial } \boldsymbol{\tau}, \quad \Leftrightarrow \tau_{ij} = e_{ijk} (\text{axial } \boldsymbol{\tau})_k, \quad (2.7)$$

80 Eq.(2.5b) may be solved for $\boldsymbol{\tau}$

$$\boldsymbol{\tau} = \frac{1}{2} \mathbf{E} \text{div } \boldsymbol{\mu} = - \text{Skw curl } \boldsymbol{\mu}^T, \quad (2.8)$$

81 that in components read $\tau_{ij} = \frac{1}{2} e_{ijk} \mu_{kl,l}$. Hence, the skew-symmetric part of the force stress
 82 tensor \boldsymbol{s} is determined by rotational equilibrium. Clearly, CE is retrieved taking $\ell = 0$, for then
 83 $\boldsymbol{\mu} = \boldsymbol{\tau} = \mathbf{0}$.

84 We now write the total energy in the sense of Eshelby [8]

$$\mathcal{L} = \int_{\mathcal{B}} \left[\frac{1}{2} \boldsymbol{\sigma} \cdot \boldsymbol{\varepsilon} + \frac{1}{2} \boldsymbol{\mu} \cdot \boldsymbol{\chi} - \boldsymbol{p} \cdot \left(\boldsymbol{\varphi} - \frac{1}{2} \text{curl } \boldsymbol{u} \right) \right] dV - \int_{\partial \mathcal{B}} (\boldsymbol{t}_n \cdot \boldsymbol{u} + \boldsymbol{q}_n \cdot \boldsymbol{\varphi}) dA, \quad (2.9)$$

having introduced the Lagrangian multiplier vector $\boldsymbol{p} = [p_i]$ to account for the constraint (2.1)
 and being \boldsymbol{n} the unit normal, in the outwards direction, to the surface element dA . Besides, we
 let the normal tensor $\mathfrak{N} = \boldsymbol{n} \otimes \boldsymbol{n}$, the projector tensor $\mathfrak{P} = \mathbf{1} - \mathfrak{N}$, and the skew tensor $\boldsymbol{P} = \mathbf{E} \boldsymbol{p}$
 associated with the vector \boldsymbol{p} thought of as an axial vector. The prescribed boundary force and
 couple vector are given by

$$\boldsymbol{t}_n = \boldsymbol{t}_n^e + \boldsymbol{\tau} \boldsymbol{n} - \frac{1}{2} \boldsymbol{n} \times \text{grad } \mu_{nn}, \quad \boldsymbol{q}_n = \mathfrak{P} \boldsymbol{q}_n^e = \boldsymbol{q}_n^e - \mu_{nn} \boldsymbol{n},$$

85 being $\boldsymbol{t}_n^e = \boldsymbol{\sigma} \boldsymbol{n}$ and $\boldsymbol{q}_n^e = \boldsymbol{\mu} \boldsymbol{n}$ the "elastic" part of the force and couple stress vector and $\mu_{nn} =$
 86 $\mathfrak{N} \cdot \boldsymbol{\mu} = \boldsymbol{n} \cdot \boldsymbol{\mu} \boldsymbol{n}$ the normal part of the couple stress. We observe that the surface integral in (2.9)

87 may be equivalently rewritten as

$$-\int_{\partial\mathcal{B}} [(\mathbf{t}_n^e + \boldsymbol{\tau}\mathbf{n}) \cdot \mathbf{u} + \mathbf{q}_n^e \cdot \boldsymbol{\varphi}] dA.$$

Indeed, recalling the vector identities

$$\operatorname{div}(\mathbf{a} \times \mathbf{b}) = \mathbf{b} \cdot \operatorname{curl} \mathbf{a} - \mathbf{a} \cdot \operatorname{curl} \mathbf{b}, \quad (2.10a)$$

$$\operatorname{div}(\phi \mathbf{b}) = \operatorname{grad} \phi \cdot \mathbf{b} + \phi \operatorname{div} \mathbf{b}, \quad (2.10b)$$

$$\operatorname{curl} \operatorname{grad} \phi = \mathbf{o}, \quad (2.10c)$$

and making use of the divergence theorem, it is easily proved that

$$\begin{aligned} -\frac{1}{2} \int_{\partial\mathcal{B}} (\mathbf{n} \times \operatorname{grad} \mu_{nn} \cdot \mathbf{u}) dA &= \frac{1}{2} \int_{\partial\mathcal{B}} (\mathbf{u} \times \operatorname{grad} \mu_{nn} \cdot \mathbf{n}) dA \\ &= \frac{1}{2} \int_{\mathcal{B}} \operatorname{div}(\mathbf{u} \times \operatorname{grad} \mu_{nn}) dV = \frac{1}{2} \int_{\mathcal{B}} (\operatorname{grad} \mu_{nn} \cdot \operatorname{curl} \mathbf{u}) dV \\ &= \int_{\mathcal{B}} (\operatorname{grad} \mu_{nn} \cdot \boldsymbol{\varphi}) dV = \int_{\mathcal{B}} \operatorname{div}(\mu_{nn} \boldsymbol{\varphi}) dV = \int_{\partial\mathcal{B}} (\boldsymbol{\varphi} \cdot \mu_{nn} \mathbf{n}) dA, \end{aligned}$$

88 having made use of Eq.(2.2). Therefore, $-\frac{1}{2} \mathbf{n} \times \operatorname{grad} \mu_{nn}$ in \mathbf{t}_n cancels out the term $\mu_{nn} \mathbf{n}$ in \mathbf{q}_n .

89 By the divergence theorem and making use of the equilibrium equations (2.5), we get

$$\mathcal{L} = - \int_{\mathcal{B}} L dV,$$

90 having introduced the Lagrangian density function L

$$L(\operatorname{grad} \mathbf{u}, \boldsymbol{\varphi}, \operatorname{grad} \boldsymbol{\varphi}, \mathbf{p}) = \frac{1}{2} \boldsymbol{\sigma} \cdot \boldsymbol{\varepsilon} + \frac{1}{2} \boldsymbol{\mu} \cdot \boldsymbol{\chi} + \mathbf{p} \cdot (\boldsymbol{\varphi} - \frac{1}{2} \operatorname{curl} \mathbf{u}), \quad (2.11)$$

that, component-wise, reads

$$L(\operatorname{grad} \mathbf{u}, \boldsymbol{\varphi}, \operatorname{grad} \boldsymbol{\varphi}, \mathbf{p}) = \frac{1}{2} \sigma_{ij} u_{(i,j)} + \frac{1}{2} \mu_{ij} \varphi_{i,j} + p_i (\varphi_i - \frac{1}{2} e_{ijk} u_{k,j}).$$

The Euler-Lagrange (E-L) equations are

$$-\frac{\partial}{\partial x_j} \frac{\partial L}{\partial u_{i,j}} + \frac{\partial L}{\partial u_i} = -\sigma_{ij,j} + \frac{1}{2} p_{m,j} e_{mji} = 0, \quad (2.12a)$$

$$-\frac{\partial}{\partial x_j} \frac{\partial L}{\partial \varphi_{i,j}} + \frac{\partial L}{\partial \varphi_i} = -\mu_{ij,j} + p_i = 0, \quad (2.12b)$$

$$\frac{\partial L}{\partial p_i} = \varphi_i - \frac{1}{2} e_{ijk} u_{k,j} = 0, \quad (2.12c)$$

that, recalling (2.7), amount to

$$-\operatorname{div}(\boldsymbol{\sigma} + \frac{1}{2} \mathbf{P}) = 0, \quad (2.13a)$$

$$\mathbf{p} - \operatorname{div} \boldsymbol{\mu} = 0, \quad (2.13b)$$

91 and the constraint (2.1). In particular, looking at (2.5b, 2.13b), we are able to give a physical

92 meaning to the Lagrange multiplier \mathbf{p}

$$\mathbf{p} = 2 \text{ axial } \boldsymbol{\tau} = 2(\tau_{23}, \tau_{31}, \tau_{12}) \quad (2.14)$$

93 wherefrom, $\frac{1}{2} \mathbf{P} = \boldsymbol{\tau}$ and Eqs.(2.13) correspond to (2.5).

94 Special care is required when dealing with the Lagrange multiplier \mathbf{p} . Indeed, acknowledging
 95 (2.2), it can be proved that $\operatorname{div} \mathbf{p} = 0$, whence

$$\mathbf{p} = -\operatorname{curl} \mathbf{h},$$

with the gauge relation $\operatorname{div} \mathbf{h} = 0$. Consequently, making use of the vector identities (2.10) alongside

$$\operatorname{curl} \mathbf{h} \cdot \operatorname{curl} \mathbf{u} = 2 \operatorname{grad} u \cdot \operatorname{Skw} \operatorname{grad} \mathbf{h},$$

we may equivalently write

$$L(\operatorname{grad} \mathbf{u}, \boldsymbol{\varphi}, \operatorname{grad} \boldsymbol{\varphi}, \mathbf{h}) = \frac{1}{2} \sigma_{ij} u_{(i,j)} + \frac{1}{2} \mu_{ij} \varphi_{i,j} - e_{kji} h_k \varphi_{i,j} + \frac{1}{2} u_{i,j} (h_{i,j} - h_{j,i}),$$

96 up to boundary terms. This observation will be used in Sec. 4 when seeking expressions for the
 97 Lagrange multiplier.

98 3. Plane strain

99 We now consider plane-strain conditions [24]

$$u_3 = \varphi_1 = \varphi_2 = 0,$$

100 by which there is no dependence of the deformation on x_3 . Thus, $\boldsymbol{\varepsilon} = \operatorname{grad}_2 \mathbf{u}$, and grad_2 is the
 101 gradient operator restricted to the co-ordinates x_α , Greek subscripts ranging in the set $\{1, 2\}$.
 102 Similarly, the constraint (2.1) reduces to the single component

$$\varphi_3 = \frac{1}{2} (u_{2,1} - u_{1,2}) = \frac{1}{2} (\mathbf{u}_{,1} \cdot \mathbf{e}_2 - \mathbf{u}_{,2} \cdot \mathbf{e}_1), \quad (3.1)$$

that immediately satisfies (2.2), while the wryness tensor (2.3) becomes

$$\boldsymbol{\chi} = \operatorname{grad}_2 \varphi_3 = \varphi_{3,1} \mathbf{e}_3 \otimes \mathbf{e}_1 + \varphi_{3,2} \mathbf{e}_3 \otimes \mathbf{e}_2,$$

103 having introduced the dyadic operator for vectors such that $(\mathbf{a} \otimes \mathbf{b})\mathbf{c} = (\mathbf{b} \cdot \mathbf{c})\mathbf{a}$, for any triple of
 104 vectors \mathbf{a} , \mathbf{b} and \mathbf{c} .

105 Within a Stroh formalism, we define the usual rank 2 matrices

$$Q_{\alpha\beta} = c_{\alpha 1 \beta 1}, \quad R_{\alpha\beta} = c_{\alpha 1 \beta 2}, \quad T_{\alpha\beta} = c_{\alpha 2 \beta 2}, \quad (3.2)$$

106 where \mathbf{Q} and \mathbf{T} are *symmetric*, alongside the symmetric matrix

$$U_{\alpha\beta} = \ell^2 g_{3\alpha 3\beta}. \quad (3.3)$$

107 Through these, we can define the elastic part of the reduced traction vectors (in the plane of strain)

$$\mathbf{t}_1^e = \mathbf{Q}\mathbf{u}_{,1} + \mathbf{R}\mathbf{u}_{,2}, \quad \mathbf{t}_2^e = \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{T}\mathbf{u}_{,2}, \quad (3.4)$$

108 and the *out-of-plane component* of the reduced couple-stress vectors \mathbf{q}_1^e and \mathbf{q}_2^e , respectively

$$\mathbf{q}_1^e = \mathbf{q}_1^e \cdot \mathbf{e}_3 = U_{11} \varphi_{3,1} + U_{12} \varphi_{3,2}, \quad \mathbf{q}_2^e = \mathbf{q}_2^e \cdot \mathbf{e}_3 = U_{21} \varphi_{3,1} + U_{22} \varphi_{3,2}. \quad (3.5)$$

109 We observe that

$$\boldsymbol{\sigma} \cdot \boldsymbol{\varepsilon} = \mathbf{t}_1^e \cdot \mathbf{u}_{,1} + \mathbf{t}_2^e \cdot \mathbf{u}_{,2},$$

110 and

$$\boldsymbol{\mu} \cdot \boldsymbol{\chi} = \boldsymbol{\mu} \cdot \operatorname{grad}_2 \varphi_3 = \mu_{31} \chi_{31} + \mu_{32} \chi_{32} = q_{13}^e \varphi_{3,1} + q_{23}^e \varphi_{3,2}.$$

Besides, specializing (2.13b), the Lagrangian multiplier \mathbf{p} has the single non-zero component

$$p_3 = \mu_{31,1} + \mu_{32,2} = q_{13,1}^e + q_{23,2}^e,$$

111 and from (2.7, 2.14) we get the single non-zero component in the skew part of the force stress
 112 tensor

$$\tau_{12} = \frac{1}{2} P_{12} = \frac{1}{2} p_3 = -\tau_{21}. \quad (3.6)$$

Then, the Lagrangian density (2.11) becomes

$$L(\mathbf{u}_{,1}, \mathbf{u}_{,2}, \varphi_3, \varphi_{3,1}, \varphi_{3,2}, p_3) = \frac{1}{2} [\mathbf{t}_1^e \cdot \mathbf{u}_{,1} + \mathbf{t}_2^e \cdot \mathbf{u}_{,2} + q_{13}^e \varphi_{3,1} + q_{23}^e \varphi_{3,2}] + p_3 \varphi_3 - \frac{1}{2} p_3 (\mathbf{u}_{,1} \cdot \mathbf{e}_2 - \mathbf{u}_{,2} \cdot \mathbf{e}_1),$$

where it is understood that the scalar product now carries over 2 components only. The conjugate momenta are

$$\frac{\partial L}{\partial \mathbf{u}_{,1}} = \mathbf{t}_1^e - \frac{1}{2} p_3 \mathbf{e}_2 = \mathbf{t}_1^e + \tau_{21} \mathbf{e}_2 = \mathbf{s}_1, \quad (3.7a)$$

$$\frac{\partial L}{\partial \mathbf{u}_{,2}} = \mathbf{t}_2^e + \frac{1}{2} p_3 \mathbf{e}_1 = \mathbf{t}_2^e + \tau_{12} \mathbf{e}_1 = \mathbf{s}_2, \quad (3.7b)$$

$$\frac{\partial L}{\partial \varphi_{3,1}} = q_{13}^e, \quad (3.7c)$$

$$\frac{\partial L}{\partial \varphi_{3,2}} = q_{23}^e, \quad (3.7d)$$

whereupon the Lagrangian may be rewritten as

$$L(\mathbf{u}_{,1}, \mathbf{u}_{,2}, \varphi_3, \varphi_{3,1}, \varphi_{3,2}, p_3) = \frac{1}{2} \mathbf{u}_{,1} \cdot \mathbf{Q} \mathbf{u}_{,1} + \mathbf{u}_{,1} \cdot \mathbf{R} \mathbf{u}_{,2} + \frac{1}{2} \mathbf{u}_{,2} \cdot \mathbf{T} \mathbf{u}_{,2} + \frac{1}{2} U_{11} \varphi_{3,1}^2 + U_{12} \varphi_{3,1} \varphi_{3,2} + \frac{1}{2} U_{22} \varphi_{3,2}^2 + p_3 \varphi_3 + \frac{1}{2} p_3 (\mathbf{u}_{,2} \cdot \mathbf{e}_1 - \mathbf{u}_{,1} \cdot \mathbf{e}_2).$$

We now come to an important juncture and treat either co-ordinate as a time-like variable, say x_2 to fix ideas. Consequently, we introduce the Legendre transformation

$$H(\mathbf{u}_{,1}, \mathbf{s}_2, \varphi_3, \varphi_{3,1}, q_{23}^e, p_3) = \mathbf{s}_2 \cdot \mathbf{u}_{,2} + q_{23}^e \varphi_{3,2} - L = \frac{1}{2} \mathbf{s}_2 \cdot \mathbf{u}_{,2} + \frac{1}{2} q_{23}^e \varphi_{3,2} - \frac{1}{2} \mathbf{t}_1^e \cdot \mathbf{u}_{,1} - \frac{1}{2} q_{13}^e \varphi_{3,1} - p_3 \varphi_3 - \frac{1}{4} p_3 (\mathbf{u}_{,2} \cdot \mathbf{e}_1 - 2 \mathbf{u}_{,1} \cdot \mathbf{e}_2),$$

113 provided that we write $\mathbf{u}_{,2}$ in terms of \mathbf{s}_2 by (3.7b) and $\varphi_{3,2}$ in terms of q_{23}^e by (3.7d). For the
114 former, making use of (3.4), we get

$$\mathbf{u}_{,2} = \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right), \quad (3.8)$$

115 assuming that \mathbf{T} is invertible, while for the latter

$$\varphi_{3,2} = U_{22}^{-1} (q_{23}^e - U_{21} \varphi_{3,1}), \quad (3.9)$$

assuming $U_{22} \neq 0$. Therefore, we can write the Hamiltonian density function (whose arguments are omitted for brevity)

$$H = \frac{1}{2} \mathbf{s}_2 \cdot \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) + \frac{1}{2} q_{23}^e \frac{q_{23}^e - U_{21} \varphi_{3,1}}{U_{22}} - \frac{1}{2} \mathbf{u}_{,1} \cdot \mathbf{Q} \mathbf{u}_{,1} - \frac{1}{2} \mathbf{u}_{,1} \cdot \mathbf{R} \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) - \frac{1}{2} U_{11} \varphi_{3,1}^2 - \frac{1}{2} U_{12} \varphi_{3,1} \frac{q_{23}^e - U_{21} \varphi_{3,1}}{U_{22}} - p_3 \varphi_3 - \frac{1}{4} p_3 \left[\mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) \cdot \mathbf{e}_1 - 2 \mathbf{u}_{,1} \cdot \mathbf{e}_2 \right],$$

which reduces to

$$H = \frac{1}{2} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) \cdot \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) + \frac{1}{2} \frac{(q_{23}^e - U_{21} \varphi_{3,1})^2}{U_{22}} - \frac{1}{2} \mathbf{u}_{,1} \cdot \mathbf{Q} \mathbf{u}_{,1} - \frac{1}{2} U_{11} \varphi_{3,1}^2 - p_3 \left(\varphi_3 - \frac{1}{2} \mathbf{u}_{,1} \cdot \mathbf{e}_2 \right). \quad (3.10)$$

116 Indeed, letting the generalized co-ordinate vector $\bar{\mathbf{q}} = (\mathbf{u}, \varphi_3)$ and the conjugate momenta $\bar{\mathbf{p}} =$
 117 (\mathbf{s}_2, q_{23}^e) , the first set of canonical equations

$$\frac{\delta H}{\delta \bar{\mathbf{p}}} = \dot{\bar{\mathbf{q}}} \quad (3.11)$$

is

$$\frac{\delta H}{\delta \mathbf{s}_2} = \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) = \mathbf{u}_{,2},$$

corresponding to (3.8), and

$$\frac{\delta H}{\delta q_{23}^e} = \frac{(q_{23}^e - U_{21} \varphi_{3,1})}{U_{22}} = \varphi_{3,2},$$

118 that matches (3.9). Likewise, the second set of canonical equations

$$\frac{\delta H}{\delta \bar{\mathbf{q}}} = -\dot{\bar{\mathbf{p}}}, \quad (3.12)$$

119 yields

$$\frac{\delta H}{\delta \mathbf{u}} = -\frac{\partial}{\partial x_1} \frac{\partial H}{\partial \mathbf{u}_{,1}} = \left[\mathbf{R} \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) + \mathbf{Q} \mathbf{u}_{,1} - \frac{1}{2} p_3 \mathbf{e}_2 \right]_{,1} = -\mathbf{s}_{2,2} \quad (3.13)$$

120 and

$$\frac{\delta H}{\delta \varphi_3} = -\frac{\partial}{\partial x_1} \frac{\partial H}{\partial \varphi_{3,1}} + \frac{\partial H}{\partial \varphi_3} = \left[\frac{U_{21}}{U_{22}} (q_{23}^e - U_{21} \varphi_{3,1}) + U_{11} \varphi_{3,1} \right]_{,1} - p_3 = -q_{23,2}^e. \quad (3.14)$$

121 Indeed, in light of Eqs.(3.6,3.7a), Eq.(3.13) is simply

$$(\mathbf{t}_1^e + \tau_{21} \mathbf{e}_2)_{,1} + \mathbf{s}_{2,2} = \mathbf{s}_{1,1} + \mathbf{s}_{2,2} = \mathbf{o},$$

122 that corresponds to (2.5a). Similarly, making use of (3.5,3.6) and (3.9), Eq.(3.14) may be rewritten
 123 as

$$q_{13,1}^e + q_{23,2}^e - 2\tau_{12} = 0,$$

which amounts to the rotational equilibrium (2.8). Thus, for a homogeneous material, we get

$$\mathbf{u}_{,2} = -\mathbf{T}^{-1} \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{T}^{-1} \phi_{,1} - \Lambda_{,1} \mathbf{T}^{-1} \mathbf{e}_1, \quad (3.15a)$$

$$\phi_{,2} = \left(\mathbf{R} \mathbf{T}^{-1} \mathbf{R}^T - \mathbf{Q} \right) \mathbf{u}_{,1} - \mathbf{R} \mathbf{T}^{-1} \phi_{,1} + \Lambda_{,1} \left(\mathbf{e}_2 + \mathbf{R} \mathbf{T}^{-1} \mathbf{e}_1 \right), \quad (3.15b)$$

$$\varphi_{3,2} = -\frac{U_{21}}{U_{22}} \varphi_{3,1} + U_{22}^{-1} \Phi_{,1}, \quad (3.15c)$$

$$\Phi_{,2} = \left(\frac{U_{21}^2}{U_{22}} - U_{11} \right) \varphi_{3,1} - \frac{U_{21}}{U_{22}} \Phi_{,1} + 2\Lambda, \quad (3.15d)$$

124 having let the stress functions $\phi = \int^{x_1} \mathbf{s}_2 d\xi_1$, $\Lambda = \frac{1}{2} \int^{x_1} p_3 d\xi_1 = \int^{x_1} \tau_{12} d\xi_1$ and $\Phi = \int^{x_1} q_{23}^e d\xi_1$,
 125 that are defined up to an arbitrary function of x_2 . In the spirit of considering x_2 a time-like
 126 variable, this is a system of ODEs in canonical form. We also observe a connection with Günther
 127 tensor potentials (2.6), namely $\phi_\alpha = -W_{\alpha 3}$ and $\Phi_{,1} = -Z_{33,1} + \phi_2$, cf. [15, Eqs(2.38-40)].

Now, we only need to dispose of the so-far undetermined Lagrange multiplier Λ . For this, we
 take the scalar product of (3.8) with \mathbf{e}_1 and make use of the constraint (3.1)

$$\mathbf{u}_{,2} \cdot \mathbf{e}_1 = -2\varphi_3 + \mathbf{u}_{,1} \cdot \mathbf{e}_2 = \mathbf{T}^{-1} \left(-\frac{1}{2} p_3 \mathbf{e}_1 - \mathbf{R}^T \mathbf{u}_{,1} + \mathbf{s}_2 \right) \cdot \mathbf{e}_1,$$

128 whereupon we find

$$\Lambda_{,1} = \zeta \left(\phi_{,1} \cdot \mathbf{f}_1 - \mathbf{u}_{,1} \cdot \mathbf{f}_2 + 2\varphi_3 \right), \quad (3.16)$$

having let the shorthand vectors

$$\mathbf{f}_1 = \mathbf{T}^{-1} \mathbf{e}_1, \quad \text{and} \quad \mathbf{f}_2 = \mathbf{e}_2 + \mathbf{R} \mathbf{f}_1,$$

129 and $\zeta^{-1} = \mathbf{T}^{-1} \mathbf{e}_1 \cdot \mathbf{e}_1 = \mathbf{f}_1 \cdot \mathbf{e}_1$. The connection (3.16) shows that, similarly to classical
 130 incompressible elasticity, the Lagrange multiplier is determined by an algebraic relation where

131 no x_2 derivative appears. Consequently, the governing equations (3.15,3.16) form a system
 132 of Differential Algebraic Equations (DAEs) in *semi-explicit form*. However, in contrast to
 133 incompressible elasticity, Eqs.(3.15d) and (3.16) indicate that a Stroh classical formulation, where
 134 the unknown vectors only appear in differential form, is not accessible.

We now show that this system of DAEs has *differential index 1*. For the sake of convenience, we let the matrices

$$\mathbf{N}_1 = -\mathbf{T}^{-1}\mathbf{R}^T, \quad (3.17a)$$

$$\mathbf{N}_2 = \mathbf{R}\mathbf{T}^{-1}\mathbf{R}^T - \mathbf{Q} = -\mathbf{R}\mathbf{N}_1 - \mathbf{Q} = \mathbf{N}_2^T, \quad (3.17b)$$

$$\mathbf{N}_3 = U_{22}^{-1} \begin{bmatrix} -U_{21} & 1 \\ U_{21}^2 - U_{22}U_{11} & -U_{21} \end{bmatrix}. \quad (3.17c)$$

Differentiating (3.16) with respect to x_2 and then making use of (3.15c), we get

$$\zeta^{-1}A_{,2} = \phi_{,2} \cdot \mathbf{f}_1 - \mathbf{u}_{,2} \cdot \mathbf{f}_2 + 2U_{22}^{-1}(-U_{21}\varphi_3 + \Phi),$$

and, by (3.15a,3.15b),

$$\begin{aligned} \zeta^{-1}A_{,2} = & \left(\mathbf{N}_2\mathbf{u}_{,1} + \mathbf{N}_1^T\phi_{,1} + A_{,1}\mathbf{f}_2 \right) \cdot \mathbf{f}_1 \\ & - \left(\mathbf{N}_1\mathbf{u}_{,1} + \mathbf{T}^{-1}\phi_{,1} - A_{,1}\mathbf{f}_1 \right) \cdot \mathbf{f}_2 + 2U_{22}^{-1}(-U_{21}\varphi_3 + \Phi), \end{aligned}$$

that provides the evolution equation for A

$$\begin{aligned} \zeta^{-1}A_{,2} = & \left(\mathbf{N}_2\mathbf{f}_1 - \mathbf{N}_1^T\mathbf{f}_2 \right) \cdot \mathbf{u}_{,1} + \left(\mathbf{N}_1\mathbf{f}_1 - \mathbf{T}^{-1}\mathbf{f}_2 \right) \cdot \phi_{,1} + 2A_{,1}\mathbf{f}_1 \cdot \mathbf{f}_2 \\ & + 2U_{22}^{-1}(-U_{21}\varphi_3 + \Phi). \end{aligned}$$

For better understanding, we adopt the convention that vectors are columns and their transpose are rows. Thus, letting the 7-component row vector

$$\boldsymbol{\xi}^T = [\mathbf{u}^T, \phi^T, A, \varphi_3, \Phi],$$

135 we finally obtain the system of first order linear PDEs

$$\frac{d\boldsymbol{\xi}}{dx_2} = \mathbf{N} \frac{d\boldsymbol{\xi}}{dx_1} + \mathbf{b}, \quad (3.18)$$

136 where we have let the 7 by 7 Stroh matrix

$$\mathbf{N} = \begin{bmatrix} \mathbf{N}_{5 \times 5} & \mathbf{O} \\ \mathbf{O} & \mathbf{N}_3 \end{bmatrix}, \quad (3.19)$$

with the 5 by 5 matrix

$$\mathbf{N}_{5 \times 5} = \begin{bmatrix} \mathbf{N}_1 & \mathbf{T}^{-1} & -\mathbf{f}_1 \\ \mathbf{N}_2 & \mathbf{N}_1^T & \mathbf{f}_2 \\ \zeta \left(\mathbf{N}_2\mathbf{f}_1 - \mathbf{N}_1^T\mathbf{f}_2 \right)^T & \zeta \left(\mathbf{N}_1\mathbf{f}_1 - \mathbf{T}^{-1}\mathbf{f}_2 \right)^T & 2\zeta\mathbf{f}_1 \cdot \mathbf{f}_2 \end{bmatrix}.$$

137 and the right hand side is a linear function of $\boldsymbol{\xi}$

$$\mathbf{b} = \begin{bmatrix} 0 \\ 0 \\ 2\zeta U_{22}^{-1}(-U_{21}\varphi_3 + \Phi) \\ 0 \\ 2A \end{bmatrix}. \quad (3.20)$$

Clearly, the Stroh (or *fundamental elasticity*) matrix (3.19) has block structure and coupling of the unknowns $\boldsymbol{\xi}_1^T = [\mathbf{u}^T, \phi^T, A]$ and $\boldsymbol{\xi}_2^T = [\varphi_3, \Phi]$ only occurs through the right hand side (3.20).

Indeed, we can write the coupled system

$$\frac{d\xi_1}{dx_2} = \mathbf{N}_{5 \times 5} \frac{d\xi_1}{dx_1} + \mathbf{L}_{5 \times 2} \xi_2, \quad (3.21a)$$

$$\frac{d\xi_2}{dx_2} = \mathbf{N}_3 \frac{d\xi_2}{dx_2} + \mathbf{L}_{2 \times 5} \xi_1, \quad (3.21b)$$

138 with

$$\mathbf{L}_{5 \times 2} = 2\zeta U_{22}^{-1} \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ -U_{21} & 1 \end{bmatrix}, \quad \mathbf{L}_{2 \times 5} = 2 \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}. \quad (3.22)$$

139 (a) Isotropic material

140 We now show that the Hamiltonian/Stroh formulation so far developed correctly represents the
141 governing equations of plane isotropic CS elasticity. These are [12,19]

$$(G + \lambda) \operatorname{grad}_2 \operatorname{div}_2 \mathbf{u} + G \Delta_2 \left[\mathbf{u} - \frac{1}{2} \ell^2 \operatorname{curl}_2 \operatorname{curl}_2 \mathbf{u} \right] = 0, \quad (3.23)$$

where $\Delta_2 \equiv \operatorname{div}_2 \operatorname{grad}_2$, while $(\operatorname{curl}_2 \mathbf{u})_\alpha = e_{\alpha\beta} u_{\beta,\alpha}$, having let the rank 2 alternating tensor $e_{\alpha\gamma}$ such that $e_{11} = e_{22} = 0$ and $e_{12} = -e_{21} = 1$. Besides, the shearing force is connected to the rotation through

$$\tau_{12} = G \ell^2 \Delta_2 \varphi_3.$$

142 Upon introducing the potentials ω, H such that

$$u_1 = \omega_{,1} + H_{,2}, \quad u_2 = \omega_{,2} - H_{,1}, \quad (3.24)$$

the governing equations (3.23) decouple as [19, Eqs.(14)]

$$(2G + \lambda) \Delta_2 \omega = 0, \quad \text{and} \quad G \Delta_2 \left(1 - \frac{1}{2} \ell^2 \Delta_2 \right) H = 0.$$

143 Indeed, $\varphi_3 = -\frac{1}{2} \Delta_2 H$ and

$$\mu_{3\alpha} = 2G \ell^2 \varphi_{3,\alpha} = -G \ell^2 \Delta_2 H_{,\alpha}. \quad (3.25)$$

144 whence we get the physical meaning of H , whose bilaplacian is related to the shearing force τ_{12} ,

$$\tau_{12} = \Lambda_{,1} = G \ell^2 \Delta_2 \varphi_3 = -\frac{1}{2} G \ell^2 \Delta_2^2 H. \quad (3.26)$$

Finally, the scalar potential is related to displacement flux

$$\operatorname{div}_2 \mathbf{u} = \Delta_2 \omega.$$

We let the matrices (3.2,3.3)

$$\mathbf{U} = 2G \ell^2 \mathbf{1}_2, \quad \mathbf{Q} = \begin{bmatrix} 2G + \lambda & 0 \\ 0 & G \end{bmatrix},$$

$$\mathbf{R} = \begin{bmatrix} 0 & \lambda \\ G & 0 \end{bmatrix}, \quad \mathbf{T} = \begin{bmatrix} G & 0 \\ 0 & 2G + \lambda \end{bmatrix},$$

where $\mathbf{1}_2$ is the rank 2 identity matrix. It easily follows that

$$\mathbf{N}_1 = - \begin{bmatrix} 0 & 1 \\ \frac{\lambda}{2G + \lambda} & 0 \end{bmatrix}, \quad \mathbf{N}_2 = -4G \begin{bmatrix} \frac{G + \lambda}{2G + \lambda} & 0 \\ 0 & 0 \end{bmatrix},$$

while $\zeta = G$,

$$\mathbf{f}_1 = G^{-1} \mathbf{e}_1, \quad \mathbf{f}_2 = 2\mathbf{e}_2.$$

The system (3.18) becomes

$$u_{1,2} = -u_{2,1} + \frac{1}{G}\phi_{1,1} - \frac{1}{G}A_{,1} \quad (3.27a)$$

$$u_{2,2} = -\frac{\lambda}{2G + \lambda}u_{1,1} + \frac{1}{2G + \lambda}\phi_{2,1}, \quad (3.27b)$$

$$\phi_{1,2} = -4G\frac{G + \lambda}{2G + \lambda}u_{1,1} - \frac{\lambda}{2G + \lambda}\phi_{2,1}, \quad (3.27c)$$

$$\phi_{2,2} = -\phi_{1,1} + 2A_{,1}, \quad (3.27d)$$

$$A_{,2} = -2Gu_{1,1} - \phi_{2,1} + \ell^{-2}\Phi, \quad (3.27e)$$

$$\varphi_{3,2} = \frac{1}{2G\ell^2}\Phi_{,1}, \quad (3.27f)$$

$$\Phi_{,2} = -2G\ell^2\varphi_{3,1} + 2A. \quad (3.27g)$$

Differentiation of Eq.(3.27d) with respect to x_1 gives

$$s_{12,1} + s_{22,2} = 2\tau_{12,1},$$

that immediately corresponds to (2.5a) in consideration of the connection $s_{21} = \sigma_{21} + \tau_{21} = s_{12} - 2\tau_{12}$. Similarly, differentiation of (3.27c) lends

$$s_{12,2} + \frac{\lambda}{2G + \lambda}s_{22,1} + 4G\frac{G + \lambda}{2G + \lambda}u_{1,11} = 0,$$

which, with a bit of algebra, corresponds to the first of Eqs.(3.23). Cross differentiation of (3.27f) and (3.27g) allows eliminating $\Phi_{,12}$ to give

$$A_{,1} = G\ell^2\Delta_2\varphi_3,$$

that matches Eq.(3.26). Besides, plugging this result in either equation lends

$$\Phi_{,12} = q_{23,2}^e = \mu_{32,2} = 2G\ell^2\varphi_{3,22},$$

145 which corresponds to (3.25).

146 4. Antiplane deformations

147 Under antiplane shear deformations, the displacement field $\mathbf{u} = (u_1, u_2, u_3)$ is completely defined
148 by the out-of-plane component $u_3 = w(x_1, x_2)$. Thus we have

$$u_1 = u_2 = \varphi_3 = 0,$$

149 and again no dependence of the deformation on x_3 . Thus, Eq.(2.1) lends the rotation $\varphi = \frac{1}{2}\text{curl}_2 w$
150 (see [19] for the definition of curl operating on a scalar field)

$$\varphi_\alpha = \frac{1}{2}e_{\alpha\gamma}w_{,\gamma}. \quad (4.1)$$

Thus, we define the 2D rotation vector

$$\boldsymbol{\varphi}^T = [\varphi_1, \varphi_2],$$

whence the curvature tensor (2.3) is immediately obtained and it is deviatoric

$$\chi_{\alpha\beta} = \varphi_{\alpha,\beta} = \frac{1}{2}e_{\alpha\gamma}w_{,\gamma\beta}, \quad \Leftrightarrow \quad \chi = \frac{1}{2} \begin{bmatrix} w_{,12} & w_{,22} \\ -w_{,11} & -w_{,12} \end{bmatrix}.$$

151 Furthermore, from (2.8) and (2.13b), we get the non-zero components of the skew force stress
152 tensor

$$\tau_{13} = -\frac{1}{2}\mu_{2\beta,\beta} = -\frac{1}{2}p_2, \quad \tau_{23} = \frac{1}{2}\mu_{1\beta,\beta} = \frac{1}{2}p_1. \quad (4.2)$$

153 The Lagrangian density (2.11) becomes

$$L(\text{grad}_2 w, \boldsymbol{\varphi}, \text{grad}_2 \boldsymbol{\varphi}, \mathbf{p}) = \frac{1}{2} [\sigma_{3\alpha} w_{,\alpha} + \mathbf{q}_1^e \cdot \boldsymbol{\varphi}_{,1} + \mathbf{q}_2^e \cdot \boldsymbol{\varphi}_{,2}] + p_1 (\varphi_1 - \frac{1}{2} w_{,2}) + p_2 (\varphi_2 + \frac{1}{2} w_{,1}). \quad (4.3)$$

The Euler-Lagrange equation associated with the variation of w reads

$$-\sigma_{3\alpha,\alpha} + \frac{1}{2} p_{1,2} - \frac{1}{2} p_{2,1} = 0,$$

154 that, by (4.2), reduces to (2.13a). Similarly, through varying $\boldsymbol{\varphi}$, we get the vector E-L equation,

$$-\mathbf{q}_{1,1} - \mathbf{q}_{2,2} + \mathbf{p} = \mathbf{o}, \quad (4.4)$$

155 that corresponds to (2.13b).

156 We now try to relate antiplane problems in CS elasticity with the theory of anisotropic
 157 Kirchhoff plates, which admits a classical Stroh formalism. If we identify the Lagrange multiplier
 158 \mathbf{p} with the shearing force for plates, by assuming $\mathbf{p} = -\text{curl}_2 h$, we immediately obtain the plate
 159 equilibrium equation $\text{div}_2 \mathbf{p} = 0$. Besides, employing the divergence theorem, (4.3) attains the
 160 alternative form

$$L_p(\text{grad}_2 w, \text{grad}_2 \boldsymbol{\varphi}, h, \text{grad}_2 h) = \frac{1}{2} [(\sigma_{3\alpha} + h_{,\alpha}) w_{,\alpha} + \mu_{\alpha\beta} \varphi_{\alpha,\beta}] + h (\varphi_{1,2} - \varphi_{2,1}), \quad (4.5)$$

161 that is formally equivalent to the Lagrangian density adopted for anisotropic plates [8, Eq.(4.15)],
 162 provided we identify $\boldsymbol{\varphi}$ with $\boldsymbol{\theta}$, $\mu_{\alpha\beta}$ with the bending moment in the plate, $M_{\alpha\beta}$, and $\sigma_{3\alpha} + h_{,\alpha}$
 163 with either normal force, $N_{1\alpha}$ or $N_{2\alpha}$, acting in the plane of the plate. Indeed, such an assumption
 164 allows to express the constraint associated with the Lagrange multiplier h in terms of "time"
 165 derivatives, i.e. they become *rheonomic*. However, to allow for a classical Stroh formulation, one
 166 feature is missing: namely, unlike in plate problems, the term $\sigma_{3\alpha} + h_{,\alpha}$ is not constitutively
 167 defined.

We proceed with the Lagrangian density (4.3) and, making use of (2.14), obtain the conjugate momenta

$$\frac{\partial L}{\partial w_{,2}} = \sigma_{32} - \frac{1}{2} p_1 = \sigma_{32} - \tau_{23} = s_{32}, \quad (4.6a)$$

$$\frac{\partial L}{\partial \boldsymbol{\varphi}_{,2}} = \mathbf{q}_2^e. \quad (4.6b)$$

168 In analogy with (3.3), we let the symmetric matrix

$$\hat{U}_{\alpha\beta} = c_{3\alpha 3\beta}, \quad (4.7)$$

169 whence

$$\sigma_{3\alpha} = \hat{U}_{\alpha\beta} w_{,\beta}, \quad (4.8)$$

170 and Eq.(4.6a) may be easily solved for $w_{,2}$

$$w_{,2} = \hat{U}_{22}^{-1} \left(s_{32} + \frac{1}{2} p_1 - \hat{U}_{21} w_{,1} \right). \quad (4.9)$$

171 For (4.6b) we need to let, in analogy with (3.2),

$$\hat{Q}_{\alpha\beta} = \ell^2 g_{\alpha 1 \beta 1}, \quad \hat{R}_{\alpha\beta} = \ell^2 g_{\alpha 1 \beta 2}, \quad \hat{T}_{\alpha\beta} = \ell^2 g_{\alpha 2 \beta 2}, \quad (4.10)$$

172 so that, paralleling (3.4),

$$\mathbf{q}_1^e = \hat{\mathbf{Q}} \boldsymbol{\varphi}_{,1} + \hat{\mathbf{R}} \boldsymbol{\varphi}_{,2}, \quad \mathbf{q}_2^e = \hat{\mathbf{R}}^T \boldsymbol{\varphi}_{,1} + \hat{\mathbf{T}} \boldsymbol{\varphi}_{,2}, \quad (4.11)$$

173 we can write

$$\boldsymbol{\varphi}_{,2} = \hat{\mathbf{T}}^{-1} \left(\mathbf{q}_2^e - \hat{\mathbf{R}}^T \boldsymbol{\varphi}_{,1} \right). \quad (4.12)$$

174 Besides, from (4.4) and the constitutive law (4.11), we get

$$\mathbf{p} = \hat{\mathbf{Q}} \boldsymbol{\varphi}_{,11} + \left(\hat{\mathbf{R}} + \hat{\mathbf{R}}^T \right) \boldsymbol{\varphi}_{,12} + \hat{\mathbf{T}} \boldsymbol{\varphi}_{,22}, \quad (4.13)$$

175 which shows that indeed $\text{div}_2 \mathbf{p} = 0$, inasmuch as (2.2) holds, as discussed in Sec.2. Recalling
176 (2.14), this implies

$$\tau_{31,2} - \tau_{32,1} = 0, \quad (4.14)$$

177 which is in fact satisfied by Eqs.(2.14) of [20].

We define the Hamiltonian density function $H = H(s_{32}, w, \varphi, \varphi_1, \mathbf{q}_2^e, \mathbf{p})$

$$\begin{aligned} H = s_{32}w, \varphi + \mathbf{q}_2^e \cdot \varphi, \varphi - L = \frac{1}{2} \hat{U}_{22}^{-1} \left(s_{32} + \frac{1}{2} p_1 - \hat{U}_{21} w, \varphi \right)^2 \\ + \frac{1}{2} \left(\mathbf{q}_2^e - \hat{\mathbf{R}}^T \varphi, \varphi_1 \right) \cdot \hat{\mathbf{T}}^{-1} \left(\mathbf{q}_2^e - \hat{\mathbf{R}}^T \varphi, \varphi_1 \right) - \frac{1}{2} \hat{U}_{11} w, \varphi_1^2 - \frac{1}{2} \varphi, \varphi_1 \cdot \hat{\mathbf{Q}} \varphi, \varphi_1 - \mathbf{p} \cdot \varphi - \frac{1}{2} p_2 w, \varphi, \end{aligned} \quad (4.15)$$

whence, from (3.11), we retrieve (4.9)

$$\frac{\delta H}{\delta s_{32}} = \hat{U}_{22}^{-1} \left(s_{32} + \frac{1}{2} p_1 - \hat{U}_{21} w, \varphi \right) = w, \varphi,$$

and (4.12)

$$\frac{\delta H}{\delta \mathbf{q}_2^e} = \hat{\mathbf{T}}^{-1} \left(\mathbf{q}_2^e - \hat{\mathbf{R}}^T \varphi, \varphi_1 \right) = \varphi, \varphi.$$

The canonical equation (3.12) gives

$$\frac{\delta H}{\delta \varphi} = \left[\hat{\mathbf{Q}} \varphi, \varphi_1 + \hat{\mathbf{R}} \hat{\mathbf{T}}^{-1} \left(\mathbf{q}_2^e - \hat{\mathbf{R}}^T \varphi, \varphi_1 \right) \right]_{,1} - \mathbf{p} = -\mathbf{q}_2^e, \varphi,$$

and

$$\frac{\delta H}{\delta w} = \left[\hat{U}_{21} \hat{U}_{22}^{-1} \left(s_{32} + \frac{1}{2} p_1 - \hat{U}_{21} w, \varphi \right) + \hat{U}_{11} w, \varphi_1 + \frac{1}{2} p_2 \right]_{,1} = -s_{32}, \varphi,$$

178 corresponding to (2.13b) and (2.13a), respectively. We introduce the stream functions, which are
179 defined up to a function of x_2 ,

$$\phi = \int^{x_1} s_{32} d\xi_1, \quad \Phi = \int^{x_1} \mathbf{q}_2^e d\xi_1, \quad \Lambda = \frac{1}{2} \int^{x_1} \mathbf{p} d\xi_1,$$

and write the first order system

$$\varphi, \varphi_2 = \hat{\mathbf{N}}_1 \varphi, \varphi_1 + \hat{\mathbf{T}}^{-1} \Phi, \varphi_1, \quad (4.16a)$$

$$\Phi, \varphi_2 = \hat{\mathbf{N}}_2 \varphi, \varphi_1 + \hat{\mathbf{N}}_1^T \Phi, \varphi_1 + 2\Lambda, \quad (4.16b)$$

$$w, \varphi_2 = -\frac{\hat{U}_{21}}{\hat{U}_{22}} w, \varphi_1 + \frac{1}{\hat{U}_{22}} \phi, \varphi_1 + \frac{1}{\hat{U}_{22}} \Lambda_{1,1} \quad (4.16c)$$

$$\phi, \varphi_2 = \left(\frac{\hat{U}_{21}^2}{\hat{U}_{22}} - \hat{U}_{11} \right) w, \varphi_1 - \frac{\hat{U}_{21}}{\hat{U}_{22}} \phi, \varphi_1 - \frac{\hat{U}_{21}}{\hat{U}_{22}} \Lambda_{1,1} - \Lambda_{2,1}, \quad (4.16d)$$

180 having let

$$\hat{\mathbf{N}}_1 = -\hat{\mathbf{T}}^{-1} \hat{\mathbf{R}}^T, \quad \hat{\mathbf{N}}_2 = -\hat{\mathbf{R}} \hat{\mathbf{N}}_1 - \hat{\mathbf{Q}}.$$

181 It only remains to determine an expression for the Lagrange multiplier \mathbf{p} , which amounts to
182 acknowledging the constraint (2.1). In fact, Eq.(4.1) allows to solve (4.16c) for $\Lambda_{1,1}$ and to dispense
183 with w, φ_1 and w, φ_2

$$\Lambda_{1,1} = 2\hat{U}_{22}\varphi_1 - 2\hat{U}_{21}\varphi_2 - \phi, \varphi_1. \quad (4.17)$$

184 In light of (4.1) and (4.8), this algebraic condition simply states that $\tau_{23} = \sigma_{32} - s_{32}$. When we plug
185 this result into (4.16d) and use (4.8), we find

$$\phi, \varphi_2 = 2\hat{U}_{11}\varphi_2 - 2\hat{U}_{12}\varphi_1 - \Lambda_{2,1} = -\sigma_{31} - \tau_{31} = -s_{31} \quad (4.18)$$

that, differentiated with respect to x_1 , gives the equilibrium equation (2.5a). To get an equation
for $\Lambda_{2,1}$, we cannot directly employ the connection $2\varphi_2 = -w, \varphi_1$, for it is algebraic. Instead, we take

advantage of $\text{div}_2 \mathbf{p} = 0$, whereby

$$\text{div}_2 \mathbf{A} = 0.$$

Mathematically, this amounts to exploiting (2.2), which is obtained cross differentiating and adding (4.1), whence a differentiation index 2 is implied. Thus, Eq.(4.17) immediately lends the *evolution equation*

$$\Lambda_{2,2} = -2\hat{U}_{22}\varphi_1 + 2\hat{U}_{21}\varphi_2 + \phi_{,1},$$

that corresponds to (4.14), integrated with respect to x_1 . In this form, the problem's variables are φ , Φ , ϕ , Λ , and they are governed by a semi-explicit system of first order DAE, the single algebraic relation being (4.17). To obtain a pure system of ODEs, an evolution equation for Λ_1 is demanded. This is obtained differentiating (4.17) with respect to x_2 and then integrating with respect to x_1

$$\Lambda_{1,2} = 2 \left[\hat{\mathbf{N}}_1 \varphi + \hat{\mathbf{T}}^{-1} \Phi \right] \cdot \left(\hat{U}_{22} \mathbf{e}_1 - \hat{U}_{21} \mathbf{e}_2 \right) - \phi_{,2} = 2\varphi \cdot \left[\hat{\mathbf{N}}_1^T \mathbf{f}_2 + \mathbf{f}_1 \right] + 2\Phi \cdot \hat{\mathbf{T}}^{-1} \mathbf{f}_2 + \Lambda_{2,1},$$

186 having made use of (4.16a,4.18) and let

$$\mathbf{f}_1 = \hat{U}_{12} \mathbf{e}_1 - \hat{U}_{11} \mathbf{e}_2, \quad \mathbf{f}_2 = \hat{U}_{22} \mathbf{e}_1 - \hat{U}_{21} \mathbf{e}_2.$$

187 Consequently, the system of DAEs has *differentiation order 3*, that is typical of constrained
188 mechanical systems. Also, we note that

$$\sigma_{31} = 2\varphi \cdot \mathbf{f}_1, \quad \sigma_{32} = 2\varphi \cdot \mathbf{f}_2. \quad (4.19)$$

We thus obtain the linear system in the variables $(\varphi, \Phi, \phi, \Lambda)$

$$\varphi_{,2} = \hat{\mathbf{N}}_1 \varphi_{,1} + \hat{\mathbf{T}}^{-1} \Phi_{,1}, \quad (4.20a)$$

$$\Phi_{,2} = \hat{\mathbf{N}}_2 \varphi_{,1} + \hat{\mathbf{N}}_1^T \Phi_{,1} + 2\Lambda, \quad (4.20b)$$

$$\phi_{,2} = -\Lambda_{2,1} - 2\varphi \cdot \mathbf{f}_1, \quad (4.20c)$$

$$\Lambda_{1,2} = \Lambda_{2,1} + 2\varphi \cdot \left[\hat{\mathbf{N}}_1^T \mathbf{f}_2 + \mathbf{f}_1 \right] + 2\Phi \cdot \hat{\mathbf{T}}^{-1} \mathbf{f}_2, \quad (4.20d)$$

$$\Lambda_{2,2} = \phi_{,1} - 2\varphi \cdot \mathbf{f}_2, \quad (4.20e)$$

189 Cross-differentiating Eqs.(4.20a,4.20b) to eliminate $\Phi_{,12}$ yields (4.13). Besides, multiplying
190 (4.20a) by $-\mathbf{R}$ and substituting in (4.20b) gives (4.4). In light of Eqs.(4.19), Eq.(4.20c) gives the
191 equilibrium equation (2.5a), while (4.20e) amounts to (4.14), both having being integrated along
192 x_1 . Finally, adding (4.20c) and (4.20d) and differentiating lends (4.6a), while cross-differentiating
193 (4.20c,4.20e) and adding lends the second order connection for ϕ

$$\Delta_2 \phi = 2\varphi_{,1} \cdot \mathbf{f}_2 - 2\varphi_{,2} \cdot \mathbf{f}_1 = \sigma_{32,1} - \sigma_{31,2}, \quad (4.21)$$

194 that supports the interpretation of ϕ as a stress function for the problem.

195 Thus, letting $\hat{\xi}^T = [\varphi^T, \Phi^T, \phi, \Lambda^T]$, we have

$$\frac{d\hat{\xi}}{dx_2} = \hat{\mathbf{N}} \frac{d\hat{\xi}}{dx_1} + \hat{\mathbf{b}}, \quad (4.22)$$

196 where we have let the Stroh matrix

$$\hat{\mathbf{N}} = \begin{bmatrix} \hat{\mathbf{N}}_1 & \hat{\mathbf{T}}^{-1} & \mathbf{o} & \mathbf{o} & \mathbf{o} \\ \hat{\mathbf{N}}_2 & \hat{\mathbf{N}}_1^T & \mathbf{o} & \mathbf{o} & \mathbf{o} \\ \mathbf{o}^T & \mathbf{o}^T & 0 & 0 & -1 \\ \mathbf{o}^T & \mathbf{o}^T & 0 & 0 & 1 \\ \mathbf{o}^T & \mathbf{o}^T & 1 & 0 & 0 \end{bmatrix}, \quad (4.23)$$

197 and the right hand side is a linear function of $\hat{\xi}$

$$\hat{b} = 2 \begin{bmatrix} \mathbf{O} & \mathbf{O} & \mathbf{o} & \mathbf{o} & \mathbf{o} \\ \mathbf{O} & \mathbf{O} & \mathbf{o} & \mathbf{e}_1 & \mathbf{e}_2 \\ -\mathbf{f}_1^T & \mathbf{o}^T & 0 & 0 & 0 \\ (\hat{\mathbf{N}}_1^T \mathbf{f}_2 + \mathbf{f}_1)^T & (\mathbf{T}^{-1} \mathbf{f}_2)^T & 0 & 0 & 0 \\ \mathbf{f}_2^T & \mathbf{o}^T & 0 & 0 & 0 \end{bmatrix} \hat{\xi}. \quad (4.24)$$

198 (a) Isotropic anti-plane deformations

199 We now show that the above canonical formulation correctly reproduces the governing equations
200 for antiplane deformations in isotropic CS media. Such framework demands

$$\epsilon = \frac{1}{2} w_{,1} \mathbf{e}_3 \otimes \mathbf{e}_1 + \frac{1}{2} w_{,2} \mathbf{e}_3 \otimes \mathbf{e}_2,$$

201 whence

$$\sigma_{31} = G w_{,1} = -2G\varphi_2, \quad \sigma_{32} = G w_{,2} = 2G\varphi_1. \quad (4.25)$$

202 In particular,

$$\sigma_{32,1} - \sigma_{31,2} = 2G \operatorname{div} \boldsymbol{\varphi} = 0, \quad (4.26)$$

and, by (4.21), ϕ turns harmonic. From (2.4), we get the curvature tensor

$$\mu_{11} = 2G\ell^2(1 + \eta)\varphi_{1,1} = G\ell^2(1 + \eta)w_{,12} = -\mu_{22}, \quad (4.27a)$$

$$\mu_{21} = 2G\ell^2(\varphi_{2,1} + \eta\varphi_{1,2}) = -G\ell^2(w_{,11} - \eta w_{,22}), \quad (4.27b)$$

$$\mu_{12} = 2G\ell^2(\eta\varphi_{2,1} + \varphi_{1,2}) = -G\ell^2(\eta w_{,11} - w_{,22}), \quad (4.27c)$$

203 whereby, from (2.13b, 2.14), we have [20, Eq.(2.14)]

$$\tau_{13} = -G\ell^2 \Delta_2 \varphi_2, \quad \tau_{23} = G\ell^2 \Delta_2 \varphi_1, \quad (4.28)$$

204 which clearly satisfy (4.14) in light of (2.2). The equilibrium equation (2.5a) reads [20, Eq.(2.15)]

$$G \left(1 - \frac{1}{2} \ell^2 \Delta_2\right) \Delta_2 w = 0, \quad (4.29)$$

205 or, equivalently, given that $\operatorname{curl}_2 \boldsymbol{\varphi} = -\frac{1}{2} \Delta_2 w$,

$$2G \operatorname{curl}_2 \boldsymbol{\varphi} - \tau_{31,1} - \tau_{32,2} = 0. \quad (4.30)$$

206 We let the vectors

$$\mathbf{f}_1^T = [0, -G], \quad \mathbf{f}_2^T = [G, 0],$$

alongside the matrices (4.7, 4.10)

$$\begin{aligned} \hat{\mathbf{U}} &= G\mathbf{1}_2, & \hat{\mathbf{Q}} &= 2G\ell^2 \begin{bmatrix} 1 + \eta & 0 \\ 0 & 1 \end{bmatrix}, \\ \hat{\mathbf{R}} &= 2G\ell^2 \begin{bmatrix} 0 & 0 \\ \eta & 0 \end{bmatrix}, & \hat{\mathbf{T}} &= 2G\ell^2 \begin{bmatrix} 1 & 0 \\ 0 & 1 + \eta \end{bmatrix}, \end{aligned}$$

where $\mathbf{1}_2$ is the rank 2 identity matrix. It easily follows that

$$\hat{\mathbf{N}}_1 = - \begin{bmatrix} 0 & \eta \\ 0 & 0 \end{bmatrix}, \quad \hat{\mathbf{N}}_2 = -2G\ell^2 \begin{bmatrix} 1 + \eta & 0 \\ 0 & 1 - \eta^2 \end{bmatrix}.$$

Eq.(4.20) gives the first order system

$$\varphi_{1,2} = -\eta\varphi_{2,1} + \frac{1}{2G\ell^2}\Phi_{1,1}, \quad (4.31a)$$

$$\varphi_{2,2} = \frac{1}{2G\ell^2(1+\eta)}\Phi_{2,1} \quad (4.31b)$$

$$\Phi_{1,2} = -2G\ell^2(1+\eta)\varphi_{1,1} + 2A_1 \quad (4.31c)$$

$$\Phi_{2,2} = -2G\ell^2(1-\eta^2)\varphi_{2,1} - \eta\Phi_{1,1} + 2A_2 \quad (4.31d)$$

$$\phi_{,2} = 2G\varphi_2 - A_{2,1} \quad (4.31e)$$

$$A_{1,2} = -2G(1+\eta)\varphi_2 + A_{2,1} + \ell^{-2}\Phi_1 \quad (4.31f)$$

$$A_{2,2} = \phi_{,1} - 2G\varphi_1. \quad (4.31g)$$

Cross-differentiating and adding Eqs.(4.31e) and (4.31g) shows that ϕ is harmonic inasmuch as (2.2) holds, which result is in line with (4.21). Consequently, letting the harmonic conjugate function ϕ^* , upon recalling that $\phi_{,2} = -\phi_{,1}^*$, we get, from (4.31e),

$$\phi^* = Gw + A_2 = \int^{x_1} s_{31} d\xi_1,$$

207 which gives to the harmonic conjugate function the role of the stress function for s_{31} .
 208 Eqs.(4.31a,4.31b) correspond to Eqs.(4.27c) and (4.27a), respectively. Eqs.(4.31c,4.31d) represent
 209 rotational equilibrium (2.13b), provided that we use (4.31a) to eliminate $\Phi_{1,1}$. Similarly, in light of
 210 (4.25) and of (2.14), Eq.(4.31e) lends translational equilibrium (4.18). Eq.(4.31g) amounts to (4.14),
 211 while (4.31f) is (4.30), having differentiated and used (4.31a) to eliminate $\ell^{-2}\Phi_{1,1}$.

212 5. Conclusions

213 We derived the Hamiltonian formalism associated with the indeterminate couple stress theory
 214 of elasticity for general anisotropic media. The Hamiltonian framework is known to lead to
 215 the celebrated Stroh formalism in classical elasticity. This canonical rewriting of the governing
 216 equations is of great theoretical and practical value, because it lends fundamental existence
 217 and uniqueness results, as well as providing a powerful tool for solving problems in generally
 218 anisotropic media. For such reasons, we extend the formalism to the couple stress theory. This
 219 is a strain gradient theory that incorporates microstructural effects in a fashion similar to lattice
 220 elasticity [16]. We show that, unlike classical and constrained elasticity, the theory does not allow
 221 for a standard Stroh formalism, owing to the nature of the internal constraint on the micro-rotation
 222 vector. Indeed, the constraint is *algebraic* and it cannot be eliminated. The resulting canonical
 223 formulation is a differential algebraic system of equations (DAE), which may be rewritten
 224 in purely differential terms by developing suitable evolution equations. However, the simple
 225 structure of classical elasticity cannot be reproduced.

226 The developed canonical system is then specialized to the case of plane and antiplane strain
 227 for couple stress anisotropic media. The antiplane framework is especially noteworthy because
 228 it admits a Lagrangian formulation that exactly matches that of flexural/extensional Kirchhoff
 229 anisotropic plates, which are amenable to a Stroh formalism. Nonetheless, the corresponding
 230 canonical system in couple stress elasticity still lacks the features of a classical Stroh formulation,
 231 because the term corresponding to the normal force in the plate is not determined constitutively,
 232 owing to the presence of tangential stresses. This notwithstanding, the Hamiltonian formalism
 233 still provides a wealth of informations, including unexpected connections which are not apparent
 234 from the standard treatment.

235 **Competing Interests.** The authors declare that they have no competing interests

236 **Funding.** AN gratefully acknowledge financial support under the H2020 MSCA RISE project EffectFact, GA
 237 101008140

240 Table of symbols

Symbol	Description	Symbol	Description
s	Cauchy stress tensor	μ	Couple-stress tensor
ϵ	Strain tensor	χ	Curvature tensor
σ	Sym part of the stress tensor	τ	Skew-sym part of the stress tensor
u	Displacement field	φ	Micro-rotation field
L	Lagrangian density	H	Hamiltonian density
N	Stroh matrix	U	Microstructure matrix (symmetric)
Q, T	Diagonal blocks in N (symmetric)	R, R^T	Off-diagonal blocks in N
e_{ijk}	Rank 3 permutation tensor	δ_{ij}	Kroneker delta tensor
e_1, e_2, e_3	Orthonormal basis vectors	Λ, G, ℓ, η	Constitutive parameters

243 References

- 244 1. VI Alshits, AN Darinskii, and J Lothe.
245 On the existence of surface waves in half-infinite anisotropic elastic media with piezoelectric
246 and piezomagnetic properties.
247 *Wave motion*, 16(3):265–283, 1992.
- 248 2. DM Barnett.
249 Bulk, surface, and interfacial waves in anisotropic linear elastic solids.
250 *International Journal of Solids and Structures*, 37(1-2):45–54, 2000.
- 251 3. DM Barnett and J Lothe.
252 Consideration of the existence of surface wave (rayleigh wave) solutions in anisotropic elastic
253 crystals.
254 *Journal of physics F: Metal physics*, 4(5):671, 1974.
- 255 4. DM Barnett and J1 Lothe.
256 Dislocations and line charges in anisotropic piezoelectric insulators.
257 *Physica status solidi (b)*, 67(1):105–111, 1975.
- 258 5. RT Edmondson and YB Fu.
259 Stroh formulation for a generally constrained and pre-stressed elastic material.
260 *International Journal of Non-Linear Mechanics*, 44(5):530–537, 2009.
- 261 6. Hui Fan and Limei Xu.
262 Love wave in a classical linear elastic half-space covered by a surface layer described by the
263 couple stress theory.
264 *Acta Mechanica*, 229(12):5121–5132, 2018.
- 265 7. YB Fu.
266 Existence and uniqueness of edge waves in a generally anisotropic elastic plate.
267 *Quarterly Journal of Mechanics and Applied Mathematics*, 56(4):605–616, 2003.
- 268 8. YB Fu.
269 Hamiltonian interpretation of the Stroh formalism in anisotropic elasticity.
270 *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*,
271 463(2088):3073–3087, 2007.
- 272 9. YB Fu and DW Brookes.
273 Edge waves in asymmetrically laminated plates.
274 *Journal of the Mechanics and Physics of Solids*, 54(1):1–21, 2006.
- 275 10. YB Fu and J Kaplunov.
276 Analysis of localized edge vibrations of cylindrical shells using the stroh formalism.
277 *Mathematics and mechanics of solids*, 17(1):59–66, 2012.
- 278 11. YB Fu, J Kaplunov, and D Prikazchikov.
279 Reduced model for the surface dynamics of a generally anisotropic elastic half-space.
280 *Proceedings of the Royal Society A*, 476(2234):20190590, 2020.

- 281 12. K F Graff and Y-H Pao.
 282 The effects of couple-stresses on the propagation and reflection of plane waves in an elastic
 283 half-space.
 284 *Journal of Sound and Vibration*, 6(2):217–229, 1967.
- 285 13. W Günther.
 286 Zur statik und kinematik des cosseratschen kontinuums.
 287 *Abh. Braunschweig. Wiss. Ges.*, 10(213):1, 1958.
- 288 14. WT Koiter.
 289 Couple-stress in the theory of elasticity.
 290 In *Proc. K. Ned. Akad. Wet.*, volume 67, pages 17–44. North Holland Pub, 1964.
- 291 15. M Lazar and H Kirchner.
 292 Cosserat (micropolar) elasticity in stroh form.
 293 *International journal of solids and structures*, 42(20):5377–5398, 2005.
- 294 16. R D Mindlin.
 295 Micro-structure in linear elasticity.
 296 *Archive for Rational Mechanics and Analysis*, 16(1):51–78, 1964.
- 297 17. RD Mindlin and HF Tiersten.
 298 Effects of couple-stresses in linear elasticity.
 299 *Archive for Rational Mechanics and analysis*, 11(1):415–448, 1962.
- 300 18. G Mishuris, A Piccolroaz, and E Radi.
 301 Steady-state propagation of a mode III crack in couple stress elastic materials.
 302 *International Journal of Engineering Science*, 61:112–128, 2012.
- 303 19. A Nobili.
 304 Asymptotically consistent size-dependent plate models based on the couple-stress theory
 305 with micro-inertia.
 306 *European Journal of Mechanics-A/Solids*, 89:104316, 2021.
- 307 20. A Nobili, E Radi, and C Signorini.
 308 A new Rayleigh-like wave in guided propagation of antiplane waves in couple stress
 309 materials.
 310 *Proceedings of the Royal Society A*, 476(2235):20190822, 2020.
- 311 21. A Nobili, E Radi, and A Vellender.
 312 Diffraction of antiplane shear waves and stress concentration in a cracked couple stress elastic
 313 material with micro inertia.
 314 *Journal of the Mechanics and Physics of Solids*, 124:663–680, 2019.
- 315 22. A Nobili and V Volpini.
 316 Microstructured induced band pattern in love wave propagation for novel nondestructive
 317 testing (ndt) procedures.
 318 *International Journal of Engineering Science*, 168:103545, 2021.
- 319 23. A Nobili, V Volpini, and C Signorini.
 320 Antiplane stoneley waves propagating at the interface between two couple stress elastic
 321 materials.
 322 *Acta Mechanica*, 232(3):1207–1225, 2021.
- 323 24. W Nowacki.
 324 Theory of asymmetric elasticity.
 325 Pergamon Press, Headington Hill Hall, Oxford OX 3 0 BW, UK, 1986., 1986.
- 326 25. N S Ottosen, M Ristinmaa, and C Ljung.
 327 Rayleigh waves obtained by the indeterminate couple-stress theory.
 328 *European Journal of Mechanics-A/Solids*, 19(6):929–947, 2000.
- 329 26. LP Solie and BA Auld.
 330 Elastic waves in free anisotropic plates.
 331 *The Journal of the Acoustical Society of America*, 54(1):50–65, 1973.
- 332 27. AN Stroh.
 333 Dislocations and cracks in anisotropic elasticity.
 334 *Philosophical magazine*, 3(30):625–646, 1958.
- 335 28. AN Stroh.
 336 Steady state problems in anisotropic elasticity.
 337 *Journal of Mathematics and Physics*, 41(1-4):77–103, 1962.
- 338 29. TCT Ting.

339 *Anisotropic elasticity: theory and applications.*
340 Number 45. Oxford University Press on Demand, 1996.

341 30. KC Wu.

342 Extension of stroh's formalism to self-similar problems in two-dimensional elastodynamics.
343 *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering*
344 *Sciences*, 456(1996):869–890, 2000.