

This is the peer reviewed version of the following article:

Harnack inequality for hypoelliptic ultraparabolic equations with a singular lower order term / Polidoro, Sergio; Ragusa, M. A.. - In: REVISTA MATEMATICA IBEROAMERICANA. - ISSN 0213-2230. - STAMPA. - 24:3(2008), pp. 1011-1046. [10.4171/RMI/565]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

27/04/2024 21:05

Harnack inequality for hypoelliptic ultraparabolic equations with a singular lower order term

Sergio Polidoro and Maria Alessandra Ragusa

Abstract

We prove a Harnack inequality for the positive solutions of ultraparabolic equations of the type

$$\mathcal{L}_0 u + \mathcal{V} u = 0,$$

where \mathcal{L}_0 is a linear second order hypoelliptic operator and \mathcal{V} belongs to a class of functions of Stummel-Kato type. We also obtain the existence of a Green function and an uniqueness result for the Cauchy-Dirichlet problem.

1. Introduction

We prove some regularity results for the solutions of the equation in \mathbb{R}^{N+1}

$$(1.1) \quad \mathcal{L}_0 u + \mathcal{V} u = 0,$$

where \mathcal{V} is a singular potential belonging to a Stummel-Kato class (see Definition 1.1 below) and \mathcal{L}_0 is a linear second order operator of the form

$$(1.2) \quad \mathcal{L}_0 = \sum_{k=1}^m X_k^2 + X_0 - \partial_t.$$

We always denote by $z = (x, t)$ the point in \mathbb{R}^{N+1} ; the X_k 's in (1.2) are smooth vector fields on \mathbb{R}^N , i.e.

$$X_k(x) = \sum_{j=1}^N a_j^k(x) \partial_{x_j}, \quad k = 0, \dots, m,$$

2000 Mathematics Subject Classification: 35K70, 35J10, 35K20, 32A37, 35B65.

Keywords: Hypoelliptic operator, Schrödinger equation, Harnack inequality, Green function.

where any a_j^k is a C^∞ function. In the sequel we also consider the X_k 's as vector fields in \mathbb{R}^{N+1} and we denote

$$(1.3) \quad Y = X_0 - \partial_t.$$

We also assume that $X_0(0) = 0$, so that

$$(1.4) \quad Y(0) = -\partial_t.$$

As we will see in Remark 3.2, this assumption is not necessary, but allows us to simplify the notations.

We say that a curve $\gamma : [0, T] \rightarrow \mathbb{R}^{N+1}$ is *L-admissible* if it is absolutely continuous and satisfies

$$\gamma'(s) = \sum_{k=1}^m \lambda_k(s) X_k(\gamma(s)) + \mu(s) Y(\gamma(s)), \quad \text{a.e. in } [0, T],$$

for suitable piecewise constant real functions $\lambda_1, \dots, \lambda_m, \mu$, with $\mu \geq 0$. We next state our main assumptions:

[H.1] there exists a homogeneous Lie group $\mathbb{G} = (\mathbb{R}^{N+1}, \circ, \delta_\lambda)$ such that

- (i) X_1, \dots, X_m, Y are left translation invariant on \mathbb{G} ;
- (ii) X_1, \dots, X_m are δ_λ -homogeneous of degree one and Y is δ_λ -homogeneous of degree two;

[H.2] for every $(x, t), (\xi, \tau) \in \mathbb{R}^{N+1}$ with $t > \tau$, there exists an *L*-admissible path $\gamma : [0, T] \rightarrow \mathbb{R}^{N+1}$ such that $\gamma(0) = (x, t)$, $\gamma(T) = (\xi, \tau)$.

Operators of this kind have been studied by Kogoj and Lanconelli in [11]. The above hypotheses and the main properties of homogeneous Lie groups will be discussed in detail in the next section, here we recall that assumptions [H.1]–[H.2] yield the well known Hörmander condition [10]:

$$(1.5) \quad \text{rank Lie}\{X_1, \dots, X_m, Y\}(z) = N + 1, \quad \text{for every } z \in \mathbb{R}^{N+1},$$

then \mathcal{L}_0 is hypoelliptic (i.e. every distributional solution to $\mathcal{L}_0 u = f$ is smooth whenever f is smooth; see, for instance, Proposition 10.1 in [11]). Hence \mathcal{L}_0 belongs to the general class of the hypoelliptic operators on homogeneous groups first studied by Folland [8]. We recall that a general theory of function spaces related to Hörmander operators has been developed by Rothschild and Stein in [24], and by Nagel, Stein and Wainger in [21]. An invariant Harnack inequality for the positive solutions of $\mathcal{L}_0 u = 0$ and a Gaussian upper estimate of its fundamental solution Γ_0 have been proved

in [11]. We also recall that Gaussian lower bounds for operators verifying assumptions [H.1]–[H.2] on Lie group of step three have been given in [22]; and one-side Liouville theorems are provided in [12].

Let us point out that several meaningful examples of operators of the form (1.2) satisfy assumptions [H.1]–[H.2]:

- heat operators on Carnot groups

$$(1.6) \quad \Delta_{\mathbb{G}} - \partial_t,$$

where $\Delta_{\mathbb{G}} = \sum_{k=1}^m X_k^2$ denotes the sub-Laplacian on a homogeneous Carnot group \mathbb{G} (see Varopoulos, Saloff-Coste and Coulhon [28]);

- heat operators *with drift* on Carnot groups

$$(1.7) \quad \Delta_{\mathbb{G}} + X_0 - \partial_t,$$

(see Alexopoulos [1]). Note that, due to our assumption [H.1], the vector field X_0 belongs to the second layer of the Lie algebra of the group);

- Kolmogorov type operators

$$(1.8) \quad \Delta_{\mathbb{R}^m} + \langle Bx, \nabla \rangle - \partial_t,$$

where $\Delta_{\mathbb{R}^m}$ is the Laplace operator on \mathbb{R}^m and B is a constant $N \times N$ real matrix (see [16] and its bibliography for a survey on known results on Kolmogorov type operators. In [18] necessary and sufficient conditions are given on the matrix B in order to satisfy assumptions [H.1]–[H.2]);

- operators on the “link of a Carnot and a Kolmogorov group”

$$(1.9) \quad \Delta_{\mathbb{G}} + \langle Bx, \nabla \rangle - \partial_t$$

here the domain of the solution is $\mathbb{R}^m \times \mathbb{R}^p \times \mathbb{R}^q \times \mathbb{R}$, \mathbb{G} is a Carnot group on $\mathbb{R}^m \times \mathbb{R}^p$ and B is a $(m+q) \times (m+q)$ matrix as in the Kolmogorov operator (1.8) (see Kogoj and Lanconelli [11, Example 9.7]). A general procedure for the construction of sequences of linked groups of dimension and step arbitrarily large is given in the paper [13].

We are concerned with the regularity of the operator

$$(1.10) \quad \mathcal{L}_{\mathcal{V}} = \mathcal{L}_0 + \mathcal{V},$$

where \mathcal{V} belongs to the following Stummel-Kato class (defined by the fundamental solution Γ_0 of \mathcal{L}_0).

Definition 1.1. Let Ω be an open subset contained in \mathbb{R}^{N+1} . A function $\mathcal{V} \in L^1(\Omega)$ belongs to the space $SK(\Omega, \mathcal{L}_0)$, if

$$(1.11) \quad \lim_{h \rightarrow 0} \eta_{\mathcal{V}}(h) = 0, \quad \lim_{h \rightarrow 0} \eta_{\mathcal{V}}^*(h) = 0,$$

where

$$(1.12) \quad \begin{aligned} \eta_{\mathcal{V}}(h) &= \sup_{(x,t) \in \Omega} \int_{(y,s) \in \Omega, \ t-h^2 < s < t} \Gamma_0(x, t, y, s) |\mathcal{V}(y, s)| dy ds, \\ \eta_{\mathcal{V}}^*(h) &= \sup_{(y,s) \in \Omega} \int_{(x,t) \in \Omega, \ s < t < s+h^2} \Gamma_0(x, t, y, s) |\mathcal{V}(x, t)| dx dt. \end{aligned}$$

We say that u is a weak solution of $\mathcal{L}_{\mathcal{V}}u = 0$ if

1. there exists $p > 1$ such that $u, X_1u, \dots, X_mu \in L_{\text{loc}}^p(\Omega)$,
2. $\mathcal{V}u \in L_{\text{loc}}^1(\Omega)$,
3. $\int_{\Omega} \sum_{k=1}^m X_k u X_k^* \varphi + \int_{\Omega} u Y^* \varphi + \int_{\Omega} u \mathcal{V} \varphi = 0$, for every $\varphi \in C_0^\infty(\Omega)$.

As in the Euclidean setting, the Stummel-Kato class can be related to the Morrey spaces $L^{p,\lambda}(\Omega, \mathbb{G})$; in Section 3 we will prove the inclusion $L^{1,\lambda}(\Omega, \mathbb{G}) \subset SK(\Omega, \mathcal{L}_0)$ for $\lambda \in]Q - 2, Q[$, where Q is the homogeneous dimension of \mathbb{G} (see Section 2 for the definitions). We also give a simple sufficient condition for the integrability of $\mathcal{V}u$: we show that, if the derivatives $X_j u, X_j X_k u$, for $j, k = 1, \dots, m$ and $Y u$ belong to $L_{\text{loc}}^1(\Omega)$ then $\mathcal{V}u \in L_{\text{loc}}^1(\Omega)$.

Our main result is an invariant Harnack inequality for the positive solutions to $\mathcal{L}_{\mathcal{V}}u = 0$. The proof of the Harnack inequality given by Kogoj and Lanconelli in [11] (for the solutions to $\mathcal{L}_0 u = 0$) is based on a mean value theorem and follows the same lines of the classical proof of the Harnack inequality for harmonic functions. That approach has been used in the study of Kolmogorov operators (1.8) by Kuptsov in [14], later by Garofalo and Lanconelli in [9] then by Lanconelli and Polidoro in [18] and relies on some accurate estimates of the derivatives $X_1 \Gamma_0, \dots, X_m \Gamma_0$ of the fundamental solution of \mathcal{L}_0 . Here we use a method based on the Green function G_0 of \mathcal{L}_0 related to suitable “cylindrical” open sets and on a pointwise lower bound for G_0 . This technique is inspired by some arguments by Safanov in [25], and used in [15] where Kusuoka and Stroock obtain Harnack inequality for solutions to certain degenerate equations. It has been also used by Fabes and Stroock in [6], [7] to study uniformly elliptic and parabolic operators with measurable coefficients and later adapted by Montanari in [20] to obtain a Harnack inequality for \mathcal{L}_0 belonging to a class of totally degenerate hypoelliptic operators. The same method has been successfully used by the authors in [23], in the study of Kolmogorov operators (1.8).

We finally recall some papers where the second order part of the operator \mathcal{L}_0 has non-smooth coefficients. We quote Sturm [27] and Zhang [29], that consider the operator (1.10) where \mathcal{L}_0 is uniformly parabolic, Citti, Garofalo and Lanconelli [5] and Lu [19], who consider the Schrödinger operator related to sum of square of Hörmander's vector fields $\mathcal{L}_\mathcal{V} = \sum_{k=1}^m X_k^2 + \mathcal{V}$, Zhang [30], who studies the analogous parabolic operator $\mathcal{L}_\mathcal{V} = \sum_{k=1}^m X_k^2 - \partial_t + \mathcal{V}$ as (1.6). Recently Bramanti and Brandolini in [4] consider operators, without potential function, of the following type: $\mathcal{L} = \sum_{i,j=1}^m a_{ij}(x) X_i X_j$, where a_{ij} belong to the Sarason class VMO . They extend to spaces of homogeneous type some regularity estimates.

We end this introduction with a short outline of this paper. In Section 2 we recall the known facts about homogeneous Lie groups and on the boundary value problems for \mathcal{L}_0 , that will be needed through in the sequel, then we state our main results. In Section 3 we discuss the main properties of the fundamental solution and of the Green function for \mathcal{L}_0 . In Section 4 we construct a Green function for $\mathcal{L}_\mathcal{V}$ by the Levi parametrix method; some L^p estimates and a pointwise lower bound for the Green function are proved. Then, in Section 5 we prove the results of this paper, in a preliminary statement only for bounded potentials \mathcal{V} , then, by a limiting argument, for every \mathcal{V} in the Stummel-Kato class.

2. Known facts and statement of main results

In this section we briefly recall the basic properties of the homogeneous Lie groups and the related homogeneous vector fields. We refer to the recent monograph [2] by Bonfiglioli, Lanconelli and Uguzzoni, for a more exhaustive treatment of that topic. We state our main results in the last part of the section.

A Lie group $\mathbb{G} = (\mathbb{R}^{N+1}, \circ)$ is said *homogeneous* if there exists a family of dilations $(\delta_\lambda)_{\lambda>0}$ of the form

$$\delta_\lambda : \mathbb{R}^{N+1} \rightarrow \mathbb{R}^{N+1}, \quad \delta_\lambda(x_1, \dots, \xi_N, t) = (\lambda^{\alpha_1} x_1, \dots, \lambda^{\alpha_N} \xi_N, \lambda^{\alpha_0} t)$$

for some positive $\alpha_1, \dots, \alpha_N, \alpha_0$, with the following property

$$(2.1) \quad \delta_\lambda(z \circ \zeta) = (\delta_\lambda z) \circ (\delta_\lambda \zeta), \quad \text{for every } z, \zeta \in \mathbb{R}^{N+1} \text{ and } \lambda > 0.$$

Hypotheses [H.1]–[H.2] imply that \mathbb{R}^N has a direct sum decomposition

$$\mathbb{R}^N = V_1 \oplus \dots \oplus V_n$$

such that, if we decompose any point $x \in \mathbb{R}^N$ as $x = x^{(1)} + \dots + x^{(n)}$ with $x^{(k)} \in V_k$, then the dilations are

$$(2.2) \quad \delta_\lambda(x^{(1)} + \dots + x^{(n)}, t) = (\lambda x^{(1)} + \dots + \lambda^n x^{(n)}, \lambda^2 t),$$

for any $\lambda > 0$. If we let $m_k = \dim V_k$, the natural number

$$Q = 2 + \sum_{k=1}^n k m_k$$

is usually called the *homogeneous dimension* of \mathbb{G} with respect to $(\delta_\lambda)_{\lambda>0}$. We also introduce the following δ_λ -homogeneous norms on \mathbb{R}^{N+1} and \mathbb{R}^N :

$$\|(x, t)\|_{\mathbb{G}} = \left(\sum_{k=1}^n |x^{(k)}|^{\frac{2n!}{k}} + |t|^{n!} \right)^{\frac{1}{2n!}} \quad |x|_{\mathbb{G}} = \left(\sum_{k=1}^n |x^{(k)}|^{\frac{2n!}{k}} \right)^{\frac{1}{2n!}}$$

($|x^{(k)}|$ is the Euclidean norm of $x^{(k)}$). We denote by

$$d(z, \zeta) = \|\zeta^{-1} \circ z\|_{\mathbb{G}}$$

the *quasi-distance* between two points $z, \zeta \in \mathbb{R}^{N+1}$, and by

$$B_r(z) = \{\zeta \in \mathbb{R}^{N+1} : d(z, \zeta) < r\}$$

the ball with center at z and radius r . Recall that there exists a positive constant c such that

$$(2.3) \quad d(z, w) \leq c(d(z, \zeta) + d(\zeta, w)), \quad d(z, w) \leq c d(w, z),$$

for every $z, \zeta, w \in \mathbb{R}^{N+1}$ (see [8], Proposition 1.4).

We also recall that, due to the fact that X_0, \dots, X_m only depend on the *space* variable x , the composition law \circ is Euclidean in the *time* variable t , i.e.

$$(2.4) \quad (x, t) \circ (y, s) = (\sigma(x, t, y, s), t + s)$$

for a suitable smooth function σ (see [11], Proposition 10.2). Moreover, since X_1, \dots, X_m and Y are homogeneous vector fields of degree 1 and 2, respectively, we have

$$(2.5) \quad \begin{aligned} ((x, t) \circ (y, s))^{(1)} &= x^{(1)} + y^{(1)}, \\ ((x, t) \circ (y, s))^{(k)} &= x^{(k)} + y^{(k)} + \sigma_k(x, t, y, s) \end{aligned}$$

for $k = 2, \dots, m$, where $\sigma_k(x, t, y, s)$ is a polynomial function that only depends on $x^{(k+1)} + \dots + x^{(m)}, t, y^{(k+1)} + \dots + y^{(m)}$ and s . As a consequence, the determinant of the Jacobian matrix of the function $z \mapsto z_0 \circ z$ equals one, thus the Lebesgue measure of \mathbb{R}^{N+1} is left-invariant under left translations, namely

$$(2.6) \quad \text{meas}(z_0 \circ E) = \text{meas}(E),$$

for every $z_0 \in \mathbb{R}^{N+1}$ and every measurable set $E \subset \mathbb{R}^{N+1}$.

Another consequence of the homogeneity of the vector fields X_1, \dots, X_m and Y is that they are of the form

$$(2.7) \quad \begin{aligned} X_k &= \sum_{j=1}^n a_{j-1}^k(x^{(1)}, \dots, x^{(j-1)}) \cdot \nabla^{(j)}, \quad k = 1, \dots, m, \\ Y &= \sum_{j=2}^n b_{j-2}(x^{(1)}, \dots, x^{(j-2)}) \cdot \nabla^{(j)} - \partial_t, \end{aligned}$$

where $\nabla^{(j)} = (0, \dots, 0, \partial_{x_1^{(j)}}, \dots, \partial_{x_{m_j}^{(j)}}, 0, \dots, 0)$ denotes the gradient with respect to the variable $x^{(j)}$ and a_j^k and b_j are δ_λ -homogeneous polynomial functions of degree j with values in V_{j+1} and V_{j+2} respectively. As a first consequence we have that $X_k^* = -X_k$ for $k = 1, \dots, m$ and $Y^* = -Y$, thus the formal adjoint of \mathcal{L}_0 is $\mathcal{L}_0^* = \sum_{k=1}^m X_k^2 - Y$.

Let us explicitly note that hypothesis [H.2] and formula (2.7) imply that, if we write \mathcal{L}_0 as

$$(2.8) \quad \mathcal{L}_0 \equiv \sum_{i,j=1}^N a_{i,j}(x) \partial_{x_i x_j} + \sum_{j=1}^N b_j(x) \partial_{x_j} - \partial_t,$$

then the $m \times m$ block matrix $(a_{i,j}(x))_{i,j=1,\dots,m}$ is constant and positive definite.

We next recall some results, due to Lanconelli and Pascucci [17], concerning the boundary value problem for \mathcal{L}_0 . Let $k \in \mathbb{N}$ and $\varepsilon > 0$ be two constants that will be chosen in the sequel. We denote

$$(2.9) \quad O = B_{\text{eucl}(ke_1, k+k\varepsilon)} \cap B_{\text{eucl}(-ke_1, k+k\varepsilon)},$$

where $B_{\text{eucl}(x,r)}$ is the Euclidean ball of \mathbb{R}^N with center at x and radius r . Moreover, for positive T we let

$$Q(T) = O \times]0, T[, \quad S = O \times \{0\}, \quad S(T) = O \times \{T\}, \quad \text{and} \quad M(T) = \partial O \times]0, T[$$

be the “unit” cylinder of \mathbb{R}^{N+1} , its lower and upper basis (resp.), and its lateral boundary. We will call *parabolic boundary* of $Q(T)$ the set

$$\partial_r Q(T) = S \cup (\partial O \times [0, T]).$$

Finally, for every positive R and for any $(\xi, \tau) \in \mathbb{R}^{N+1}$, we set

$$Q_R(\xi, \tau, T) = (\xi, \tau) \circ \delta_R(Q(R^{-2}T)),$$

and, analogously,

$$\begin{aligned} M_R(\xi, \tau, T) &= (\xi, \tau) \circ \delta_R(M(R^{-2}T)), \quad S_R(\xi, \tau) = (\xi, \tau) \circ \delta_R(S), \\ S_R(\xi, \tau, T) &= (\xi, \tau) \circ \delta_R(S(R^{-2}T)), \quad \partial_r Q_R(\xi, \tau, T) = (\xi, \tau) \circ \delta_R(\partial_r Q(R^{-2}T)) \end{aligned}$$

(note that, by (2.2) and (2.4), T is the true height of the sets $Q_R(\xi, \tau, T)$, $M_R(\xi, \tau, T)$ and $\partial_r Q_R(\xi, \tau, T)$, and $S_R(\xi, \tau, T) = Q_R(\xi, \tau, T) \cap \{(x, t) : t = \tau + T\}$). We also remark that, by (2.2) and (2.6), we have

$$\text{meas}(Q_R(\xi, \tau, R^2T)) = R^Q \text{meas}(Q(T)).$$

Moreover

$$(2.10) \quad \text{meas}(S_R(\xi, \tau)) = R^{Q-2} \text{meas}(S),$$

where, with a slight abuse of notations, $\text{meas}(S_R(\xi, \tau))$ is the N -dimensional measure of the set $S_R(\xi, \tau)$ and, obviously,

$$(2.11) \quad \text{meas}(Q_R(\xi, \tau, R^2T)) = T R^Q \text{meas}(S).$$

Consider the Cauchy-Dirichlet problem in the unit cylinder

$$(2.12) \quad \begin{cases} \mathcal{L}_0 u = f & \text{in } Q(T) \\ u = 0 & \text{in } \partial_r Q(T) \end{cases}$$

with $f \in C_0^\infty(Q(T))$. As noticed before, the $m \times m$ block matrix $(a_{i,j}(x))_{i,j=1,\dots,m}$ in (2.8) is constant and positive definite so that, in particular, $a_{11} > 0$. Then, by Proposition 2.4 and Theorem 2.5 in [17]¹ there exists a positive ε in the definition of O such that the Dirichlet problem (2.12) has a unique (classical) solution $u \in C(Q(T) \cup \partial_r Q(T)) \cap C^\infty(Q(T))$ (in the sequel ε in the definition of O will be always chosen as above).

We say that $G_0 : (Q(T) \cup \partial_r Q(T)) \times Q(T) \rightarrow \mathbb{R}$ is a Green function for $Q(T)$ if, for every $f \in C_0(S_R)$, the function

$$u(z) = - \int_{Q(T)} G_0(z, \zeta) f(\zeta) d\zeta$$

is solution of the Cauchy-Dirichlet problem (2.12). In [17], Theorem 2.7 it is proved that a Green function G_0 exists and is smooth out of the diagonal of the set $(Q(T) \cup \partial_r Q(T)) \times Q(T)$; $G_0(x, t, \xi, \tau) \geq 0$; for any $(x, t), (\xi, \tau) \in \mathbb{R}^{N+1}$, $G_0(x, t, \xi, \tau) = 0$ if, $t \leq \tau$.

¹in [17] it is assumed that \mathcal{L}_0 is the heat operator out of a compact set of \mathbb{R}^{N+1} . \mathcal{L}_0 can be suitably modified outside $Q(T)$ in order to fulfill such a requirement.

The function $G_0^*(z, \zeta) \equiv G_0(\zeta, z)$ is a Green function for the adjoint operator \mathcal{L}_0^* . For every positive R and for any $(\xi, \tau) \in \mathbb{R}^{N+1}$ the function $G_0((\xi, \tau) \circ \delta_R(\zeta), (\xi, \tau) \circ \delta_R(z))$ is a Green function for the set $Q_R(\xi, \tau, T)$. The Green function can be characterized as

$$(2.13) \quad G_0(x, t, y, \tau) = \Gamma_0(x, t, y, \tau) - h(x, t, y, \tau),$$

where $h(\cdot, \cdot, y, 0)$ is the solution of the boundary value problem

$$(2.14) \quad \begin{cases} \mathcal{L}_0 u = 0 & \text{in } Q(T) \\ u = 0 & \text{in } M(T) \\ u = \Gamma_0(\cdot, \cdot, y, 0) & \text{in } S \end{cases}$$

The existence of a generalized solution h can be proved by a standard method, based on the elliptic regularization procedure. An universal barrier at every point of $M(T) \cup S$ has been constructed in the proof of Theorem 2.5 in [17], then h attains the boundary data by continuity. Moreover, by the hypoellipticity of \mathcal{L}_0 , h is a smooth classical solution to $\mathcal{L}_0 u = 0$ in $Q(T)$. Since $G_0(\zeta, \cdot)$ is a Green function for the adjoint operator \mathcal{L}_0^* , we also have that h is smooth for $(x, t) \neq (y, 0)$. By the minimum principle it plainly follows $h \geq 0$, then

$$(2.15) \quad G_0(x, t, y, s) \leq \Gamma_0(x, t, y, s), \quad \text{for every } (x, t), (y, s) \in \mathbb{R}^{N+1}.$$

We finally note that, for any $\varphi \in C_0(\mathbb{R}^N)$, the function

$$u(x, t) = \int_{\mathbb{R}^N} \Gamma_0(x, t, y, 0) \varphi(y) dy$$

is a classical solution to the Cauchy problem $\mathcal{L}_0 u = 0$ in $\mathbb{R}^N \times \mathbb{R}^+$, $u(x, 0) = \varphi(x)$; as a consequence, for every $\varphi \in C_0(S)$, the function

$$v(x, t) = \int_S G_0(x, t, y, 0) \varphi(y) dy$$

is a classical solution to the Cauchy-Dirichlet problem $\mathcal{L}_0 u = 0$ in $Q(T)$, $u = \varphi$ in S and $u \equiv 0$ in $M(T)$.

We next state the main results of this note. For every $R, T > 0$ and $(\xi, \tau) \in \mathbb{R}^{N+1}$, consider the Cauchy-Dirichlet problem

$$(2.16) \quad \begin{cases} \mathcal{L}_v u = f & \text{in } Q_R(\xi, \tau, T) \\ u = 0 & \text{in } \partial_r Q_R(\xi, \tau, T) \end{cases}$$

with $f \in C_0(Q_R(\xi, \tau, T))$.

We say that u is a weak solution of (2.16) if it is a weak solution to $\mathcal{L}_V u = f$ in $Q_R(\xi, \tau, T)$, it belongs to $C(Q_R(\xi, \tau, T) \cup \partial_r Q_R(\xi, \tau, T))$ and attains the boundary data by continuity. We say that

$$G : (Q_R(\xi, \tau, T) \cup \partial_r Q_R(\xi, \tau, T)) \times Q_R(\xi, \tau, T) \rightarrow \mathbb{R}$$

is a Green function for (2.16) if $G(\cdot, w)$ is a weak solution to (2.16), for every $w \in Q_R(\xi, \tau, T)$.

Theorem 2.1. *The Cauchy-Dirichlet problem (2.16) has a unique weak solution u . Moreover a Green function G for $Q_R(\xi, \tau, T)$ exists, and the function $G^*(\zeta, z) = G(z, \zeta)$ is a Green function for the adjoint operator \mathcal{L}_V^* .*

Before stating our second result, we introduce two further notations. Let us consider the cylinder $Q_R(\xi, \tau, R^2)$ and, for every $\alpha, \beta, \gamma, \delta \in]0, 1[$: $\alpha < \beta < \gamma$, let us set

$$\begin{aligned} Q^- &= \{(x, t) \in Q_{\delta R}(\xi, \tau, R^2) : \tau + \alpha R^2 \leq t \leq \tau + \beta R^2\}, \\ Q^+ &= \{(x, t) \in Q_{\delta R}(\xi, \tau, R^2) : \tau + \gamma R^2 \leq t\}. \end{aligned}$$

Theorem 2.2. (Harnack). *Let $\mathcal{V} \in SK(\Omega, \mathcal{L}_0)$. Then there exist two constants $R_0 > 0$ and $\delta_0 \in]0, 1[$ such that, for every $Q_R(\xi, \tau, R^2) \subset\subset \Omega$, with $R \leq R_0$ and Q^+, Q^- as above, with $\delta \in]0, \delta_0[$, we have*

$$\sup_{Q^-} u \leq M \inf_{Q^+} u,$$

for every positive weak solution u of $\mathcal{L}_V u = 0$. Here M is a positive constant that depends on η_V, η_V^* and on the constants $\alpha, \beta, \gamma, \delta$.

Proposition 2.3. *If u is a weak solution of $\mathcal{L}_V u = 0$ in Ω , with $\mathcal{V} \in SK(\Omega, \mathcal{L}_0)$, then u is continuous. Moreover, for any $\beta \in]0, 1[$ there exists a positive constant C_β , dependent only on \mathcal{L}_0 and β , such that*

$$|u(z) - u(z_0)| \leq (C_\beta d(z, z_0)^\beta + 2\eta_V(5c^2 d(z, z_0)^{1-\beta})) \sup_{B_{4r}(z_0)} |u|$$

for every $z_0 \in \Omega$, $r \in]0, 1[$ such that $B_{4r}(z_0) \subset \Omega$ and for every $z \in B_{r_\beta}(z_0)$ (where $r_\beta = r^{\frac{1}{1-\beta}}$ and c is the constant in (2.3)).

If $\mathcal{V} \in L^{1,\lambda}(\Omega, \mathbb{G})$ with $\lambda \in]Q - 2, Q[$, then

$$|u(z) - u(z_0)| \leq C_\beta (1 + \|\mathcal{V}\|_{L^{1,\lambda}(\Omega, \mathbb{G})}) \sup_{B_{4r}(z_0)} |u| \cdot d(z, z_0)^\alpha,$$

where $\alpha = \min\{\beta, (1 - \beta)(\lambda - Q + 2)\}$.

3. Preliminary results

In this section we recall some result about the fundamental solution and to the Green function G_0 for operators satisfying assumptions [H.1]–[H.2]; we then prove a lower bound for G_0 . We end the section with some remarks on the Stummel-Kato class $SK(\Omega, \mathcal{L}_0)$.

In [11], Kogoj and Lanconelli prove the existence of a fundamental solution $\Gamma_0(z, \zeta)$ for the operators \mathcal{L}_0 satisfying conditions [H.1]–[H.2]. The main properties of Γ_0 are analogous to the properties of the heat kernel: Γ_0 is smooth in $\{(z, \zeta) \in \mathbb{R}^{N+1} \times \mathbb{R}^{N+1} : z \neq \zeta\}$; $\Gamma_0(x, t, \xi, \tau) \geq 0$; for any $(x, t), (\xi, \tau) \in \mathbb{R}^{N+1}$, $\Gamma_0(x, t, \xi, \tau) > 0$ if, and only if, $t > \tau$.

Γ_0 is invariant with respect to the translations of \mathbb{G} :

$$\Gamma_0(z, \zeta) = \Gamma_0(\zeta^{-1} \circ z, 0) \equiv \Gamma_0(\zeta^{-1} \circ z), \quad \text{for every } z, \zeta \in \mathbb{R}^{N+1},$$

and it is δ_λ -homogeneous of degree $2 - Q$ with respect to the dilations of \mathbb{G} :

$$(3.1) \quad \Gamma_0(\delta_\lambda(z)) = \lambda^{2-Q} \Gamma_0(z) \quad \text{for every } z \in \mathbb{R}^{N+1}, \lambda > 0;$$

as a consequence we have that

$$\lim_{|z| \rightarrow \infty} \Gamma_0(z) = 0; \quad \limsup_{z \rightarrow 0} \Gamma_0(z) = +\infty \quad \text{and} \quad \Gamma_0 \in L^1_{\text{loc}}(\mathbb{R}^{N+1}).$$

For every $\varphi \in C_0^\infty(\mathbb{R}^{N+1})$ and $z \in \mathbb{R}^{N+1}$ we have

$$(3.2) \quad \mathcal{L}_0 \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) \varphi(\zeta) d\zeta = -\varphi(z), \quad \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) \mathcal{L}_0 \varphi(\zeta) d\zeta = -\varphi(z),$$

and $\mathcal{L}_0 \Gamma_0(z, \cdot) = -\delta_z$ (the Dirac measure centered at z). Moreover

$$\int_{\mathbb{R}^N} \Gamma_0(x, t) dx = 1, \quad \text{for every } t > 0.$$

The function $\Gamma_0^*(z, \zeta) \equiv \Gamma_0(\zeta, z)$ is the fundamental solution of the adjoint operator \mathcal{L}_0^* .

Since Γ_0 is a δ_λ -homogeneous functions of degree $-Q + 2$ and the derivatives $X_j \Gamma_0$, for $j = 1, \dots, m$, are δ_λ -homogeneous functions of degree $-Q + 1$, from the general theory of function spaces on homogeneous Lie groups (see for instance Folland [8, Proposition (1.15)]; see also Rothschild and Stein [24] for a more developed analysis of differential operators on Lie groups) it follows that there exist a positive constant \tilde{C} such that, for every

$z_1, z_2, \zeta \in \mathbb{R}^{N+1}$, with $d(z_1, \zeta) \geq 2d(z_1, z_2)$ we have

$$\begin{aligned}
 |\Gamma_0(z_1, \zeta) - \Gamma_0(z_2, \zeta)| &\leq \tilde{C} \frac{d(z_1, z_2)}{d(z_1, \zeta)^{Q-1}}, \\
 (3.3) \quad |X_j \Gamma_0(z_1, \zeta) - X_j \Gamma_0(z_2, \zeta)| &\leq \tilde{C} \frac{d(z_1, z_2)}{d(z_1, \zeta)^Q}, \\
 |X_j^{(\zeta)} \Gamma_0(z_1, \zeta) - X_j^{(\zeta)} \Gamma_0(z_2, \zeta)| &\leq \tilde{C} \frac{d(z_1, z_2)}{d(z_1, \zeta)^Q},
 \end{aligned}$$

for $j = 1, \dots, m$ (the notation $X_j^{(\zeta)}$ means that the vector field X_j acts on the variable ζ). Moreover, if we set for $f \in L^p(\mathbb{R}^{N+1})$

$$(3.4) \quad T_f(z) = \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) f(\zeta) d\zeta$$

we have (see [8, Theorem (5.14)]):

i) if $1 < p < \frac{Q}{2}$, then $T_f \in L^q(\mathbb{R}^{N+1})$, for $\frac{1}{q} = \frac{1}{p} - \frac{2}{Q}$, and

$$(3.5) \quad \|T_f\|_q \leq C_p \|f\|_p;$$

ii) if $p > \frac{Q}{2}$, then

$$(3.6) \quad |T_f(z_1) - T_f(z_2)| \leq C_p d(z_1, z_2)^\alpha \|f\|_p, \quad \text{for every } z_1, z_2 \in \mathbb{R}^{N+1}$$

for some positive constant C_p and $\alpha = \min \left\{ 1, 2 - \frac{Q}{p} \right\}$.

Finally, for $j = 1, \dots, m$ we have

$$(3.7) \quad X_j T_f(z) = \int_{\mathbb{R}^{N+1}} X_j \Gamma_0(z, \zeta) f(\zeta) d\zeta$$

and, analogously,

i) if $1 < p < Q$, then $X_j T_f \in L^q(\mathbb{R}^{N+1})$, with $\frac{1}{q} = \frac{1}{p} - \frac{1}{Q}$, and

$$(3.8) \quad \|X_j T_f\|_q \leq C_p \|f\|_p;$$

ii) if $p > Q$, then

$$(3.9) \quad |X_j T_f(z_1) - X_j T_f(z_2)| \leq C_p d(z_1, z_2)^\alpha \|f\|_p, \quad \text{for } \alpha = 1 - \frac{Q}{p}.$$

Note that, by (3.5), formula (3.2) extends to

$$(3.10) \quad \mathcal{L}_0 \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) f(\zeta) \psi(\zeta) d\zeta = -f(z) \psi(z),$$

for any $f \in L^p_{\text{loc}}(\mathbb{R}^{N+1})$, with $1 < p < \frac{Q}{2}$, and any cut-off function ψ . We also recall that T_f is continuous from $L^1(\mathbb{R}^{N+1})$ to $L^{\frac{Q-2}{Q}}_{\text{weak}}(\mathbb{R}^{N+1})$; more specifically, there exists a positive constant C such that

$$\text{meas}\left\{z \in \mathbb{R}^{N+1} : |T_f(z)| > \alpha\right\} \leq \left(\frac{C}{\alpha}\right)^{\frac{Q-2}{Q}} \|f\|_{L^1(\mathbb{R}^{N+1})}, \quad \text{for every } \alpha > 0$$

(see [8, Prop. 1.10]) hence we will also use formula (3.10) for $f \in L^1_{\text{loc}}(\mathbb{R}^{N+1})$.

We next prove a lower bound for the Green function G_0 for \mathcal{L}_0 :

Proposition 3.1. *For any positive R and T and every $(\xi, \tau) \in \mathbb{R}^{N+1}$ and $\alpha \in]0, 1[$ there exist $\delta_0, \varepsilon \in]0, 1[$ such that*

$$G_0(x, t, y, \tau) \geq \frac{2\varepsilon}{\text{meas}(S_R(\xi, \tau))}$$

for every $\delta \in]0, \delta_0]$, $y \in S_{\delta R}(\xi, \tau)$ and $(x, t) \in Q_{\delta R}(\xi, \tau, T)$, such that $t \geq \tau + \alpha T$.

Proof. Thanks to the invariance of the operator with respect to the translations and the dilations of the Lie group \mathbb{G} , it is not restrictive to assume $(\xi, \tau) = (0, 0)$ and $R = 1$; we also denote $S = S_1(0, 0)$. Aiming to prove that $G_0(0, t, 0, 0) > 0$, for every $t \in]0, T]$, we recall (2.13). We first note that h is a bounded function in the set $\{(x, t, 0, 0) \in \overline{Q(T)} \times \{(0, 0)\}\}$. On the other hand $\Gamma_0(0, t) = t^{-\frac{Q-2}{2}} \Gamma_0(0, 1)$ by (3.1), then

$$G_0(0, t, 0, 0) = \Gamma_0(0, t) - h(0, t, 0, 0) \rightarrow +\infty$$

as $t \rightarrow 0+$. Then $G_0(0, t, 0, 0) > 0$ for any positive small t . Since $G_0 \geq 0$ by the Bony's maximum principle ([3, Theorem 3.2]) $G_0(0, t, 0, 0) > 0$ for $t \in]0, T]$. In order to prove our claim we let

$$\varepsilon = \frac{1}{4} \text{meas}(S) \min_{[\alpha T, T]} G_0(0, t, 0, 0);$$

it is not restrictive to suppose $\varepsilon < 1$. Since G_0 is a continuous function, there exists $\delta_0 \in]0, 1[$ such that

$$G_0(x, t, y, 0) \geq \frac{2\varepsilon}{\text{meas}(S)};$$

for every $(x, t) \in Q_\delta(0, 0, T)$, such that $t \geq \alpha T$ and $y \in S_\delta(0, 0)$, with $\delta \in]0, \delta_0]$. This proves the claim for $(\xi, \tau) = (0, 0)$ and $R = 1$. The result in the general case follows by using the invariance with respect to the Lie group structure. ■

Remark 3.2. In the above proof we used the assumption (1.4) in order to apply the Bony's strong maximum principle along the segment $\{(0, t) : 0 \leq t \leq T\}$, which is a trajectory of the vector field Y .

If we remove the requirement (1.4), then the hypothesis [H.2] and (2.7) yield $Y(0) = b_0 \cdot \nabla^{(2)} - \partial_t$, for some constant vector b_0 belonging to the second layer of the Lie algebra of \mathbb{G} . In this case, it is possible to adapt the proof of Proposition 3.1 by using the change of variable $(x, t) \mapsto (x - tb_0, t)$, however the assertion has to be stated according to the appropriate geometry.

We end this section with some further remarks about our definition of the Stummel-Kato class. We first recall the upper gaussian estimate for the fundamental solution provided by Kogoj and Lanconelli (see (5.1) in [11]), that allows us to establish whether a given function \mathcal{V} does satisfy condition (1.11): for every $t > 0, x \in \mathbb{R}^N$

$$(3.11) \quad \Gamma_0(x, t) \leq \frac{C}{t^{\frac{Q-2}{2}}} \exp\left(-\frac{|x|_{\mathbb{G}}}{Ct}\right)$$

for some positive constant C .

We next observe that, unlike in the usual definition of the Stummel-Kato class, in formula (1.12) we integrate \mathcal{V} on an unbounded set. A definition more similar to that one of the elliptic case should be given in terms of the following functions

$$(3.12) \quad \begin{aligned} \tilde{\eta}_{\mathcal{V}}(h) &= \sup_{(x,t) \in \Omega} \int_{\Omega \cap Q_h(x,t,h^2)} \Gamma_0(x,t,y,s) |\mathcal{V}(y,s)| dy ds, \\ \tilde{\eta}_{\mathcal{V}}^*(h) &= \sup_{(y,s) \in \Omega} \int_{\Omega \cap Q_h^*(y,s,h^2)} \Gamma_0(x,t,y,s) |\mathcal{V}(x,t)| dx dt; \end{aligned}$$

however, it turns out that $\tilde{\eta}_{\mathcal{V}}$ and $\tilde{\eta}_{\mathcal{V}}^*$ define the same class as $\eta_{\mathcal{V}}$ and $\eta_{\mathcal{V}}^*$.

Remark 3.3. We have that

$$\begin{aligned} \lim_{h \rightarrow 0} \eta_{\mathcal{V}}(h) = 0 &\iff \lim_{h \rightarrow 0} \tilde{\eta}_{\mathcal{V}}(h) = 0; \\ \lim_{h \rightarrow 0} \eta_{\mathcal{V}}^*(h) = 0 &\iff \lim_{h \rightarrow 0} \tilde{\eta}_{\mathcal{V}}^*(h) = 0. \end{aligned}$$

One of the two implications is an easy consequence of the inequalities $\tilde{\eta}_{\mathcal{V}}(h) \leq \eta_{\mathcal{V}}(h)$ and $\tilde{\eta}_{\mathcal{V}}^*(h) \leq \eta_{\mathcal{V}}^*(h)$. The other one easily follows from the homogeneity of Γ_0 , with respect to the dilation of the Lie group, and from the absolute continuity of the integral.

We next compare the spaces $SK(\Omega, \mathcal{L}_0)$ and the following Morrey spaces $L^{p,\lambda}(\Omega, \mathbb{G})$

Definition 3.4. Let Ω be an open subset of \mathbb{R}^{N+1} and let $p, \lambda \in \mathbb{R}$ be such that $1 \leq p < \infty$ and $0 \leq \lambda \leq Q$. We say that a function $f \in L^p_{\text{loc}}(\Omega)$ belongs to the Morrey space $L^{p,\lambda}(\Omega, \mathbb{G})$ if $\|f\|_{L^{p,\lambda}(\Omega, \mathbb{G})} < \infty$, where

$$\|f\|_{L^{p,\lambda}(\Omega, \mathbb{G})} = \left(\sup_{r>0, z \in \Omega} \frac{1}{r^\lambda} \int_{\Omega \cap B_r(z)} |f(w)|^p dw \right)^{\frac{1}{p}}.$$

Although the class $SK(\Omega, \mathcal{L}_0)$ and the spaces $L^{p,\lambda}(\Omega, \mathbb{G})$ are defined analogously to the classic ones, we observe some substantial differences between them. In the case of elliptic equations we have

$$(3.13) \quad L^{1,\lambda}(\Omega) \subseteq SK(\Omega) \subseteq L^{1,\mu}(\Omega), \quad 0 < \mu \leq n - 2 < \lambda < n.$$

An analogous result is true for the sum of the squares of the Hörmander fields, however in the case of parabolic (and degenerate parabolic) operators, we can prove the first inclusion, but the second one seems false (see Example 2.10 in [23]).

Proposition 3.5. We have

$$L^{1,\lambda}(\Omega, \mathbb{G}) \subseteq SK(\Omega, \mathcal{L}_0), \quad \text{for every } \lambda \in]Q - 2, Q[.$$

Proof. By using the homogeneity of the fundamental solution Γ_0 we find

$$(3.14) \quad \int_{\Omega \cap Q_h(x, t, h^2)} \Gamma_0(x, t, w) |\mathcal{V}(w)| dw \leq c_\lambda h^{\lambda - Q + 2} \frac{1}{h^\lambda} \int_{\Omega \cap B_h(x, t)} |\mathcal{V}(w)| dw,$$

for every $\mathcal{V} \in L^{1,\lambda}(\Omega, \mathbb{G})$, and by Remark 3.3 this inequality yields the desired inclusion. \blacksquare

Since we are concerned with weak solutions to $\mathcal{L}_\nu u = 0$, we need a sufficient condition for the requirement $\mathcal{V}u \in L^1_{\text{loc}}$. We recall that, in the case of uniformly elliptic operators, $\mathcal{V}u \in L^1_{\text{loc}}(\Omega)$ provided that u belongs to the space $H^1_{\text{loc}}(\Omega)$ (see Schechter [26]) and a similar condition holds for the sum of squares of Hörmander vector fields (see [5]). Here we prove that $\mathcal{V}u$ is locally integrable when u belongs to the Sobolev-Folland-Stein space $W^{2,1}(\Omega, \mathcal{L}_0)$, namely if the following norm

$$\|u\|_{W^{2,1}(\Omega, \mathcal{L}_0)} = \|u\|_{L^1(\Omega)} + \sum_{j=1}^m \|X_j u\|_{L^1(\Omega)} + \sum_{i,j=1}^m \|X_i X_j u\|_{L^1(\Omega)} + \|Y u\|_{L^1(\Omega)}$$

is finite.

Lemma 3.6. *If $u \in W_{\text{loc}}^{2,1}(\Omega, \mathcal{L}_0)$ and H, K are two compact sets such that $K \subset\subset H \subset \Omega$, then there exists a positive constant C , dependent only on H, K and $\mathcal{V} \in SK(\Omega, \mathcal{L}_0)$, such that*

$$(3.15) \quad \int_K |\mathcal{V}(z)u(z)|dz \leq C\|u\|_{W^{2,1}(H, \mathcal{L}_0)}.$$

Proof. We first claim that, for every $v \in C_0^\infty(\Omega)$, we have

$$(3.16) \quad \int_\Omega |\mathcal{V}(z)v(z)|dz \leq C_0\|v\|_{W^{2,1}(\Omega, \mathcal{L}_0)},$$

where C_0 is a positive constant dependent only on \mathcal{V} and on the support of v . Indeed, if we denote by H the support of v then

$$\begin{aligned} \int_\Omega |\mathcal{V}(z)v(z)|dz &\leq \int_H |\mathcal{V}(z)| \left(\int_H \Gamma_0(z, \zeta) |\mathcal{L}_0 v(\zeta)| d\zeta \right) dz \\ &\leq \int_H |\mathcal{L}_0 v(\zeta)| d\zeta \sup_{\eta \in H} \left(\int_H |\mathcal{V}(z)| \Gamma_0(z, \eta) dz \right) \leq \eta_{\mathcal{V}}^*(c_H) \|v\|_{W^{2,1}(\Omega, \mathcal{L}_0)} \end{aligned}$$

where $c_H = \max\{|t - \tau| : (x, t), (\xi, \tau) \in H\}$. This proves (3.16). The thesis follows from a standard density argument. \blacksquare

4. The Green function for $\mathcal{L}_{\mathcal{V}}$

In this section we use the *parametrix method* to prove the existence of a Green function G for the operator $\mathcal{L}_{\mathcal{V}}$, related to any given cylinder $Q_R(\xi, \tau, T)$. We construct G as a perturbation of G_0 :

$$G(z, w) = G_0(z, w) + \int_{Q_R} G_0(z, \eta) \Phi(\eta, w) d\eta,$$

for some unknown function Φ . A formal argument, based on the fact that $\mathcal{L}_0 G_0(z, w) = -\delta_w(z)$ and on the requirement that $\mathcal{L}_{\mathcal{V}} G(z, w) = -\delta_w(z)$ leads to the following Volterra equation for Φ

$$\Phi(z, \zeta) = \mathcal{V}(z)G_0(z, \zeta) + \int_{Q_R} \mathcal{V}(z)G_0(z, \eta) \Phi(\eta, \zeta) d\eta;$$

The successive approximation method then gives:

$$(4.1) \quad G(z, w) = G_0(z, w) + \sum_{k=1}^{\infty} J_k(z, w),$$

where

$$(4.2) \quad \begin{aligned} J_1(z, w) &= \int_{Q_R(\xi, \tau, T)} G_0(z, \eta) \mathcal{V}(\eta) G_0(\eta, w) d\eta \\ J_{k+1}(z, w) &= \int_{Q_R(\xi, \tau, T)} G_0(z, \eta) \mathcal{V}(\eta) J_k(\eta, w) d\eta. \end{aligned}$$

We will prove that these integrals J_k are well defined, then the L^p convergence of the series and we finally show that G is a Green function for $\mathcal{L}_\mathcal{V}$. Aiming to unify the notations, in the sequel we will denote $J_0 = G_0$ so that

$$J_1(z, w) = \int_{Q_R(\xi, \tau, T)} G_0(z, \eta) \mathcal{V}(\eta) J_0(\eta, w) d\eta.$$

Lemma 4.1. *The functions in (4.2) belong to $L^p(Q_R(\xi, \tau, T))$ for every $p \in [1, \frac{Q}{Q-2})$ and there exists a positive constant c_p such that*

$$(4.3) \quad \begin{aligned} \|J_k(z, \cdot), L^p(Q_R(\xi, \tau, T))\| &\leq c_p \eta_\mathcal{V}^*(T)^k, \\ \|J_k(\cdot, w), L^p(Q_R(\xi, \tau, T))\| &\leq c_p \eta_\mathcal{V}(T)^k, \end{aligned}$$

for every $w, z \in Q_R(\xi, \tau, T)$. Moreover, $J_k(x, t, y, s) = 0$ for every $t \leq s$. We can also write J_{k+1} as

$$(4.4) \quad J_{k+1}(z, w) = \int_{Q_R(\xi, \tau, T)} J_k(z, \eta) \mathcal{V}(\eta) G_0(\eta, w) d\eta.$$

Proof. We let $\tilde{\mathcal{V}}(\eta) = |\mathcal{V}(\eta)|$ and define \tilde{J}_k , by using formulas (4.2) with $\tilde{\mathcal{V}}$. Note that $\eta_\mathcal{V}(T) = \eta_{\tilde{\mathcal{V}}}(T)$ and $\eta_\mathcal{V}^*(T) = \eta_{\tilde{\mathcal{V}}}^*(T)$, then $\tilde{\mathcal{V}} \in SK(Q_T, \mathcal{L}_0)$ if and only if $\mathcal{V} \in SK(Q_T, \mathcal{L}_0)$. We first prove the inequalities in (4.3) for the non-negative functions \tilde{J}_k , the required estimates will follow from the trivial inequality $|J_k| \leq \tilde{J}_k$.

Due to the fact that every \tilde{J}_k is non-negative, (4.4) is immediate. In order to prove the L^p estimates for \tilde{J}_k we note that

$$(4.5) \quad \begin{aligned} \int_{Q_R(\xi, \tau, T)} G_0(z, \eta) |\mathcal{V}(\eta)| d\eta &\leq \int_{Q_R(\xi, \tau, T)} \Gamma_0(z, \eta) |\mathcal{V}(\eta)| d\eta = \eta_\mathcal{V}(T), \\ \int_{Q_R(\xi, \tau, T)} G_0(\eta, w) |\mathcal{V}(\eta)| d\eta &\leq \int_{Q_R(\xi, \tau, T)} \Gamma_0(\eta, w) |\mathcal{V}(\eta)| d\eta = \eta_\mathcal{V}^*(T), \end{aligned}$$

since $G_0 \leq \Gamma_0$.

We next define the sequences:

$$s_k = \sup_{z \in Q_R(\xi, \tau, T)} \int_{Q_R(\xi, \tau, T)} \tilde{J}_k(z, \eta) |\mathcal{V}(\eta)| d\eta,$$

$$s_k^* = \sup_{\eta \in Q_R(\xi, \tau, T)} \int_{Q_R(\xi, \tau, T)} |\mathcal{V}(z)| \tilde{J}_k(z, \eta) dz,$$

and we prove the following inequalities

$$(4.6) \quad s_k \leq \eta_{\mathcal{V}}(T)^{k+1}, \quad s_k^* \leq \eta_{\mathcal{V}}^*(T)^{k+1}$$

by induction on k . For $k = 1$ we have

$$s_1 \leq \sup_{z \in Q_R(\xi, \tau, T)} \int_{Q_R(\xi, \tau, T)} G_0(z, \zeta) |\mathcal{V}(\zeta)| \cdot$$

$$\cdot \left(\sup_{w \in Q_R(\xi, \tau, T)} \int_{Q_R(\xi, \tau, T)} G_0(w, \eta) |\mathcal{V}(\eta)| d\eta \right) d\zeta \leq \eta_{\mathcal{V}}^2(T),$$

by (4.5). The same argument and (4.2) give

$$s_{k+1} \leq s_k \eta_{\mathcal{V}}(T),$$

for any $k > 1$, then the first inequality in (4.6) is proved. The proof of the second one is analogous.

To obtain the L^p estimate for \tilde{J}_k we set, for $p \in \left[1, \frac{Q}{Q-2}\right)$:

$$\mathcal{T} = \left\{ \varphi \in C_0^\infty(Q_R(\xi, \tau, T)) : \varphi \geq 0, \quad \|\varphi\|_{L^{p'}(Q_R(\xi, \tau, T))} \leq 1 \right\}.$$

For any $\varphi \in \mathcal{T}$, we have

$$\int_{Q_R(\xi, \tau, T)} \tilde{J}_{k+1}(z, w) \varphi(w) dw = \int_{Q_R(\xi, \tau, T)} \tilde{J}_k(z, \eta) |\mathcal{V}(\eta)| \left(\int_{Q_R(\xi, \tau, T)} G_0(\eta, w) \varphi(w) dw \right) d\eta$$

$$\leq c_p \|\varphi\|_{L^{p'}(Q_R(\xi, \tau, T))} s_k \leq c_p \eta_{\mathcal{V}}^*(T)^{k+1},$$

by (4.4) and (4.6), where

$$c_p = \sup_{\eta \in Q_R(\xi, \tau, T)} \|\Gamma_0(\eta, \cdot)\|_{L^p(Q_R(\xi, \tau, T))}.$$

Thus

$$\|\tilde{J}_k(z, \cdot)\|_{L^p(Q_R(\xi, \tau, T))} = \sup_{\varphi \in \mathcal{T}} \int_{Q_R(\xi, \tau, T)} \tilde{J}_k(z, w) \varphi(w) dw \leq c_p \eta_{\mathcal{V}}^*(T)^k$$

and the first inequality in (4.3) holds for every $k \in \mathbb{N}$. In the same way we obtain the second one. Since $|J_k(z, w)| \leq \tilde{J}_k(z, w)$, the estimates (4.3) and the identity (4.4) also hold for every J_k , and the lemma is completely proved. ■

Proposition 4.2. *Let $T > 0$ be such that $\eta_{\mathcal{V}}(T) < 1$ and $\eta_{\mathcal{V}}^*(T) < 1$. Then*

i) for every $p \in [1, \frac{Q}{Q-2})$ the series introduced in (4.1) converges in $L^p(Q_R(\xi, \tau, T))$ and there exists a positive constant c_p such that

$$\|G(z, \cdot)\|_{L^p(Q_R(\xi, \tau, T))} \leq c_p \sum_{k=0}^{\infty} \eta_{\mathcal{V}}(T)^k; \quad \|G(\cdot, w)\|_{L^p(Q_R(\xi, \tau, T))} \leq c_p \sum_{k=0}^{\infty} \eta_{\mathcal{V}}^*(T)^k$$

ii) $G(x, t, y, s) = 0$ for $t \leq s$;

iii) the derivatives

(4.7)

$$\begin{aligned} X_j G(\cdot, w) &= X_j G_0(\cdot, w) + \sum_{k=1}^{\infty} \int_{Q_R(\xi, \tau, T)} X_j G_0(\cdot, \eta) \mathcal{V}(\eta) J_k(\eta, w) d\eta, \\ X_j G(z, \cdot) &= X_j G_0(z, \cdot) + \sum_{k=1}^{\infty} \int_{Q_R(\xi, \tau, T)} J_k(z, \eta) \mathcal{V}(\eta) X_j G_0(\eta, \cdot) d\eta \end{aligned}$$

are defined as elements of the space $L_{\text{loc}}^p(Q_R(\xi, \tau, T))$ for any $p \in [1, \frac{Q}{Q-1})$ and, for every compact set $K \subset Q_R(\xi, \tau, T)$, there exists a positive constant \bar{c}_p such that

$$\|X_j G(\cdot, w)\|_{L^p(K)} \leq \bar{c}_p \sum_{k=0}^{\infty} \eta_{\mathcal{V}}(T)^k, \quad \|X_j G(z, \cdot)\|_{L^p(K)} \leq \bar{c}_p \sum_{k=0}^{\infty} \eta_{\mathcal{V}}^*(T)^k,$$

for $j = 1, \dots, m$;

iv) for every $(x, t) \in Q_R(\xi, \tau, T)$,

$$\int_{S_R(\xi, \tau)} |G(x, t, y, \tau)| dy \leq \sum_{k=1}^{\infty} \eta_{\mathcal{V}}(T)^k; \quad \int_{S_R(\xi, \tau, T)} |G(y, \tau+T, x, t)| dy \leq \sum_{k=1}^{\infty} \eta_{\mathcal{V}}^*(T)^k;$$

v) for every $z \in Q_R(\xi, \tau, T)$, we have

$$\int_{Q_R(\xi, \tau, T)} |G(z, w) \mathcal{V}(w)| dw \leq \sum_{k=1}^{\infty} \eta_{\mathcal{V}}(T)^k, \quad \int_{Q_R(\xi, \tau, T)} |\mathcal{V}(\zeta) G(\zeta, z)| d\zeta \leq \sum_{k=1}^{\infty} \eta_{\mathcal{V}}^*(T)^k.$$

Proof. Assertions i) and ii) are direct consequences of Lemma 4.1. In order to prove iii), we show that the series

$$\sum_{k=1}^{\infty} \int_{Q_R(\xi, \tau, T)} X_j G_0(\cdot, \eta) \mathcal{V}(\eta) J_k(\eta, w) d\eta,$$

is convergent in $L_{\text{loc}}^p(Q_R(\xi, \tau, T))$.

Let K be a compact subset of $Q_R(\xi, \tau, T)$; for any $\varphi \in \mathcal{T}$, such that $\text{supp}(\varphi) \subset K$, we have

$$\begin{aligned} & \int_K \left(\int_{Q_R(\xi, \tau, T)} X_j G_0(z, \eta) \mathcal{V}(\eta) J_k(\eta, w) d\eta \right) \varphi(z) dz \\ & \leq \bar{c}_p \|\varphi\|_{L^{p'}(K)} \sup_{w \in Q_R(\xi, \tau, T)} \int_{Q_R(\xi, \tau, T)} |\mathcal{V}(\eta)| \tilde{J}_k(\eta, w) d\eta \\ & \leq \bar{c}_p \eta_{\mathcal{V}}^*(T)^{k+1} \end{aligned}$$

by (4.6), where

$$\bar{c}_p = \sup_{\eta \in Q_R(\xi, \tau, T)} \|X_j G_0(\cdot, \eta)\|_{L^p(K)}.$$

Hence

$$\begin{aligned} & \left\| \int_{Q_R(\xi, \tau, T)} X_j G_0(\cdot, \eta) \mathcal{V}(\eta) J_k(\eta, w) d\eta \right\|_{L^p(K)} \\ & = \sup_{\varphi \in \mathcal{T}} \int_K \left(\int_{Q_R(\xi, \tau, T)} X_j G_0(z, \eta) \mathcal{V}(\eta) J_k(\eta, w) d\eta \right) \varphi(z) dz \\ & \leq \bar{c}_p \eta_{\mathcal{V}}^*(T)^{k+1}. \end{aligned}$$

This proves the first identity in (4.7) and the estimate

$$\left\| X_j G(z, \cdot) \right\|_{L^p(K)} \leq \bar{c}_p \sum_{k=0}^{\infty} \eta_{\mathcal{V}}^*(T)^k.$$

The same argument gives the second identity and the corresponding estimate.

In order to prove *iv*), we note that, for every $k \in \mathbb{N}$,

$$\begin{aligned} & \left| \int_{S_R(\xi, \tau)} J_k(x, t, y, \tau) dy \right| \\ & \leq \left| \int_{Q_R(\xi, \tau, T)} J_{k-1}(x, t, \eta) \mathcal{V}(\eta) \left(\int_{S_R(\xi, \tau)} G_0(\eta, y, \tau) dy \right) d\eta \right| \\ & \leq \eta_{\mathcal{V}}^k(T), \end{aligned}$$

by (4.6). This proves the first estimate, the proof of the second one is analogous.

Finally, *v*) is an immediate consequence of (4.6). This concludes the proof of Proposition 4.2. ■

Corollary 4.3. *The function G defined in (4.1) is solution, in the distribution sense, of $\mathcal{L}_V G(\cdot, \zeta) = -\delta_\zeta$, $\mathcal{L}_V^* G(z, \cdot) = -\delta_z$. Namely: $G(\cdot, \zeta)$, $X_j G(\cdot, \zeta) \in L^p(Q_R(\xi, \tau, T))$, for some $p > 1$ and for $j = 1, \dots, m$, $GV \in L^1(Q_R(\xi, \tau, T))$ and $\forall \varphi \in C_0^\infty(Q_R(\xi, \tau, T))$, we have*

$$\int_{Q_R(\xi, \tau, T)} \left(\sum_{j=1}^m X_j G(z, \zeta) X_j \varphi(z) + G(z, \zeta) Y \varphi(z) - G(z, \zeta) \mathcal{V}(z) \varphi(z) \right) dz = \varphi(\zeta),$$

$$\int_{Q_R(\xi, \tau, T)} \left(\sum_{j=1}^m X_j G(z, \zeta) X_j \varphi(\zeta) - G(z, \zeta) Y \varphi(\zeta) - G(z, \zeta) \mathcal{V}(\zeta) \varphi(\zeta) \right) d\zeta = \varphi(z).$$

Proof. Since G_0 is the Green function of \mathcal{L}_0 , we have

$$\int_{Q_R(\xi, \tau, T)} \left(\sum_{j=1}^m X_j G_0(z, \zeta) X_j \varphi(z) + G_0(z, \zeta) Y \varphi(z) \right) dz = \varphi(\zeta),$$

for every $\varphi \in C_0^\infty(Q_R(\xi, \tau, T))$. For any $k \in \mathbb{N}$, we multiply the above identity by $\mathcal{V}(\zeta) J_{k-1}(\zeta, w)$ and integrate on $Q_R(\xi, \tau, T)$; we find

$$\int_{Q_R(\xi, \tau, T)} \left(\sum_{j=1}^m X_j \varphi(z) \int_{Q_R(\xi, \tau, T)} X_j G_0(z, \zeta) \mathcal{V}(\zeta) J_{k-1}(\zeta, w) d\zeta + \right.$$

$$\left. Y \varphi(z) \int_{Q_R(\xi, \tau, T)} G_0(z, \zeta) \mathcal{V}(\zeta) J_{k-1}(\zeta, w) d\zeta \right) dz = \int_{Q_R(\xi, \tau, T)} \varphi(\zeta) \mathcal{V}(\zeta) J_{k-1}(\zeta, w) d\zeta$$

and the first identity follows from the definition (4.1), (4.2) and from (4.7). In analogue way we can proceed for the second equality. \blacksquare

Proposition 4.4. *Let $T > 0$ be such that $\eta_V(T) < 1$ and $\eta_V^*(T) < 1$. Then, for any $(\xi, \tau) \in \mathbb{R}^{N+1}$ the function G defined by (4.1) is the Green function for the Cauchy-Dirichlet problem (2.16) related to $Q_R(\xi, \tau, T)$.*

Moreover $G^(w, z) = G(z, w)$ is the Green function for the Cauchy-Dirichlet problem*

$$(4.8) \quad \begin{cases} \mathcal{L}_V^* v = g & \text{in } Q_R(\xi, \tau, T) \\ v = 0 & \text{in } \partial_r^* Q_R(\xi, \tau, T) \end{cases}$$

with $g \in C_0(Q_R(\xi, \tau, T))$, namely the function

$$v(y, s) = - \int_{Q_R(\xi, \tau, T)} G^*(y, s, z) g(z) dz$$

is a weak solution to $\mathcal{L}_V^ v = g$ in $Q_R(\xi, \tau, T)$ and attains the boundary data by continuity (in (4.8) $\partial_r^* Q_R(\xi, \tau, T) = (\xi, \tau) \circ \delta_R(\partial_r^* Q(R^{-2}T))$, where $\partial_r^* Q(T) = S(T) \cup (\partial O \times [0, T])$).*

Proof. As said in Section 2, G is a Green function for the Cauchy-Dirichlet problem (2.16) if, for any $f \in C_0(Q_R(\xi, \tau, T))$, the function

$$u(z) = - \int_{Q_R(\xi, \tau, T)} G(z, \zeta) f(\zeta) d\zeta$$

is a weak solution to $\mathcal{L}_V u = f$ in $Q_R(\xi, \tau, T)$ and attains the boundary data by continuity. The fact that u solves $\mathcal{L}_V u = f$ is a direct consequence of Corollary 4.3.

In order to verify that u continuously vanishes at $\partial_r Q_R(\xi, \tau, T)$ we first note that

$$\mathcal{L}_0 u(z) = f(z) - \mathcal{V}(z)u(z),$$

then

$$u(z) = - \int_{Q_R(\xi, \tau, T)} G_0(z, \eta) f(\eta) d\eta + \int_{Q_R(\xi, \tau, T)} G_0(x, t, \eta) \mathcal{V}(\eta) u(\eta) d\eta,$$

for every $z \in Q_R(\xi, \tau, T)$. Since the function

$$u_0(z) = - \int_{Q_R(\xi, \tau, T)} G_0(z, \eta) f(\eta) d\eta$$

is a solution to the boundary value problem (2.12) (which is related to \mathcal{L}_0) it is known that it continuously vanishes at $\partial_r Q_R(\xi, \tau, T)$. Hence, we have to show that

$$(4.9) \quad \lim_{(x,t) \rightarrow (x_0, t_0)} \int_{Q_R(\xi, \tau, T)} G_0(x, t, \eta) \mathcal{V}(\eta) u(\eta) d\eta = 0,$$

for every $(x_0, t_0) \in \partial_r Q_R(\xi, \tau, T)$.

In order to prove (4.9) we observe that u is a bounded function, by Proposition 4.2 (i). Let us first consider a point $(x_0, t_0) \in S_R(\xi, \tau)$. Since \mathcal{V} belongs to the Stummel-Kato class, we have

$$\left| \int_{Q_R(\xi, \tau, T)} G_0(x, t, \eta) \mathcal{V}(\eta) u(\eta) d\eta \right| \leq \|u\|_\infty \eta_V(t) \rightarrow 0 \quad \text{as } t \rightarrow 0^+.$$

This proves that $u(x, t) \rightarrow 0$ as $(x, t) \rightarrow (x_0, t_0)$, for any $(x_0, t_0) \in S_R(\xi, \tau)$.

We next consider a point $(x_0, t_0) \in M_R(\xi, \tau, T)$. For every positive ε there exists a $\delta > 0$ such that

$$\left| \int_{(y,s) \in Q_R(\xi, \tau, T): t_0 - \delta < s < t_0} G_0(x, t, y, s) \mathcal{V}(y, s) u(y, s) dy ds \right| \leq \|u\|_\infty \eta_V(\delta) < \varepsilon$$

$\forall (x, t) \in Q_R(\xi, \tau, T)$, since $\mathcal{V} \in SK(\Omega, \mathcal{L}_0)$. Moreover there exists a positive constant \tilde{H} such that $G_0(x, t, y, s) \leq \tilde{H}$, for every $(x, t), (y, s) \in Q_R(\xi, \tau, T)$ such that $s < t_0 - \delta$ and $t > t_0 - \delta/2$. Hence

$$\lim_{(x,t) \rightarrow (x_0, t_0)} \int_{\eta \in Q_R(\xi, \tau, T): s < t_0 - \delta} G_0(x, t, \eta) \mathcal{V}(\eta) u(\eta) d\eta = 0,$$

that proves (4.9). This completes the proof that $u(x, t) \rightarrow 0$ as $(x, t) \rightarrow (x_0, t_0)$ for every $(x_0, t_0) \in \partial_r Q_R(\xi, \tau, T)$, thus G is a Green function for $\mathcal{L}_{\mathcal{V}}$ in $Q_R(\xi, \tau, T)$.

The proof that G^* is a Green function for $\mathcal{L}_{\mathcal{V}}^*$ in $Q_R(\xi, \tau, T)$ is analogous and will be omitted. \blacksquare

We next prove a lower bound for G analogous to Proposition 3.1.

Proposition 4.5. *For every $\alpha_0 \in]0, 1[$ there exist $\varepsilon, \delta_0 \in]0, 1[$, and $R_0 > 0$ such that, if $R \in]0, R_0]$, $\delta \in]0, \delta_0]$, and G is the Green function related to $Q_R(\xi, \tau, R^2)$ then*

$$G(x, t, y, \tau) \geq \frac{\varepsilon}{\text{meas}(S_R(\xi, \tau))}$$

for every $y \in S_{\delta R}(\xi, \tau)$ and for every $(x, t) \in Q_{\delta R}(\xi, \tau, R^2)$, such that $t \geq \tau + \alpha_0 R^2$

Proof. We claim that there exists a positive constant c such that

$$(4.10) \quad |J_k(x, t, \bar{x}, \bar{t})| \leq \frac{c}{(t - \bar{t})^{\frac{Q-2}{2}}} (\eta_{\mathcal{V}}(t - \bar{t}) + \eta_{\mathcal{V}}^*(t - \bar{t}))^k,$$

for every $k \in \mathbb{N}$ and any $(x, t), (\bar{x}, \bar{t}) \in Q_R(\xi, \tau, R^2)$. As a consequence, from (4.1) and Proposition 3.1 we get

$$\begin{aligned} G(x, t, y, \tau) &\geq G_0(x, t, y, \tau) - \frac{c}{(t - \tau)^{\frac{Q-2}{2}}} \sum_{k=1}^{\infty} (\eta_{\mathcal{V}}(t - \tau) + \eta_{\mathcal{V}}^*(t - \tau))^k \\ &\geq \frac{2\varepsilon}{\text{meas}(S_R(\xi, \tau))} - \frac{c}{(t - \tau)^{\frac{Q-2}{2}}} \sum_{k=1}^{\infty} (\eta_{\mathcal{V}}(t - \tau) + \eta_{\mathcal{V}}^*(t - \tau))^k, \end{aligned}$$

for any $y \in S_{\delta R}(\xi, \tau)$ and for every $(x, t) \in Q_{\delta R}(\xi, \tau, R^2)$, such that $t \geq \tau + \alpha_0 R^2$. Moreover, since $\text{meas}(S_R(\xi, \tau)) = R^{Q-2} \text{meas}(S)$ and $\alpha_0 R^2 \leq t - \tau \leq R^2$, the above inequality gives

$$G(x, t, y, \tau) \geq \frac{2\varepsilon}{\text{meas}(S_R(\xi, \tau))} - \frac{c'}{\text{meas}(S_R(\xi, \tau))} \sum_{k=1}^{\infty} (\eta_{\mathcal{V}}(t - \tau) + \eta_{\mathcal{V}}^*(t - \tau))^k,$$

for some positive constant c' . The claim then follows by choosing R_0 suitably small.

We next prove (4.10) by induction. We first recall (3.11), then

$$(4.11) \quad G_0(x, t, y, s) \leq \Gamma_0(x, t, y, s) \leq \frac{C}{(t-s)^{\frac{Q-2}{2}}},$$

for every $(x, t), (y, s) \in Q_R(\xi, \tau, R^2)$. Since

$$\begin{aligned} J_1(x, t, \bar{x}, \tau) &= \int_{\tau}^{\frac{t+\tau}{2}} \int_{S_R(\xi, \tau, s)} G_0(x, t, y, s) \mathcal{V}(y, s) G_0(y, s, \bar{x}, \tau) dy ds + \\ &+ \int_{\frac{t+\tau}{2}}^t \int_{S_R(\xi, \tau, s)} G_0(x, t, y, s) \mathcal{V}(y, s) G_0(y, s, \bar{x}, \tau) dy ds, \end{aligned}$$

we have

$$\begin{aligned} |J_1(x, t, \bar{x}, \tau)| &\leq \frac{C}{\left(\frac{t-\tau}{2}\right)^{\frac{Q-2}{2}}} \int_{\tau}^{\frac{t+\tau}{2}} \int_{S_R(\xi, \tau, s)} |\mathcal{V}(y, s)| \Gamma_0(y, s, \bar{x}, \tau) dy ds \\ &+ \frac{C}{\left(\frac{t-\tau}{2}\right)^{\frac{Q-2}{2}}} \int_{\frac{t+\tau}{2}}^t \int_{S_R(\xi, \tau, s)} \Gamma_0(x, t, y, s) |\mathcal{V}(y, s)| dy ds, \end{aligned}$$

so that (4.10) follows for $k = 1$.

For $k > 1$ we argue analogously: we write

$$\begin{aligned} J_{k+1}(x, t, \bar{x}, \tau) &= \int_{\tau}^{\frac{t+\tau}{2}} \int_{S_R(\xi, \tau, s)} J_k(x, t, y, s) \mathcal{V}(y, s) G_0(y, s, \bar{x}, \tau) dy ds \\ &+ \int_{\frac{t+\tau}{2}}^t \int_{S_R(\xi, \tau, s)} J_k(x, t, y, s) \mathcal{V}(y, s) G_0(y, s, \bar{x}, \tau) dy ds, \end{aligned}$$

and we use (4.6) in the second integral. This completes the proof. \blacksquare

5. Proof of the main results

In this section we prove the main results of this paper. As said in the introduction, the main difficulty is in the fact that \mathcal{V} is unbounded, then we cannot rely on the usual maximum principle. To overcome this problem, we first prove Proposition 2.3 and an uniqueness result for *bounded* solutions, then we prove the Harnack inequality (Theorem 2.2) for a bounded function \mathcal{V} , with the constant M depending on $\eta_{\mathcal{V}}$ and $\eta_{\mathcal{V}}^*$, but not on the L^∞ norm of \mathcal{V} . We finally remove the hypotheses of boundedness from u and \mathcal{V} , by using a technique due to Zhang [29]. We consider the sequence of operators

$$(5.1) \quad \mathcal{L}_{\mathcal{V}_m} = \mathcal{L}_0 + \mathcal{V}_m,$$

where

$$\mathcal{V}_m(x, t) = \begin{cases} -m & \text{if } \mathcal{V}(x, t) \leq -m \\ \mathcal{V}(x, t) & \text{if } -m < \mathcal{V}(x, t) < m, \\ m & \text{if } \mathcal{V}(x, t) \geq m. \end{cases}$$

and we approximate the solution u to $\mathcal{L}_{\mathcal{V}} = 0$ by a sequence u_m of solutions to $\mathcal{L}_{\mathcal{V}_m} = 0$. Since $\eta_{\mathcal{V}_m}(T) \leq \eta_{\mathcal{V}}(T)$, and $\eta_{\mathcal{V}_m}^*(T) \leq \eta_{\mathcal{V}}^*(h)$, the Harnack inequality for bounded solutions extends to u .

Lemma 5.1. *If u is a bounded weak solution of $\mathcal{L}_{\mathcal{V}}u = 0$ in Ω , with $\mathcal{V} \in SK(\Omega, \mathcal{L}_0)$, then u is continuous. Moreover, for any $\beta \in]0, 1[$ there exists a positive constant C_β , dependent only on \mathcal{L}_0 and β , such that*

$$|u(z) - u(z_0)| \leq (C_\beta d(z, z_0)^\beta + 2\eta_{\mathcal{V}}(5c^2 d(z, z_0)^{1-\beta})) \sup_{B_{4r}(z_0)} |u|$$

for every $z_0 \in \Omega$, $r \in]0, 1[$ such that $B_{4r}(z_0) \subset \Omega$ and for every $z \in B_{r_\beta}(z_0)$ (where $r_\beta = r^{\frac{1}{1-\beta}}$ and c is the constant in (2.3)).

If $\mathcal{V} \in L^{1,\lambda}(\Omega, \mathbb{G})$ with $\lambda \in]Q - 2, Q[$, then

$$|u(z) - u(z_0)| \leq C_\beta (1 + \|\mathcal{V}\|_{L^{1,\lambda}(\Omega, \mathbb{G})}) \sup_{B_{4r}(z_0)} |u| \cdot d(z, z_0)^\alpha,$$

where $\alpha = \min \{\beta, (1 - \beta)(\lambda - Q + 2)\}$.

Proof. Let $z_0 \in \Omega$, $r \in (0, 1)$ be such that $B_{4r}(z_0) \subset \Omega$ and let $z \in B_{r_\beta}(z_0)$. We choose $\varrho = 2d(z_0, z)^{1-\beta}$ and a function $\varphi \in C_0^\infty(B_{2\varrho}(z_0))$ such that $\varphi \equiv 1$ in $B_\varrho(z_0)$ and that $|X_j \varphi| \leq \frac{c}{\varrho}$, $|X_i X_j \varphi| \leq \frac{c}{\varrho^2}$, for $i, j = 1, \dots, m$ and $|Y \varphi| \leq \frac{c}{\varrho^2}$, for some positive constant c only depending on the operator \mathcal{L}_0 .

Since $\varrho \leq 2r$, we have $B_{2\varrho}(z_0) \subset B_{4r}(z_0) \subset \Omega$ and $(\varphi u): B_{2\varrho}(z_0) \rightarrow \mathbb{R}$ satisfies

$$\mathcal{L}_0(\varphi u) = \sum_{j=1}^m X_j^2(\varphi u) + Y(\varphi u) = \varphi \mathcal{L}_0 u + u \mathcal{L}_0 \varphi + 2 \sum_{j=1}^m X_j \varphi X_j u.$$

By the representation formula (3.10) we have that

$$\begin{aligned} (5.2) \quad u(z) &= - \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) \mathcal{L}_0 \varphi(\zeta) u(\zeta) d\zeta - 2 \sum_{j=1}^m \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) \langle X_j \varphi, X_j u \rangle d\zeta \\ &\quad + \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) \mathcal{V}(\zeta) u(\zeta) \varphi(\zeta) d\zeta \\ &= A_1(z) + A_2(z) + A_3(z), \quad \forall z \in B_\varrho(z_0). \end{aligned}$$

We explicitly remark that $d(z_0, \zeta) \geq 2d(z_0, z)$, for every $\zeta \in B_{2\varrho}(z_0) \setminus B_\varrho(z_0)$. From the first inequality in (3.3) we estimate the two terms in A_1 as follows

$$\begin{aligned} |A_1(z) - A_1(z_0)| &\leq \int_{\mathbb{R}^{N+1}} |\Gamma_0(z, \zeta) - \Gamma_0(z_0, \zeta)| \cdot |\mathcal{L}_0 \varphi(\zeta) u(\zeta)| d\zeta \\ &\leq \tilde{C} \sup_{B_{2\varrho}(z_0)} |u \mathcal{L}_0 \varphi| \int_{B_{2\varrho}(z_0) \setminus B_\varrho(z_0)} \frac{d(z_1, z_2)}{d(z_1, \zeta)^{Q-1}} d\zeta \leq C_\beta d(z, z_0)^\beta \sup_{B_{2\varrho}(z_0)} |u|, \end{aligned}$$

for some positive constant C_β depending on \tilde{C} in (3.3), on $\mathcal{L}_0 \varphi$ and on β .

We next consider A_2 . We integrate by parts

$$\begin{aligned} A_2(z) &= 2 \sum_{j=1}^m \int_{\mathbb{R}^{N+1}} X_j^{(\zeta)} (\Gamma_0(z, \zeta) X_j \varphi(\zeta)) u(\zeta) d\zeta \\ &= 2 \sum_{j=1}^m \int_{\mathbb{R}^{N+1}} \Gamma_0(z, \zeta) X_j^2 \varphi(\zeta) u(\zeta) d\zeta + 2 \sum_{j=1}^m \int_{\mathbb{R}^{N+1}} X_j^{(\zeta)} \Gamma_0(z, \zeta) X_j \varphi(\zeta) u(\zeta) d\zeta, \end{aligned}$$

(as in (3.3), the notation $X_j^{(\zeta)}$ means that the vector field X_j acts on the variable ζ). We then estimate the first sum by the same argument as A_1 ; for the second one we use the third inequality in (3.3).

We finally consider A_3 . Let us first observe that, in view of (2.3), we have $d(z, \zeta) \leq c^2 (d(z, z_0) + 2\varrho) \leq 5c^2 d(z_0, z)^{1-\beta}$ for every $\zeta \in \text{supp}(\varphi)$, and $z \in B_r(z_0)$. Moreover, if $\zeta = (\xi, \tau)$ and $z = (x, t)$, then $|t - \tau| \leq d^2(z, \zeta)$, so that

$$|A_3(z)| \leq \sup_{B_{2\varrho}(z_0)} |u| \int_{B_{2\varrho}(z_0)} \Gamma_0(z, \zeta) |\mathcal{V}(\zeta)| d\zeta \leq \sup_{B_{2\varrho}(z_0)} |u| \cdot \eta_\mathcal{V} (5c^2 d(z_0, z)^{1-\beta}),$$

for every $z \in B_\varrho(z_0)$. This proves the first claim of Lemma 5.1. The second assertion directly follows from Proposition 3.5 (see (3.14)). ■

We next prove a uniqueness result for Cauchy-Dirichlet problem (2.16).

Lemma 5.2. *If u is a bounded solution to the problem*

$$(5.3) \quad \begin{cases} \mathcal{L}_\mathcal{V} u = 0 & \text{in } Q_R(\xi, \tau, T) \\ u = 0 & \text{in } \partial_r Q_R(\xi, \tau, T), \end{cases}$$

then $u \equiv 0$.

Proof. By the maximum principle, if u and v are weak solutions of the problem

$$\begin{cases} \mathcal{L}_0 u = f & \text{in } Q_R(\xi, \tau, T) \\ u = 0 & \text{in } \partial_r Q_R(\xi, \tau, T) \end{cases}$$

with $f \in L^1(Q_R(\xi, \tau, T))$, then $u \equiv v$.

Hence, if u is a solution of the non-homogeneous problem

$$\begin{cases} \mathcal{L}_0 u + \mathcal{V}u = f & \text{in } Q_R(\xi, \tau, T) \\ u = 0 & \text{in } \partial_r Q_R(\xi, \tau, T) \end{cases}$$

with $f \in L^1(Q_R(\xi, \tau, T))$, then u is almost everywhere equal to

$$(5.4) \quad v(z) = \int_{Q_R(\xi, \tau, T)} G_0(z, \zeta) (\mathcal{V}(\zeta)u(\zeta) - f(\zeta)) d\zeta.$$

Suppose now that u is a solution of the homogeneous problem (5.3). We then have

$$u(x, t) = \int_{Q_R(\xi, \tau, T)} G_0(x, t, y, s) \mathcal{V}(y, s) u(y, s) dy ds.$$

for every $(x, t) \in Q_R(\xi, \tau, h)$. Then recalling that $G_0(x, t, y, s) = 0$ for $t \leq s$ we have for $t < \tau + \delta$

$$\|u\|_{L^\infty(Q_R(\xi, \tau, \delta))} \leq \eta_V(\delta) \|u\|_{L^\infty(Q_R(\xi, \tau, \delta))}.$$

Thus, if we choose δ such that $\eta_V(\delta) < 1$, we have $u \equiv 0$ in $Q_R(\xi, \tau, \delta)$. We then conclude the proof by iterating this method. \blacksquare

Arguing as above, we can easily prove the following property.

Remark 5.3. If the function \mathcal{V} is bounded and u is a solution to the problem

$$\begin{cases} \mathcal{L}_\mathcal{V} u = f & \text{in } Q_R(\xi, \tau, T) \\ u = 0 & \text{in } M_R(\xi, \tau, T) \\ u = g & \text{in } S_R(\xi, \tau) \end{cases}$$

for some $f \in L^1(Q_R(\xi, \tau, T))$ and $g \in C_0(S_R(\xi, \tau))$, then

$$u(z) = \int_{S_R(\xi, \tau)} G(z, y, \tau) g(y) dy - \int_{Q_R(\xi, \tau, T)} G(z, \eta) f(\eta) d\eta.$$

In order to state our next result, we introduce some further notations. For a given $(\xi, \tau) \in \mathbb{R}^{N+1}$ and $R > 0$, we set

$$Q_R^* = Q_R^*(\xi, \tau, R^2) = Q_R(\xi^*, \tau^*, R^2), \quad \text{where } (\xi^*, \tau^*) = (\xi, \tau) \circ (0, -R^2).$$

Note that $\tau^* = \tau - R^2$ by (2.4), then we may consider Q_R^* as the cylinder whose *upper basis* is centered at (ξ, τ) . We also set

$$M(R) = \sup_{Q_R^*} u, \quad m(R) = \inf_{Q_R^*} u \quad \text{Osc}(u, \xi, \tau, R) = M(R) - m(R).$$

Lemma 5.4. *Let $u \geq 0$ be a bounded solution of $\mathcal{L}_V u = 0$ in Q_R^* . Then there exist $\delta, \varrho \in (0, 1)$ and a positive R_0 , which depend on η_V and \mathcal{L}_0 , such that*

$$\text{Osc}(u, \xi, \tau, \delta R) \leq \varrho M(R)$$

for every $R \in]0, R_0]$.

Proof. The method is inspired by that in [29] (and has been used in [23]). Let ε, δ and R_0 be as in Proposition 4.5, and set

$$\mathcal{S} = \left\{ (x, \tau^*) \in S_R(\xi^*, \tau^*) : u(x, \tau^*) \geq \frac{M(R) + m(R)}{2} \right\},$$

Consider two possibilities.

Case 1: $\text{meas}(\mathcal{S}) \geq \frac{1}{2} \text{meas}(S_R(\xi^*, \tau^*))$.

Define the function

$$v(z) = \int_{S_R(\xi^*, \tau^*)} G_0(z, y, \tau^*) (u(y, \tau^*) - m(R)) dy + \int_{Q_R^*} G_0(z, \zeta) \mathcal{V}(\zeta) u(\zeta) d\zeta,$$

and note that it is a solution to

$$\begin{cases} \mathcal{L}_0 v = -\mathcal{V}u & \text{in } Q_R(\xi^*, \tau^*, R^2) \\ v = u - m(R) & \text{in } S_R(\xi^*, \tau^*) \\ v \leq u - m(R) & \text{in } M_R(\xi^*, \tau^*, R^2). \end{cases}$$

The function $u - m(R)$ is non-negative in Q_R^* and $\mathcal{L}_0(u - m(R)) = -\mathcal{V}u$, then, by the comparison principle, we find

$$u(z) - m(R) \geq \int_{S_R(\xi^*, \tau^*)} G_0(z, y, \tau^*) (u(y, \tau^*) - m(R)) dy + \int_{Q_R^*} G_0(z, \zeta) \mathcal{V}(\zeta) u(\zeta) d\zeta$$

for almost every $z \in Q_R^*$. We next apply Proposition 3.1 with $T = R^2$, and we obtain

$$\begin{aligned} \int_{S_R(\xi^*, \tau^*)} G_0(z, y, \tau^*) (u(y, \tau^*) - m(R)) dy &\geq \int_{\mathcal{S}} G_0(z, y, \tau^*) (u(y, \tau^*) - m(R)) dy \\ &\geq \frac{M(R) - m(R)}{2} \int_{\mathcal{S}} G_0(z, y, \tau^*) dy \geq \frac{M(R) - m(R)}{2} \int_{\mathcal{S}} \frac{\varepsilon}{\text{meas}(S_R(\xi^*, \tau^*))} dy \\ &\geq \frac{\varepsilon}{4} (M(R) - m(R)), \quad \text{for every } z \in Q_{\delta R}^*(\xi, \tau, (\delta R)^2). \end{aligned}$$

On the other hand, we have

$$\left| \int_{Q_R^*} G_0(z, \zeta) \mathcal{V}(\zeta) u(\zeta) d\zeta \right| \leq M(R) \eta_V(R^2),$$

where the integral is sufficiently small provided that we fix R_0 such that $\eta_{\mathcal{V}}(R^2) \leq \frac{\varepsilon}{8}$ for any $R \in]0, R_0]$. Observing that

$$m(\delta R) - m(R) \geq \frac{\varepsilon}{4}(M(R) - m(R)) - \frac{\varepsilon}{8}M(R)$$

it follows

$$M(\delta R) - m(\delta R) \leq \left(1 - \frac{\varepsilon}{4}\right)(M(R) - m(R)) + \frac{\varepsilon}{8}M(R) \leq \left(1 - \frac{\varepsilon}{8}\right)M(R).$$

This concludes the the proof in the first case, since $\varepsilon \in]0, 1[$.

Case 2: $\text{meas}(\mathcal{S}) \leq \frac{1}{2} \text{meas}(S_R(\xi^*, \tau^*))$. In this case we set

$$w(z) = \int_{S_R(\xi^*, \tau^*)} G_0(z, y, \tau^*)(M(R) - u(y, \tau^*))dy + \int_{Q_R^*} G_0(z, \zeta) \mathcal{V}(\zeta)u(\zeta)d\zeta.$$

Following the method used in Case 1 we find

$$\begin{aligned} M(R) - u(z) &\geq \int_{S_R(\xi^*, \tau^*) \setminus \mathcal{S}} G_0(z, y, \tau^*)(M(R) - u(y, \tau^*))dy - \frac{\varepsilon}{8}M(R) \\ &\geq \frac{\varepsilon}{4}(M(R) - m(R)) - \frac{\varepsilon}{8}M(R), \text{ for a. e. } z \in Q_{\delta R}^*(\xi, \tau, (\delta R)^2) \end{aligned}$$

and then

$$M(\delta R) - m(\delta R) \leq \left(1 - \frac{\varepsilon}{8}\right)M(R).$$

The proof of Lemma 5.4 is then accomplished. ■

Proposition 5.5. *Let R_0 and δ_0 as in Proposition 4.5. Let $u \geq 0$ be a solution of $\mathcal{L}_{\mathcal{V}}u = 0$ in Ω , $Q_R(\xi, \tau, R^2) \subset\subset \Omega$, with $R \leq R_0$, and let \mathcal{V} a bounded function. Then, for every $\alpha, \beta, \gamma, \delta \in]0, 1[$ such that $\alpha < \beta < \gamma$ and $\delta < \delta_0$ there exists a positive M that depends on $\eta_{\mathcal{V}}, \eta_{\mathcal{V}}^*$ and on the constants $\alpha, \beta, \gamma, \delta$, but does not depend on the norm $\|\mathcal{V}\|_{L^\infty}$, such that*

$$\sup_{Q^-} u \leq M \inf_{Q^+} u.$$

Proof. We first note that the boundedness of \mathcal{V} yields the continuity of u , by the representation formula (3.10) and a standard bootstrap argument. Then there exists $(\bar{x}, \bar{t}) \in \overline{Q^+}$ such that $u(\bar{x}, \bar{t}) = \min_{\overline{Q^+}} u$. It is not restrictive to assume $u(\bar{x}, \bar{t}) = 1$.

Following the line of the proof of Theorem 5.4 in [7], we consider, for every $r \in [0, \beta R^2]$, the following function

$$v(x, t) = \int_{S_R(\xi, \tau, r)} G(x, t, y, r)u(y, r)dy, \quad \forall (x, t) \in Q_R(\xi, \tau, R^2).$$

By the comparison principle (recall that \mathcal{V} is bounded and that $u \geq 0$) we obtain $u(x, t) \geq v(x, t)$, for every $(x, t) \in Q_R(\xi, \tau, R^2)$, then

$$(5.5) \quad u(\bar{x}, \bar{t}) \geq \int_{S_R(\xi, \tau, r)} G(\bar{x}, \bar{t}, y, r) u(y, r) dy.$$

Let us fix $\delta' = \frac{\delta + \delta_0}{2}$ and consider, for any $\lambda > 0$, the set

$$\mathcal{S}(r, \lambda) = \{y \in S_{\delta' R}(\xi, \tau, r) : u(y, r) \geq \lambda\}.$$

Then inequality (5.5) and Proposition 4.5 (with $\alpha_0 = \gamma - \beta$) imply that

$$(5.6) \quad 1 = u(\bar{x}, \bar{t}) \geq \int_{\mathcal{S}(r, \lambda)} G(\bar{x}, \bar{t}, y, r) u(y, r) dy \geq \frac{\lambda \varepsilon \text{meas}(\mathcal{S}(r, \lambda))}{\text{meas}(S_R(\xi, \tau))}.$$

We set

$$(5.7) \quad K = \frac{1}{2} \left(1 + \frac{1}{\varrho}\right) \quad r(\lambda) = \frac{R}{\delta} \left(\frac{4}{\varepsilon \lambda (1 - \varrho)}\right)^{\frac{1}{Q-2}}$$

where ϱ is the constant in Lemma 5.4, and we note that

$$Q_{\delta r(\lambda)}^*(\xi, \tau, (\delta r(\lambda))^2) \cap S_R(\xi, \tau, r) = S_{\delta r(\lambda)}(\zeta, \tau)$$

for every $r \in [t - (\delta r(\lambda))^2, t]$. Then

$$(5.8) \quad \begin{aligned} \text{meas}(Q_{\delta r(\lambda)}^*(\xi, \tau, (\delta r(\lambda))^2) \cap S_R(\xi, \tau, r)) &= \text{meas}(S_{\delta r(\lambda)}(\zeta, \tau)) \\ &\quad (\text{by the analogous of (2.10) for the } N\text{-dimensional measure}) \\ &= (\delta r(\lambda))^{Q-2} \cdot \text{meas}(S) = \frac{4R^{Q-2}}{\varepsilon \lambda (1 - \varrho)} \cdot \text{meas}(S). \end{aligned}$$

We next prove the following statement. Let $\lambda > 0$ and $(x, t) \in Q_{\delta' R}(\xi, \tau, R^2)$ with $t \leq \tau + \beta R^2$ be such that $u(x, t) \geq \lambda$ and that

$$Q_{r(\lambda)}^*(x, t, r(\lambda)^2) \subset Q_{\delta' R}(\xi, \tau, R^2).$$

Then there exists $(x', t') \in Q_{r(\lambda)}^*(x, t, r(\lambda)^2)$ such that $u(x', t') \geq K\lambda$.

Indeed, from (5.6) it follows that

$$\text{meas}\left(\mathcal{S}\left(r, \frac{\lambda}{2}(1 - \varrho)\right)\right) \leq \frac{2R^{Q-2}}{\lambda \varepsilon (1 - \varrho)} \text{meas}(S)$$

so that, by (5.8), there is a $(\xi', \tau') \in Q_{\delta r(\lambda)}^*(\xi, \tau, (\delta r(\lambda))^2) \cap S_R(\xi, \tau, r)$ such that $u(\xi', \tau') < \frac{\lambda}{2}(1 - \varrho)$. Our claim then follows from Lemma 5.4.

We next show that there exists a positive constant M such that $u(x, t) \leq M$ for every $(x, t) \in Q^-$. The thesis then follows, since $u(\bar{x}, \bar{t}) = \min_{\overline{Q^+}} u = 1$. Suppose, by contradiction, that there were a $z_0 \in Q^-$ such that $u(z_0) > M$. Then, repeating the arguments used above to obtain $u(x', t') \geq K\lambda$, there exists a sequence (z_j) such that

$$u(z_j) \geq M K^j, \quad z_{j+1} \in Q_{r_j}^*(z_j, r_j^2), \quad \text{where } r_j = r(M K^j),$$

provided that

$$(5.9) \quad Q_{r_j}^*(z_j, r_j^2) \subset Q_{\delta'R}(\xi, \tau, R^2), \quad \text{for every } j \in \mathbb{N}.$$

In order to prove (5.9) we note that

$$d(z_{j+1}, z_j) \leq c_0 r_j = c_0 \frac{R}{\delta} \left(\frac{4}{\varepsilon M(1-\varrho)K^j} \right)^{\frac{1}{Q-2}}$$

where $c_0 = \max_{z \in Q_1^*(0,0,1)} d(z, (0,0))$ (recall (2.1)). Hence

$$d(z_j, z_0) \leq c_0 \frac{R}{\delta} \left(\frac{4}{\varepsilon M(1-\varrho)K^j} \right)^{\frac{1}{Q-2}} \sum_{i=1}^{\infty} K^{-\frac{i}{Q-2}},$$

so that we can choose a positive M , that depends on α, δ, δ_0 but does not depend on R , such that (5.9) holds. Hence the sequence $u(z_j)$ is unbounded and we get a contradiction with the continuity of u . This accomplishes the proof. \blacksquare

In a similar way it is true the next result for the adjoint operator.

Remark 5.6. Let $v \geq 0$ be a solution of $\mathcal{L}_{\mathcal{V}}^* v = 0$ in $Q_R(\xi, \tau, R^2)$, where \mathcal{V} is a bounded function and it is in the class $SK(Q_R(\xi, \tau, R^2), \mathcal{L}_0)$. Then

$$\sup_{Q^+} v \leq M \inf_{Q^-} v,$$

for some positive constant M depending on $\eta_{\mathcal{V}}, \eta_{\mathcal{V}}^*$ and on the constants $\alpha, \beta, \gamma, \delta$, but that does not depend on $\|\mathcal{V}\|_{L^\infty}$.

Lemma 5.7. Let u be a solution of $\mathcal{L}_{\mathcal{V}} u = 0$ in Ω . Then, for any $z_0 \in \Omega$ there exists a compact neighborhood K of z_0 such that $K \subset \Omega$ and that u is the limit in $L^1(K)$ of a sequence $(u_m)_{m \in \mathbb{N}}$, where every u_m satisfies

$$\mathcal{L}_{\mathcal{V}_m} u_m = 0 \text{ in } K.$$

Moreover, for every compact set $H \subset \text{int}(K)$, there exists a positive constant c_H such that

$$(5.10) \quad |u_m(z)| \leq c_H \quad \forall z \in H, \quad \forall m \in \mathbb{N}.$$

Proof . Consider a cylindrical set $Q_R(\xi, \tau, T)$ such that $\overline{Q_R(\xi, \tau, T)} \subset \Omega$, and suppose that $\eta_V(T) < 1$ and $\eta_V^*(T) < 1$. Let also consider a text function $\varphi \in C_0^\infty(Q_R(\xi, \tau, T))$ such that $\varphi \equiv 1$ in a compact neighborhood K of z_0 such that $K \subset Q_R(\xi, \tau, T)$. We have

$$\mathcal{L}_{\mathcal{V}_m}(u\varphi) = \varphi\mathcal{L}_0u + u\mathcal{L}_0\varphi + 2 \sum_{j=1}^m X_j\varphi X_ju + u\varphi\mathcal{V}_m =$$

(recalling that u is solution of $\mathcal{L}_0u + \mathcal{V}u = 0$)

$$= u\mathcal{L}_0\varphi + 2 \sum_{j=1}^m X_j\varphi X_ju + (\mathcal{V}_m - \mathcal{V})u\varphi.$$

In the sequel we will set $f = 2 \sum_{j=1}^m X_j\varphi X_ju + u\mathcal{L}_0\varphi$. We also consider the Green function G_m related to $\mathcal{L}_{\mathcal{V}_m}$ and set

$$(5.11) \quad u_m(z) = - \int_{Q_R(\xi, \tau, T)} G_m(z, \zeta) f(\zeta) d\zeta.$$

We have

$$\begin{cases} \mathcal{L}_m(u_m - \varphi u) = -(\mathcal{V}_m - \mathcal{V})\varphi u & \text{in } Q_R(\xi, \tau, T) \\ u_m - \varphi u = 0 & \text{in } \partial_r Q_R(\xi, \tau, T), \end{cases}$$

and the function $(\mathcal{V}_m - \mathcal{V})\varphi u$ belongs to $L^1(Q_R(\xi, \tau, T))$, then, by Remark 5.3, we find

$$(u_m - \varphi u)(z) = \int_{Q_R(\xi, \tau, T)} G_m(z, \zeta) (\mathcal{V}_m - \mathcal{V})\varphi u(\zeta) d\zeta.$$

We next integrate over $Q_R(\xi, \tau, T)$ and use property (i) of Proposition 4.2, for $p = 1$. We obtain

$$\|u_m - \varphi u\|_{L^1(Q_R(\xi, \tau, T))} \leq c_1 \|(\mathcal{V}_m - \mathcal{V})\varphi u\|_{L^1(Q_R(\xi, \tau, T))},$$

for some constant c_1 that does not depend on m . On the other hand

$$|(\mathcal{V}_m(\zeta) - \mathcal{V}(\zeta))\varphi(\zeta)u(\zeta)| \leq |\mathcal{V}(\zeta)\varphi(\zeta)u(\zeta)|,$$

for almost every $\zeta \in Q_R(\xi, \tau, T)$ and the function $\mathcal{V}\varphi u \in L^1(Q_R(\xi, \tau, T))$, then

$$\lim_{m \rightarrow \infty} \|u_m - \varphi u\|_{L^1(Q_R(\xi, \tau, T))} = 0.$$

This proves the first claim.

We next prove (5.10). We set

$$\tilde{H} = \text{supp}((X_1\varphi)^2 + \cdots + (X_m\varphi)^2 + (Y\varphi)^2)$$

and note that $f(\zeta) = 0$ for every $\zeta \notin \tilde{H}$.

We next prove that there exists a positive constant \tilde{c} , that depends on H and \tilde{H} , but does not depend on m , such that

$$(5.12) \quad G_m(z, \zeta) \leq \tilde{c} \quad \text{for every } z \in H, \zeta \in \tilde{H}.$$

As a consequence, by formula (5.11) we obtain

$$|u_m(z)| \leq \int_{\tilde{H}} G_m(z, \zeta) f(\zeta) d\zeta \leq \tilde{c} \|f\|_{L^1(Q_R(\xi, \tau, T))},$$

for every $z \in H$, and the proof is concluded, since

$$\|f\|_{L^1(Q_R(\xi, \tau, T))} \leq k \left(\sum_{j=1}^m \|X_j u\|_{L^1(Q_R(\xi, \tau, T))} + \|u\|_{L^1(Q_R(\xi, \tau, T))} \right),$$

for a positive constant k that only depends on φ and on \mathcal{L}_0 .

We prove (5.12) by using the Harnack inequality stated in Remark 5.6. Let z be a point of H . For every $\zeta \in \tilde{H}$ we consider a cylindrical open set $\tilde{Q} \subset Q_R(\xi, \tau, T)$ such that $\zeta \in \tilde{Q}^+$ and $\tilde{Q} \cap H = \emptyset$. Since $G_m(z, \cdot)$ is a positive solution to $\mathcal{L}_{\mathcal{V}_m}^* v = 0$, by Remark 5.6 we have

$$\sup_{\tilde{Q}^+} G_m(z, \cdot) \leq M \inf_{\tilde{Q}^-} G_m(z, \cdot),$$

for some positive constant M that does not depend on m . On the other hand

$$\text{meas}(\tilde{Q}^-) \inf_{\tilde{Q}^-} G_m(z, \cdot) \leq \int_{Q_R(\xi, \tau, T)} G_m(z, \zeta) d\zeta \leq c_1 \sum_{k=0}^{\infty} \eta_{\mathcal{V}}(T)^k,$$

where c_1 is the constant appearing in the statement (i) of Proposition 4.2. Thus $G_m(z, \zeta) \leq \tilde{k}$, for every $z \in H$ and $\zeta \in \tilde{Q}^+$, where the constant \tilde{k} depends on M, c_1 and $\eta_{\mathcal{V}}(T)$. The estimate (5.12) then follows from a standard covering argument for the compact set \tilde{H} . This completes the proof. \blacksquare

Proof of Proposition 2.3. Let u be a solution of the equation $\mathcal{L}_{\mathcal{V}} u = 0$ in Ω . By Lemma 5.7, u is the limit, in $L_{\text{loc}}^1(\Omega)$ of a sequence of bounded functions $(u_m)_{m \in \mathbb{N}}$ such that $\mathcal{L}_{\mathcal{V}_m} u_m = 0$ in a suitable compact set $K \subset \Omega$. We then apply Lemma 5.1 to every function u_m , then there exists a subsequence $(u_{m_k})_{k \in \mathbb{N}}$ that converges uniformly to u in K . Thus the estimate of the modulus of continuity stated in Lemma 5.1 extends to u . This completes the proof. \blacksquare

Proof of Theorem 2.1. Let $Q_R(\xi_0, \tau_0, T)$ be any cylindrical set. If $\eta_V(T) < 1$ and $\eta_V^*(T) < 1$, the result immediately follows from Proposition 4.2.

If otherwise $\eta_V(T) \geq 1$, or $\eta_V^*(T) \geq 1$, we choose $h > 0$ such that $\eta_V(h) < 1$, $\eta_V^*(h) < 1$. Consider the cylinders

$$\begin{aligned} Q^{(s)}(T_0) &= O \times]s, s + T_0[, & Q_R^{(s)}(\xi_0, \tau_0, T_0) &= (\xi_0, \tau_0) \circ \delta_R Q^{(s)}(R^{-2}T_0) \\ S^{(s)} &= O \times \{s\}, & S_R^{(s)}(\xi_0, \tau_0) &= (\xi_0, \tau_0) \circ \delta_R S^{(s)}, \end{aligned}$$

and let $G^{(s)}$ denote the Green function related to $Q_R^{(s)}(\xi_0, \tau_0, h)$ (we can employ the argument used in Proposition 4.2 without any change). We then extend the definition of G given in Proposition 4.2 as follows: for every $(x, t) \in Q_R(\xi_0, \tau_0, T)$ such that $s + h < t \leq s + 2h$, we set

$$G(x, t, y, s) = \int_{S_R^{(s)}(\xi_0, \tau_0)} G^{(s+h)}(x, t, w, s + h) G^{(s)}(w, s + h, y, s) dw.$$

It is easy to verify that G is a Green function for the set $Q_R(\xi_0, \tau_0, 2h)$ and that $G^*(\zeta, z) = G(z, \zeta)$ is a Green function for the adjoint operator \mathcal{L}_V^* . For $(x, t) \in Q_R(\xi_0, \tau_0, T)$ such that $s + 2h < t \leq s + 4h$ we repeat the above argument and define the Green function in the set $Q_R(\xi_0, \tau_0, 4h)$ as

$$G(x, t, y, s) = \int_{S_R^{(s)}(\xi_0, \tau_0)} G^{(s+2h)}(x, t, w, s + 2h) G^{(s)}(w, s + 2h, y, s) dw.$$

After a finite number of iterations we obtain a Green function for the $Q_R(\xi_0, \tau_0, T)$. This completes the proof. ■

Proof of Theorem 2.2. As in Proposition 2.3, we obtain the result by using Lemma 5.7 and Proposition 5.5. ■

Acknowledgments. We wish to thank E. Lanconelli for his interest in this problem and for many helpful discussions.

References

- [1] ALEXOPOULOS, G. K.: Sub-Laplacians with drift on Lie groups of polynomial volume growth. *Mem. Amer. Math. Soc.* **155**, no. 739, (2002).
- [2] BONFIGLIOLI, A., LANCONELLI, E. AND UGUZZONI, F.: *Stratified Lie groups and potential theory for their sub-Laplacians*. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [3] BONY, J. M.: Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés. *Ann. Inst. Fourier (Grenoble)* **19** (1969), 277–304.
- [4] BRAMANTI, M. AND BRANDOLINI, L.: Estimates of BMO type for singular integrals on spaces of homogeneous type and applications to hypoelliptic PDEs. *Rev. Mat. Iberoamericana* **21** (2005), no. 2, 511–556.

- [5] CITTI, G., GAROFALO, N. AND LANCONELLI, E.: Harnack's inequality for sum of squares of vector fields plus a potential. *Amer. J. Math.* **115** (1993), no. 3, 699–734.
- [6] FABES, E. B. AND STROOCK, D. W.: The L^p –integrability of Green's functions and fundamental solutions for elliptic and parabolic equations. *Duke Math. J.* **51** (1984), no. 4, 997–1016.
- [7] FABES, E. B. AND STROOCK, D. W.: A new proof of Moser's parabolic Harnack inequality using the old ideas of Nash. *Arch. Rational Mech. Anal.* **96** (1986), no. 4, 327–338.
- [8] FOLLAND, G. B.: Subelliptic estimates and function spaces on nilpotent Lie groups. *Ark. Mat.* **13** (1975), no. 2, 161–207.
- [9] GAROFALO, N. AND LANCONELLI, E.: Level sets of the fundamental solution and Harnack inequality for degenerate equations of Kolmogorov type. *Trans. Amer. Math. Soc.* **321** (1990), no. 2, 775–792.
- [10] HÖRMANDER, L.: Hypoelliptic second order differential equations. *Acta Math.* **119** (1967), 147–171.
- [11] KOGOJ, A. AND LANCONELLI, E.: An invariant Harnack inequality for a class of hypoelliptic ultraparabolic equations. *Med. J. Math.* **1** (2004), no. 1, 51–80.
- [12] KOGOJ, A. AND LANCONELLI, E.: One-side Liouville theorems for a class of hypoelliptic ultraparabolic equations. In *Geometric Analysis of PDE and Several Complex Variables*, 305–312. *Contemp. Math.* **368**. Amer. Math. Soc., Providence, RI, 2005.
- [13] KOGOJ, A. AND LANCONELLI, E.: Link of groups and homogeneous Hörmander operators. *Proc. Amer. Math. Soc.* **135** (2007), no. 7, 2019–2030.
- [14] KUPTSOV, L. P.: On parabolic means. *Dokl. Acad. Nauk SSSR* **252** (1980), no. 2, 296–301.
- [15] KUSUOKA, S. AND STROOCK, D. W.: Applications of the Malliavin calculus, III. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **34** (1987), no. 2, 391–442.
- [16] LANCONELLI, E., PASCUCCI, A. AND POLIDORO, S.: Linear and nonlinear ultraparabolic equations of Kolmogorov type arising in diffusion theory and in finance. In *Nonlinear problems in mathematical physics and related topics, II*, 243–265. *Int. Math. Ser.* **2**. Kluwer/Plenum, New York, 2002.
- [17] LANCONELLI, E. AND PASCUCCI, A.: On the fundamental solution for hypoelliptic second order partial differential equations with non-negative characteristic form. *Ricerche Mat.* **48** (1999), no. 1, 81–106.
- [18] LANCONELLI, E. AND POLIDORO, S.: On a class of hypoelliptic evolution operators. *Rend. Sem. Mat. Univ. Politec. Torino* **52** (1994), no. 1, 29–63.
- [19] LU, G.: On Harnack's inequality for a class of strongly degenerate Schrödinger operators formed by vector fields. *Differential Integral Equations* **7** (1994), no. 1, 73–100.
- [20] MONTANARI, A.: Harnack inequality for totally degenerate Kolmogorov-Fokker-Planck operators. *Boll. Un. Mat. Ital. (7)* **10** (1996), no. 4, 903–926.

- [21] NAGEL, A., STEIN, E. M. AND WAINER, S.: Balls and metrics defined by vector fields. I. Basic properties. *Acta Math.* **155** (1985), no. 1-2, 103–147.
- [22] PASCUCI, A. AND POLIDORO, S.: Harnack inequalities and Gaussian estimates for a class of hypoelliptic operators. *Trans. Amer. Math. Soc.* **358** (2006), no. 11, 4873–4893.
- [23] POLIDORO, S. AND RAGUSA, M. A.: A Green function and regularity results for an ultraparabolic equation with a singular potential. *Adv. Differential Equations* **7** (2002), no. 11, 1281–1314.
- [24] ROTHSCCHILD, L. P. AND STEIN, E. M.: Hypoelliptic differential operators and nilpotent groups. *Acta Math.* **137** (1977), no. 3-4, 247–320.
- [25] SAFANOV, M. N.: Harnack’s inequality for elliptic equations and the Hölder property of their solutions. *J. Soviet Mathematics* **21** (1983), 851–863.
- [26] SCHECHTER, M.: *Spectra of partial differential operators*. North-Holland Series in Applied Mathematics and Mechanics **14**. North-Holland Publishing Co., Amsterdam, 1971.
- [27] STURM, K. T.: Harnack’s inequality for parabolic operators with singular low order terms. *Math. Z.* **216** (1994), no. 4, 593–611.
- [28] VAROPOULOS, N. TH., SALOFF-COSTE, L. AND COULHON, T.: *Analysis and geometry on groups* Cambridge Tracts in Mathematics **100**. Cambridge University Press, Cambridge, 1992.
- [29] ZHANG, Q.: On a parabolic equation with a singular lower order term. *Trans. Amer. Math. Soc.* **348** (1996), no. 7, 2811–2844.
- [30] ZHANG, Q.: On a parabolic equation with a singular lower order term, II. The Gaussian bounds. *Indiana Univ. Math. J.* **46** (1997), no. 3, 989–1020.

Recibido: 15 de julio de 2007

Revisado: 8 de octubre de 2007

Sergio Polidoro
Dipartimento di Matematica Pura e Applicata
Università degli Studi di Modena e Reggio Emilia
Via G. Campi 213/b
41100 Modena-Italy
`sergio.polidoro@unimore.it`

Maria Alessandra Ragusa
Dipartimento di Matematica
Università di Catania
Viale Andrea Doria, 6
95125 Catania, Italy
`maragusa@dmf.unict.it`