

This is the peer reviewed version of the following article:

Satellite-detected tropospheric nitrogen dioxide and spread of SARS-CoV-2 infection in Northern Italy / Filippini, T.; Rothman, K. J.; Goffi, A.; Ferrari, F.; Maffei, G.; Orsini, N.; Vinceti, M.. - In: SCIENCE OF THE TOTAL ENVIRONMENT. - ISSN 0048-9697. - 739:(2020), pp. 140278-.. [10.1016/j.scitotenv.2020.140278]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

02/05/2024 17:33

Satellite-detected tropospheric nitrogen dioxide and spread of SARS-CoV-2 infection in Northern Italy

Tommaso Filippini, Kenneth J. Rothman, Alessia Goffi, Fabrizio Ferrari, Giuseppe Maffei, Nicola Orsini, Marco Vinceti



PII: S0048-9697(20)33799-2

DOI: <https://doi.org/10.1016/j.scitotenv.2020.140278>

Reference: STOTEN 140278

To appear in: *Science of the Total Environment*

Received date: 2 May 2020

Revised date: 13 June 2020

Accepted date: 15 June 2020

Please cite this article as: T. Filippini, K.J. Rothman, A. Goffi, et al., Satellite-detected tropospheric nitrogen dioxide and spread of SARS-CoV-2 infection in Northern Italy, *Science of the Total Environment* (2020), <https://doi.org/10.1016/j.scitotenv.2020.140278>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Satellite-detected tropospheric nitrogen dioxide and spread of SARS-CoV-2 infection in Northern Italy

Authors: Tommaso Filippini¹, Kenneth J. Rothman^{2,3}, Alessia Goffi⁴, Fabrizio Ferrari⁴, Giuseppe Maffei⁴, Nicola Orsini⁵, Marco Vinceti^{1,2}

Affiliations:

¹Environmental, Genetic and Nutritional Epidemiology Research Center (CREAGEN), Section of Public Health, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy;

² RTI Health Solutions, Research Triangle Park, NC, USA;

³ Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA;

⁴Terraria, Milan, Italy;

⁵Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

Corresponding author:

Marco Vinceti, Dipartimento di Scienze Biomediche, Metaboliche e Neuroscienze – UNIMORE, Via Campi 287, 41125 Modena (Italy) – E-mail marco.vinceti@unimore.it

Abstract

Following the outbreak of Severe Acute Respiratory Syndrome CoronaVirus 2 (SARS-CoV-2) last December 2019 in China, Italy was the first European country to be severely affected, with the first local case diagnosed on 20 February 2020. The virus spread quickly, particularly in the North of Italy, with three regions (Lombardy, Veneto and Emilia-Romagna) being the most severely affected. These three regions accounted for more than 80% of SARS-CoV-2 positive cases when the tight lockdown was established (March 8). These regions include one of Europe's areas of heaviest air pollution, the Po valley. Air pollution has been recently proposed as a possible risk factor of SARS-CoV-2 infection, due to its adverse effect on immunity and to the possibility that

polluted air may even carry the virus. We investigated the association between air pollution and subsequent spread of the SARS-CoV-2 infection within these regions. We collected NO₂ tropospheric levels using satellite data available on the European Space Agency before the lockdown. Using a multivariable restricted cubic spline regression model, we compared NO₂ levels with SARS-CoV-2 infection prevalence rate at different time points after the lockdown, namely March 8, 22 and April 5, in the 28 provinces of Lombardy, Veneto and Emilia Romagna. We found little association of NO₂ levels with SARS-CoV-2 prevalence up to about 130 µmol/m², while a positive association was evident at higher levels at each time point. Notwithstanding the limitations of the use of aggregated data, these findings lend some support to the hypothesis that high levels of air pollution may favor the spread of the SARS-CoV-2 infection.

Keywords: covid-19; coronavirus; air pollution; nitrogen dioxide; Sentinel-5P; public health.

1. Introduction

The first indigenous case of COVID-19 in Italy was diagnosed on February 20, 2020. Despite the isolation of potential hotspots in eleven affected municipalities ('red areas') and two national lockdowns established on February 23 and March 8-9, the outbreak quickly spread throughout Northern Italy and particularly in the Lombardy, Veneto and Emilia-Romagna regions (Gabutti et al., 2020; Ministry of Health, 2020). The later and more intensive lockdown, which started on March 8 in almost the entire territory of these three regions and on March 9 over all of Italy, was effective in slowing and curbing the infection, with a flattening and then a reversal of the outbreak beginning 9 days later. Further restrictions were established on March 22, including closure of all non-essential activities and prohibition of movements out of the municipality of residence.

Northern Italy is the most heavily industrialized and polluted area of Italy (EEA, 2019). The rapid and more intense spread of COVID-19 in northern Italy, following the initial outbreak in densely populated and polluted Wuhan, China, prompted the hypothesis that the spread of the

SARS-CoV-2 infection and possibly the severity of the associated COVID-19 disease was related to high air pollution levels (Coccia, 2020; Conticini et al., 2020; RIAS, 2020; SIMA, 2020; Watts and Kommenda, 2020).

Though there is some evidence that air pollution may enhance the susceptibility to viral infections including that by SARS-CoV-2 (Barcelo, 2020; Chen et al., 2010; Domingo and Rovira, 2020; Peng et al., 2020; Tsatsakis et al., 2020), the connection is still speculative. Transmission of airborne infections such as that due to SARS-CoV-2 occurs mainly through droplets and aerosols in the vicinity of infected individuals (Gabutti et al., 2020), but airborne particles might serve as carriers for pathogens (Setti et al., 2020a; SIMA, 2020; Zhao et al., 2019). Another possible mechanism linking air pollution to SARS-CoV-2 infection and COVID-19 might be higher susceptibility to respiratory viral diseases induced by air pollutants, through damage to respiratory tissue, immune dysregulation, overexpression of inflammatory cytokines and chemokines leading to innate immune system hyper-activation (Conticini et al., 2020) or ACE-2 receptor overexpression (Frontera et al., 2020).

Among the most commonly investigated air contaminants, nitrogen dioxide (NO_2) is a secondary pollutant formed by combustion processes, primarily from road transport (41%), energy production (22%), energy use from households and commercial activities (13%), and from industry (13%) (EEA, 2020). NO_2 is used for air pollution monitoring, with a threshold yearly average of $40 \mu\text{g}/\text{m}^3$ according to both European and WHO standards (EEA, 2019). In 2017 around 7% of the European urban population was exposed to concentrations above the annual EU limit (EEA, 2019). Traffic-related NO_2 exposure has been shown to be associated with increased risk of asthma and decreased lung function (Bowatte et al., 2017; Cai et al., 2017; McCreanor et al., 2007) and also rhinitis in adult population (Cesaroni et al., 2008; de Marco et al., 2002). In addition, high NO_2 levels has been associated with increased mortality for all causes, cardiovascular and respiratory

mortality (Brunekreef et al., 2009; Eum et al., 2019; Hoek et al., 2013), and also pneumonia in older adults (Eum et al., 2019). With reference to SARS-CoV-2 infection, a recent spatial analysis has highlighted that up to 78% of COVID-19 deaths occurred in the five European areas located in Italy and Spain also showing the highest NO₂ levels, thus indicating a possible contribution to fatality by NO₂ long-term exposure (Ogen, 2020). In addition, a recent Chinese study showed a positive association between short-term exposure to air pollutants (i.e. particulate matter, sulfur dioxide, carbon monoxide, ozone and nitrogen dioxide) and newly diagnosed COVID-19 confirmed cases (Zhu et al., 2020), and other studies suggested a positive association with poor Air Quality Index (Li et al., 2020; Zoran et al., 2020).

The aim of our study was to investigate the relation between levels of air pollution in Northern Italy, assessed through remote-sensing satellite information on tropospheric NO₂, and subsequent spread of the SARS-CoV-2 infection using publicly available data on newly-diagnosed infected cases.

2. Methods

2.1. Study area

We studied the Lombardy, Veneto and Emilia-Romagna regions of Northern Italy, which have a population of 19.4 million people, around 30% of Italian population, and accounted for 82% of Italian cases of SARS-CoV-2 infection on March 8, 2020 (Ministry of Health, 2020) with more severe forms of the disease requiring intensive care unit admission (Frontera et al., 2020). These regions contribute about 40% of the gross national product (ISTAT, 2020b), and are also characterized by the highest number of workers in the international export business sector (ISTAT, 2020a).

2.2. Health endpoints

The number of new daily SARS-CoV-2 positive cases was available by region and province of diagnosis from the from national governmental website (CPD, 2020). From these data we computed prevalence rates, using population data for the study provinces from the Italian National Institute of Statistic website (ISTAT, 2020b).

2.3. Environmental exposure assessment

To assess the environmental pollution characterizing the study area before onset of the outbreak, we used daily information for the 1-23 February period from the Copernicus Open Access Hub-Sentinel-5P mission. Through this source, the European Space Agency (ESA) makes satellite data publicly available for the entire Europe (ESA, 2020). We focused on the geolocated (with a spatial resolution of $7 \times 7 \text{ km}^2$) tropospheric column of nitrogen dioxide reported by ESA Sentinel-5P, made available after Near Real Time (NRT) processing (typically 3 hours from sensing), and, after addition, Offline (OFFL) processing (5 days after sensing time) (Eskes et al., 2019).

We collected all satellite images related to the area of interest (the provinces of the three study regions) for each day of the analysis period, and we computed population-weighted average of NO_2 values (in $\mu\text{mol}/\text{m}^2$) for each province. Data about NO_2 were considered missing for one or more days when satellite coverage in those days decreased below 30% of the spatial units (provinces) for cloud-covered scenes or snow/ice on the surface influencing satellite image reliability (Eskes et al., 2019). We also validated satellite data through measured ground-level NO_2 concentration by monitoring stations, finding a high correlation between satellite and measured provincial average NO_2 , with Pearson correlation coefficients ranging from 0.34 (Sondrio) to 0.81 (Cremona).

To control for population mobility, a possible confounding factor, we collected data on mobile phone daily movements through anonymous data of the SIM cards of residents (approximately 27 million people) processed and made available by Teralytics (Polzer, 2020). Information on the position of mobile telephones is available through the Call Detail Records (CDR), from which transport models can describe daily trip chains. We used daily movements in Italy available from February 1 through March 27 divided by the population in each province. We also retrieved temperature data publicly available from the European Centre for Medium-Range Weather Forecasts website (ECMWF, 2020), and relative humidity that has been calculated using environmental temperature and dewpoint temperature according previous method (Lawrence, 2005). We also considered as a potential confounder the presence of international airports in the study provinces for which the traffic flow was higher than 100,000 passengers in January 2020 (ASSAEROPORTI, 2020). This criterion included airports in Milan, Bergamo, Varese in Lombardy, Venice, Verona and Treviso in Veneto, and Bologna in Emilia-Romagna.

2.4. Data analysis

To assess the average amount of air pollution preceding the onset of the SARS-CoV-2 outbreak, we modeled the time-series of daily NO₂ tropospheric levels from February 1st to 24th using a linear regression taking into account heteroskedasticity and autocorrelation up to 7 days using a Newey–West estimator (Newey and West, 1987). We also modeled these values after the lockdowns. We then examined the relation between NO₂ levels before February 24th, using the estimated NO₂ levels on February 12, which is the midpoint of the period before the lockdown, and SARS-CoV-2 infection prevalence rates at several time points: 14 days (corresponding to the period between outbreak onset and the establishment of the tight lockdown), 28 and 42 days, corresponding to March 8, March 22, and April 5. We used restricted cubic spline regression

analysis fitting a model with three fixed knots (10th, 50th and 90th percentiles) according to Harrell's method (Harrell, 2001) to assess the shape of the relation.

We also collected information on other possible confounding factors, including population density and age distribution in 2019, percentage of population commuting daily for work or school, percentage of single-member families, and percentage of dwellings with one resident only based on 2011 census data, the latest available at provincial level (ISTAT, 2020b).

We performed the analysis within a multivariable model adjusting for population density (number of inhabitants per km²), an index indicating age of the population (ratio between resident population aged ≥ 65 years and those aged ≤ 14 years), people mobility (measured using number of movements of mobile telephones before the lockdown), temperature (°C), relative humidity in the three subsequent periods (in percentage), and airport presence (yes/no). We alternatively further added to the model the percentage of population commuting daily outside the municipality of residence for work or study, single member families, or dwellings occupied by only one resident.

3. Results

Figure 1 shows the average tropospheric levels of NO₂ in study provinces before the disease outbreak, and in the subsequent periods during the spread of SARS-CoV-2 infection, when NO₂ levels steadily decreased. In particular, before the outbreak the average NO₂ levels were high in Lombardy (193 $\mu\text{mol}/\text{m}^2$), Milan (238 $\mu\text{mol}/\text{m}^2$) and Monza/Brianza (243 $\mu\text{mol}/\text{m}^2$), and they fell below 100 $\mu\text{mol}/\text{m}^2$ in all investigated provinces in the period following the tightest mobility restrictions.

Table 1 reports the province-specific prevalence rates of the SARS-CoV-2 infection in the three follow-up time points considered in the study, i.e. the one corresponding to the institution of the tight lockdown (March 8), and the next two 2-week periods, along with corresponding NO₂ pollution levels before and after the detection of the outbreak (February 24).

We first fit a spline curve to the crude data. There was an increase in prevalence rates can be noted at low levels of NO₂, with a plateau above 150 µmol/m² (Figure S1). In multivariable analysis adjusting for population density, elderly population, airport presence, people mobility before February 24, and temperature and relative humidity in the two weeks before prevalence assessment (Figure 2), NO₂ levels in the pre-outbreak period were not meaningfully associated with COVID-19 prevalence rates at any of the three follow-up time points, up to approximately 130 µmol/m² of NO₂ levels. Above 130 µmol/m² of NO₂ levels, however, there was a higher prevalence of COVID-19 with increasing levels of air pollution, with a similar curve relating air pollution to prevalence for each of the three time periods. Further addition of possible confounding factors, such as percentage of daily commuters outside the municipality of residence (Figure S2), percentage of single-member families (Figure S3), and percentage of dwellings occupied by only one resident (Figure S4), did not appreciably change the results. In these analyses, the limited precision inherent in these ecological data are reflected in the broad confidence bands.

4. Discussion

In this study, we found a positive association between levels of NO₂ levels and subsequence prevalence of SARS-CoV-2 positivity in Northern Italy, though this occurred only at high levels of NO₂. This relation is consistent with severe outdoor air pollution enhancing the spread of the SARS-CoV-2 virus, possibly by altering the immunological status and thus increasing individual susceptibility to infectious diseases (Ibironke et al., 2019; Marchini et al., 2020; Rivas-Santiago et al., 2015; Williams et al., 2011). The area of the study comprises the densest industrial, trading and agricultural zone in Italy, with an orography that favors the stagnation of pollutants. This region has seen some of the worst air pollution in Europe (EEA, 2019; Mazzola et al., 2010). Within this area, high levels of air pollution have been associated with increased mortality and

hospitalization for both cardiovascular and respiratory diseases (Carugno et al., 2016; Fattore et al., 2011).

A positive association between spread of the SARS-CoV-2 outbreak in Italy and excesses of particulate matter at monitoring stations has also been reported (CPD, 2020; Setti et al., 2020b), and in the US an increase of $1 \mu\text{g}/\text{m}^3$ in particulate matter ($\text{PM}_{2.5}$) has been associated with a 15% increase in mortality from COVID-19 (Wu et al., 2020). High air pollution levels could also have favored the spread of another coronavirus-induced disease, SARS, based on the positive correlation between air pollution and its lethality in Chinese provinces (Cui et al., 2003). In addition, a study carried out in Milan metropolitan area (Lombardy region) found a positive association between COVID-19 new daily cases and particulate matter levels, with Pearson correlation coefficient between average and maximum PM_{10} levels and daily new cases of 0.35 and 0.51, respectively (Zoran et al., 2020). We used tropospheric NO_2 levels as a proxy for overall air pollution. This metric was closely associated with ground levels of this contaminant in a validation study in the study area as well as in other European urban areas (Ialongo et al., 2020; Lorente et al., 2019).

In vitro studies have shown that NO_2 can alter expression and synthesis of pro-inflammatory mediators in human bronchial cells (Bayram et al., 2001; Blomberg et al., 1997; Blomberg et al., 1999; Devalia et al., 1993; Mirowsky et al., 2016), thus increasing inflammation and hyperresponsiveness of epithelial cells. Also, both direct cytotoxicity and cytokine-mediated has been reported after NO_2 exposure of epithelial cells in the lung (Ayyagari et al., 2004; Persinger et al., 2002). Overall, these observations provide some biological plausibility for the association we and other detected. In addition, NO_2 levels are likely to mirror the air levels of other contaminants, such as particulate matter (Bigi et al., 2012), which also has deleterious effects on the immunity (Blomberg, 2000; Glencross et al., 2020).

Despite biologic plausibility, this study has important limitations. Most important is the potential for ecologic bias, because we used aggregated data. With aggregated data we could not take into account individual risk factors that affected the spread of infection, nor control directly for confounding at the individual level. Factors such as working in health care delivery, travel to China or other infected countries, residence in nursing homes, and individual contact with infected relatives, friends, or workmates were not controlled and differences in the distributions of these factors may have influenced the results. A further limitation was the inability to examine other clinical endpoints of COVID-19, but information on hospitalizations, deaths, and other outcomes were not publicly available at the provincial level. Finally, the effect estimates we computed were imprecise, as reflected in broad confidence intervals. Although these limitations preclude strong inferences from these ecologic data, our findings appear to confirm an association between heavy air pollution and spread of the SARS-CoV-2 virus.

Declarations of interest: none

Abbreviation list:

CDR: Call Detail Records

ESA: European Space Agency

NO₂: nitrogen dioxide

NRT: Near Real Time processing

OFFL: Offline processing

SARS-CoV-2: Severe Acute Respiratory Syndrome CoronaVirus 2

References

- ASSAEROPORTI. January 2020 statistics, Associazione Italiana Gestori di Aeroporti, 2020. http://assaeroporti.com/statistiche_202001/ (Accessed: 12 June 2020)
- Ayyagari VN, Januszkiewicz A, Nath J. Pro-inflammatory responses of human bronchial epithelial cells to acute nitrogen dioxide exposure. *Toxicology* 2004; 197: 149-64. doi: 10.1016/j.tox.2003.12.017

- Barcelo D. An environmental and health perspective for COVID-19 outbreak: Meteorology and air quality influence, sewage epidemiology indicator, hospitals disinfection, drug therapies and recommendations. *J Environ Chem Eng* 2020; 104006. doi: 10.1016/j.jece.2020.104006
- Bayram H, Sapsford RJ, Abdelaziz MM, Khair OA. Effect of ozone and nitrogen dioxide on the release of proinflammatory mediators from bronchial epithelial cells of nonatopic nonasthmatic subjects and atopic asthmatic patients in vitro. *J Allergy Clin Immunol* 2001; 107: 287-94. doi: 10.1067/mai.2001.111141
- Bigi A, Ghermandi G, Harrison RM. Analysis of the air pollution climate at a background site in the Po valley. *J Environ Monit* 2012; 14: 552-63. doi: 10.1039/c1em10728c
- Blomberg A. Airway inflammatory and antioxidant responses to oxidative and particulate air pollutants - experimental exposure studies in humans. *Clin Exp Allergy* 2000; 30: 310-7. doi: 10.1046/j.1365-2222.2000.00814.x
- Blomberg A, Krishna MT, Bocchino V, Biscione GL, Shute JK, Kelly FJ, et al. The inflammatory effects of 2 ppm NO₂ on the airways of healthy subjects. *Am J Respir Crit Care Med* 1997; 156: 418-24. doi: 10.1164/ajrccm.156.2.9612042
- Blomberg A, Krishna MT, Helleday R, Soderberg M, Ledin MC, Kelly FJ, et al. Persistent airway inflammation but accommodated antioxidant and lung function responses after repeated daily exposure to nitrogen dioxide. *Am J Respir Crit Care Med* 1999; 159: 536-43. doi: 10.1164/ajrccm.159.2.9711068
- Bowatte G, Erbas B, Lodge CJ, Knibbs LD, Gurrin LC, Marks GB, et al. Traffic-related air pollution exposure over a 5-year period is associated with increased risk of asthma and poor lung function in middle age. *Eur Respir J* 2017; 50. doi: 10.1183/13993003.02357-2016
- Brunekreef B, Beelen R, Hoek G, Schouten L, Bausch-Goldbohm S, Fischer P, et al. Effects of long-term exposure to traffic-related air pollution on respiratory and cardiovascular mortality in the Netherlands: the NLCS-AIR study. *Res Rep Health Eff Inst* 2009; 5-71; discussion 73-89
- Cai Y, Zijlema WL, Doiron D, Blangiardo M, Burton PR, Fortier I, et al. Ambient air pollution, traffic noise and adult asthma prevalence: a BioSHaRE approach. *Eur Respir J* 2017; 49. doi: 10.1183/13993003.02127-2015
- Carugno M, Consonni D, Randi G, Catelan D, Grisotto L, Bertazzi PA, et al. Air pollution exposure, cause-specific deaths and hospitalizations in a highly polluted Italian region. *Environ Res* 2016; 147: 415-24. doi: 10.1016/j.envres.2016.03.003
- Cesaroni G, Badaloni C, Porta D, Forastiere F, Perucci CA. Comparison between various indices of exposure to traffic-related air pollution and their impact on respiratory health in adults. *Occup Environ Med* 2008; 65: 683-90. doi: 10.1136/oem.2007.037846
- Chen PS, Tsai FT, Lin CK, Yang CY, Chan CC, Young CY, et al. Ambient influenza and avian influenza virus during dust storm days and background days. *Environ Health Perspect* 2010; 118: 1211-6. doi: 10.1289/ehp.0901782
- Coccia M. Factors determining the diffusion of COVID-19 and suggested strategy to prevent future accelerated viral infectivity similar to COVID. *Sci Total Environ* 2020; 729: 138474. doi: 10.1016/j.scitotenv.2020.138474
- Conticini E, Frediani B, Caro D. Can atmospheric pollution be considered a co-factor in extremely high level of SARS-CoV-2 lethality in Northern Italy? *Environ Pollut* 2020; 261: 114465. doi: 10.1016/j.envpol.2020.114465
- CPD. Italian Civil Protection Department: COVID-19 data, 2020. <https://github.com/pcm-dpc/COVID-19> (Accessed: 12 June 2020)
- Cui Y, Zhang ZF, Froines J, Zhao J, Wang H, Yu SZ, et al. Air pollution and case fatality of SARS in the People's Republic of China: an ecologic study. *Environ Health* 2003; 2: 15. doi: 10.1186/1476-069X-2-15

- de Marco R, Poli A, Ferrari M, Accordini S, Giammanco G, Bugiani M, et al. The impact of climate and traffic-related NO₂ on the prevalence of asthma and allergic rhinitis in Italy. *Clin Exp Allergy* 2002; 32: 1405-12. doi: 10.1046/j.1365-2745.2002.01466.x
- Devalia JL, Campbell AM, Sapsford RJ, Rusznak C, Quint D, Godard P, et al. Effect of nitrogen dioxide on synthesis of inflammatory cytokines expressed by human bronchial epithelial cells in vitro. *Am J Respir Cell Mol Biol* 1993; 9: 271-8. doi: 10.1165/ajrcmb/9.3.271
- Domingo JL, Rovira J. Effects of air pollutants on the transmission and severity of respiratory viral infections. *Environ Res* 2020; 187: 109650. doi: 10.1016/j.envres.2020.109650
- ECMWF. ERA5 dataset European Centre for Medium-Range Weather Forecasts, 2020. <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5> (Accessed: 12 June 2020)
- EEA. EEA Report No 10/2019 - Air quality in Europe 2019. In: European Environment Agency, editor, Copenhagen, Denmark, 2019. <https://www.eea.europa.eu/publications/air-quality-in-europe-2019> (Accessed: 12 June 2020)
- EEA. The European environment state and outlook 2020. In: Agency EE, editor, 2020. <https://www.eea.europa.eu/publications/soer-2020> (Accessed: 12 June 2020)
- ESA. Sentinel-5P data. European Space Agency, 2020. https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-5P (Accessed: 12 June 2020)
- Eskes H, van Geffen J, Boersma F, Eichmann K-U, Apituley A, Pedergnana M, et al. Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Nitrogen dioxide. In: Institute RNM, editor, 2019. <https://sentinel.esa.int/documents/247904/2474726/Sentinel-5P-Level-2-Product-User-Manual-Nitrogen-Dioxide> (Accessed: 12 June 2020)
- Eum KD, Kazemiparkouhi F, Wang B, Manjourides J, Pun V, Pavlu V, et al. Long-term NO₂ exposures and cause-specific mortality in American older adults. *Environ Int* 2019; 124: 10-15. doi: 10.1016/j.envint.2018.12.060
- Fattore E, Paiano V, Borgini A, Tittarelli A, Bertoldi M, Crosignani P, et al. Human health risk in relation to air quality in two municipalities in an industrialized area of Northern Italy. *Environ Res* 2011; 111: 1321-7. doi: 10.1016/j.envres.2011.06.012
- Frontera A, Cianfanelli L, Vlachos K, Landoni G, Cremona G. Severe air pollution links to higher mortality in COVID-19 patients: The "double-hit" hypothesis. *J Infect* 2020. doi: 10.1016/j.jinf.2020.05.031
- Gabutti G, d'Anchera E, Sandri F, Savio M, Stefanati A. Coronavirus: Update Related to the Current Outbreak of COVID-19. *Infect Dis Ther* 2020. doi: 10.1007/s40121-020-00295-5
- Glencross DA, Ho TR, Camina N, Hawrylowicz CM, Pfeffer PE. Air pollution and its effects on the immune system. *Free Radic Biol Med* 2020. doi: 10.1016/j.freeradbiomed.2020.01.179
- Harrell FE. Regression modeling strategies: with applications to linear models, logistic and ordinal regression, and survival analysis. New York: Springer-Verlag, 2001.
- Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B, et al. Long-term air pollution exposure and cardio- respiratory mortality: a review. *Environ Health* 2013; 12: 43. doi: 10.1186/1476-069X-12-43
- Ialongo I, Virta H, Eskes H, Hovila J, Douros J. Comparison of TROPOMI/Sentinel-5 Precursor NO₂ observations with ground-based measurements in Helsinki. *Atmos Meas Tech* 2020; 13: 205-218. doi: 10.5194/amt-13-205-2020
- Ibironke O, Carranza C, Sarkar S, Torres M, Choi HT, Nwoko J, et al. Urban Air Pollution Particulates Suppress Human T-Cell Responses to Mycobacterium Tuberculosis. *Int J Environ Res Public Health* 2019; 16. doi: 10.3390/ijerph16214112

- ISTAT. [Coeweb - Statistiche del commercio estero]. Italian National Institute of Statistics, 2020a. <https://www.coeweb.istat.it> (Accessed: 12 June 2020)
- ISTAT. Data from the Italian National Institute of Statistics, 2020b. <https://www.istat.it/it/popolazione-e-famiglie> (Accessed: 12 June 2020)
- Lawrence MG. The Relationship between Relative Humidity and the Dewpoint Temperature in Moist Air: A Simple Conversion and Applications. *Bulletin of the American Meteorological Society* 2005; 86: 225-234. doi: 10.1175/BAMS-86-2-225
- Li H, Xu XL, Dai DW, Huang ZY, Ma Z, Guan YJ. Air Pollution and temperature are associated with increased COVID-19 incidence: a time series study. *Int J Infect Dis* 2020. doi: 10.1016/j.ijid.2020.05.076
- Lorente A, Boersma KF, Eskes HJ, Veefkind JP, van Geffen J, de Zeeuw MB, et al. Quantification of nitrogen oxides emissions from build-up of pollution over Paris with TROPOMI. *Sci Rep* 2019; 9: 20033. doi: 10.1038/s41598-019-56428-5
- Marchini T, Zirlik A, Wolf D. Pathogenic role of air pollution particulate matter in cardiometabolic disease: Evidence from mice and humans. *Antioxid Redox Signal* 2020. doi: 10.1089/ars.2020.8096
- Mazzola M, Lanconelli C, Lupi A, Busetto M, Vitale V, Tomasi C. Columnar aerosol optical properties in the Po Valley, Italy, from MFRSR data. *Journal of Geophysical Research: Atmospheres* 2010; 115. doi: 10.1029/2009JD013310
- McCreanor J, Cullinan P, Nieuwenhuijsen MJ, Stewart-Evans J, Malliarou E, Jarup L, et al. Respiratory effects of exposure to diesel traffic in persons with asthma. *N Engl J Med* 2007; 357: 2348-58. doi: 10.1056/NEJMoa071535
- Ministry of Health. Official Italian statistics on SARS-CoV-2, March 8, 2020, 2020. http://www.salute.gov.it/imgs/C_17_pagineAree_5351_0_file.pdf (Accessed: 12 June 2020)
- Mirowsky JE, Dailey LA, Devlin RB. Differential expression of pro-inflammatory and oxidative stress mediators induced by nitrogen dioxide and ozone in primary human bronchial epithelial cells. *Inhal Toxicol* 2016; 28: 374-82. doi: 10.1080/08958378.2016.1185199
- Newey WK, West KD. A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica* 1987; 55: 703-708. doi: 10.2307/1913610
- Ogen Y. Assessing nitrogen dioxide (NO₂) levels as a contributing factor to coronavirus (COVID-19) fatality. *Sci Total Environ* 2020; 726: 138605. doi: 10.1016/j.scitotenv.2020.138605
- Peng L, Zhao X, Tao Y, Mi S, Huang J, Zhang Q. The effects of air pollution and meteorological factors on measles cases in Lanzhou, China. *Environ Sci Pollut Res Int* 2020. doi: 10.1007/s11356-020-07903-4
- Persinger RL, Poynter ME, Ckless K, Janssen-Heininger YM. Molecular mechanisms of nitrogen dioxide induced epithelial injury in the lung. *Mol Cell Biochem* 2002; 234-235: 71-80. doi: 10.1023/A:1015973530559
- Polzer G. [La mappa della nostra era glaciale: così il coronavirus ha congelato l'Italia]. *La Repubblica*, Italy, 2020. <https://lab.gedidigital.it/repubblica/2020/cronaca/coronavirus-mappa-italia-impatto-sulla-mobilita/?ref=RHPPLF-BH-I252606083-C8-P3-S1.8-T1> (Accessed: 12 June 2020)
- RIAS. [Inquinamento atmosferico e COVID-19, Rete Italiana Ambiente e Salute]. *SCI RE*, 2020. <https://www.scienzainrete.it/articolo/inquinamento-atmosferico-e-covid-19/rete-italiana-ambiente-e-salute/2020-04-13> (Accessed: 12 June 2020)
- Rivas-Santiago CE, Sarkar S, Cantarella Pt, Osornio-Vargas A, Quintana-Belmares R, Meng Q, et al. Air pollution particulate matter alters antimycobacterial respiratory epithelium innate immunity. *Infect Immun* 2015; 83: 2507-17. doi: 10.1128/IAI.03018-14

- Setti L, Passarini F, De Gennaro G, Barbieri P, Perrone MG, Borelli M, et al. SARS-Cov-2RNA found on particulate matter of Bergamo in Northern Italy: First evidence. *Environ Res* 2020a; 188: 109754. doi: 10.1016/j.envres.2020.109754
- Setti L, Passarini F, De Gennaro G, Barbieri P, Perrone MG, Piazzalunga A, et al. The Potential role of Particulate Matter in the Spreading of COVID-19 in Northern Italy: First Evidence-based Research Hypotheses. *medRxiv* 2020b: 2020.04.11.20061713. doi: 10.1101/2020.04.11.20061713
- SIMA. Position Paper Particulate Matter and COVID-19 Italian Society of Environmental Medicine (SIMA), 2020. http://www.simaonlus.it/wpsima/wp-content/uploads/2020/03/COVID_19_position-paper_ENG.pdf (Accessed: 12 June 2020)
- Tsatsakis A, Petrakis D, Nikolouzakakis TK, Docea AO, Calina D, Vinceti M, et al. COVID-19, an opportunity to reevaluate the correlation between long-term effects of anthropogenic pollutants on viral epidemic/pandemic events and prevalence. *Food Chem Toxicol* 2020; 141: 111418. doi: 10.1016/j.fct.2020.111418
- Watts J, Kommenda N. Coronavirus pandemic leading to huge drop in air pollution. *The Guardian*, 2020. https://www.theguardian.com/environment/2020/mar/23/coronavirus-pandemic-leading-to-huge-drop-in-air-pollution?CMP=Share_iOSApp_Other (Accessed: 26 April 2020)
- Williams L, Ulrich CM, Larson T, Wener MH, Wood B, Chen-Levy Z, et al. Fine particulate matter (PM_{2.5}) air pollution and immune status among women in the Seattle area. *Arch Environ Occup Health* 2011; 66: 155-65. doi: 10.1080/19338244.2010.539636
- Wu X, Nethery RC, Sabath BM, Braun D, Dominici F. Exposure to air pollution and COVID-19 mortality in the United States. *medRxiv* 2020: 2020.04.05.20054502. doi: 10.1101/2020.04.05.20054502
- Zhao Y, Richardson B, Takle E, Chai L, Schmitt D, Xin H. Airborne transmission may have played a role in the spread of 2015 highly pathogenic avian influenza outbreaks in the United States. *Sci Rep* 2019; 9: 11755. doi: 10.1038/s41598-019-47788-z
- Zhu Y, Xie J, Huang F, Cao L. Association between short-term exposure to air pollution and COVID-19 infection: Evidence from China. *Sci Total Environ* 2020; 727: 138704. doi: 10.1016/j.scitotenv.2020.138704
- Zoran MA, Savastru RS, Savastru DM, Tautan MN. Assessing the relationship between surface levels of PM_{2.5} and PM₁₀ particulate matter impact on COVID-19 in Milan, Italy. *Sci Total Environ* 2020; 738: 139825. doi: 10.1016/j.scitotenv.2020.139825

Table 1. Number of total SARS-CoV-2 positive cases and prevalence rate on March 8, March 22, and April 5 and predicted NO₂ tropospheric levels ($\mu\text{mol}/\text{m}^2$) before the lockdown, in the subsequent periods after partial lockdown (February 24-March 8), after full lockdown (March 8-March 22), and after March 22.

	Population at Jan 1, 2019 ^a	Total cases			Prevalence rate (per 100,000)			NO ₂ levels ($\mu\text{mol}/\text{m}^2$)			
		Mar 8	Mar 22	Apr 5	Mar 8	Mar 22	Apr 5	Before Feb 24	After Feb 24		
									Feb 24-Mar 8	Mar 8-Mar 22	After Mar 22
Lombardy	10060574	4189	27206	50455	41.6	270.4	501.5	198	126	98	72
Bergamo (BG)	1114590	997	6296	9712	89.5	557.7	871.4	199	124	93	59
Brescia (BS)	1265954	501	5317	9340	39.6	420.0	737.8	169	120	91	61
Como (CO)	599204	27	512	1384	4.5	85.4	231	163	101	95	64
Cremona (CR)	358955	665	2895	4233	185.3	806.5	1179.3	177	107	84	60
Lecco (LC)	337380	53	872	1678	15.7	258.5	497.4	151	92	81	48
Lodi (LO)	230198	853	1772	2255	370.6	769.8	979.6	173	98	83	74
Mantua (MN)	412292	56	905	3046	13.6	219.5	348.5	159	105	74	53
Milan (MI)	3250315	406	5096	11230	12.5	156.9	345.5	245	157	111	98
Monza/Brianza (MB)	873935	59	1108	3046	6.8	126.8	348.5	255	153	138	80
Pavia (PV)	545888	243	1306	2619	44.5	239.2	479.8	147	98	72	50
Sondrio (SO)	181095	6	205	591	3.3	113.2	326.3	33	25	22	29
Varese (VA)	890768	32	386	1191	3.6	43.3	133.7	166	99	92	60
Veneto	4905854	670	5122	11226	13.7	104.4	228.8	136	94	75	50
Belluno (BL)	202950	23	226	538	11.3	111.4	265.1	50	56	31	29
Padua (PD)	937908	255	1277	2744	27.2	136.2	292.6	147	88	84	55
Rovigo (RO)	234937	5	76	186	2.1	32.3	79.2	118	68	56	42
Treviso (TV)	887806	126	935	1712	14.2	105.3	192.8	108	96	59	42
Venezia (VE)	853338	126	732	1425	14.8	85.8	167	126	94	66	46
Verona (VR)	926497	63	1046	2688	6.8	112.9	290.1	181	114	97	59
Vicenza (VI)	862418	50	631	1647	5.8	73.2	191	141	97	84	54
Emilia-Romagna	4459477	1180	7555	17089	26.5	168.4	383.2	109	81	66	38
Bologna (BO)	1014619	62	674	2521	6.1	66.4	248.5	109	86	71	37
Ferrara (FE)	345691	6	150	488	1.7	43.4	141.2	104	71	55	39
Forlì-Cesena (FC)	394627	15	329	977	3.8	83.4	247.6	80	61	51	29
Modena (MO)	705393	97	1010	2609	13.8	143.2	369.9	117	95	70	42
Parma (PR)	451631	276	1209	2275	61.1	267.7	503.7	121	87	72	42
Piacenza (PC)	287152	528	1765	2892	183.9	614.7	1007.1	140	113	90	58
Ravenna (RA)	389456	13	309	708	3.3	79.3	181.8	95	67	60	31
Reggio Emilia (RE)	531891	70	1167	3066	13.2	219.4	576.4	118	83	68	43
Rimini (RN)	339017	113	942	1553	33.3	277.9	458.1	90	53	40	23
Italy	60359546	7375	59138	128948	12.2	98.0	213.6				

^aMost recent data available from Italian National Institute of Statistic (ISTAT, 2020b).

Figure 1. Northern Italy study area showing levels of NO₂ tropospheric levels ($\mu\text{mol}/\text{m}^2$) before the first lockdown (A), and in the subsequent periods February 24-March 8 (B), March 8-March 22 (C), and after March 22 (D).

Figure 2. Restricted cubic spline regression analysis between NO₂ tropospheric levels ($\mu\text{mol}/\text{m}^2$) before the spread of the outbreak and SARS-CoV-2 positivity prevalence (cases per 100,000) in the three periods after the lockdown dates (A: February 24-March 8; B: March 8-March 22; C: March 22-April 5). Results presented SARS-CoV-2 infection prevalence rate (solid line) with 95% confidence interval (dash lines) in a multivariable model adjusted for population density, an index indicating age of the population, people mobility measured from telephone movements before the lockdown, temperature ($^{\circ}\text{C}$) and relative humidity in the three subsequent periods, and airport presence. Shaded circles are weighted on number of cases corresponding to the prevalence rates at each time point.

Declaration of competing interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof

Author statement

MV and TF designed the original study, and with KJR analyzed and interpreted the data, and wrote the manuscript. NO designed and carried out data analysis with TF. AG, FF and GM downloaded and processed environmental data. All authors read and approved the final manuscript.

Journal Pre-proof

Graphical abstract

Highlights

- Air pollution might increase susceptibility to SARS-CoV-2 infection
- We examined infection prevalence in the most affected regions in Northern Italy
- Satellite-detected tropospheric nitrogen dioxide was used to assess air pollution
- High NO₂ levels were associated with spread of the infection
- A causal role remains speculative, given ecologic biases and uncontrolled confounding





A**B****C****D**

Figure 1




Figure 2