

Article

The cold chain and the COVID-19 pandemic: an unusual increase in histamine content in fish samples collected in Southern Italy during lockdown

Francesco Giuseppe Galluzzo¹, Gaetano Cammilleri^{1,*}, Antonello Cicero¹, Licia Pantano¹, Andrea Pulvirenti², Andrea Macaluso¹, Nicola Cicero³, Vittorio Calabrese⁴ and Vincenzo Ferrantelli¹

¹Istituto Zooprofilattico Sperimentale della Sicilia, Palermo, Italy; ²Dipartimento di Scienze della Vita, Università degli studi di Modena e Reggio Emilia, Modena, Italy; ³Dipartimento SASTAS, Università degli studi di Messina, Messina, Italy and ⁴Dipartimento di Scienze biomediche e biotecnologiche, Università degli studi di Catania, Catania, Italy

*Correspondence to: Gaetano Cammilleri, Istituto Zooprofilattico Sperimentale della Sicilia, via Gino Marinuzzi 3, 90129, Palermo, Italy. E-mail: gaetano.cammilleri86@gmail.com

Received 21 July 2021; Revised 8 November 2021; Editorial decision 9 November 2021.

Abstract

Objectives: We analysed 900 samples of fresh (250) and processed (650) fish products collected in Sicily (Southern Italy) in 2020 during the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic (hereafter: COVID-19).

Materials and methods: The samples were divided temporally based on four phases relating to the various restrictions imposed by the Italian government in this period. The validated method of ultrahigh-performance liquid chromatography combined with a diode array detector was then employed for the analysis.

Results: The samples collected during the Phase I lockdown period and after it had ended (Phase II) revealed significant increases in the mean histamine levels: (41.89±87.58) mg/kg and (24.91±76.76) mg/kg, respectively. The 11 (1.3% of the total) fresh fish samples that were identified as being non-compliant with Regulation (EC) No.2073/2005 were only found during these two periods. All the processed samples were always compliant. The histamine values decreased as the restrictions eased, achieving a mean value of (11.16±9.3) mg/kg (Phase III).

Conclusions: There was an increase in the incidence of fish samples that were non-compliant with Regulation (EC) No.2073/2005 compared to previous surveillance data. These results provide a first report on the effect of lockdown measures on food safety and the cold chain. Our findings must cause food safety operators to intensify their controls over fresh fish products in such periods to safeguard consumer health. Further studies are required to evaluate whether the same trend would be observed with other food contaminants.

Keywords: Histamine; cold-chain; food safety; COVID-19.

Introduction

In December 2019, China reported a cluster of cases of pneumonia with an unknown cause that would later be designated as COVID-19 (coronavirus disease 2019). The disease is caused by the acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and has caused significant harm to the global population, including in the form of major socioeconomic damage (Chamola et al., 2020). Two months after the first case was reported, the World Health Organization (WHO) declared a public health emergency of international concern (Wu and McGoogan, 2020). The first case of COVID-19 reported in Italy was in January 2020; just one month later, more than 300 cases were recorded (Santacroce et al., 2020; WHO, 2020). A number of restrictions were imposed by the Italian government to contain the spread of the disease in the country. A national lockdown was initially put in place from 9 March to 3 May 2020 (Official Journal of The Republic of Italy, 2020a). In this period, known as Phase I, people could only move across regions for health or business purposes. All schools were closed and only grocery stores and shops selling other essential items remained open. During Phase II (4 May–14 July 2020) and Phase III (15 June–7 October 2020) there was a gradual loosening of the containment measures in correspondence with the downwards trend of the epidemic up to 5 November 2020.

After this date, the government introduced dynamic restrictions to contain the infection that were based on 21 parameters such as the number of cases and the capacity of intensive care units to admit patients.

The measures first adopted even included a suspension of the activities of the country's accommodation, catering and travel sectors. This, along with clear changes in the ways consumers ate, shopped and interacted with food, caused significant damage to the fishing industry (Cavallo et al., 2020). Generally, there was an increase in the consumption of comfort and unhealthy food, which was associated with the negative emotions experienced due to the pandemic (Ben Hassen et al., 2020). Furthermore, the rise in online food shopping led to problems obtaining delivery slots (Hobbs, 2020). In the south of Italy, local markets were closed to the general public during Phases I and II, leading to a surplus of unsold fish products. The changes put in place because of COVID-19 may have caused less attention to be paid to the health aspects of the food sold during the lockdown (Aday and Aday, 2020), probably due to the extended food storage dates introduced at this time.

Histamine is a biogenic amine (BA) that can be found in fish and fish products. It is produced due to the decarboxylation of histidine, a reaction catalysed by histidine decarboxylase, which is found in some bacterial species belonging to genera including *Morganella*, *Klebsiella*, *Photobacterium*, and *Vibrio* (Ababouch et al., 1991; Bjornsdottir et al., 2009; Hwang et al., 2020; Wang et al., 2020). The fish species with high histidine levels belong to the Scombridae (*Scomber scombrus*, *Thunnus thynnus*), Clupeidae (*Sardina pilchardus*, *Clupea harengus*) and *Engraulis* (*Engraulis encrasicolus*) genera (Colombo et al., 2018). The histamine levels in fish-based products can rise if they are subjected to poor storage conditions (Lehane and Olley, 2000). Consequently, it can be used as both a valuable marker of quality and an indicator of the freshness of such items (Bodmer et al., 1999; Sánchez-Pérez et al., 2018). Histamine plays numerous roles in the human body, but an increased level in the blood can produce several symptoms through its actions with histamine receptors (Schnedl et al., 2019). Scombroid poisoning is one of the most common forms of intoxication due to the histamine in fish products (Feng et al., 2016). The symptoms of this type of food poisoning are often associated with seafood allergies, and treatment is similarly linked (Hungerford,

2010). Histamine levels in fish products are regulated by European Commission (EC) Regulation No.2073/2005 (EC, 2005). The maximum levels permitted are 200 mg/kg and 400 mg/kg for fresh and processed fish products, respectively. Histamine cannot be degraded with standard cooking methods, meaning that prevention measures are crucial (Chung et al., 2017). A cold chain is an essential method for preventing histamine formation. High temperatures can increase its presence in food irreversibly (Hattori and Seifert, 2017), and the production of histidine decarboxylase cannot be averted at low temperatures (European Food Safety Authority, 2015). The goal of the current study was to evaluate whether the restrictions imposed during the COVID-19 pandemic had an impact on the presence of histamine in the fresh and processed fish products sold by supermarkets and street vendors in Southern Italy.

Materials and Method

Reagents and standards

Histamine dihydrochloride (99%), acetonitrile, potassium monophosphate, perchloric acid, sodium 1-decanesulfonate and potassium hydrogen phosphate trihydrate were purchased from Sigma-Aldrich (Amsterdam, The Netherlands). All chemicals and solvents were of analytical grade. Ultrapure water used for analysis was obtained from a Millipore purification system (Millipore, Burlington, MA, USA). Standards solutions of 5, 20, 40, 80, and 120 mg/L were made by diluting a 1000 mg/L histamine standard solution (Sigma-Aldrich, Amsterdam, The Netherlands) with deionized water.

Sampling plan and sample collection

In 2020, we collected 900 fish samples (250 fresh and 650 processed) at random from supermarkets and street vendors in Sicily (Southern Italy). All of the processed samples were produced in Italy from March to December 2020, as reported on the relevant labels. The samples were grouped for the statistical analysis according to the restrictions introduced by the Italian government due to COVID-19. Four different phases were then identified based on these same restrictions.

Phase I refers to the national lockdown that lasted from 9 March to 3 May and it was during this period that the severest restrictions were in place. Few types of economic activity were permitted (pharmacies, para-pharmacies, and grocery stores), public events were forbidden and restaurants were closed. Phase II was subsequently established by the Italian government and covered the date range from 4 May to 14 June. Takeaway food was available and public parks were open in this period. The downwards trend of the epidemiological curve led to Phase III, during which the government left only minor restrictions in place. This stage lasted from 15 June to 7 October and allowed public events for up to 200 attendees indoors and the reopening of cultural and social centers. Phase IV (8 October–December 2020) was characterized by the start of a second wave of cases. Consequently, dynamic restrictions were announced for 5 November onwards, the extent of which depended on the evolution of certain epidemiological parameters.

Eighty-six samples were collected during the lockdown (Phase I), 207 in Phase II, 399 in Phase III, and 208 in Phase IV. They belonged to six species: *Thunnus thynnus* (bluefin tuna; $n=401$); *Sardina pilchardus* (sardines; $n=367$); *Engraulis encrasicolus* (anchovies; $n=62$); *Scomber scombrus* (mackerel; $n=71$); *Coryphaena hippurus* (mahi-mahi; $n=9$); and *Thunnus alalunga* (albacore; $n=2$). The data are summarized in Table 1. The fresh fish samples were put in sterile bags and transported at $(+4\pm 1)$ °C in a refrigerated vehicle with a temperature control system. They were then stored at (-20 ± 1) °C

Table 1. List of fresh and processed samples collected divided by pandemic phases

Sample	Type	Phase I		Phase II		Phase III		Phase IV	
		Street vendor	Supermarket	Street vendor	Supermarket	Street vendor	Supermarket	Street vendor	Supermarket
Fresh									
Anchovies	Whole	—	—	—	—	4	6	4	5
Tuna	Fillets	16	20	28	26	25	25	14	13
Mackerel	Fillets	—	—	—	—	5	4	—	—
Mackerel	Whole	—	—	5	4	—	—	4	15
Sardines	Fillets	—	—	5	4	8	10	—	—
Fresh in total		16	20	38	34	42	45	22	33
Processed									
Albacore	Marinated	—	2	—	—	—	—	—	—
Anchovies	In oil	2	3	—	—	9	9	—	—
Anchovies	Salted	3	2	—	—	4	5	—	—
Anchovies	Marinated	4	2	—	—	—	—	—	—
Tuna	In oil	3	3	—	—	30	24	38	70
Tuna	Salted	—	8	3	6	—	—	—	27
Tuna	Mixed	—	—	—	—	—	4	—	—
Mackerel	In oil	—	2	—	9	—	9	—	9
Mackerel	Salted	—	3	—	—	—	—	—	—
Mackerel	Marinated	—	2	—	—	—	—	—	—
Mahi-mahi	In oil	—	—	6	3	—	—	—	—
Sardines	In oil	—	—	—	63	55	90	—	9
Sardines	Salted	1	—	23	13	33	30	—	—
Sardines	Marinated	—	4	—	9	—	—	—	—
mixed fishes	In oil	—	—	—	—	3	7	—	—
mixed fishes	surimi	—	6	—	—	—	—	—	—
Processed in total		13	37	32	103	134	178	38	115
Total		86		207		399		208	

Values indicate the number of samples collected in the specific period.

Table 2. Results of the validation process of the UHPLC-DAD method

Histamine level (mg/kg)	Observed concentration (mg/kg) ^a	RSD (%)	Recovery (%)
100	104.0±0.9	0.9	104
200	200±3	1.4	100
400	401±2	0.5	100

DAD, diode array detector; RSD, relative standard deviation; SD, standard deviation; UPLC, ultra-high performance liquid chromatography.

^amean±SD (n=10).

before extraction for the ultra-high performance liquid chromatography (UHPLC) analysis, which was conducted on the collection day. Salt and oil in the processed samples were removed before the analysis.

Extraction procedure

The extraction of the samples was carried out according to Cicero *et al.* (2020a). Briefly, 10 g of samples were homogenized and placed in a 50-mL tube with the addition of 10 mL of perchloric acid aqueous solution (6%). The sample was vortexed for 1 min. Next, 30 mL of deionized water were added to the sample and vortexed for 1 min. The solution was centrifuged at 5098g for 10 min, then at ambient temperature, the supernatant was transferred to a 50-mL flask and made up to volume with deionized water. A total of 1 mL was filtered on a 0.45-µm microfilter and put in vials for the UHPLC analysis. Each sample was analysed in duplicate.

UHPLC-DAD analysis

The analysis was conducted on an Agilent 1290 UHPLC with an ultraviolet/diode array detector (UV/DAD; Agilent Technologies, Santa Clara, CA, USA). A supelcosil LC-ABZ (15 cm×4.6 mm, diameter=5 mm) was used for the separation. Acetonitrile and an aqueous phosphate buffer solution at pH 6.9 were used as mobile phase (15:85, in volume). The chromatography conditions and instrumental parameters were set according to Cicero *et al.* (2020b). Briefly, the injection volume was 20 µL, the flow rate was 1.2 mL/min at room temperature and the detector wavelength was set at 210 nm for a runtime of only 6 min. The method involved an isocratic elution using a mobile phase A consisting of the phosphate buffer aqueous solution at pH 6.9 and a mobile phase B consisting of acetonitrile (85:15, in volume). The method was validated according to the ISO/IEC 17025:2017 (<https://www.iso.org/directives-and-policies.html>). The method showed a limit of quantification (LOQ) of 7.2 mg/kg and a limit of detection (LOD) of 2.2 mg/kg. The linearity of the method was calculated by linear regression of the areas obtained from the analysis, in triplicate, of histamine calibration standards solutions, accepting a determination coefficient (R^2)>0.999. The recovery and relative standard deviation (RSD) parameters were determined by spiking blank tuna samples at three concentration levels (100, 200, and 400 mg/kg), performing 10 replicates for each level. The results are summarized in Table 2.

Data collection and statistical analysis

All data were collected and elaborated with R version 4.0.2 (General Public License). Results under the LOQ of the method were considered as the LOQ value (Helsel, 2005) for the statistical analysis.

The packages used for R analysis and data visualization were: Rcmdr (Rcmdr), ggbiplot2 and pplotly (Kabacoff, 2011; Sievert, 2020). Data were not normally distributed (Shapiro–Wilk Test P -value<0.05); therefore, a Kruskal–Wallis test was carried out to verify differences between sampling periods for each sample type. Post-hoc analysis was performed with Dunn’s test (Dunn, 1964) and adjusted with the Benjamini–Hochberg method (Benjamini and Hochberg, 1995) to determine differences between groups. Only fresh tuna samples revealed the presence of histamine and statistical analysis among different phases was carried out taking these samples only into account.

Results and Discussion

Histamine content and general consideration

The results of the analysis are set out in Figure 1 and Table 3. Histamine was detected in 47 fresh tuna samples (5.00%) at levels between 15.07 mg/kg and 596.69 mg/kg. About 1.22% of the samples were over the limits imposed by Regulation (EC) No.2073/2005 (200 mg/kg for fresh fish products), which is comparable to the multiannual studies conducted in Italy (Cicero et al., 2020b; Lo Magro et al., 2020). The non-compliant histamine levels were only found in the fresh tuna samples obtained from street vendors. This confirms that there is a high incidence of this BA forming in this type of product due to the high levels of free histidine in their tissues. High amounts of histamine are due to time and/or temperature abuses during handling and storage (Lo Magro et al., 2020). The street vendors in our study were displaying the sampled fresh tuna on ice, but the FDA (U.S. Food and Drug Administration)-recommended visual checks of the condition of ice around a product were not being made, suggesting an absence of time/temperature controls (FDA, 2021). Furthermore, as histamine formation can be affected by evisceration, the high concentrations of this BA in our samples suggest that the fresh tuna was stored for a prolonged period without this occurring. In a recent study on eviscerated

yellowfin tuna stored at 30 °C for 12 hours, the maximum histamine level achieved was 2400 mg/kg (Benner et al., 2009). Unfortunately, the tuna samples we collected had all been filleted, and so it was not possible to determine the time that had elapsed between evisceration and the fish being offered for sale on the market. In contrast to the fresh tuna, no histamine was found in the samples of processed tuna and other processed fish products, confirming that the Italian fish-processing industries had continued to comply with mandated hazard analyses and critical control-point procedures, even during the lockdown.

Histamine content and pandemic period

Phase I (9 March to 3 May 2020)

The samples from Phase I were the least compliant with Regulation (EC) No.2073/2005, the incidence of non-compliant samples being 4.65%. This figure increased to 11.11% if we only considered the fresh samples. Four fresh tuna fillets contained histamine levels above the EC limit, with a concentration range from 250.02 to 596.69 mg/kg. All the non-compliant samples were collected from street vendors between 17 March and 31 March 2020. The statistical analysis for the fresh tuna revealed significant differences between the first and the other phases examined in the study (Kruskal–Wallis chi-squared=73.081; P <0.05). There may be a variety of reasons for these results, including in relation to the containment measures imposed during Phase I and the radical change in consumer attitudes that occurred in this period (Aday and Aday, 2020). This phase saw panic buying (Hossain, 2020) and a preference for frozen meals (Galanakis et al., 2021). The convenience during lockdown of processed food with extended use-by dates meant that it was favoured over highly perishable items like fresh fish (Cavallo et al., 2020). In addition, the lack of slots available for the delivery of online food orders (Hobbs, 2020) and the consumer’s reduced purchasing power also increased the demand for processed food (Khan and Moverley Smith, 2020). As an example, the period 17 February–15 March,




Figure 1. Scatterplot of the samples analysed sorted by period and phase of sampling ($n=900$).

Table 3. Histamine contents found in the fish samples analysed (expressed as mg/kg)

Sample	Phase I	Phase II	Phase III	Phase IV
Fresh				
Analysed	36	72	87	55
>LOD ^a	27	9	8	—
Non-compliant	4 (11.11%)	7 (9.72%)	—	—
Mean±SD	85.91±123.30	52.51±26.15	14.63±19.62	—
Processed				
Analysed	50	135	312	135
>LOD ^a	—	—	—	—
Non-compliant	—	—	—	—
Mean±SD	—	—	—	—
Total				
Analysed	86	207	399	208
>LOD ^a	27	9	8	—
Non-compliant	4 (4.65%)	7 (3.38%)	—	—
Mean±SD	41.89±87.58	24.91±76.76	11.16±9.30	—

The observation '>LOD' (limit of detection) refers to the number of samples with detectable histamine levels. The term 'non-compliant' indicates samples that reached histamine levels above the limits imposed by the EU Regulation (200 mg/kg for fresh fish, 400 mg/kg for processed fish). The percentage of non-compliant on the total of the sample analysed is given in parentheses. The term 'analysed' indicates the total number of samples subjected to UHPLC-DAD analysis.

DAD, diode array detector; EU, European Union; SD, standard deviation; UPLC, ultra-high performance liquid chromatography.

^aLOD is 2.2 mg/kg.

2020 saw the sales of canned tuna increase by +36% compared to the same dates in 2019. Furthermore, Phase I had included a ban on fishing and aquaculture activities until 26 March 2020 to protect the health of local fisherman, a consequence of which was a reduced supply (Official Journal of The Republic of Italy, 2020b).

Along with the closure of restaurants, the factors referred to above led to a decline in the demand for fresh fish (D'Oronzio *et al.*, 2020). Consequently, there was initial overproduction during the first lockdown: fresh fish went unsold and fish market operators and street vendors, probably due to liquidity issues, may have decided to keep their fresh products for longer in uncontrolled storage conditions. The result may have been less compliance with permitted histamine levels, as reflected in our samples (Mattioli *et al.*, 2020; Swinnen and McDermott, 2020). Furthermore, flawed mitigation measures agreed by those involved in fishing caused an increase in the amount of unsold fish products (D'Oronzio *et al.*, 2020). Activities began to recover slightly in April 2020, albeit with an ongoing significant reduction in the fishing effort (in terms of days and hours).

These conditions could explain the absence of non-compliant samples during April despite the first lockdown period. The incidence of non-compliant samples found during the first lockdown period was higher than over 5 years of studies conducted in Southern Italy (Muscarella *et al.*, 2013; Piersanti *et al.*, 2014; Cicero *et al.*, 2020b), suggesting an alarming situation during this period. The food chain is susceptible to alteration in food demand and offer (Aday and Aday, 2020) and food fraud could occur more easily in lockdown due to the fact that there were less food inspections and less governance (Brooks *et al.*, 2021).

No histamine was detected in the processed samples, confirming that the fish processing industry in Italy continued to comply with regulations and operate good practices.

Phase II (4 May–14 June 2020)

Samples from seven tuna fillets were non-compliant with EU legislation during Phase II (4 May–14 July 2020), with a mean histamine content of (407.63±139.78) mg/kg and a range between 215.76 mg/kg and 565.21 mg/kg. The percentage of non-compliant samples in Phase II was 9.72% if only the fresh samples were considered. All of

these samples were collected from street vendors. Phase II was characterized by less restrictive regulations than Phase I (17 May 2020).

This slight decrease in the number of non-compliant samples can be linked to both the fact that there was only a gradual loosening of the COVID-19 restrictions and the simultaneous reorganization of the surveillance procedures on food safety imposed after the national lockdown. Several food fraud cases were brought against street vendors in Sicily. In the majority, fresh fish was displayed on ice, which provides only partial coverage of an item and exposes the rest of it to ambient temperatures. Consequently, the high incidence of non-compliant samples in our study suggests an absence of time/temperature controls of the fish products examined, leading to microbial growth and histamine production.

In this period, restaurants could sell food to take away and via home deliveries. They were able to also serve food at the table from 1 June 2020, but with limitations in place relating to social distancing. In concert with this, there was a gradual increase in the consumption of fish products.

Phases III–IV (15 June–December 2020)

No non-compliant samples were found from the period 15 June to December 2020, despite an increase in the number we collected in Phase III ($n=399$). June 2020 onwards saw an increase in the domestic consumption of fresh fish (D'Oronzio *et al.*, 2020; EUMOFA, 2020a). The Hotelier–Restaurant–Café (HORECA) channels were also open in this period, leading to a rise in demand. In Sicily, local restaurants predominantly buy their fresh fish from local and street vendors, who have an important sales channel as a result. However, the increased demand in this phase was not matched with an adequate supply. This was related to the size of Sicily's fishing vessels, 80% of which are less than 10 m in length, meaning that they could not be used because of the government's social distancing requirements (D'Oronzio *et al.*, 2020).

Furthermore, the fear of a production surplus, as occurred in Phase I, caused the sector to reduce its fishing activities, giving rise to conditions that were the complete opposite of those seen in the first lockdown period (EUMOFA, 2020b). Phases III, IV, and V were therefore characterized by a significant reduction in the availability

of fresh fish. Most of the fish caught were sold, and so the sector did not need to adopt inadequate storage practices that could compromise the health of consumers. The restrictions imposed by the government and the closure of the HORECA channels thus seem to have been the key factors undermining the safety of the fresh fish products sold.

Conclusions

This study examines the impact of the restrictions imposed by the Italian government during the COVID-19 pandemic on the safety of fresh fish products. The results reveal a relationship between the extent of these measures and the incidence of samples that did not comply with regulations regarding histamine levels. Furthermore, our findings show that large-scale distributors of processed fish products maintained high hygiene standards compared to street vendors of fresh fish. The latter's non-compliance with safety standards during the lockdown is alarming and should lead to food safety inspectors intensifying their efforts to ensure that controls on fresh fish products are in place in similar circumstances in the future, using targeted strategies capable of safeguarding health. The limitations of this research are the small data pool and the imbalance of the sample types analysed for each period. Additional studies are therefore required.

Author Contributions

Francesco Giuseppe Galluzzo: Conceptualization, Methodology and data curation; Gaetano Cammilleri: Investigation and data curation; Antonello Cicero: Formal Analysis and Methodology; Licia Pantano: Formal Analysis and data curation; Andrea Pulvirenti: Writing, Review and Editing; Andrea Macaluso: Investigation and Supervision; Antonio Vella: Visualization and Supervision; Vittorio Calabrese: Validation and Supervision; Nicola Cicero: Review and editing; Vincenzo Ferrantelli: Project Administration, Writing, Review and Editing.

Funding

This research received no external funding.

Conflict of Interest

The authors declare no conflict of interest.

References

- Ababouch, L., Afilal, M. E., Rhafiri, S., et al. (1991). Identification of histamine-producing bacteria isolated from sardine (*Sardina pilchardus*) stored in ice and at ambient temperature (25 °C). *Food Microbiology*, 8(2): 127–136.
- Aday, S., Aday, M. S. (2020). Impact of COVID-19 on the food supply chain. *Food Quality and Safety*, 4(4): 167–180.
- Ben Hassen, T., El Bilali, H., Allahyari, M. S. (2020). Impact of COVID-19 on food behavior and consumption in Qatar. *Sustainability*, 12(17): 6973.
- Benjamini, Y., Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B*, 57(1): 289–300.
- Benner, Jr. R. A., Staruszkiewicz, W. F., Conrad, S. M., et al. (2009). Biogenic amine production in yellowfin tuna (*Thunnus albacares*) under controlled decomposition conditions. 2009 Annual Meeting of the International Association for Food Protection, 12–15 July 2009, Grapevine, TX, USA, Poster presentation 1–28.
- Bjornsdottir, K., Bolton, G. E., McClellan-Green, P. D., et al. (2009). Detection of Gram-negative histamine-producing bacteria in fish: a comparative study. *Journal of Food Protection*, 72(9): 1987–1991.
- Bodmer, S., Imark, C., Kneubühl, M. (1999). Biogenic amines in foods: histamine and food processing. *Inflammation Research*, 48(6): 296–300.
- Brooks, C., Parr, L., Smith, J. M., et al. (2021). A review of food fraud and food authenticity across the food supply chain, with an examination of the impact of the COVID-19 pandemic and Brexit on food industry. *Food Control*, 130: 108171.
- Cavallo, C., Sacchi, G., Carfora, V. (2020). Resilience effects in food consumption behaviour at the time of Covid-19: perspectives from Italy. *Heliyon*, 6(12): e05676.
- Chamola, V., Hassija, V., Gupta, V., et al. (2020). A comprehensive review of the COVID-19 pandemic and the role of IoT, Drones, AI, blockchain, and 5G in managing its impact. *IEEE Access*, 8: 90225–90265.
- Chung, B. Y., Park, S. Y., Byun, Y. S., et al. (2017). Effect of different cooking methods on histamine levels in selected foods. *Annals of Dermatology*, 29(6): 706–714.
- Cicero, A., Cammilleri, G., Galluzzo, F. G., et al. (2020a). Histamine in fish products randomly collected in Southern Italy: a 6-year study. *Journal of Food Protection*, 83(2): 241–248.
- Cicero, A., Galluzzo, F. G., Cammilleri, G., et al. (2020b). Development of a rapid and eco-friendly UHPLC analytical method for the detection of histamine in fish products. *International Journal of Environmental Research and Public Health*, 17(20): 7453.
- Colombo, F. M., Cattaneo, P., Confalonieri, E., et al. (2018). Histamine food poisonings: a systematic review and meta-analysis. *Critical Reviews in Food Science and Nutrition*, 58(7): 1131–1151.
- D'Oronzio, M. A., Di Paolo, I., Diglio, G., et al. (2020). The Covid-19 Emergency and the Italian Fish Sector: Impact and Answers [Online]. https://www.crea.gov.it/documents/68457/0/L%E2%80%99Emergenza+COVID-19+e+il+settore+ittico+italiano_Impatti_e_risposte_Gen_2021.pdf/8da26719-4390-808d-f403-fd7f1d710942?t=1611327296230. Accessed on 8 April 2021. (in Italian)
- Dunn, O. J. (1964). Multiple comparisons using rank sums. *Technometrics*, 6(3): 241–252.
- EC (European Commission). (2005). Commission Regulation (EC) No. 2073/2005 of 15 November 2005 on Microbiological Criteria for Foodstuffs [Online]. <http://data.europa.eu/eli/reg/2005/2073/oj/eng>. Accessed on 27 August 2020.
- EUMOFA (European Market Observatory for Fisheries and Aquaculture Products). (2020a). Studies and reports [Online]. <https://www.eumofa.eu/it/market-analysis>. Accessed on 29 April 2021. (in Italian)
- EUMOFA (European Market Observatory for Fisheries and Aquaculture Products). (2020b). Monthly Highlights, No.9. 2020 Sep 7 [Online]. <https://www.eumofa.eu/it/market-analysis>. Accessed on 19 April 2021. (in Italian)
- European Food Safety Authority. (2015). Scientific and technical assistance on the evaluation of the temperature to be applied to pre-packed fishery products at retail level. *EFSA Journal*, 13(7): 4162.
- Feng, C., Teuber, S., Gershwin, M. E. (2016). Histamine (Scombroid) fish poisoning: a comprehensive review. *Clinical Reviews in Allergy & Immunology*, 50(1): 64–69.
- FDA (U.S. Food and Drug Administration). (2021). Fish and Fishery Products Hazards and Controls Guidance, Fourth Edition – June 2021. FDA, Silver Spring, MD, USA.
- Galanakis, C. M., Rizou, M., Aldawoud, T. M. S., et al. (2021). Innovations and technology disruptions in the food sector within the COVID-19 pandemic and post-lockdown era. *Trends in Food Science & Technology*, 110: 193–200.
- Hossain. (2020). In Pictures: panic-buying around the world amid fears over coronavirus outbreak [Online]. The Straits Times. <https://www.straitstimes.com/multimedia/photos/in-pictures-panic-buying-around-the-world-amid-fears-over-coronavirus-outbreak>. Accessed on 20 July 2021.
- Hattori, Y., Seifert, R., (Eds). (2017). Histamine and Histamine Receptors in Health and Disease. 1st Edition. Springer, Cham, Switzerland, pp. 222–224.
- Helsel, D. R. (2005). Nondetects and Data Analysis Statistics for Censored Environmental Data [Online]. <https://www.cabdirect.org/cabdirect/abstract/20053102639>. Accessed on 12 July 2021.

- Hobbs, J. E. (2020). Food supply chains during the COVID-19 pandemic. *Canadian Journal of Agricultural Economics*, 68(2): 171–176.
- Hungerford, J. M. (2010). Scombroid poisoning: a review. *Toxicon*, 56(2): 231–243.
- Hwang, C. C., Kung, H. F., Lee, Y. C., *et al.* (2020). Histamine fish poisoning and histamine production by *Raoultella ornithinolytica* in Milkfish Surimi. *Journal of Food Protection*, 83(5): 874–880.
- Kabacoff, R. (2011). *R in Action: Data Analysis and Graphics with R*. Manning, Shelter Island, NY, USA, pp. 227–426.
- Khan, M. A., Moverley Smith, J. E. (2020). “Covibesity,” a new pandemic. *Obesity Medicine*, 19: 100282.
- Lehane, L., Olley, J. (2000). Histamine fish poisoning revisited. *International Journal of Food Microbiology*, 58(1–2): 1–37.
- Lo Magro, S., Summa, S., Iammarino, M., *et al.* (2020). A 5-years (2015–2019) control activity of an EU Laboratory: contamination of histamine in fish products and exposure assessment. *Applied Sciences*, 10(23): 8693.
- Mattioli, A. V., Sciomer, S., Cocchi, C., *et al.* (2020). Quarantine during COVID-19 outbreak: changes in diet and physical activity increase the risk of cardiovascular disease. *Nutrition, Metabolism, and Cardiovascular Diseases*, 30(9): 1409–1417.
- Muscarella, M., Lo Magro, S., Campaniello, M., *et al.* (2013). Survey of histamine levels in fresh fish and fish products collected in Puglia (Italy) by ELISA and HPLC with fluorimetric detection. *Food Control*, 31: 211–217.
- Official Journal of The Republic of Italy. (2020a). Decree of the President of the Council of Ministers April 26, 2020 [Online]. <https://www.gazzettaufficiale.it/eli/id/2020/04/27/20A02352/sg>. Accessed on 16 December 2020. (in Italian)
- Official Journal of The Republic of Italy. (2020b). Decree of the President of the Council of Ministers 11 June 2020 [Online]. <https://www.gazzettaufficiale.it/eli/id/2020/06/11/20A03194/sg>. Accessed on 16 December 2020. (in Italian)
- Piersanti, A., Tavoloni, T., Lestingi, C., *et al.* (2014). High-throughput histamine analysis approach in an official control laboratory: analytical methods and four years fish products results. *Food Chemistry*, 153: 437–443.
- Sánchez-Pérez, S., Comas-Basté, O., Rabell-González, J., *et al.* (2018). Biogenic amines in plant-origin foods: are they frequently underestimated in low-histamine diets? *Foods*, 7(12): 205.
- Santacroce, L., Charitos, I. A., Del Prete, R. (2020). COVID-19 in Italy: an overview from the first case to date. *Electronic Journal of General Medicine*, 17(6): em235.
- Schnedl, W. J., Lackner, S., Enko, D., *et al.* (2019). Evaluation of symptoms and symptom combinations in histamine intolerance. *Intestinal Research*, 17(3): 427–433.
- Sievert, C. (2020). Interactive Web-Based Data Visualization with R, Plotly, and Shiny. CRC Press, Boca Raton, FL, USA.
- Swinnen, J., McDermott, J. (2020). COVID-19 and Global Food Security [Online]. International Food Policy Research Institute, Washington, DC, USA. <https://ebrary.ifpri.org/digital/collection/p15738coll2/id/133762>. Accessed on 8 January 2021.
- Wang, D., Yamaki, S., Kawai, Y., *et al.* (2020). Histamine production behaviors of a psychrotolerant histamine-producer, *Morganella psychrotolerans*, in various environmental conditions. *Current Microbiology*, 77(3): 460–467.
- WHO (World Health Organization). (2020). Coronavirus disease 2019 (COVID-19) Situation Report - 37 [Online]. <https://www.who.int/publications/m/item/situation-report---37>. Accessed on 1 April 2021.
- Wu, Z. Y., McGoogan, J. M. (2020). Characteristics of and important lessons from the Coronavirus Disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*, 323(13): 1239–1242.