

This is the peer reviewed version of the following article:

Natural complementary feeds in gestating and lactating sows: effects on performance and metabolic profile / A., Beltrami; Ielo, Maria Cristina; Volpelli, Luisa Antonella; Minelli, Giovanna; LO FIEGO, Domenico Pietro. - In: ITALIAN JOURNAL OF ANIMAL SCIENCE. - ISSN 1828-051X. - ELETTRONICO. - 12:(2013), pp. 30-30. (Intervento presentato al convegno ASPA 20th Congress tenutosi a Bologna nel 11-13 June 2013).

Page Press

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

05/05/2024 20:19

(Article begins on next page)



Abstract N.0028

Dear Professor **Luisa Antonella Volpelli**,

here are the details of your abstract as they were saved on Mar 14, 2013.

Nutrition and feeding

NATURAL COMPLEMENTARY FEEDS IN GESTATING AND LACTATING SOWS: EFFECTS ON PERFORMANCE AND METABOLIC PROFILE

Andrea Beltrami*#, Maria Cristina Ielo§, Luisa Antonella Volpelli*§, Giovanna Minelli*§, Domenico Pietro Lo Fiego*§

**Dipartimento di Scienze della Vita, Università di Modena e Reggio Emilia, Italy; #METHODO CHEMICALS, Novellara (RE), Italy; §BIOGEST-SITEIA, Università di Modena e Reggio Emilia, Italy*
Corresponding author: volpelli@unimore.it

Pregnancy and lactation cause high metabolic effort and high cellular metabolism in sows; the contemporary presence of stressors may lead to ROS (Reactive Oxygen Species) production, which may overcome the action of the physiological anti-oxidants. The consequences may be DNA, protein and PUFA oxidation, peroxides production, ROS release in blood, and finally oxidative stress. Liver function, sow immune defense and fertility, immunity transfer to piglets and their vitality may be unbalanced.

In this work, two natural complementary feeds (GOLD and GOLD EPS), aiming to reduce oxidative stress, were used at 0.3% in the diet of lactating (trial L, 83 sows in total, 3 groups: control, GOLD, GOLD EPS) and gestating-lactating (trial GL, 105 sows, 2 groups; control, GOLD) sows. GOLD contains: brewer's yeast and organic selenium from *Saccharomyces cerevisiae*; betaine, phosphatidylcholine, RRR- α -tocopherol. GOLD EPS also contains *Silybum marianum* extract. Collected data were: sows' feed intake and post-weaning performance; number of newborn and weaned piglets, mortality, individual growing; liver metabolic profile, oxidative stress indicators; blood, colostrum and milk IgG.

In trial L, higher weights were recorded for GOLD and GOLD EPS piglets (7352 and 7329 g at weaning) in comparison with control (7077 g); a trend to post-weaning fertility improvement for GOLD EPS sows were also observed (new successful farrowing: 62, 68 and 77 % for control, GOLD and GOLD EPS).

In trial GL, GOLD piglets were slightly heavier at birth and less affected by diarrhea. A positive increase of serum anti-oxidant power (OXY) was observed at half pregnancy in GOLD sows, followed by a decrease at farrowing. GOLD sows also showed a constant decrease of bilirubin. Serum IgG, after an increase at farrowing (control 14.4; GOLD 16.4 g/l), showed a decrease in GOLD sows at the end of lactation, when IgG level increased in piglets (control 4.6; GOLD 7.8 g/l): this might indicate a



higher transfer from mother to piglets.

Notwithstanding the difficulty of on-field conditions, these trials provide partially good results about the performance of sows fed the tested products.

Acknowledgments

Thanks are due to METHODO CHEMICALS for the support to the research.

Presentation: Oral *Publishable:* YES