

This is a pre print version of the following article:

A previously unreported function of beta1B integrin isoform in caspase-8-dependent integrin-mediated keratinocyte death / Lotti, Roberta; Marconi, Alessandra; Truzzi, Francesca; Dallaglio, Katiuscia; Gemelli, Claudia; R. G., Borroni; Palazzo, Elisabetta; Pincelli, Carlo. - In: JOURNAL OF INVESTIGATIVE DERMATOLOGY. - ISSN 0022-202X. - STAMPA. - 130:(2010), pp. 2569-2577. [10.1038/jid.2010.195]

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

25/05/2024 12:55

(Article begins on next page)

## **A novel function of $\beta_1B$ integrin isoform in caspase-8-dependent integrin-mediated keratinocyte death**

Roberta Lotti<sup>1,3</sup>, Alessandra Marconi<sup>1,3</sup>, Francesca Truzzi<sup>1</sup>, Katuscia Dallaglio<sup>1</sup>, Claudia Gemelli<sup>2</sup>, Riccardo Giovanni Borroni<sup>1</sup>, Elisabetta Palazzo<sup>1</sup> and Carlo Pincelli<sup>1</sup>.

<sup>1</sup>Institute of Dermatology, and <sup>2</sup>Department of Biomedical Sciences, School of Biosciences and Biotechnologies, University of Modena and Reggio Emilia, 41100 Modena, Italy.

<sup>3</sup>These two authors contributed equally to this work.

### **Corresponding Author:**

Carlo Pincelli, MD

Via del Pozzo, 71

4100 Modena, Italy

Tel +39 059 422 2931

Fax +39 059 422 4271

[carlo.pincelli@unimore.it](mailto:carlo.pincelli@unimore.it)

### **Short title:**

Pro-apoptotic role of  $\beta_1B$  integrin

### **Abbreviation list:**

shFADD: short-hairpin-RNA-Fas-Associated-Death-Domain; IMD: integrin-mediated death; DISC: death-inducing signaling complex; SCC: Squamous-cell carcinoma.

## Abstract

Integrins regulate adhesive cell-matrix interactions and mediate survival signals. On the other hand, unligated or free cytoplasmic fragments of integrins induce apoptosis in many cell types (integrin-mediated death). We have previously shown that  $\beta_1$  integrins expression protects keratinocyte stem cells from anoikis, while the role of the  $\beta_1B$  integrin isoform has never been clarified. Here we report that suspended keratinocytes undergo apoptosis via the activation of caspase-8, independently of Fas/Fas Ligand system. Indeed, anti- $\beta_1$  integrin neutralizing antibodies induced apoptosis in short-hairpin-RNA-Fas-Associated-Death-Domain treated cells. Moreover, before and during suspension, caspase-8 directly associated with  $\beta_1$  integrin, that in turn internalized and progressively degraded, shedding the cytoplasmic domain.  $\beta_1B$  was expressed only in the cytoplasm in a perinuclear fashion and remained unaltered during suspension. At 24 hrs, as  $\beta_1A$  located close to the nucleus,  $\beta_1B$  co-localized with  $\beta_1A$  and co-immunoprecipitated with caspase-8. Caspase-8 was activated earlier in  $\beta_1B$  integrin transfected keratinocytes, and these cells underwent a higher rate of apoptosis than mock cells. By contrast, caspase-8 was not activated in siRNA  $\beta_1B$  transfected cells. These results indicate that when  $\beta_1A$  is unligated,  $\beta_1B$  is responsible for “integrin-mediated death” in human keratinocytes.

## Introduction

$\beta_1$  integrin is abundantly expressed in keratinocytes and is critical for keratinocyte adhesion to type IV collagen (Fleischmajer et al, 1998; Tiberio et al, 2002). A  $\beta_1B$  variant has also been isolated, in which a unique 12-amino-acid sequence replaces the last 21 COOH-terminal amino acids of wild-type  $\beta_1$  integrin (Altruda et al, 1990). Although  $\beta_1B$  integrin is expressed not only in keratinocytes, but also in liver and in skeletal and cardiac muscles (Balzac et al, 1993), its functional properties remain largely unknown.  $\beta_1$  integrins mediate binding to the extracellular matrix (ECM) and transduce an intracellular signal promoting cell survival in many tissues and cells, including keratinocytes (Howlett et al, 1995; Marconi et al, 2004). Cells denied adhesion because of integrin ligand deprivation undergo a form of cell death named anoikis (Frisch and

Francis, 1994). While anoikis occurs as the consequence of loss of integrin-mediated attachment, a form of integrin-mediated death (IMD) has been proposed, whereby integrins in the antagonized or unligated state, can promote apoptosis independently of cell detachment (Stupack et al, 2001; Brassard et al, 1999). In most instances, anoikis process involves the extrinsic apoptotic pathway that is typically triggered by death receptor ligands, which in turn recruit the adaptor protein FADD. The death receptor-bound FADD associates with caspase-8 in the death-inducing signaling complex (DISC) leading to caspase autoactivation (Muzio et al, 1998; Marconi et al, 2004) (Figure S1). In IMD, integrins seem to cluster with caspase-8 without the involvement of the DISC (Stupack et al, 2001), although the actual mechanisms underlying this process remain to be clarified. In this study, we demonstrate that  $\beta_1$ B integrin clusters with unligated wild-type  $\beta_1$  integrin and (pro-)caspase-8 to induce IMD in human keratinocytes, providing evidence for a previously unidentified role of this integrin isoform.

## Results

**Death receptors are not involved in anoikis.** To test the role of  $\beta_1$  integrin and caspase-8 during anoikis, cells were suspended and treated with a neutralizing anti- $\beta_1$  integrin antibody. We show that suspension induces keratinocyte apoptosis that is significantly augmented by blocking  $\beta_1$  integrin, indicating that this integrin plays a role in keratinocyte anoikis (Figure S1a) (Marconi et al, 2004). We also show that caspase-8 is activated early in keratinocyte anoikis (Figure S1b), suggesting the involvement of death receptors in this process. To evaluate whether death receptors are actually responsible for caspase-8 activation and anoikis in our system, we evaluated the modulation of Fas/FasL in suspended keratinocytes and after treatment with anti- $\beta_1$  integrin. During a 0 to 18 hr time course, no up-regulation of either Fas (Figure 1a) or FasL (Figure 1b) was detected in both suspended and anti- $\beta_1$  integrin-treated keratinocytes. In addition, no FasL was released by suspended keratinocytes up to 3 hrs (Figure 1c). To definitely rule out the involvement of death receptors in keratinocyte anoikis, we retrovirally infected HaCaT keratinocytes with short-hairpin-RNA-Fas-Associated-Death-Domain (shFADD). Western blot

analysis showed reduced levels of FADD in shRNA FADD-infected keratinocytes, as compared to mock cells (Figure 2a). We first tested the capability of shFADD to prevent apoptosis. As expected, TRAIL induced apoptosis in mock but not in shFADD keratinocytes (Figure 2b). By contrast, caspase-8 was activated in shFADD as well as in mock keratinocytes treated with anti- $\beta_1$  integrin, as shown by the appearance of the active fragment (23 kD) (Figure 2c). Furthermore, both shFADD and mock keratinocytes showed the same percent of sub G1 peak positive cells in presence or in absence of anti- $\beta_1$  integrin. At 18 hrs, while keratinocytes reached high degree of anoikis and caspase-8 was fully activated, blocking FADD did not prevent cell death in keratinocytes (Figure 2d).

**Direct interaction of  $\beta_1$  integrin and caspase-8 in adherent and suspended cells.** If keratinocytes undergo anoikis without the involvement of FADD, we reasoned that there must be a direct interaction between  $\beta_1$  integrin and caspase-8. To determine whether caspase-8 and  $\beta_1$  integrin interact in keratinocytes, adherent cells were double stained with anti- $\beta_1$  integrin and anti-caspase-8 antibodies. Subconfluent keratinocytes expressed  $\beta_1$  integrin both at the membrane and in the cytoplasm, while caspase-8 displayed a diffuse cytosolic localization. Double staining revealed that  $\beta_1$  integrin and caspase-8 co-localize in the cytoplasm (Figure 3a). Consequently, we investigated the interaction between caspase-8 and  $\beta_1$  integrin in keratinocyte anoikis. Western blotting on lysates from suspended keratinocytes showed the appearance of caspase-8 active fragment (18 kDa) starting at 1 hr (Figure 3b). The same lysates were immunoprecipitated with anti- $\beta_1$  integrin antibody. Procaspase-8 and  $\beta_1$  integrin co-immunoprecipitated at 0 hrs, while procaspase-8 progressively decreased, indicating activation, up to 24 hrs. Furthermore, suspension affected  $\beta_1$  integrin conformation starting at 1 hr, and precipitated protein decreased at 6 hrs, eventually disappearing at 24 hrs. FADD did not immunoprecipitate with  $\beta_1$  integrin-procaspase-8 complex, further confirming that death receptors are not involved in keratinocyte anoikis (Figure 3c). Taken together, these results clearly indicate that  $\beta_1$  integrin and caspase-8 directly interact, and that  $\beta_1$  integrin is altered during anoikis.

**$\beta_1$  integrin is internalized and degraded during suspension.** As  $\beta_1$  integrin is progressively altered during keratinocyte suspension, we wanted to investigate the fate and the function of  $\beta_1$  integrin in this system. To this end, we incubated keratinocytes with neutralizing anti- $\beta_1$  integrin antibody and performed anoikis assay. Confocal images demonstrate that  $\beta_1$  integrin was partially internalized after 30 min and almost completely at 2 hrs (Figure 4a). Indeed, as suspension proceeded, cell-to-cell contacts progressively decreased along with the internalization of integrin, in agreement with the observation that cells detached from each other are more prone to this kind of cell death (Frisch and Francis, 1994). Moreover, in suspended cells, the extracellular portion of  $\beta_1$  integrin was progressively degraded up to 24 hrs, as shown by the use of an antibody directed against this integrin domain (Figure 4b). When suspended keratinocytes were blotted using an antibody raised against the full-length  $\beta_1A$  integrin, a band of approximately 75 kD was detected between 2 and 24 hrs (Figure 4b), representing the cytoplasmic fragment of  $\beta_1A$  integrin, likely derived from the degradation of the extracellular domain.  $\beta_1B$  differs from wild-type  $\beta_1$  integrin ( $\beta_1A$ ) in the cytoplasmic domain (Altruda et al, 1990) and is mainly detected within the endoplasmic reticulum (Balzac et al, 1993; Kee et al, 2000). In agreement with previous works, we show that  $\beta_1A$  variant was mostly expressed at the cell membrane. By contrast, we detected  $\beta_1B$  form at the cytoplasmic level in a perinuclear fashion (Figure S2a). Given the difference in the cytoplasmic tails, we used specific antibodies recognizing the different  $\beta_1$  integrin domains. Although both  $\beta_1A$  and  $\beta_1B$  were expressed in human keratinocytes, the content of  $\beta_1A$  protein was markedly higher than that of  $\beta_1B$  integrin (Figure S2b). The role of this integrin form in keratinocytes is not fully understood. It has been proposed that  $\beta_1B$  acts as a dominant negative regulator of cell adhesion, by interfering with focal contact formation, thus decreasing adhesiveness (Retta et al, 1998). To better understand the role of this  $\beta_1$  integrin variant, we suspended human keratinocytes and blotted lysates using an antibody raised against  $\beta_1B$  integrin isoform. Unlike  $\beta_1A$ , no protein modifications of  $\beta_1B$  were observed up to 24 hrs (Figure 4b), indicating that no degradation of this variant occurs during suspension.

**$\beta_1A$  and  $\beta_1B$  integrins cluster and immunoprecipitate with caspase-8 during suspension.**

Confocal images of cells at 0 hr suspension show that  $\beta_1A$  expression was strictly localized at the cell membrane, while active caspase-8 was still undetectable. At 1 hr, caspase-8 became activated, while  $\beta_1A$  began to internalize. After 24 hrs suspension, caspase-8 was clearly detected with a cytoplasmic punctate pattern. At this time,  $\beta_1A$  integrin was also observed in perinuclear location (Figure 5a). On the other hand, at 0 hr,  $\beta_1B$  was predominantly expressed at the perinuclear level. This expression pattern remained unchanged up to 24 hrs, when caspase-8 was fully activated (Figure 5b). These results suggest that during keratinocyte suspension,  $\beta_1A$  integrin internalizes and locates close to the nucleus, likely co-localizing with  $\beta_1B$ . As  $\beta_1A$  and  $\beta_1B$  integrins co-localize with caspase-8 at the cytoplasmic level, we analyzed whether caspase-8 co-immunoprecipitates with either integrin isoforms. At 0 hr, we observed a weak band indicating a small amount of procaspase-8 bound to both  $\beta_1$  integrin variants. After 18 hrs suspension, binding of  $\beta_1A$  or  $\beta_1B$  to caspase-8 was markedly enhanced. As negative controls, no precipitates were observed when lysates were omitted nor when primary antibody was replaced by total rabbit serum (Figure 5c).

**Pro-apoptotic role of  $\beta_1B$  integrin.** To definitely assess the pro-apoptotic role of  $\beta_1B$  integrin, keratinocytes were transfected with a full-length  $\beta_1B$  expressing vector.  $\beta_1B$  was up-regulated at 24 hrs (Figure 6a), and it remained overexpressed until 48 hrs, both at the mRNA and the protein level (Figure 6a and b).  $\beta_1B$  integrin expression was strikingly increased at the intracellular level in transfected cells up to 48 hrs (Figure 6c). Moreover, keratinocytes that expressed high levels of  $\beta_1B$  integrin spontaneously detached from the culture dishes, as compared to mock cells (Figure 6e) and to keratinocytes transfected with a full-length  $\beta_1A$  expressing vector where adhesion and morphology were not affected (Figure 6d and e). Before inducing suspension, keratinocytes underwent cell death, as it appeared from the significant increase in the percent of TUNEL positive cells 24 and 48 hrs after transfection with  $\beta_1B$ , but not with  $\beta_1A$  (Figure 6f). When

keratinocytes were suspended, activation of caspase-8 at 0 hr occurred only in  $\beta_1B$  integrin transfected cells. During suspension, activation of caspase-8 was observed earlier in  $\beta_1B$  integrin overexpressing cells than in mock cells, as shown by the appearance of the active fragment at 3 and 6 hrs (Figure 6g). On the other hand, caspase-8 activation was not detected in keratinocytes transfected with siRNA  $\beta_1B$  at 12 hrs (Figure 6h). Finally, percentage of cells in subG1 peak at 12 hrs is higher in scramble than in siRNA  $\beta_1B$  transfected keratinocytes (Figure 6i).

## Discussion

In this study, we report that the beta 1 B isoform cooperates with antagonized beta1 A integrin to induce a form of integrin-mediated cell death in human keratinocytes, independently of death receptors. Death-domain-containing proteins trigger anoikis in epithelial cells (Frisch, 1999; Rytömaa et al, 1999), and anoikis is mediated by Fas/FasL and by the activation of caspase-8 in endothelial cells (Aoudjit and Vuori, 2001). Furthermore, FADD plays a critical role in detachment-induced apoptosis (Frisch, 1999), and epithelial cell lines expressing a dominant-negative form of FADD are protected from anoikis (Rytömaa et al, 1999). By contrast, we demonstrate that, while blocking beta 1 integrin activates caspase-8, death receptors are not involved in this system. Indeed, either cell suspension or anti-beta1 integrin failed to up-regulate Fas and FasL, and induced apoptosis in shFADD infected keratinocytes, with an early activation of caspase-8. This is consistent with previous work demonstrating that caspase-8 activation can occur in a manner independent of FADD (Stupack et al, 2001). Since keratinocytes undergo anoikis following the early activation of caspase-8, without the involvement of FADD, one might postulate that caspase-8 is activated downstream of caspase-3, in agreement with previous reports of death receptor-independent apoptosis (Ryu et al, 2005). Alternatively, caspase-8 activation could occur downstream of caspase-9 as an amplification loop of the intrinsic apoptotic pathway (Grossmann et al, 2001). However, both explanations have been ruled out by our previous study where we

reported that caspase-9 as well as executioner caspases are activated downstream of caspase-8 during keratinocyte anoikis (Marconi et al, 2004).

Given the exclusion of death receptors from the apoptotic pathway, we pursued the hypothesis of a direct interaction between integrins and caspase-8. Indeed, this study provides evidence that  $\beta_1$  integrin and caspase-8 co-localize in the cytoplasm and co-immunoprecipitate even before detachment. Immediately after suspension, caspase-8 is activated, while integrin is internalized and degraded. This suggests that keratinocyte suspension promotes  $\beta_1$  integrin fragmentation of the extracellular domain. Interestingly, this fragment was previously shown to induce apoptosis via caspase-8 activation in myocytes (Menon et al, 2006) and  $\beta_5$  cytoplasmic domain is proapoptotic in squamous cell carcinoma cell lines (Janes and Watt, 2004). Association of integrins with caspase-8 has been previously reported by Stupack and coworkers who, at variance with our study, observed a direct interaction of caspase-8 with  $\beta_3$  integrin only in cell lines adherent to an inappropriate ligand (Stupack et al, 2001). Furthermore,  $\alpha_v\beta_3$  integrin has been recently reported to associate with caspase-8 in osteoclasts plated on a non-specific matrix, and procaspase-8 was only detected in suspended cells (Zhao et al, 2005). To our knowledge, the present study describes a previously unreported form of IMD in normal human keratinocytes. However, at variance with the above publications, we show that a direct association between procaspase-8 and  $\beta_1$  integrin occurs in keratinocytes still adherent to the proper ligand. Thus in keratinocytes,  $\beta_1$  integrin-procaspase-8 complex exists in a “pre-apoptotic” state until a stimulus triggers IMD, further confirming that integrins themselves can play a proapoptotic role (Giancotti et al, 1990; Plath et al, 2000).

We also show that  $\beta_1B$  isoform plays a critical role in IMD.  $\beta_1B$  integrin is mostly expressed in the cytoplasm and, unlike  $\beta_1$  integrin, does not undergo degradation during suspension, while caspase-8 is activated. Moreover,  $\beta_1A$  and  $\beta_1B$  integrins cluster at the perinuclear level and immunoprecipitate with caspase-8 during suspension. A direct association between caspase-3 and  $\alpha_5\beta_1$  integrin was detected in non-adherent rat fibroblasts (Rajeswari et al, 2006), but, at variance with our study, activation of the caspase was observed only when the complex was

translocated to the cell membrane, and not in the cytoplasm. While the study did not address the signal allowing translocation, we hypothesize that, in our system, the  $\beta_1B$  variant located in the cytoplasm cooperates with the cytoplasmic fragment of  $\beta_1A$ . This integrin fragment derives from the degradation of  $\beta_1A$  integrin, and proves the unligated state of this integrin. At the cytoplasmic level, the two integrin isoforms enhance the recruitment of caspase-8 and induce cell death, in good agreement with previous results on the proapoptotic role of integrins (Kozlova et al, 2001).

The pro-apoptotic role of  $\beta_1B$  integrin was definitely shown both in cells overexpressing the cDNA and in cells where expression of the isoform was silenced by  $\beta_1B$  siRNA. It appears that  $\beta_1B$  is responsible for keratinocyte apoptosis even before cell suspension, with the early activation of caspase-8. These results indicate that the expression of  $\beta_1B$  integrin is sufficient to induce IMD in human keratinocytes, even when cells are adherent to the appropriate ligand. We postulate that unligated  $\beta_1A$  integrin is internalized, recruits procaspase-8 and clusters with  $\beta_1B$  integrin that in turn is ligated to procaspase-8 at perinuclear level. Caspase-8 is then activated, leading to the initiation of the extrinsic apoptotic pathway.

Squamous-cell carcinomas (SCCs) are tumors that arise in multilayered epithelia such as the epidermis, esophagus, cervix and the oral cavity. While the tumors differ according to tissue of origin in their frequency, aetiology and prognosis, they are all characterized by the alteration of integrins. In the epidermis, SCC may originate from keratinocyte stem cells (Zhang et al, 2005), which are protected from anoikis by virtue of the high levels of  $\beta_1$  integrin (Tiberio et al, 2002). SCCs are characterized by perturbation of integrin expression and extensive loss of the basement membrane (Ferreira et al, 2009; Savoia et al, 1994). Keratinocyte stem cells detach from the basement membrane, proliferate and may reattach to an inadequate location to eventually metastasize. We speculate that  $\beta_1B$  integrin works as a “sensor” of the detachment from the proper ligand and stimulates apoptosis to prevent spread of SCC, thus acting as a “fail-safe” device.

## **Materials and Methods**

### **DNA constructs**

Retroviral constructs pSUPER.Retro and pSUPER.Retro.shFADD were a kind gift from S.M. Frisch (West Virginia University, USA). A full-length  $\beta_1B$  or  $\beta_1A$  integrin cDNA were generated by RT-PCR and inserted in the pcDNA3.1/V5-His using the TOPO TA Expression kit (Invitrogen Corporation, Carlsbad, CA, USA).

### **Cell culture and infection**

Normal human keratinocytes were obtained from foreskin and cultured as described previously (Pincelli et al, 1997). The spontaneously transformed keratinocyte line HaCaT was kindly provided by Dr. N. Fusenig (DKFZ Heidelberg) and cultivated as already described (Boukamp et al, 1988). Packaging lines for the described retroviral constructs were generated by transinfection in the ecotropic Phoenix and amphotropic GP+envAm12 cells, as previously described (Grande et al, 1999). HaCat cells were transduced by two cycles of infection (6 hrs each) with viral supernatant in presence of polybrene (8  $\mu$ g/ml). Transduced cells were selected in presence of 1  $\mu$ g/ml puromycin (Sigma, St Louis, MO). Anoikis assay was performed as previously described (Marconi et al, 2004).

### **Cell culture and transfection**

HaCaT cells were transfected with described pcDNA3.1 plasmids with Lipofectamine 2000 and OptiMEM (Invitrogen) as suggested in the datasheet. Subconfluent normal human keratinocytes were double transfected with  $\beta_1B$  integrin siRNA (Sense: CAAAGUGGCUUAUAAAGUAUU; Antisense: 5'-PUACUUUAUAAGCCACUUUGUU; Thermo Scientific) with Lipofectamine 2000 and OptiMEM (Invitrogen) as suggested in the datasheet. Protein levels were detected by

Western Blotting as described below. Anoikis assay was performed as previously described (Marconi et al, 2004).

### **Antibodies**

Rabbit polyclonal antibodies used were: anti-caspase 8 (AR-18; gift of J.C. Reed, The Burnham Institute, USA); anti- $\beta_1$  integrin (M-106; Santa Cruz Biotechnology, Inc.); anti- $\beta_1A$  and anti- $\beta_1B$  antisera (gift from G. Tarone, University of Turin, Italy). Mouse monoclonal antibodies used were: neutralizing anti- $\beta_1$  integrin (Lia 1/2; Immunotech, Marseile, France); anti- $\beta_1$  integrin (K20; Santa Cruz Biotechnology, Inc.); anti-caspase 8 (Ab-3; Calbiochem, Darmstadt, Germany); anti-active caspase 8 (C15; Alexis Biochemicals); anti-FADD (A66-2; BD PharMingen); anti- $\beta$  actin (AC-15; Sigma). Secondary antibodies: Alexa Fluor 546 and 488-conjugated goat IgGs (Invitrogen); HRP-conjugated goat IgGs were also used (BioRad, Hercules, CA).

### **FACS, flow cytometry**

For anoikis (sub G1 peak positive), cells were suspended in hypotonic fluorochrome solution: propidium iodide (PI) 50 mg/ml, 0.1% sodium citrate, 0.1% Triton X-100; labeled at 4°C for at least 15 minutes and then analyzed using Epics XL flow cytometer (Epics XL, Coulter, Hialeah, FL, USA). Plots were analyzed by WinMDI 2.8 software.

Both cells undergone anoikis or treated with PVlgG, as a positive control for Fas Ligand expression, were analyzed for extracellular antigen expression. Cells were labelled with monoclonal anti-human Fas antibody (clone UB2; Immunotech) and with biotin-coniugated rabbit anti-mouse immunoglobulin (Dako, Copenaghen, Denmark), or with biotin conjugated anti-human FasL monoclonal antibody (Clone NOK-1; BDPharMingen) diluted in PBS with 5% FBS. Cells were then labelled with streptavidin/RPE-Cy5 (Dako) and analyzed with Epics XL flow cytometer (Coulter). Plots were analyzed by WinMDI 2.8 software.

### **ELISA**

To measure the release of soluble FasL during anoikis, we performed a two-site enzyme linked

immunoassay ELISA (MBL, Nagoya, Japan) on culture media according the manufacturer's instructions.

### **MTT assay**

MTT assay was performed in selected infected keratinocyte. After treatment with TRAIL (kindly provided by Dr. Martin Leverkus, Magdeburg, Germany) and CHX 1 $\mu$ g/ml, cells were incubated with MTT (Sigma) as previously described (Truzzi et al, 2008). Results are expressed as the mean  $\pm$  SD of three different experiments. Student's T-test was used for comparison of the means.

### **Co-immunoprecipitation and immunoblotting**

Lysates were processed for IP and WB as described previously (Marconi et al, 2004) or with lysis buffer (50 mM Tris-HCl, pH 8.5, 150 mM NaCl, 1% deoxycolate, 1% Triton X-100, 0.1% SDS, 0.2% sodium azide and inhibitor cocktail [Roche]). Proteins were immunoprecipitated with anti- $\beta_1$  integrin (K20) or anti- $\beta_1$  antisera. Samples were resolved by 7 or 12% SDS-PAGE and immunoblotted as described previously (Marconi et al, 2004).

### **Immunofluorescence analysis**

Sub-confluent cells were fixed in ice-cold methanol or PFA 4% and incubated with monoclonal anti- $\beta_1$  integrin antibody and the appropriate secondary antibody. Cells were then labelled with polyclonal anti-caspase 8 antibody and appropriate secondary antibody. Alternatively cells were first stained with rabbit anti- $\beta_1$  integrin variants antisera and with mouse anti-active caspase-8. Micrographs were taken on a Confocal Scanning Laser Microscopy (Leica TCS4D) (Leica, Exton,PA). In selected cases, a series of images of the same x/y plane was taken and the reconstruction made on the z-axis.

### **Internalization assay**

Keratinocytes were pre-incubated with neutralizing anti- $\beta_1$  integrin antibody and anoikis assay was performed. At each time-point, cells were fixed in PFA 4% and immunofluorescence performed with the addition of the secondary antibody.

### **RT-PCR**

Total RNA was extracted from transfected cells and RT-PCR was performed as previously described (Truzzi et al, 2008). Nucleotide sequences of the  $\beta_1$ B integrin oligomers used (MWG Biotech, Ebersberg, Germany) were:  $\beta_1$ B-DP: 5'-gggaacaacgagggtcatggttcat-3';  $\beta_1$ B-RP: 5'-ttataagccacttgcttttgatg-3'.

### **TUNEL assay**

Cells were fixed in PFA 4% both 24 and 48 hrs after transfection. Then cells were stained with the In Situ Cell Death Detection Kit (Roche Diagnostics, Basel, Switzerland) as recommended by manufacturer. Positive cells were analyzed as detailed previously (Truzzi et al, 2008).

### **Conflict of Interest**

The authors declare no conflict of interest.

### **Acknowledgments**

We thank Prof. G. Tarone for the critical reading of this manuscript. This work was partially supported by a grant from Ministry of Instruction, University and Research 20077FPJ22.

## References

Altruda F, Cervella P, Tarone G, et al. (1990). A human integrin beta 1 subunit with a unique cytoplasmic domain generated by alternative mRNA processing. *Gene* 95: 261-266.

Aoudjit F and Vuori K. (2001). Matrix attachment regulates Fas-induced apoptosis in endothelial cells: a role for c-flip and implications for anoikis. *J Cell Biol* 152: 633-643.

Arredondo J, Chernyavsky AI, Karaouni A, et al. (2005). Novel mechanisms of target cell death and survival and of therapeutic action of IVIg in Pemphigus. *Am J Pathol* 167:1531-44.

Balzac F, Belkin AM, Kotliansky VE, et al. (1993). Expression and functional analysis of a cytoplasmic domain variant of the beta 1 integrin subunit. *J Cell Biol* 121: 171-178.

Boukamp P, Petrussevska RT, Breitkreutz D, et al. (1988). Normal keratinization in a spontaneously immortalized aneuploid human keratinocyte cell line. *J Cell Biol* 106: 761-771.

Brassard DL, Maxwell E, Malkowski M, et al. (1999). Integrin alpha(v)beta(3)-mediated activation of apoptosis. *Exp Cell Res* 251: 33-45.

Ferreira M, Fujiwara H, Morita K, et al. (2009). An activating beta1 integrin mutation increases the conversion of benign to malignant skin tumors. *Cancer Res* 69: 1334-1342.

Fleischmajer R, Perlish JS, MacDonald ED, et al. (1998). There is binding of collagen IV to beta 1 integrin during early skin basement membrane assembly. *Ann N Y Acad Sci* 857: 212-227.

Frisch SM. (1999). Evidence for a function of death-receptor-related, death-domain-containing proteins in anoikis. *Curr Biol* 9: 1047-1049.

Frisch SM and Francis H. (1994). Disruption of epithelial cell-matrix interactions induces apoptosis. *J Cell Biol* 124: 619-626.

Giancotti FG, and Ruoslahti E. (1990). Elevated levels of the alpha 5 beta 1 fibronectin receptor suppress the transformed phenotype of Chinese hamster ovary cells. *Cell* 60: 849-859.

Grande A, Piovani B, Aiuti A, et al. (1999). Transcriptional targeting of retroviral vectors to the erythroblastic progeny of transduced hematopoietic stem cells. *Blood* 93: 3276-3285.

Grossmann J, Walther K, Artinger M, et al. (2001). Apoptotic signaling during initiation of detachment-induced apoptosis ("anoikis") of primary human intestinal epithelial cells. *Cell Growth Differ* 12: 147-155.

Howlett AR, Bailey N, Damsky C, et al. (1995). Cellular growth and survival are mediated by beta 1 integrins in normal human breast epithelium but not in breast carcinoma. *J Cell Sci* 108: 1945-1957.

Janes SM, and Watt FM. (2004). Switch from alphavbeta5 to alphavbeta6 integrin expression protects squamous cell carcinomas from anoikis. *J Cell Biol* 166: 419-431.

Kee WJ, Li ER, Watt FM. (2000). beta1B integrin subunit contains a double lysine motif that can cause accumulation within the endoplasmic reticulum. *J Cell Biochem* 78: 97-111.

Kozlova NI, Morozevich GE, Chubukina AN, et al. (2001). Integrin alphavbeta3 promotes anchorage-dependent apoptosis in human intestinal carcinoma cells. *Oncogene* 20: 4710-4717.

Marconi A, Atzei P, Panza C, et al. (2004). FLICE/caspase-8 activation triggers anoikis induced by beta1-integrin blockade in human keratinocytes. *J Cell Sci* 117: 5815-5823.

Menon B, Krishnamurthy P, Kaverina E, et al. (2006). Expression of the cytoplasmic domain of beta1 integrin induces apoptosis in adult rat ventricular myocytes (ARVM) via the involvement of caspase-8 and mitochondrial death pathway. *Basic Res Cardiol* 101: 485-493.

Muzio M, Stockwell BR, Stennicke HR, et al. (1998). An induced proximity model for caspase-8 activation. *J Biol Chem* 273: 2926-2930.

Plath T, Detjen K, Welzel M, et al. (2000). A novel function for the tumor suppressor p16(INK4a): induction of anoikis via upregulation of the alpha(5)beta(1) fibronectin receptor. *J Cell Biol* 150: 1467-1478.

Pincelli C, Haake AR, Benassi L, et al. (1997). Autocrine nerve growth factor protects human keratinocytes from apoptosis through its high affinity receptor (TRK): a role for BCL-2. *J. Invest. Dermatol* 109: 757-764.

Rajeswari J, and Pande G. (2006). Direct association between caspase 3 and alpha5beta1 integrin and its role during anoikis of rat fibroblasts. *Cell Biol Int* 30: 963-969.

Retta SF, Balzac F, Ferraris P, et al. (1998). beta1-integrin cytoplasmic subdomains involved in dominant negative function. *Mol Biol Cell* 9: 715-731.

Rytömaa M, Martins LM, Downward J. (1999). Involvement of FADD and caspase-8 signalling in detachment-induced apoptosis. *Curr Biol* 9: 1043-1046.

Ryu HY, Emberley JK, Schlezinger JJ, et al. (2005). Environmental chemical-induced bone marrow B cell apoptosis: death receptor-independent activation of a caspase-3 to caspase-8 pathway. *Mol Pharmacol* 68: 1087-1096.

Savoia P, Cremona O, Trusolino L, et al. (1994). Integrins and basement membrane proteins in skin carcinomas. *Pathol Res Pract* 190: 950-954.

Stupack DG, Puente XS, Boutsaboualoy S, et al. (2001). Apoptosis of adherent cells by recruitment of caspase-8 to unligated integrins. *J Cell Biol* 155: 459-470.

Tiberio R, Marconi A, Fila C, et al. (2002). Keratinocytes enriched for stem cells are protected from anoikis via an integrin signaling pathway in a Bcl-2 dependent manner. *FEBS Lett* 524: 139-144.

Truzzi F, Marconi A, Lotti R, et al. (2008). Neurotrophins and their receptors stimulate melanoma cell proliferation and migration. *J Invest Dermatol* 128: 2031-2040.

Zhang W, Hanks AN, Boucher K, et al. (2005). UVB-induced apoptosis drives clonal expansion during skin tumor development. *Carcinogenesis*. 26: 249-57.

Zhao H, Ross FP, Teitelbaum SL. (2005). Unoccupied alpha(v)beta3 integrin regulates osteoclast apoptosis by transmitting a positive death signal. *Mol Endocrinol* 19: 771-780.

## Figure Legends

**Figure 1.** Fas/FasL modulation in keratinocyte anoikis. **(a)** Fas expression. Keratinocytes were suspended in polypropylene tubes in presence or absence of anti- $\beta_1$  integrin neutralizing antibody. Cells were then incubated with anti-Fas antibody and analyzed by flow cytometry. **(b)** FasL expression. Keratinocytes were treated as before, incubated with anti-FasL antibody and analyzed by flow cytometry. Pemphigus vulgaris immunoglobulins G (PVIgG) were used as a positive control stimulus for FasL (Arredondo et al, 2005). **(c)** sFasL release. Keratinocytes were kept in suspension in polypropylene tubes for different time-points and culture media were collected. Soluble FasL in culture media was analyzed by ELISA. PVIgG were used as a positive control stimulus for FasL.

**Figure 2.** FADD is not required for keratinocyte anoikis. **(a)** FADD protein expression in infected HaCaT keratinocytes. Proliferating HaCaT cells were retrovirally infected with FADD shRNA (HaCaT shFADD) or with control shRNA (HaCaT mock). Cells were selected in presence of puromycin and lysed. Lysates were run on SDS-acrylamide gel and blotted onto nitrocellulose membrane. Blot was incubated against anti-FADD antibody.  $\beta$ -actin was used as loading control. **(b)** shFADD prevents TRAIL-induced apoptosis in keratinocytes. Infected HaCaT cells were treated with increasing doses of TRAIL and viability was analyzed by MTT assay after 24 hrs. **(c)** Caspase-8 activation in shFADD cells. Retrovirally infected HaCaT cells were suspended and treated with anti- $\beta_1$  integrin neutralizing antibody. Cells were lysed at different time points, and lysates were immunoblotted with anti-caspase-8 antibody.  $\beta$ -actin was used as loading control. **(d)** shFADD cells show the same percent of sub G1 peak positive cells. Mock and shFADD keratinocytes were kept in suspension in polypropylene tubes in presence or absence of anti- $\beta_1$  integrin antibody. After 18 hrs cells were stained in PI-solution. Sub-G1 peak positive cells (M1) were analyzed by flow cytometry. Data represent the mean from three independent experiments.

**Figure 3.**  $\beta_1$  integrin associates with caspase-8. **(a)**  $\beta_1$  integrin co-localizes with caspase-8. Subconfluent normal human keratinocytes were fixed in PFA 4%. Cells were immunostained with anti- $\beta_1$  integrin and anti-procaspase-8 antibodies. Bottom panel represents the merge. Cells were analyzed by confocal microscopy. Bar, 15  $\mu\text{m}$ . **(b)** Early activation of caspase-8 during anoikis. Normal human keratinocytes were trypsinized (0 hr) and kept in suspension in polypropylene tubes for different time-points. At the end of single time-point cells were lysed and lysates were immunoblotted with anti-procaspase-8 antibody.  $\beta$ -actin was used as loading control. **(c)**  $\beta_1$  integrin immunoprecipitates with procaspase-8. Keratinocytes were treated as in (b). Lysates were immunoprecipitated with anti- $\beta_1$  integrin antibody. Immunoprecipitates were blotted with anti- $\beta_1$  integrin, anti-procaspase-8 and anti-FADD antibodies. Control 1, immunoprecipitate with anti- $\beta_1$  integrin antibody without lysates; control 2, immunoprecipitate with a non-keratinocyte antigen antibody (anti-vimentin). Control samples were run simultaneously in a different gel.

**Figure 4.**  $\beta_1$  integrin is internalized and degraded during keratinocyte suspension. **(a)**  $\beta_1$  integrin internalization. Keratinocytes were incubated with the neutralizing anti- $\beta_1$  integrin antibody for 20 min at 4°C. Cells were then suspended for different time points, fixed in PFA 4% and analyzed by confocal microscopy. Images outside the x/y planes represent the z-line projection of the cross-identified area. Bar, 8,5  $\mu\text{m}$ . **(b)**  $\beta_1$  integrin degradation. Keratinocytes were suspended in polypropylene tubes and lysed at different time points. Lysates were probed against  $\beta_1$  integrin, using an antibody directed to the extracellular portion. The same lysates were blotted with anti- $\beta_1\text{A}$  and  $\beta_1\text{B}$  integrin antisera.  $\beta$ -actin was used as loading control.

**Figure 5.**  $\beta_1$  integrin variants immunoprecipitate with caspase-8 during suspension. **(a-b)**  $\beta_1\text{A}$  and  $\beta_1\text{B}$  integrin localization during caspase-8 activation. Keratinocytes were suspended in polypropylene tubes and fixed in PFA 4% at different time points. Cells were spun and immunostained at room temperature with anti- $\beta_1\text{A}$  (a) and anti- $\beta_1\text{B}$  integrin (b) antisera and anti-active caspase-8 antibody (a,b). Cells were analyzed by confocal microscopy. (a) Bar 21  $\mu\text{m}$ . (b)

Bar 17  $\mu\text{m}$ . (c)  $\beta_1$  integrin variants immunoprecipitate with procaspase-8. Keratinocytes were suspended in polypropylene tubes and lysed after 18 hrs of suspension. Lysates were alternatively immunoprecipitated with anti- $\beta_1\text{B}$  integrin, anti- $\beta_1\text{A}$  integrin or anti-extracellular domain of  $\beta_1$  integrin antibodies. Immunoprecipitates were blotted with anti-procaspase-8. Control 1: immunoprecipitate without lysates; Control 2: immunoprecipitate with total rabbit serum.

**Figure 6.**  $\beta_1\text{B}$  integrin is pro-apoptotic in keratinocytes. (a) HaCaT cells were transfected with a full-length  $\beta_1\text{B}$  integrin-containing vector (pcDNA3.1/ $\beta_1\text{B}$ ) or with the empty vector (pcDNA3.1). mRNA was extracted from cells after 24 and 48 hrs of transfection,  $\beta_1\text{B}$  integrin expression was analyzed by RT-PCR.  $\beta$ -actin was used as loading control. (b) HaCaT cells were transfected as in (a). Cells were lysed and protein extract were blotted with anti- $\beta_1\text{B}$  integrin antibody.  $\beta$ -actin is the loading control. (c) HaCaT cells were transfected as before. Keratinocytes were fixed in PFA 4% after 24 and 48 hrs of transfection. Cells were spinned and stained with anti- $\beta_1$  integrin antibody and analyzed at confocal microscopy. Images outside the x/y planes represent the z-line projection of the cross-identified area. Bar 12  $\mu\text{m}$ . (d) HaCaT cells were transfected with a full-length  $\beta_1\text{A}$  integrin-containing vector (pcDNA3.1/ $\beta_1\text{A}$ ). Cells were lysed and protein extracts were blotted with anti- $\beta_1\text{A}$  integrin antibody.  $\beta$ -actin is the loading control. (e) Micrograph pictures of transfected cells after 48 hrs of transfection. Bar 50  $\mu\text{m}$  (f) HaCaT cells transfected with  $\beta_1\text{B}$ ,  $\beta_1\text{A}$  or empty vector were stained with TUNEL assay 24 and 48 hrs after transfection. (g)  $\beta_1\text{B}$  transfected HaCaT cells were trypsinized 48 hrs after transfection and kept in suspension in polypropylene tubes for different time-points. Cells were lysed and lysates were blotted against anti-caspase-8 antibody.  $\beta$ -actin was used as loading control. (h) Primary human keratinocytes were transfected with siRNA  $\beta_1\text{B}$  or scramble and kept in suspension in polypropylene tubes for 12 hrs. Cell lysates were blotted against anti- $\beta_1\text{B}$  integrin or anti-caspase-8 antibody.  $\beta$ -actin was used as loading control. (i) In the same experiment, cells were stained in PI-solution. Sub-G1 peak positive cells (M1) were analyzed by flow cytometry. Data represent the mean from three independent experiments.

## Supplementary Materials