

ABSTRACT VOLUME

GeoTirol 2016

Annual Meeting DGGV | 25–28 September 2016

Innsbruck | Austria




Photo: Martin Reiser



Annual Meeting



hosting:

25-29/09/2016 INNSBRUCK



PAN GEO
AUSTRIA

Bodenseetagung

© BODENSEETAGUNG OG, Metznere / Skayne

Recommended Citation

Ortner, H. [Ed.] (2016): Abstract Volume of GeoTirol2016 - Annual Meeting of DGGV and PAN-GEO Austria, 25-28. September 2016, Innsbruck.

Disclaimer and Copyright

Each author is responsible for the content of his or her abstract.

Imprint**Publisher**

Institute of Geology
University of Innsbruck
Innrain 52f
6020 Innsbruck

Editor

Hugo Ortner

SILURIAN ANOXIC EVENTS AT THE CELLON SECTION (AUSTRIA) THROUGH AN ICHNOFABRIC EYE

BAUCON, Andrea* (1); FERRETTI, Annalisa (1); SCHOENLAUB, Hanspeter (2)

1: University of Modena, Italy; 2: Geopark Carnic Alps, Austria

andrea@tracemaker.com

ichnology, ichnofabric, anoxic events, Lau event, Chondrites

The Ordovician and Silurian periods are characterized by major biotic and geochemical events, among which large carbon isotope excursions associated with mass extinctions and anoxia. The Cellon section (Austria) is an ideal case study to understand the dynamics of these events. In fact, it represents not only a reference section for the Silurian of the world, but also it provides an almost continuous sedimentary record ranging from the Upper Ordovician to the Lower Devonian.

The goal of this study is to understand the relationship between the mentioned biotic and geochemical events and organism-substrate interactions. To this aim, the ichnofabric approach – considering the aspects of sediment texture imparted by biogenic activity – is applied to the Cellon Section. Results show three oxygen-related ichnofabrics:

The *Thalassinoides* ichnofabric is characterized by intense bioturbation (bioturbation index 5-6) and the dominant presence of *Thalassinoides* burrows. Because of the high bioturbation index, this ichnofabric is interpreted to reflect oxygenated condition of bottom waters.

The Chondrites ichnofabric is characterized by low bioturbation (BI 0-1) and the monoichnospecific presence of Chondrites. Because of the chemosymbiotic, sulphide-mining strategy of the Chondrites producers, this ichnofabric is interpreted to represent dysoxic conditions.

The unbioturbated ichnofabric is characterized by laminated, unbioturbated (BI 0) fabrics. The absence of benthic activity suggest widespread anoxic conditions.

Data interpretation suggest that the onset of anoxic events are characterized by a relatively abrupt transition from sediment-feeding strategies (*Thalassinoides* ichnofabric) to chemosymbiotic ones (Chondrites).




Photo: Robbie Shone