

# Cytogenetic and molecular analysis of the holocentric chromosomes of the potato aphid *Macrosiphum euphorbiae* (Thomas, 1878)

Valentina Monti<sup>1,2</sup>, Gian Carlo Manicardi<sup>2</sup>, Mauro Mandrioli<sup>1</sup>

**1** Dipartimento di Biologia, Università di Modena e Reggio Emilia, Via Campi 213/D, 41125 Modena, Italy

**2** Dipartimento di Scienze Agrarie e degli Alimenti, Università di Modena e Reggio Emilia, Via Amendola 2, 42100 Reggio Emilia, Italy

Corresponding author: *Mauro Mandrioli* (mauro.mandrioli@unimo.it)

---

Academic editor: *Valentina Kuznetsova* | Received 17 June 2010 | Accepted 19 July 2010 | Published 24 August 2011

**Citation:** Monti V, Manicardi GC, Mandrioli M (2011) Cytogenetic and molecular analysis of the holocentric chromosomes of the potato aphid *Macrosiphum euphorbiae* (Thomas, 1878). *Comparative Cytogenetics* 5(3): 163–172. doi: 10.3897/CompCytogen.v5i3.1724

---

## Abstract

Cytogenetic and molecular investigations on the holocentric chromosomes of the aphid *Macrosiphum euphorbiae* (Thomas, 1878) have been carried out using silver staining and C-banding (followed by chromomycin A<sub>3</sub> and DAPI staining) in order to improve our knowledge about the structure of aphid chromosomes. The C-banding pattern is peculiar since only the two X chromosomes and a single pair of autosomes presented heterochromatic bands. Silver staining and FISH with the 28S rDNA probe localized the rDNA genes on one telomere of each X chromosome that were also brightly fluorescent after chromomycin A<sub>3</sub> staining of C-banded chromosomes, whereas all other heterochromatic bands were DAPI positive. Interestingly, a remarkable nucleolar organizing region (NOR) heteromorphism was present making the two X chromosomes easily distinguishable. Southern blotting and FISH assessed the presence of the (TTAGG)<sub>n</sub> repeat at the ends of all the *M. euphorbiae* chromosomes. Karyotype analysis showed that all males possessed the X chromosome with the larger amount of rDNA suggesting a non-Mendelian inheritance of the two X chromosomes.

## Keywords

aphid, holocentric chromosomes, telomeres, heterochromatin, NOR heteromorphism

## Introduction

In the last decades classic and molecular cytogenetics provided an integrated approach for the structural, functional and evolutionary analysis of aphid holocentric chromosomes (Blackman 1980, Manicardi et al. 1998, Mandrioli et al. 1999a, 1999b, Manicardi et al. 2002, Mandrioli et al. 2011).

Interest in aphid cytogenetics is mostly due to the holocentric/holokinetic structure of their chromosomes that present a diffused centromeric activity (Wrensch et al. 1994). This particular type of chromatin organization has been described in almost all the eukaryotic taxa examined so far, with the exception of echinoderms and chordates (Wrensch et al. 1994).

Aphids, in view of the ease with which mitotic chromosome can be obtained from embryonic tissues, represent an ideal model to better understand the architecture of holocentric chromosomes, and to work out the differences/similarities with monocentric ones (Manicardi et al. 2002). At the same time, a cytogenetic analysis of aphids is very helpful since the description of species-specific chromosomal markers could make easier the identification of species that is, at present, quite difficult (Rakauskas 1998, Blackman 1980).

Aphid X chromosomes have been studied with great attention since they present several structural constraints (Manicardi et al. 2002). In particular, the X chromosomes showed a large amount of heterochromatin and possessed the rDNA cluster located at one telomere in almost all the species studied at a cytogenetic level (Mandrioli et al. 1999a, b, Manicardi et al. 2002). Furthermore, aphid X chromosomes seem to be more stable than autosomes since, when fragmentations occur, they mostly affected autosomes leaving X chromosomes usually as the longest in aphid karyotype (Khuda-Buksh and Datta 1981, Blackman 1987, Hales 1989).

In order to better understand X chromosome evolution in aphids, we decided to carry out a cytogenetic analysis of the holocentric chromosomes of the aphid *Macrosiphum euphorbiae* (Thomas, 1878), an important pest of several crops, belonging to a genus that has been up to date scarcely studied at a cytogenetic level (Blackman 1980).

## Material and methods

Specimens of *Macrosiphum euphorbiae* were collected on *Bellis perennis* (Linnaeus, 1753) in Modena (Italy) and maintained at 22°C with 16:8 hours light/darkness on *B. perennis* plants. Male aphids were obtained by exposing parthenogenetic females to short photoperiods (8:16 hours light/darkness) according to Crema (1979).

Chromosome preparations from 150 parthenogenetic females were made by spreading embryo cells, as described by Manicardi et al. (1996). Male chromosomes have been obtained by squash preparation of 30 embryos as reported by Manicardi et al. (1991).

C banding treatment was performed according to the technique of Sumner (1972). C banded chromosomes were stained with DAPI according to Donlon and Magenis (1983) and with chromomycin A<sub>3</sub> (CMA<sub>3</sub>) as described in Schweizer (1976). NOR regions were labelled by silver staining following the technique of Howell and Black (1980).

DNA extraction from aphid embryos was performed as described in Mandrioli et al. (1999a).

The 28S rDNA probe was obtained by PCR amplification of a 400 bp long fragment of the 28S rDNA gene using the two primers, F (5'-AACAAACAACCGA-TACGTTCCG) and R (5'-CTCTGTCCGTTTACAACCGAGC), designed according to the insect 28S rRNA sequences available in GenBank. The amplification mix contained 100 ng genomic DNA, 1 mM of each primer, 200 mM dNTPs and 2 U of DyNAzyme II polymerase (Finnzymes Oy). Amplification was performed using a Hybaid thermal-cycler at an annealing temperature of 60°C for 1 min with an extension time of 1 min at 72°C.

In order to test the presence of the telomeric (TTAGG)<sub>n</sub> repeat, a probe was obtained by PCR amplification using the two primers F (TTAGG)<sub>5</sub> and R (CCTAA)<sub>5</sub> in the absence of template, as described by Ijdo et al. (1991).


The telomeric and 28S rDNA probes were labelled using the PCR DIG labelling mix (Roche) according to the Roche protocols.

Southern blotting and fluorescent *in situ* hybridization (FISH) were made as described by Mandrioli et al. (1999a). FISH slides were observed using a Zeiss Axioplan epifluorescence microscope equipped with a 100 W mercury light source. Photographs of the fluorescent images were taken using a CCD camera (Spot, Digital Instrument, Madison, USA) and using the Spot software supplied with the camera and processed using Adobe Photoshop (Adobe Systems, Mountain View, CA).

## Results

The parthenogenetic females of *M. euphorbiae* showed a chromosome number of 2n=10 (Fig. 1). C banding followed by CMA<sub>3</sub> staining showed a bright fluorescence exclusively limited to one telomere of the two longest chromosomes (Fig. 1a) that, on the basis of the comparison with male karyotype, have been identified as X chromosomes. DAPI staining showed a large heterochromatic band at the opposite end of the X chromosomes in respect to the GC-rich CMA<sub>3</sub>-stained telomere. A second heterochromatic band was observed at one telomere of the autosome pair 2 (Fig. 1b).

The overlapping between CMA<sub>3</sub> areas and rDNA genes has been confirmed by both silver staining (Fig. 1c) and FISH with the 28S rDNA probe (Fig. 1d) showing an exclusive localization of the rDNA genes on *M. euphorbiae* X chromosomes. Silver staining, FISH and CMA<sub>3</sub> staining demonstrated the remarkable occurrence of heteromorphism between homologous NORs on X chromosomes (Figs. 1a, c, d) allowing us to distinguish the two X chromosomes. The same heteromorphism was also evident in interphase nuclei (Fig. 1a).



**Figure 1a–h.** *M. euphorbiae* chromosomes, stained with CMA<sub>3</sub> (**a**) and DAPI (**b**) after C banding, showing heterochromatin on one telomere of the two X chromosomes and on autosome pair 2. Silver staining (**c**) and FISH with a 28S rDNA probe (**d**) evidenced heteromorphous NORs located at one telomere of each X chromosome that was also fluorescent after CMA<sub>3</sub> staining, as summarized in the panel (**e**) Southern blotting (**f**) after digestion with *Xho*I of DNA samples of *A. pisum* **1**, *M. euphorbiae* **2** and *D. melanogaster* **3** together with FISH (**g**) assessed that the telomeric sequence (TTAGG)<sub>n</sub> constitute each chromosomal end of the *M. euphorbiae* chromosomes. CMA<sub>3</sub> staining of male chromosomes showed that all the male plates present the X chromosome with the larger NOR (**h**) Arrows indicate X chromosomes. Arrowheads indicate NORs. Asterisks evidence the presence of heteromorphous nucleoli in the *M. euphorbiae* nuclei. Bar = 10 μm.

The presence of the (TTAGG)<sub>n</sub> repeat has been evaluated by Southern blotting and FISH. Southern blotting revealed a diffuse smear of hybridization (Fig. 1f), whereas FISH experiments with the telomeric (TTAGG)<sub>n</sub> probe showed bright FITC-fluorescent spots at the ends of all chromosomes (Fig. 1g). In the interphase nuclei of *M. euphorbiae*, telomeres clustered in few highly fluorescent foci (Fig. 1g).

The male karyotype consisted of 8 autosomes and one X chromosome only (Fig. 1h). Interestingly, all the analysed males possessed the X chromosome with the larger NOR.

## Discussion

Currently, more than 4000 aphid species have been described, but the chromosome number has been reported only for about 500 of them (Blackman 1980). Furthermore, less than 10% of the described aphid karyotypes have been deeply studied at a cytogenetic level, even if aphids are intriguing model in animal cytogenetics in view of the holocentric nature of their chromosomes (Manicardi et al. 2002).

According to our results, *M. euphorbiae* has a chromosome number of  $2n=10$ , which represents the typical diploid chromosome number reported in the literature for species of this genus (Blackman 1980).

C banding carried out on *M. euphorbiae* mitotic chromosomes revealed that heterochromatin was not equilocated on each chromosome, but limited to telomeric regions of the two X chromosomes and to autosome pair 2. A preferential storage of heterochromatin on X chromosomes has been previously observed in almost all the aphid species cytogenetically studied to date with the exception of *Diuraphis noxia* (Mordvilko ex Kurdjumov, 1913) (Novotna et al. 2011), but C-bands on a single autosome pair were reported only in *Acyrtosiphon pisum* (Harris, 1776) (Bizzaro et al. 2000) and *Brevicoryne brassicae* (Linnaeus, 1758) (Giannini et al. 2003).

The different responses to  $CMA_3$  and DAPI staining after C banding point out a DNA composition heterogeneity of *M. euphorbiae* heterochromatin. Indeed, GC-rich NOR-associated heterochromatin differs from all other heterochromatic bands that are made by AT-rich DNAs. This pattern of heterochromatin heterogeneity seems to be a general characteristic of aphid chromatin since it has been described in all species investigated so far at a cytogenetic level (Manicardi et al. 2002).

Contrary to the protocol followed for several aphid species, in order to induce a clear-cut banding pattern on *M. euphorbiae* chromosomes, we modified the usual C-banding procedure making a 7 minutes long barium hydroxide treatment. Difficulties in obtaining clear-cut C-banding have been recently reported in the aphid *D. noxia* (Novotna et al. 2011). Interestingly, both these species belong to the tribe Macrosiphini, whereas most of the species with a clear cut banding were Aphidini. Further studies on the tribe Macrosiphini could shed light on the peculiarities of their chromatin organization, which are at the basis of this different banding propensity.

Southern blot experiments with the  $(TTAGG)_n$  probe showed smears in *M. euphorbiae* genome suggesting that its telomeres are composed of TTAGG repeats. This result has been confirmed by FISH experiments that clearly showed a hybridization signal on each telomere, whereas no evidence of any interstitial labelling has been observed demonstrating that the TTAGG repeats are restricted to the terminal regions of all aphid chromosomes. The presence of the  $(TTAGG)_n$  repeat in *M. euphorbiae* further support the hypothesis that this telomeric sequence is common in aphids (Bizzaro

et al. 2000, Monti et al. 2011), so that its absence in *D. noxia* (Novotna et al. 2011) could be considered an exception, as previously reported in other insects (Frydrychová et al. 2004, Lukhtanov and Kuznetsova 2010).

In interphase nuclei of most organisms the telomeric regions are situated in an ordered fashion with an association to the nuclear matrix and clustering at least in some stage of cell life (Palladino et al. 1993, Luderus et al. 1996, Pryde et al. 1997). Accordingly to previous results in aphids (Monti et al. 2011), telomeres appeared clustered into few foci in the *M. euphorbiae* nuclei and were not located mainly near the nuclear periphery as reported in other insects such as *Drosophila melanogaster* (Meigen, 1830) (Hochstrasser et al. 1986) and the cabbage moth *Mamestra brassicae* (Linnaeus, 1758) (Mandrioli 2002).

NOR number and position have been frequently reported as highly variable in insects, where rDNA genes have been frequently found also on autosomes or only on autosomes as generally reported in Lepidoptera and Psylloidea (Maryńska-Nadachowska et al. 1992, Maryńska-Nadachowska and Grozeva 2001, Nguyen et al. 2010), including several species with multiple NORs (e.g. Postiglioni and Brum-Zorrilla 1988, Juan et al. 1993). Silver staining of *M. euphorbiae* mitotic metaphases revealed two dots located on one telomere of each X chromosome. This seems to be a highly conservative characteristic of aphid chromosomes, since the same NOR localization has been described in almost all the aphids species studied to date (Kuznetsova and Gandrabur 1991, Kuznetsova and Maryńska-Nadachowska 1993, Blackman and Spence 1996, Manicardi et al. 1998, 2002), with the unique exception of *Schoutedenia ralumensis* (Rübsaamen, 1905) and *Maculolachnus submacula* (Walker, 1848) that present autosomal NORs and *Amphorophora idaei* (Borner, 1839) showing interstitial NORs on X chromosomes (Blackman 1987).

In several aphid species silver staining revealed the occurrence of an appreciable level of heteromorphism between homologous NORs due to a different distribution of rDNA genes between the two X chromosomes, but, in all the previous studies, different levels of heteromorphism have been observed both at inter- and intra-individual levels (Blackman and Spence 1996, Mandrioli et al. 1999b; Manicardi et al. 1998, 2002). Contrarily to previous observations, all the observed *M. euphorbiae* plates presented an X chromosome with a NOR region larger than the homologue allowing us to clearly differentiate the two sex chromosomes. In view of this stable chromosomal marker we performed experiments of male determination in order to evaluate if during this process both the X chromosomes have the same chances of being inherited in males.

All the parthenogenetic eggs during the prophase present two X chromosomes linked by NORs (Schrader 1940, Orlando 1974, Hales and Mitler 1983, Blackman and Hales 1986). However, in eggs developing as females, the connection is quickly lost, but in male generating eggs the X chromosomes remain attached by sticky NORs and undergo a sort of non-canonical reductional division (Blackman and Hales 1986). At the end of this peculiar division the egg has one X chromosome only and it is determined as a male.

The observation that all the *M. euphorbiae* male metaphases had an X chromosome with a large NOR evidenced a selective bias favouring X chromosomes with larger number of rDNA genes. This fact could be due by a non-random elimination of one X chromosome during male determination process or by the early abortion of embryos containing an X chromosome with few rDNA genes. Contrarily to what has been observed in *Megoura viciae* (Buckton, 1876) (Crema 1981), we never observed the presence of aborted eggs among the developing male embryos of *M. euphorbiae* so that we hypothesized that the bias occurred during the male determination process which ended with a selective elimination of the X chromosome bearing few rDNA genes.

The phenomenon of biased inheritance of X chromosomes in aphids seems to be controversial among aphids since observations on *Sitobion fragariae* (Walker, 1848), using an X-linked polymorphic microsatellite marker, suggested that X chromosome loss during male determination was random (Wilson et al. 1997), whereas the presence of strong biases in the transmission of sex chromosomes has been reported in *Rhopalosiphum padi* (Linnaeus, 1758) (Franz et al., 2005)

On the basis of our results on *M. euphorbiae*, we suggest that the presence of an unequal distribution of rDNA between the two X chromosomes could affect the attachment of the X chromosome by sticky NORs favouring the loss of the X chromosome with few rDNA genes and a not random inheritance of the X chromosomes. On the contrary, aphid specimens with an equal distribution of rDNA genes could undergo a random loss of the X chromosomes during male determination. On the basis of our hypothesis, therefore, NOR heteromorphism does not inhibit male determination (as previously reported by Blackman and Spence 1996), but affects the inheritance of the X chromosomes making it non random.

## Acknowledgements

This work is supported by the grant “F.A.R.” from the University of Modena and Reggio Emilia (M.M.).

## References

- Bizzaro D, Mandrioli M, Zanotti M, Giusti M, Manicardi GC (2000) Chromosome analysis and molecular characterization of highly repeated DNAs in the aphid *Acyrtosiphon pisum* (Aphididae, Hemiptera). *Genetica* 108: 197–202. doi: 10.1023/A:1004128314728
- Blackman RL (1980) Chromosome numbers in the Aphididae and their taxonomic significance. *Systematic Entomology* 5: 7–25. doi: 10.1111/j.1365-3113.1980.tb00393.x
- Blackman RL (1987) Reproduction, cytogenetics and development. In: Minsk AK, Harrewijn P (Eds) *Aphids: their biology, natural enemies and control*. Vol A. Elsevier, Amsterdam, 163–195.

- Blackman RL, Hales DF (1986) Behaviour of the X chromosomes during growth and maturation of parthenogenetic eggs of *Amphorophora tuberculata* (Homoptera, Aphididae) in relation to sex determination. *Chromosoma* 94: 59–64. doi: 10.1007/BF00293530
- Blackman RL, Spence JM (1996) Ribosomal DNA is frequently concentrated on only one X chromosome in permanently apomictic aphids, but this does not inhibit male determination. *Chromosome Research* 4: 314–320. doi: 10.1007/BF02263684
- Crema R (1979) Egg viability and sex determination in *Megoura viciae* (Homoptera, Aphididae). *Entomologia Experimentalis & Applicata* 26: 152–156. doi: 10.1111/j.1570-7458.1979.tb02910.x
- Crema R (1981) Lethal genes in a parthenogenetic strain of *Megoura viciae*. *Bollettino Zoologico* 48: 37–89.
- Donlon TA, Magenis RE (1983) Methyl green is a substitute for distamycin A in the formation of distamycin A/DAPI C-bands. *Human Genetics* 65: 144–146. doi: 10.1007/BF00286651
- Frantz A, Plantagenest M, Bonhomme J, Prunier N, Simon JC (2005) Strong biases in the transmission of sex chromosomes in the aphid *Rhopalosiphum padi*. *Genetical Research* 85: 111–117. doi: 10.1017/S0016672305007482
- Frydrychová R, Grossmann P, Trubac P, Vítková M, Marec F (2004) Phylogenetic distribution of TTAGG telomeric repeats in insects. *Genome* 47: 163–178. doi: 10.1139/g03-100
- Giannini S, Bizzaro D, Mandrioli M, Manicardi GC (2003) Cytogenetic analysis on the holocentric chromosomes of the cabbage aphid *Brevicoryne brassicae*. *Caryologia* 56: 143–147.
- Hales DF, Mitler TE (1983) Precocene causes male determination in the aphid *Myzus persicae*. *Journal of Insect Physiology* 29: 819–823. doi: 10.1016/0022-1910(83)90147-6
- Hales D (1989) The chromosomes of *Scoutedenia lutea* (Homoptera, Aphididae, Greenidinae) with an account of meiosis in the male. *Chromosoma* 98: 295–300. doi: 10.1007/BF00327316
- Hochstrasser M, Mathog D, Saumweber H, Sedat JW (1986) Spatial organization of chromosomes in the salivary gland nuclei of *Drosophila melanogaster*. *Journal of Cell Biology* 102: 112–115. doi: 10.1083/jcb.102.1.112
- Howell WM, Black DA (1980) Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: a 1 step-method. *Experientia* 36: 1014–1015. doi: 10.1007/BF01953855
- Ijdo JW, Wells RA, Baldini A, Reeders ST (1991) Improved telomere detection using a telomere repeat probe (TTAGGG)<sub>n</sub> generated by PCR. *Nucleic Acids Research* 19: 4780. doi: 10.1093/nar/19.17.4780
- Juan C, Pons J, Petitpierre E (1993) Localization of tandemly repeated DNA sequences in beetle chromosomes by fluorescent *in situ* hybridization. *Chromosome Research* 1: 167–174. doi: 10.1007/BF00710770
- Khuda-Buksh AR, Datta S (1981) A preliminary note on the mitotic chromosome aberrations in embryos of whole-body X-irradiated parthenogenetic viviparous *Aphis gossypii* (Homoptera, Aphididae). *Chromosome Information Service* 30: 4–5.
- Kuznetsova V, Gandrabur SI (1991) Nucleolar organizing regions in aphid chromosomes. *Cytologia* 33: 41–47.

- Kuznetsova VG, Maryńska-Nadachowska A (1993) C-heterochromatin and NORs in X-chromosomes of aphid *Dysaphis anthrisci* Born. Karyosystematics of Invertebrate Animals 2: 16–18.
- Luderus ME, van Steensel B, Chong L, Cremers FF, de Lange T (1996) Structure, subnuclear distribution and nuclear matrix association of the mammalian telomeric complex. Journal of Cell Biology 135: 867–881. doi: 10.1083/jcb.135.4.867
- Lukhtanov VA, Kuznetsova VG (2010) What genes and chromosomes say about the origin and evolution of insects and other arthropods. Russian Journal of Genetics 46: 1115–1121. doi: 10.1134/S1022795410090279
- Mandrioli M (2002) Cytogenetic characterization of telomeres in the holocentric chromosomes of the lepidopteran *Mamestra brassicae*. Chromosome Research 9: 279–286. doi: 10.1023/A:1016515607278
- Mandrioli M, Azzoni P, Lombardo G, Manicardi GC (2011) Composition and epigenetic markers of heterochromatin in the aphid *Aphis nerii* (Hemiptera: Aphididae). Cytogenetics and Genome Research 133: 67–77. doi: 10.1159/000323510
- Mandrioli M, Bizzaro D, Gionghi D, Bassoli L, Manicardi GC, Bianchi U. (1999a) Molecular cytogenetic characterization of a highly repeated DNA sequence in the peach potato aphid *Myzus persicae*. Chromosoma 108: 436–442. doi: 10.1007/s004120050395
- Mandrioli M, Bizzaro D, Giusti M, Manicardi GC, Bianchi U (1999b) The role of rDNA genes in X chromosomes association in the aphid *Acyrtosiphon pisum*. Genome 42: 381–386.
- Manicardi GC, Bizzaro D, Galli E, Bianchi U (1996) Heterochromatin heterogeneity in the holocentric X chromatin of *Megoura viciae* (Homoptera, Aphididae). Genome 39: 465–470. doi: 10.1139/g96-059
- Manicardi GC, Gautam DC, Bizzaro D, Guicciardi E, Bonvicini Pagliai AM, Bianchi U (1991) Chromosome banding in Aphids: G, C, *AluI*, and *HaeIII* banding patterns in *Megoura viciae* (Homoptera, Aphididae). Genome 34: 661–665. doi: 10.1139/g91-101
- Manicardi GC, Mandrioli M, Bizzaro D, Bianchi U (1998) Silver staining as a new banding technique to identify aphid chromosomes. Chromosome Research 6: 55–57. doi: 10.1023/A:1009270525511
- Manicardi GC, Mandrioli M, Bizzaro D, Bianchi U (2002) Cytogenetic and molecular analysis of heterochromatic areas in the holocentric chromosomes of different aphid species. In: Sobti RG, Obe G, Athwal RS (Eds) Chromosome Structure and Function, 47–56.
- Maryńska-Nadachowska A, Warchalowska-Sliwa E, Kuznetsova VG (1992) Karyotype of *Psyllina* (Homoptera). I. New data and check-list. Folia biologica 40: 15–25.
- Maryńska-Nadachowska A, Grozeva S (2001) Chromosome banding in *Cacopsylla mali* (Schmidberger) and *C. sorbi* (Linnaeus) (Homoptera, Psyllidae) with polymorphic sex chromosomes. Folia biologica 49: 157–161.
- Monti V, Giusti M, Bizzaro D, Manicardi GC, Mandrioli M (2011) Presence of a functional (TTAGG)<sub>n</sub> telomere-telomerase system in aphids. Chromosome Research, on line first.
- Nguyen P, Sahara K, Yoshido A, Marec F (2010) Evolutionary dynamics of rDNA clusters on chromosomes of moths and butterflies (Lepidoptera). Genetica 138: 343–354. doi: 10.1007/s10709-009-9424-5

- Novotná J, Havelka J, Stary P, Koutecky P, Vitková M (2011) Karyotype analysis of the Russian aphid *Diuraphis noxia* (Hemiptera: Aphididae) reveals a large X chromosome with rRNA and histone gene families. *Genetica* 139: 281–289. doi: 10.1007/s10709-011-9546-4
- Orlando E (1974) Sex determination in *Megoura viciae* Bukton (Homoptera, Aphididae). *Monitore Zoologico Italiano* 8: 61–70.
- Palladino F, Laroche T, Gilson A, Axelrode A, Pillus L, Gasser SM (1993) SIR3 and SIR4 proteins are required for the positioning and integrity of yeast telomeres. *Cell* 75: 543–555. doi: 10.1016/0092-8674(93)90388-7
- Postiglioni A, Brum-Zorrilla N (1988) Non-relationship between nucleolus and sex chromosome XY in *Chelymormpha variabilis* Boheman (Coleoptera: Chrysomelidae). *Genetica* 77: 137–141. doi: 10.1007/BF00057764
- Pryde FE, Gorham HC, Louis EJ (1997) Chromosome ends: all the same under their caps. *Current Opinion in Genetics & Development* 7: 822–828. doi: 10.1016/S0959-437X(97)80046-9
- Rakauskas R (1998) What is the (aphid) species? In: Nieto Nafria JM, AGF Dixon (Eds) *Aphids in natural and managed ecosystems*. Universidad de Leon (Secretariado de Publicaciones), Lein, 451–455.
- Schrader F (1940) Touch and go pairing in chromosomes. *Proceedings of the National Academy of Science USA* 26: 634–636. doi: 10.1073/pnas.26.11.634
- Schweizer D (1976) Reverse fluorescent chromosome banding with chromomycin and DAPI. *Chromosoma* 58: 307–324. doi: 10.1007/BF00292840
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Experimental Cell Research* 75: 304–306. doi: 10.1016/0014-4827(72)90558-7
- Wilson ACC, Sunnucks P, Hales D (1997) Random loss of X chromosome at male determination in an aphid, *Sitobion* near *fragariae*, detected using an X-linked polymorphic microsatellite marker. *Genetical Research* 69: 233–236. doi: 10.1017/S0016672397002747
- Wrench DL, Ketheley JB, Norton RA (1994) Cytogenetic of holokinetic chromosomes and inverted meiosis: keys to the evolutionary success of mites with generalization on eukaryotes. In: Houck MA (Ed) *Mites. Ecological and evolutionary analysis of life-history patterns*. Chapman & Hall, New York, London, 282–343.

Copyright of Comparative Cytogenetics is the property of Pensoft Publishers and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.