

Promoter methylation and downregulated expression of the *TBX15* gene in ovarian carcinoma

GAIA GOZZI^{1,2*}, SONIA T. CHELBI^{2,3*}, PAOLA MANNI¹, LOREDANA ALBERTI², SERGIO FONDA¹, SARA SAPONARO^{1,2}, LUCA FABBIANI⁴, FRANCESCO RIVASI⁴, JEAN BENHATTAR^{2,5} and LORENA LOSI¹

¹Pathological Anatomy Unit, Department of Life Sciences, University of Modena and Reggio Emilia, I-41124 Modena, Italy;

²University Institute of Pathology, Lausanne University Hospital, 1011 Lausanne; ³Department of Biochemistry,

University of Lausanne, 1066 Epalinges, Switzerland; ⁴Department of Diagnostic Medicine, Clinic and Public Health,

University of Modena and Reggio Emilia, I-41124 Modena, Italy; ⁵Biopath Lab,

Medical Biology and Pathology, 1006 Lausanne, Switzerland

Received April 23, 2015; Accepted June 29, 2016

DOI: 10.3892/ol.2016.5019

Abstract. *TBX15* is a gene involved in the development of mesodermal derivatives. As the ovaries and the female reproductive system are of mesodermal origin, the aim of the present study was to determine the methylation status of the *TBX15* gene promoter and the expression levels of *TBX15* in ovarian carcinoma, which is the most lethal and aggressive type of gynecological tumor, in order to determine the role of *TBX15* in the pathogenesis of ovarian carcinoma. This alteration could be used to predict tumor development, progression, recurrence and therapeutic effects. The study was conducted on 80 epithelial ovarian carcinoma and 17 control cases (normal ovarian and tubal tissues). *TBX15* promoter methylation was first determined by pyrosequencing following bisulfite modification, then by cloning and sequencing, in order to obtain information about the epigenetic haplotype. Immunohistochemical analysis was performed to evaluate the correlation between the methylation and protein expression levels. Data revealed a statistically significant increase of the *TBX15* promoter region methylation in 82% of the tumor samples and in various histological subtypes. Immunohistochemistry showed an inverse correlation between methylation levels and the expression of the *TBX15* protein. Furthermore, numerous tumor samples displayed varying degrees of intra-tumor heterogeneity. Thus, the present study determined that ovarian carcinoma typically expresses low levels of *TBX15* protein, predominantly due to an epigenetic mechanism. This

may have a role in the pathogenesis of ovarian carcinoma independent of the histological subtype.

Introduction

Ovarian cancer is the fourth most common type of cancer in the female genital tract, but has the highest mortality rate of all gynecological tumors. Ovarian carcinoma accounts for >90% of cases of ovarian cancer and may arise from relatively pluripotent cells of the coelomic epithelium or 'modified mesothelium' that covers the ovary (1). The coelomic epithelium is the mesodermal-derived epithelial lining of the intraembryonic coelom that invaginates to give rise to the müllerian ducts, the primordia for the epithelia of the fallopian tube, the endometrium and the endocervix (1). There is now compelling evidence that certain carcinomas can also originate from the fallopian tubes and the endometriosis (1,2). Ovarian carcinoma are a heterogeneous group of neoplasms that exhibit a wide range of tumor morphologies and clinical manifestations (2). According to the World Health Organization classification, the most common tumor subtypes are serous, endometrioid and mucinous, and are characterized by their morphological resemblance to various mucosal tissues of the female reproductive tract (3). Cytogenetic and molecular analysis indicates that multiple genetic and epigenetic changes are involved in the pathogenesis of ovarian carcinoma (2,4); however, it remains unclear how these alterations lead to the development of ovarian cancer. Better understanding of the molecular mechanisms responsible for the different types of ovarian cancer may improve diagnosis and treatment according to the biological characteristics of the cancer.

T-box genes constitute a family encoding transcription factors characterized by a highly conserved DNA-binding region, the T-box. These genes have a role in regulating the proliferation and differentiation of tissue-specific stem and progenitor cells during organogenesis (5,6). Previous studies have indicated that altered T-box gene expression may be associated with human cancer (7-10). For example, *TBX15*, a member of the T-box family, appears to be involved in mesodermal differentiation (11,12), putatively through its interaction with co-repressors of the Groucho family (13)

Correspondence to: Professor Lorena Losi, Pathological Anatomy Unit, Department of Life Sciences, University of Modena and Reggio Emilia, 71 Via del Pozzo, I-41124 Modena, Italy
E-mail: lorena.losi@unimore.it

*Contributed equally

Key words: *TBX15*, ovarian carcinoma, DNA methylation, immunohistochemistry, epigenetics, pyrosequencing

and, therefore, modulation of the Notch, Wingless/Wnt and decapentaplegic/bone morphogenetic protein/transforming growth factor- β signaling pathways (14). Furthermore, *TBX15* hypermethylation was observed in prostate carcinoma and may be correlated to a negative prognosis (15,16).

As stated, the *TBX15* gene is involved in mesodermal differentiation. Furthermore, the ovary and the female reproductive system are of mesodermal origin. Therefore, the present study analyzed the *TBX15* gene, in particular the epigenetic alteration of the *TBX15* gene promoter, in ovarian carcinoma in order to investigate its role in the pathogenesis of this neoplasia. This alteration could be used to predict tumor development, progression, recurrence and therapeutic effects. First, the percentage of DNA methylation at CpG sites was determined by bisulfite-pyrosequencing, as well as by cloning and sequencing, to refine the methylation pattern at the level of individual DNA molecules. Second, immunohistochemistry was performed to determine the correlation between DNA methylation and expression of the *TBX15* protein. Differences between the subtypes of ovarian carcinoma, and a possible correlation to pathological parameters and prognosis were also investigated.

Materials and methods

Patients and tissues. The present study was conducted on tissues obtained from ovaries removed surgically from women with ovarian carcinoma who were not subjected to preoperative chemotherapy. For control cases, 17 normal ovarian and tubal tissues were obtained from women treated surgically for benign gynecological conditions with no evidence of cancer (median age, 58 years). The 80 tumor samples included 29 serous, 23 mucinous and 28 endometrioid carcinomas, the three most frequent subtypes of carcinoma. All tissues were obtained via total abdominal hysterectomy and bilateral salpingo-oophorectomy performed between January 1995 and April 2010 at Pathological Anatomy Unit of the University of Modena and Reggio Emilia (Modena, Italy). For cases of mucinous carcinoma, a careful evaluation of the clinical history, in search of a primary mucinous carcinoma at another site, was performed, and these types of patients were excluded. Furthermore, cases with immunohistochemical positivity for CDX2, CEA or CK20, or negativity for CA125 were excluded from the study. The clinical and pathological features of patients are summarized in Table I. The stage was established according to the International Federation of Gynaecology and Obstetrics criteria (17). The pathological grade was specified according to the Silverberg grading system (18) for all tumors excluding serous tumors. For serous tumors, low and high grade were determined according to the proposed 2-tier grading system (19). Recurrences, metastasis and survival were established by the investigation of clinical files and pathological reports of patients. The study was approved by ethics committee of The University of Modena and Reggio Emilia and written informed consent was obtained from all patients.

Tissues were formalin-fixed and paraffin-embedded, sectioned at 3-4 μ m, stained with hematoxylin and eosin, and reviewed by two pathologists (Professor Lorena Losi and Professor Francesco Rivasi) in order to confirm the histopathological diagnosis.

Microdissection, DNA extraction and bisulfite conversion. Deparaffinized and rinsed sections (OTTIX and alcohol; Diapath S.P.A, Bergamo, Italy) were stained in 0.1% toluidine blue for 30 sec, washed, air-dried and manually microdissected using a scalpel blade under a microscopic by a pathologist (Professor Lorena Losi), in order to enrich the samples to at least 75% tumor cells. DNA extraction was performed using a Maxwell 16 FFPE Tissue LEV DNA Purification kit (Promega Corporation, Madison, WI, USA), according to the manufacturer's instructions and 100 ng genomic DNA was subjected to bisulfite conversion using the EpiTect Bisulfite kit (Qiagen, Hilden, Germany), according to the manufacturer's protocol.

Methylation analysis by pyrosequencing. Bisulfite-modified genomic DNA was PCR-amplified using specific primers covering two consecutive regions of the human *TBX15* promoter (2.8 kb upstream of the transcription start site of NM_152380.2), as previously described (20). The primers were as follows: Forward, 5'-ATGGGATAGTATAATTGATTTGGAATTT-3' and reverse, 5'-AAAAACCTTTCACCCCA TAA-3' for the upstream region (Amp1); and forward, 5'-GGTATTGGGGTAAGAGGAGA-3' and reverse, 5'-ACCACACAA AACTCCCTTTAT-3' for the downstream region (Amp2). Both reverse primers were biotinylated at the 5'-end. PCR was performed using 3 μ l of DNA, MgSO₄ at a final concentration of 3 mM and each primer at a final concentration of 0.25 μ M using 0.1 U of Platinum Taq DNA Polymerase (Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA). PCR was performed for a total of 41 cycles at an annealing temperature of 60°C. The primers of the reference gene GAPDH were: Forward, 5'-AAGGTGAAGGTCCGAGTCAAC-3' and reverse, 5'-GAGTTAAAAGCAGCCCTGGTG-3'.

The DNA methylation status of the two regions was assessed by pyrosequencing using the automated system PyroMark Q24 (Qiagen). Sequencing primers were designed using the PyroMark Assay Design SW software (version 2.0; Qiagen), as follows: Upstream, 5'-GAGGGAGTGGATTTT-3' and downstream, 5'-GGAAGTTTAGATTTTATATTTGTGA-3'. The results were analyzed using Pyromark Q24 software (version 2.0.6; Qiagen) to estimate the percentage of methylation at each CpG position for Amp1 (n=1,..,9) and Amp2 (n=1,..,10). Samples were categorized as follows: <40%, low level of methylation; 40-70%, medium level of methylation; and >70% very high level of methylation.

Methylation analysis by cloning and sequencing. The PCR products (Amp1 and Amp2) were ligated into a pGEM-T Vector (Promega Corporation), according to the manufacturer's protocol. The ligation products were transformed into chemically competent JM109 bacteria (Promega Corporation), according to the manufacturer's protocol. In total, 100 μ l mixture was plated on agar plates containing ampicillin, X-gal and isopropyl thio- β -D-galactoside for blue-white screening. The plates were incubated at 37°C overnight. Finally, DNA was extracted from 15-20 bacterial colonies, and subjected to PCR amplification of Amp1 and Amp2 followed by analysis by pyrosequencing. Cloning-sequencing data was then analyzed using BiQ Analyzer software (<http://biq-analyzer.bioinf.mpi-inf.mpg.de/>) to generate corresponding lollipop methylation diagrams (21).

Table I. Clinical and pathological features of ovarian carcinoma.

| Neoplasia | Patients, n | Median age, years | Histological grade | | | | | Stage | | | |
|------------------------|-------------|-------------------|--------------------|------|----|----|-----|-------|----|-----|----|
| | | | Low | High | I | II | III | I | II | III | IV |
| Serous carcinoma | 29 | 57 | 5 | 24 | - | - | - | 6 | 5 | 17 | 1 |
| Mucinous carcinoma | 23 | 59 | - | - | 13 | 4 | 6 | 19 | 1 | 3 | 0 |
| Endometrioid carcinoma | 28 | 61 | - | - | 14 | 8 | 6 | 22 | 3 | 3 | 0 |

Immunohistochemical analysis. Immunohistochemical analysis of *TBX15* was conducted on all 80 formalin-fixed, paraffin-embedded neoplastic tissue samples and all control samples. Immunohistochemistry was performed on a representative tumor sample in which, in addition to neoplasia, there was a marginal region of normal ovarian tissue, which was sectioned at 4 μ m. The normal controls are referred to tissues of different patients without ovarian carcinomas. Immunohistochemistry was performed using a *TBX15* rabbit anti-human polyclonal (C-terminus) primary antibody (catalog no., LS-C81114-100; dilution, 1:100; LifeSpan Biosciences, Inc., Seattle, WA, USA) with a BenchMark XT automated staining system (Ventana, Strasbourg, France), using 3,3'-diaminobenzidine as the chromogen. The secondary antibody was included in the Ultraview kit (catalog no., 05269806001; Roche Diagnostics Spa, Milan, Italy), which was used according to the manufacturer's protocol. At the end of reaction, slides were counterstained with hematoxylin. Primary antibody incubation was 37°C for 1 h.

Statistical analysis. Explorative and inferential data analysis was performed using R program version 3.2.2 (www.R-project.org/) with the following added packages: Dplyr, reshape2, data.table, extraGrid and ggplot2. One-way analysis of variance (ANOVA) followed by Tukey's honest significant difference (HSD) multiple comparisons of means were applied to analyze the difference in mean methylation between histotypes in all CpG_n sites, for Amp1 (n=1,...,9) and Amp2 (n=1,...,10) regions. Immunohistochemistry results were analyzed by two methods: i) Results were primarily analyzed by linear regression between mean methylation of pathological histotypes over all CpG sites and corresponding immunohistochemistry for Amp1, Amp2, and the mean of Amp1 and Amp2; ii) results were also analyzed by the χ^2 test to verify possible associations between pathological features (grade and stage), patient status (overall and disease-free survival) and the methylation values of pathological histotypes (split in two groups by thresholds of $\leq 40\%$ and $> 40\%$ methylation). Data was analyzed from three independent experiments. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Quantitative analysis of *TBX15* promoter methylation at multiple CpG sites by pyrosequencing. Methylation levels of 19 CpG sites within the 5'-region of the *TBX15* gene were analyzed by bisulfite pyrosequencing in 80 ovarian tumor

samples and 17 control samples. In normal ovarian tissues, a mean methylation level range of 10-30% was observed in all the 17 samples at all 19 CpG sites analyzed; this is considered to be a low level of methylation (Fig. 1A and B). For both *TBX15* promoter regions investigated (Amp1 and Amp2), a significant increase in mean methylation (range, 30-80%) was observed in the different groups of ovarian cancer compared with controls (N) at all the 19 CpG sites, as shown from Tukey's HSD post-hoc comparisons of means (Fig. 2A and B). For the upstream region of the promoter (Amp1), the endometrioid carcinoma samples showed a higher mean percentage of methylation than the serous and mucinous carcinoma samples for all the analyzed CpG positions; in particular, with the methylation values for CpGs 1, 2, 3, 4, 5, 6, 7, 9 of the serous histotype (E-S, y-axis; Fig. 2A) and CpGs 1, 4 and 5 of the mucinous histotype reached statistical significance compared with endometrioid carcinoma (E-M, y-axis; $P < 0.05$; Fig. 2A). By contrast, no significant difference was noted when comparing serous and mucinous cases. Similarly, the ANOVA and Tukey's HSD analysis revealed differences in the methylation profile of the downstream region (Amp2) between the three tumor groups. Again, endometrioid carcinomas showed the highest mean percentage of methylation. When considering the CpG positions individually, mucinous tumors presented significantly lower methylation levels at CpG sites 5 and 7 compared with serous carcinomas, whereas for endometrioid methylation was significantly increased at CpG position 1 compared with both mucinous and serous carcinomas (mucinous, $P = 0.008$; serous, $P = 0.040$; Fig. 2B). Considering all the 19 CpG sites (9 and 10 CpG sites for the Amp1 and Amp2, respectively), the mean \pm standard deviation methylation was $19.21 \pm 2.95\%$ for the control group, while the means were 50.56 ± 16.28 , 50.48 ± 15.08 and $63.27 \pm 15.56\%$ for serous, mucinous and endometrioid subtypes, respectively (Fig. 3). Unlike control samples, tumor samples exhibited a wider distribution of *TBX15* promoter methylation percentages, ranging from cases showing a methylation percentage similar to control samples to cases with methylation reaching almost 100%. Notably, when calculating the median of the methylation percentage for each group, the values were close to the mean of each group (49.7, 50.9 and 63.5% for serous, mucinous and endometrioid subtypes, respectively), demonstrating that the distribution of the observed methylation percentages was considerably homogenous within each group (Fig. 3).

***TBX15* promoter methylation patterns of individual clones.** To assess the methylation profile of the *TBX15* promoter at the single molecule level, a cloning-sequencing procedure





Figure 1. Methylation percentage of each CpG site within the (A) Amp1 and (B) Amp2 regions of the *TBX15* promoter gene. For both *TBX15* promoter regions investigated (Amp1 and Amp2), a meaningful increase in methylation was observed in the different groups of ovarian carcinoma (S, M and E) compared with the controls (N) at all 19 CpG sites, 9 for Amp1 and 10 for Amp2. Box plots indicate the first and third quartiles, median, and upper and lower quartiles (whiskers). Outliers are indicated by empty circles and mean values are indicated by black circles. Data are also presented as mean \pm SE. SE, standard error; N, normal; S, serous; M, mucinous; E, endometrioid.

was performed on the Amp1 and Amp2 PCR products. The control cases, with a mean percentage of methylation of ~20%, exhibited methylation in only a small number of CpG sites in the Amp1 and Amp2 regions. Indeed, the mean methylation of the Amp1 and Amp2 regions was 15, 20 and 17% for N5, N6 and N17, respectively, (Fig. 4).

For tumor samples, both regions (Amp1 and Amp2) were found to be either fully (or almost fully) methylated or contained only a small number of methylated CpG sites, as in normal tissues. Therefore, at the single molecule level, the *TBX15* promoter exhibited either low or high methylation in ovarian carcinoma samples. When tumor samples were analyzed more in detail, a marked association was observed

between the mean methylation level measured by pyrosequencing and the percentage of molecules with either fully (or almost fully) methylated Amp1 and Amp2 regions. This observation is illustrated in Fig. 4, with three serous tumor samples that exhibited different mean levels of *TBX15* methylation. Similar results were obtained with mucinous and endometrioid carcinomas (data not shown). For the S2 tumor sample with strong hypermethylation (80 and 91% for Amp1 and Amp2, respectively), as expected, almost all the molecules were methylated in almost all CpG sites. The S1 sample, which had an intermediate level of methylation (76 and 53% for Amp1 and Amp2, respectively), had a mixture of molecules, some with sporadic methylation, such as in the controls, and

A



B




Figure 2. Results of one way analysis of variance followed by Tukey's HSD multiple comparisons test. Difference of % methylation values (x-axis) and corresponding 95% CI (horizontal bars) between normal and histotype group means are shown for all the possible histotype couples (y-axis) in CpG sites (y-axis) for (A) Amp1 and (B) Amp2. The vertical dashed line indicates zero difference. CI crossing dashed line shows non significant difference for the corresponding couple of patients type. All histotype couples with a non-zero crossing CI present a significant % methylation difference with $P < 0.05$ in a range of $< 0.005 - 0.031$ for AMP1 and $< 0.005 - 0.046$ for AMP2. The shorter the distance of CI from zero, the larger the P-value. HSD, honest significant difference; N, normal; S, serous; M, mucinous; E, endometrioid; CI, confidence interval.

some molecules that were heavily or even fully methylated (Fig. 4). For the S3 serous sample with mild methylation (37 and 13% for Amp1 and Amp2, respectively), the situation was similar to that in the controls except that the number of methylated CpG sites was marginally higher. According to all

the results obtained, hypermethylation of the *TBX15* promoter should be considered when mean methylation of at least 40% is observed in the Amp1 or Amp2 region. In the current study, pyrosequencing revealed that hypomethylation occurred in all 17 control ovary samples and in 15 ovarian carcinoma samples.




Figure 3. Mean DNA methylation of the Amp1 and Amp2 regions of the *TBX15* promoter gene. The mean DNA methylation of each case is represented by a triangle. The tumor samples (S, M and E) showed a wider distribution of *TBX15* promoter methylation percentages, ranging from methylation percentage similar to that in control samples, to cases with methylation close to 100%. Box plots indicate the first and third quartiles, median, and upper and lower quartiles (whiskers). Outliers are indicated by empty circles and mean values are indicated by black circles. Data are also presented as mean \pm standard deviation. S, serous; M, mucinous; E, endometrioid.




Figure 4. Lollipop methylation diagram showing DNA methylation of the Amp1 and Amp2 regions of the *TBX15* promoter gene at the molecular level. Amplification, bisulfite-pyrosequencing and cloning of the Amp1 and Amp2 regions was performed on genomic DNA extracted from control ovarian (N5, N6, N17) and carcinoma (S1, S2, S3) samples. A small number of methylated CpG sites (black circles) were observed in control tissues, whereas hypermethylation was observed in a large proportion of S1 and S2 samples. N, normal; S, serous; M, mucinous; E, endometrioid.

TBX15 hypermethylation was observed in 65/80 (82%) cases of ovarian cancer, in particular in 83, 74 and 88% of serous, mucinous and endometrioid carcinomas, respectively.

Immunohistochemical expression of *TBX15*. To determine if *TBX15* promoter methylation could be a mechanism for gene

silencing, *TBX15* protein expression was analyzed by immunohistochemistry. In the control cases, *TBX15* protein was present in the nucleus of the ovarian mesothelium, stromal cells and vascular smooth muscle cells (Fig. 5A). Immunohistochemical positivity was also present in the epithelium of the fallopian tubes.




Figure 5. Immunohistochemical expression of the TBX15 protein in normal and neoplastic ovarian tissue samples. (A) Normal ovarian tissue with positive staining of TBX15 in the nucleus of the ovarian mesothelium and in cortical stromal cells (magnification, x10). Endometrioid carcinomas with different levels of *TBX15* promoter methylation: (B) 20%, (C) 72% and (D) 53% (magnification, x20). 3,3'-diaminobenzidine and hematoxylin staining.




Figure 6. Linear regression between promoter methylation and IHC of TBX15 for Amp1, Amp2 and the mean of Amp1 and Amp2. The linear regression performed in ovarian tumor samples shows an inverse correlation between the level of *TBX15* promoter methylation and the expression of the TBX15 protein, as determined by IHC ($r=-0.7$ in Amp1, $r=-0.8$ in Amp2, $r=-0.83$ for the mean between the two regions; $P<0.01$ for both line coefficients, in all three lines). S, serous; M, mucinous; E, endometrioid; IHC, immunohistochemistry.

For all the neoplastic samples with a low level of methylation (<40%), the presence of TBX15 expression was observed in almost all the tumor cells (Fig. 5B). By contrast, an absence of TBX15 protein expression was observed in almost all the tumor cells in samples with a very high level of methylation (>70%; Fig. 5C). Notably, in samples with a medium level of methylation (40-70%), only some focal regions exhibited TBX15 protein expression (Fig. 5D). Therefore, in ovarian tumor samples, an inverse correlation was observed between

the level of *TBX15* promoter methylation and the expression of the TBX15 protein ($r=-0.7$ for Amp1; $r=-0.8$ for Amp2; $r=-0.83$ for the mean between the two regions; $P<0.01$; Fig. 6). The two regions, Amp1 and Amp2, present a comparable correlation coefficient, suggesting that both regions may be involved in the silencing of the gene.

Correlation of TBX15 methylation with clinical and pathological parameters. The possible association of methylation

results with pathological and clinical features, such as grade, stage, overall survival and disease-free survival, was investigated. When cases with >40% mean methylation were considered to be methylated, the χ^2 test showed that all considered parameters were independent of *TBX15* methylation status (data not shown).

Discussion

In the current study, the methylation status of the *TBX15* promoter gene and its protein expression were investigated in order to determine its role in the pathogenesis of ovarian carcinoma according to its implication in mesodermal tissues, such as the ovary and the other organs of the reproductive system. In normal ovarian tissues, the *TBX15* methylation level was low and varied between 10-30% at a single CpG site. Notably, hypermethylation of the *TBX15* promoter was observed in 82% of the total ovarian carcinoma samples, with similar percentages in various histological subtypes.

The analysis of *TBX15* expression by immunohistochemistry confirmed the correlation between promoter methylation and the loss of *TBX15* expression in ovarian carcinoma. Therefore, DNA methylation may be the primary mechanism responsible for *TBX15* gene silencing in ovarian carcinoma. Intratumor heterogeneity was noted by immunohistochemistry and was confirmed by methylation analysis in approximately half of the ovarian cancer cases. In the present study, the analyzed tumor samples were enriched to at least 75% tumor cells by manual microdissection. The mean methylation level of tumor samples ranged between 30 and 80%, which indicates that, for some of the carcinoma samples, only certain tumor cells harbored a hypermethylated *TBX15* promoter. Additional information regarding the methylation status was obtained by bisulfite sequencing after the cloning of PCR products, which revealed that, in all normal ovarian tissues and in ~15% of cancer tissues, CpG methylation was randomly distributed over the region and no fully methylated molecules were detected. By contrast, in neoplastic samples, the mean increase in methylation compared with the controls resulted in fully or almost fully methylated molecules. Overall, methylation analysis revealed heterogeneity at CpG sites and between individual molecules. The presence of hypomethylated and hypermethylated clones in various ovarian tumor samples may be associated with the presence of normal stromal cells, tumor cells at various stages of progression towards tumorigenesis, or populations of tumor cells subject to clonal evolution. This observation is in accordance with genetic or epigenetic intratumoral heterogeneity, as previously observed in colorectal (22), breast (23) and ovarian (24) cancer.

To date, the role of *TBX15* in neoplastic diseases remains unknown; however, a previous study identified aberrant methylation of *TBX15* in prostate carcinoma, highlighting its possible role as a prognostic marker. Indeed, *TBX15* methylation was found to be associated with the pathological stage and Gleason score in prostate carcinoma (16), suggesting that *TBX15* may be useful as a methylation marker in pre- and post-treatment clinical evaluations. In the present study, no correlation was identified between *TBX15* methylation and the stage or grade of ovarian carcinoma. Furthermore, the overall and disease-free survival rates did not appear to be

associated with the methylation and immunohistochemical data. This lack of correlation may be due to the presence of intratumor heterogeneity in ovarian tumors, which confounds the validation of single biomarkers, or, more likely, because this epigenetic alteration occurs in the majority of ovarian carcinoma cases. It would be useful for future studies to determine whether hypermethylation of *TBX15* could be used to predict tumor development, progression and therapeutic effects in ovarian carcinoma.

A previous study observed that specific T-box genes promote epithelial-mesenchymal transition (EMT) in neoplastic cell lines and even in certain neoplasia. This process involves events that convert adherent epithelial cells into individual migratory cells that can invade the extracellular matrix; such cells may be critical for cancer progression and the development of metastasis (25). Furthermore, an association between cells undergoing EMT and cells with stem cell-like properties in cancer (cancer stem cells) has been noted (26,27). A member of the T-box family, Brachyury, which appears to be regulated by the β -catenin oncogene, can induce the expression of stemness markers, such as NANOG, CD133, CD166 and CD44, in a subpopulation of colorectal cancer cells that mimic invasive front mesenchymal-like cells (28). This transcription factor may be useful in predicting the outcome of chemotherapy and radiotherapy, as cancer stem cells are resistant to these treatments, and could become a target for specific therapies. A similar course may also occur in ovarian carcinoma and *TBX15* may have a role in this process.

In conclusion, the data obtained in the present study supports the hypothesis that epigenetic alteration of the *TBX15* gene is implicated in the pathogenesis of the majority of cases of ovarian carcinoma. The current data also suggests that *TBX15*, which is implicated in the development of mesoderm-derived organs, such as the female reproductive system, may have a role in the neoplastic ovarian process, independent of the histological subtype. Therefore, hypermethylation of *TBX15* may represent a potential biomarker for early detection, progression and response to treatments with a significant impact on reducing the mortality of this disease.

Acknowledgements

The authors thank Professor Melanie Cripps for the English revision of the manuscript. Financial support was partially provided by Banca Popolare dell'Emilia Romagna (Modena, Italy).

References

1. Auersperg N: Ovarian surface epithelium as a source of ovarian cancers: Unwarranted speculation or evidence-based hypothesis? *Gynecol Oncol* 130: 246-251, 2013.
2. Kurman RJ and Shih IeM: The origin and pathogenesis of epithelial ovarian cancer: A proposed unifying theory. *Am J Surg Pathol* 34: 433-443, 2010.
3. Kurman RJ, Carcangiu ML, Herrington CS and Young RH (eds): WHO Classification of Tumours of Female Reproductive Organs. 4th edition. IARC, Lyon, 2014.
4. Gloss BS and Samimi G: Epigenetic biomarkers in epithelial ovarian cancer. *Cancer Lett* 342: 257-263, 2014.
5. Showell C, Binder O and Conlon FL: T-box genes in early embryogenesis. *Dev Dyn* 229: 201-218, 2004.

6. Takashima Y and Suzuki A: Regulation of organogenesis and stem cell properties by T-box transcription factors. *Cell Mol Life Sci* 70: 3929-3945, 2013.
7. Dorfman DM, Hwang ES, Shahsafaei A and Glimcher LH: T-bet, a T-cell-associated transcription factor, is expressed in a subset of B-cell lymphoproliferative disorders. *Am J Clin Pathol* 122: 292-297, 2004.
8. Dorfman DM, Hwang ES, Shahsafaei A and Glimcher LH: T-bet, a T cell-associated transcription factor, is expressed in Hodgkin's lymphoma. *Hum Pathol* 36: 10-15, 2005.
9. Ito A, Asamoto M, Hokaiwado N, Takahashi S and Shirai T: Tbx3 expression is related to apoptosis and cell proliferation in rat bladder both hyperplastic epithelial cells and carcinoma cells. *Cancer Lett* 219: 105-112, 2005.
10. Rowley M, Grothey E and Couch FJ: The role of Tbx2 and Tbx3 in mammary development and tumorigenesis. *J Mammary Gland Biol Neoplasia* 9: 109-118, 2004.
11. Kispert A, Herrmann BG, Leptin M and Reuter R: Homologs of the mouse Brachyury gene are involved in the specification of posterior terminal structures in *Drosophila*, *Tribolium* and *Locusta*. *Genes Dev* 8: 2137-2150, 1994.
12. Vidricaire G, Jardine K and McBurney MW: Expression of the Brachyury gene during mesoderm development in differentiating embryonal carcinoma cell cultures. *Development* 120: 115-122, 1994.
13. Farin HF, Bussen M, Schmidt MK, Singh MK, Schuster-Gossler K and Kispert A: Transcriptional repression by the T-box proteins Tbx18 and Tbx15 depends on Groucho corepressors. *J Biol Chem* 282: 25748-25759, 2007.
14. Buscarlet M and Stifani S: The 'Marx' of Groucho on development and disease. *Trends Cell Biol* 17: 353-361, 2007.
15. Kron K, Pethe V, Briollais L, Sadikovic B, Ozcelik H, Sunderji A, Venkateswaran V, Pinthus J, Fleshner N, van der Kwast T and Bapat B: Discovery of novel hypermethylated genes in prostate cancer using genomic CpG island microarrays. *PLoS One* 4: e4830, 2009.
16. Kron K, Liu L, Trudel D, Pethe V, Trachtenberg J, Fleshner N, Bapat B and van der Kwast T: Correlation of ERG expression and DNA methylation biomarkers with adverse clinicopathologic features of prostate cancer. *Clin Cancer Res* 18: 2896-2904, 2012.
17. Prat J: FIGO committee on gynecologic oncology: Staging classification for cancer of the ovary, fallopian tube, and peritoneum. *Int J Gynaecol Obstet* 124: 1-5, 2014.
18. Silverberg SG: Histopathologic grading of ovarian carcinoma: A review and proposal. *Int J Gynecol Pathol* 19: 7-15, 2000.
19. Malpica A, Deavers MT, Lu K, Bodurka DC, Atkinson EN, Gershenson DM and Silva EG: Grading ovarian serous carcinoma using a two-tier system. *Am J Surg Pathol* 28: 496-504, 2004.
20. Chelbi ST, Doridot L, Mondon F, Dussour C, Rebourcet R, Busato F, Gascoin-Lachambre G, Barbaux S, Rigourd V, Mignot TM, *et al*: Combination of promoter hypomethylation and PDX1 overexpression leads to TBX15 decrease in vascular IUGR placentas. *Epigenetics* 6: 247-255, 2011.
21. Bock C, Reither S, Mikeska T, Paulsen M, Walter J and Lengauer T: BiQ analyzer: Visualization and quality control for DNA methylation data from bisulfite sequencing. *Bioinformatics* 21: 4067-4068, 2005.
22. Losi L, Baisse B, Bouzourene H and Benhattar J: Evolution of intratumoral genetic heterogeneity during colorectal cancer progression. *Carcinogenesis* 26: 916-922, 2005.
23. Moelans CB, de Groot JS, Pan X, van der Wall E and van Diest PJ: Clonal intratumor heterogeneity of promoter hypermethylation in breast cancer by MS-MLPA. *Mod Pathol* 27: 869-874, 2014.
24. Woloszynska-Read A, Mhawech-Fauceglia P, Yu J, Odunsi K and Karpf AR: Intertumor and intratumor NY-ESO-1 expression heterogeneity is associated with promoter-specific and global DNA methylation status in ovarian cancer. *Clin Cancer Res* 14: 3283-3290, 2008.
25. Imajyo I, Sugiura T, Kobayashi Y, Shimoda M, Ishii K, Akimoto N, Yoshihama N, Kobayashi I and Mori Y: T-box transcription factor Brachyury expression is correlated with epithelial-mesenchymal transition and lymph node metastasis in oral squamous cell carcinoma. *Int J Oncol* 41: 1985-1995, 2012.
26. Mani SA, Guo W, Liao MJ, Eaton EN, Ayyanan A, Zhou AY, Brooks M, Reinhard F, Zhang CC, Shipitsin M, *et al*: The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell* 133: 704-715, 2008.
27. Blick T, Hugo H, Widodo E, Waltham M, Pinto C, Mani SA, Weinberg RA, Neve RM, Lenburg ME and Thompson EW: Epithelial mesenchymal transition traits in human breast cancer cell lines parallel the CD44(hi)/CD24 (lo/-) stem cell phenotype in human breast cancer. *J Mammary Gland Biol Neoplasia* 15: 235-252, 2010.
28. Sarkar D, Shields B, Davies ML, Müller J and Wakeman JA: BRACHYURY confers cancer stem cell characteristics on colorectal cancer cells. *Int J Cancer* 130: 328-337, 2012.