

This is a pre print version of the following article:

“Quarantine during COVID-19 outbreak: changes in Diet and physical activity increase the risk of cardiovascular disease” / Mattioli, Anna Vittoria; Sciomer, Susanna; Cocchi, Camilla; Maffei, Silvia; Gallina, Sabina. - In: NMCD. NUTRITION METABOLISM AND CARDIOVASCULAR DISEASES. - ISSN 0939-4753. - 30:9(2020), pp. 1409-1417. [10.1016/j.numecd.2020.05.020]

Terms of use:


The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

07/08/2024 13:43

Journal Pre-proof

“Quarantine during COVID-19 outbreak: changes in Diet and physical activity increase the risk of cardiovascular disease”

Anna Vittoria Mattioli, Susanna Sciomer, Camilla Cocchi, Silvia Maffei, Sabina Gallina



PII: S0939-4753(20)30213-1

DOI: <https://doi.org/10.1016/j.numecd.2020.05.020>

Reference: NUMECD 2318

To appear in: *Nutrition, Metabolism and Cardiovascular Diseases*

Received Date: 22 April 2020

Revised Date: 19 May 2020

Accepted Date: 20 May 2020

Please cite this article as: Mattioli AV, Sciomer S, Cocchi C, Maffei S, Gallina S, “Quarantine during COVID-19 outbreak: changes in Diet and physical activity increase the risk of cardiovascular disease”, *Nutrition, Metabolism and Cardiovascular Diseases*, <https://doi.org/10.1016/j.numecd.2020.05.020>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 The Italian Diabetes Society, the Italian Society for the Study of Atherosclerosis, the Italian Society of Human Nutrition and the Department of Clinical Medicine and Surgery, Federico II University. Published by Elsevier B.V. All rights reserved.

**“Quarantine during COVID-19 outbreak: changes in Diet and physical activity
increase the risk of cardiovascular disease”.**

Anna Vittoria Mattioli^a, Susanna Sciomer^b, Camilla Cocchi^c, Silvia Maffei^d, Sabina Gallina^e

Affiliation:

a) Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Italy

b) Department of Cardiovascular, Respiratory Nephrological, Anesthesiological and Geriatric Sciences, Sapienza University, Italy

c) Istituto Nazionale per le Ricerche Cardiovascolari, U.O. Modena, Italy

d) Cardiovascular and Gynaecological Endocrinology Unit, Fondazione G. Monasterio CNR-Regione Toscana, Pisa, Italy;

e)) Department of Neuroscience, Imaging and Clinical Sciences, University of Chieti-Pescara, Chieti, Italy

Running title: quarantine and cardiovascular risk

Address for correspondence:

Prof Anna Vittoria Mattioli,
Surgical, Medical and Dental Department of Morphological Sciences related to Transplant,
Oncology and Regenerative Medicine
University of Modena and Reggio Emilia,
Via del pozzo, 71 41100 Modena (Italy)
E-mail: annavittoria.mattioli@unimore.it

Word count

Main text: 4040 words

Abstract: 233 words

Tables: 2

References: 86

Conflicting Interests: None Declared

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors

Author contribution. AVM, SS, SM,SG conceived of the idea at the basis of the article, AVM, CC, SG developed the different part of the manuscript, AVM, SS, CC, SM, SG performed the final supervision. All authors contributed to and approved the final manuscript.

Orcid:

Anna Vittoria Mattioli: 0000-0003-1487-9530

Abstract

Aims. CoV-19/SARS-CoV-2 is a highly pathogenic virus that is causing a global pandemic with a high number of deaths and infected people. To contain the diffusion of infection, several Governments have enforced restrictions on outdoor activities or even collective quarantine on the population. The present commentary briefly analyzes the effects of quarantine on lifestyle, including nutrition and physical activity and the impact of new technologies in dealing with this situation.

Data Synthesis. Quarantine is associated with stress and depression leading to unhealthy diet and reduced physical activity. A diet poor in fruit and vegetables is frequent during isolation, with a consequent low intake of antioxidants and vitamins. However, vitamins have recently been identified as a principal weapon in the fight against the Cov-19 virus. Some reports suggest that Vitamin D could exert a protective effect on such infection. During quarantine, strategies to further increase home-based physical activity and to encourage adherence to a healthy diet should be implemented. The WHO has just released guidance for people in self-quarantine, those without any symptoms or diagnosis of acute respiratory illness, which provides practical advice on how to stay active and reduce sedentary behaviour while at home.

Conclusions. Quarantine carries some long-term effects on cardiovascular disease, mainly related to unhealthy lifestyle and anxiety. Following quarantine, a global action supporting healthy Diet and physical activity is mandatory to encourage people to return to a good lifestyle routine.

Key-words: quarantine; COVID-19; stress; lifestyle; gender; physical activity; vitamin D

76 The emergence of novel coronavirus, officially known as Severe Acute Respiratory
77 Syndrome-Coronavirus-2 (SARS-CoV-2), has presented an important challenge for
78 healthcare systems across the world. Rapid transmission is due to high infectivity, capacity
79 to be transmitted even during asymptomatic phase and relatively low virulence. [1]
80 On March 12, 2020 the WHO defined the COVID-19 infection as pandemic. [2]
81 Quarantine and isolation are two measures that can prevent, or minimize, the impact of
82 infectious disease outbreaks. In public health practice, “quarantine” refers to the
83 separation of persons (or communities) who have been exposed to an infectious disease.
84 “Isolation,” in contrast, applies to the separation of persons who are known to be infected.
85 [3]
86 The word quarantine comes from the Italian quarantena, meaning "forty days", used in the
87 14th-15th-century Venetian language and designating the period that all ships were
88 required to be isolated before passengers and crew could go ashore during the Black Death
89 plague epidemic [4].
90 In the Modern Era there are several examples of Government imposed quarantine or travel
91 bans. i.e. at least 18 U.S. states quarantined people returning from West Africa during the
92 2014 Ebola outbreak. [3] Recently, the Italian Government among others has enforced
93 quarantine on the population to contain the diffusion of the COVID-19 virus. The previous
94 experience of the SARS outbreak showed the efficacy of timely quarantine and isolation
95 measures. [5,6] Singapore and Hong Kong, both of which had severe acute respiratory
96 syndrome (SARS) epidemics in 2002–03, provide hope and many lessons to other
97 countries. Today, quarantine and social distancing reduced transmission of COVID-19
98 infection by about 60% in China [7]. However, a further peak will likely occur when
99 restrictive measures to avoid major economic impact are relaxed. [7] Nevertheless,
100 quarantine is an unpleasant experience: with loss of freedom, uncertainty over disease
101 status, and boredom it can affect the health status of subjects. Therefore, the potential

benefits of mandatory mass quarantine need to be weighed carefully against the possible long-term negative effects on health, i.e. cardiovascular risk burden, and mental disease. [8,9]

Quarantine induces anxiety and stress. Western health care systems have been built around the concept of patient-centered care, but such a pandemic requires a change of perspective toward a concept of community-centered care.[10] This increases anxiety in people who have concerns about their own health. Survey studies on subjects who had been quarantined reported a high prevalence of psychological distress and disorder symptoms. These included emotional disturbance, depression, stress, low mood, irritability, insomnia, post-traumatic stress [11-14].

During outbreaks of infection people are likely to experience fear of falling sick or dying themselves, feelings of helplessness, and stigma [13-14]. During one influenza outbreak, around 10% to 30% of the general public were very or fairly worried about the possibility of contracting the virus. With the closure of schools and business, negative emotions experienced by individuals increased. [15-16]

The present commentary briefly analyzes the effects of quarantine on lifestyle, including nutrition and physical activity and the impact of new technologies in dealing with this situation.

Effects of stress and anxiety on lifestyle during quarantine

Sudden catastrophic events i.e. earthquake are associated with an increase in sudden cardiac deaths, and an overall increase in death from atherosclerotic and ischemic heart disease. [17]

The reasons for this increase in cardiac events after catastrophic events include an acute increase in sympathetic nervous activity and catecholamines.

127 In the case of quarantine people suffered from a chronic increase of sympathetic nervous
128 activity leading to a chronic negative effect on heart and vessels. Social isolation and
129 loneliness are associated with a very high risk of mortality and the development of major
130 chronic disease. [18] Specifically, social isolation is associated with an increased risk of
131 mortality in patients with CVD. [19]

132 The acute stress response of an integrated cascade of physiological reactions has been well
133 described [20,21] On the contrary, less is known about how chronic stress responses
134 convert to pathological changes over time, contributing to the development and
135 progression of cardiovascular disease. [21]

136 The great majority of data come from studies evaluating personal stress (i.e. the death of a
137 close person) or categories of persons (i.e. workers during major economic recession).

138 Little is known about outcome after a period of quarantine and outbreaks, even though all
139 agree this corresponds to a period of high stress. Stress-related changes in the
140 sympathetic–parasympathetic balance and the neuroendocrine dysregulation involving the
141 hypothalamus–pituitary– adrenal responses might adversely affect the cardiovascular
142 system both by accelerating the atherosclerotic process and by precipitating the occurrence
143 of a cardiovascular event. Repeated or chronic exposure to stress facilitates the progression
144 of atherosclerosis. Adrenaline and noradrenaline increase the heart rate and decrease
145 heart rate variability, optimize blood flow to muscle tissues, and elevate core body
146 temperature. The sympathetic nervous system has direct cardiostimulatory effects
147 (chronotropy and inotropy via β_1 -adrenergic receptors) and pressor effects (via
148 α_1 -adrenergic receptors) and also affects metabolism (promotes insulin resistance and
149 lipolysis) and the immune system [22,23,24]

150 Moreover, unhealthy behaviors and economic crisis contribute to increased health risk,
151 with socially isolated and lonely individuals having less favorable lifestyles. [25,26]

152 An analysis of UK Biobank indicated that health behavior accounted for more than 30% of
153 the excess risk of mortality attributed to social isolation and loneliness over a 6.5 year
154 follow-up period [27]

155 The main consequence of stress related to quarantine is a change in lifestyle and
156 nutritional habits. (Table 1) Changes in nutritional habits can be due to: 1. reduced
157 availability of goods, 2. limited access to food caused by restricted store opening hours, 3.
158 switch to unhealthy food.

159 Analysis performed following the Ebola outbreak quarantine showed that having
160 inadequate food and water supplies was a source of frustration and emotional stress. The
161 WHO and Liberia's Ministry of Health found that during the outbreak many families in
162 quarantine did not receive food supplies. In some cases even access to water and sanitation
163 facilities could not be guaranteed for people in quarantine. These conditions do not seem
164 compatible with quarantine or isolation being forms of "easy rescue". In addition, Ebola-
165 exposed families who were subject to quarantine measures in West Africa during the 2014-
166 15 outbreak suffered significantly from stigmatization and loss of livelihoods, as well as
167 possible increased exposure. [6,28,29]

168 Due to anxiety surrounding future food shortage, people purchase packaged and long-life
169 food rather than fresh food. Foods with long shelf-life shorten the life line owing to their
170 salt, sugar or trans-fat content. [30]

171 This leads to an unhealthy diet poor in antioxidants food i.e. fresh fruit and vegetables that
172 would increase oxidative stress and inflammation. [31,33]

173 The modern diet era began with the primate evolution when primate biology lost its ability
174 to synthesize vitamin C consequent to a healthy consumption of fruits [33]

175 Over the last 150 years, refining, hydrogenation, salting and frying were adopted as
176 methods of prolonging the shelf life of food substances. Refined sugar with added fat
177 increased the energy density of the food materials by 6 times. [34]. These dietary changes

178 generated a major mismatch between the genetic structure of man and what he could
179 metabolize [35]. Over the last 40 years dietary guidelines and population strategies
180 together with good public education and simultaneous efforts directed at other lifestyle
181 issues, such as promoting healthy Diet (i.e. the Mediterranean Diet or Dash Diet),
182 improving physical activity and reducing tobacco consumption, have notably reduced
183 cardiovascular mortality. [32,35-37]

184 During quarantine our diet takes a step back from being a healthy diet rich in fresh food to
185 one containing foods with a long shelf-life.

186 In addition, for many people, the typical response to chronic stressful situations is not to
187 avoid food but possibly to seek out and consume energy-dense foods [38-41].

188 Anxiety, depression, uneasiness, and anger are emotions that commonly accompany
189 chronic stress [42]. The responses to acute or chronic stress also include a number of
190 modifying behaviors such as alcohol consumption, smoking, and eating. [40,42]

191 When individuals respond to stress by eating more, anecdotal evidence suggests the foods
192 selected are typically high in sugar and fat. [42] This desire to consume a specific kind of
193 food is defined as “food craving”. Food craving is a multidimensional experience as it
194 includes cognitive (e.g., thinking about food), emotional (e.g., desire to eat or changes in
195 mood), behavioral (e.g., seeking and consuming food), and physiological (e.g., salivation)
196 aspect [43, 44]

197 The craving for carbohydrates encourages the production of serotonin which has a positive
198 effect on mood. This effect on mood is proportional to glycemic index of foods. [43]

199 Muscogiuri and colleagues recently pointed out that quarantine-related stress translates
200 into sleep disturbances that further worsen stress and increase food craving. [43]

201 They reported on importance of consuming foods containing or promoting the synthesis of
202 serotonin and melatonin at dinner. (e.g. roots, leaves, fruits and seeds such as almonds,

203 bananas, cherries and oats) In addition, milk and dairy products are the main sources of
204 sleep-inducing tryptophan amino acid, a precursor of serotonin and melatonin.
205 Tryptophan is involved in the regulation of satiety and calorie intake via serotonin which
206 mainly reduces carbohydrate and fat intake and inhibits neuropeptide Y. [43] Dairy
207 products such as yogurt could also increase the activity of natural killer cells and reduce
208 the risk of respiratory infections suggesting a potential function protection against SARS-
209 CoV-2 disease. This unhealthy nutritional habit may contribute to excess energy intakes
210 and weight gain, increasing the risk of developing obesity. [45,46] Obesity is associated
211 with chronic inflammation, and it is a strong risk factor of heart disease, diabetes, and lung
212 disease that have been demonstrated to increase the risk for more serious complications of
213 CoVID-19 [47]

214

215 Torres and coworkers identified that people cope with stress by eating and drinking in an
216 attempt to feel better (“stress-related eating”). These stress-driven eaters and drinkers
217 were more likely to consume unhealthy foods such as snacks, hamburgers, soda cola, and
218 chocolate regularly and to drink wine and spirits more frequently. In addition the lack of
219 emotional support from friends and relatives was predictive of stress driven eating and
220 drinking behaviors [42,48]. During quarantine, stress-driven eaters would easily switch
221 from a healthy diet to an unhealthy one. In addition, the increase in macronutrient intake
222 could also be accompanied by micronutrient deficiency. During quarantine the Diet is poor
223 in fresh fruit and vegetables. Fruits and vegetables are rich in micronutrients and
224 antioxidants. [49] Micronutrients include vitamins that act as antioxidants by reducing the
225 inflammatory response and improving the immune response. Anti-oxidants increase the
226 number of T-cell subsets, enhance lymphocyte response to mitogen, increased interleukin-
227 2 production, and potentiated natural killer cell activity. This would affect cardiovascular
228 risk mainly in high-risk patients. Therefore, during this period it is essential to follow a

229 balanced nutritional diet that includes a high amount of antioxidants and vitamins. In
 230 some cases it may be helpful to take vitamin and minerals supplements. However, despite
 231 epidemiologic studies have reported that high dietary intake of foods rich in vitamin E,
 232 vitamin C, and β - carotene, have been inversely associated with the incidence of CAD,
 233 different results are obtained with vitamin and antioxidant supplementations. [50]
 234 Several reasons have been proposed to explain this lack of results obtained with
 235 antioxidants supplements: define the optimal dosage, use the appropriate vitamin isomer,
 236 interference or competition between vitamins, inter-individual variation to response to
 237 vitamin. [50]
 238 The very recent literature focused on the potential beneficial effect of Vit. D
 239 supplementation on patients with COVID-19.

240 **Vitamin D and COVID-19**

241 From recent journal literature, it is known that COVID-19 infection is associated with the
 242 increased production of pro-inflammatory cytokines, and C-reactive protein [51,52].
 243 Antioxidants and vitamins exert protective effects against infection and inflammation.
 244 Some research suggested the efficacy of vitamin supplements to prevent COVID 19
 245 infections. A letter from Panarese and coworkers suggested that vitamin D deficiency may
 246 also contribute to airway/gastrointestinal infectious illnesses. [53] Vitamin D has immune-
 247 modulatory properties, that include down-regulation of pro-inflammatory cytokines. [54-
 248 56] It is possible that the protective effect of vitamin D against COVID-19 is related to
 249 suppression of cytokine response and reduced severity/risk for ARDS. In addition, a meta-
 250 analysis shows that regular oral vitamin D2/D3 intake (in doses up to 2000 IU/d without
 251 additional bolus), is safe and protective against acute respiratory tract infection, especially
 252 in subjects with vitamin D deficiency. [56] The elderly display a very high prevalence of
 253 hypovitaminosis D, especially during the winter. [53] It therefore seems plausible that

254 Vitamin D prophylaxis may contribute to reducing the severity of illness caused by SARS-
255 CoV-2, particularly in settings where hypovitaminosis D is frequent, including people
256 currently living in Northern countries. [54] Regardless of age, ethnicity, and latitude;
257 recent data showed that 40% of the Europeans are vitamin D deficient (25(OH)D levels
258 <50 nmol/L), and 13% are severely deficient (25(OH)D <30 nmol/L). It is known that
259 severe vitamin D deficiency dramatically increases the risk of mortality, infections, and
260 many other diseases. [57]

261 In a retrospective multicentre study of 212 cases with laboratory-confirmed infection of
262 SARS-CoV-2, an increase in serum 25(OH)D level in the body could either improve clinical
263 outcomes or mitigate worst (severe to critical) outcomes, while a decrease in serum
264 25(OH)D level in the body could worsen clinical outcomes of COVID-2019 patients.
265 [58]

266 In this patient's population, those with 25OHD >75 nmol/l had lower symptoms than
267 those with lower 25OHD.

268 A recent review suggested using vitamin D loading doses of 200,000–300,000 IU in
269 50,000-IU capsules to reduce the risk and severity of COVID-19. Grant and coworkers
270 suggested that higher vitamin D doses and 25OHD concentrations would be better for
271 prevention and probably, reduce the risk of influenza and COVID-19 incidence and death.
272 [59] Several reviews consider the ways in which vitamin D reduces the risk of viral
273 infections. [56,59,60] Vitamin D has many mechanisms by which it reduces the risk of
274 microbial infection and death. Vitamin D helps maintain tight junctions, gap junctions,
275 and adherens junctions (e.g., by E-cadherin) fighting against the action of viruses to
276 disturb junction integrity, increasing infection by the virus and other microorganisms. [61]

277 Vitamin D also enhances cellular immunity by reducing the cytokine storm induced by the
278 innate immune system. The latter generates both pro-inflammatory and anti-inflammatory
279 cytokines in response to viral and bacterial infections, as observed in COVID-19 patients.

[51] Vitamin D can reduce the production of pro-inflammatory Th1 cytokines, such as tumor necrosis factor α and interferon γ . [62] Food rich in vitamin D are: salmon, sardines, cod liver oil, canned tuna, egg yolks, mushrooms, and meat. [63] Fish are poor represented in the diet typically consumed by the elderly people living in the North of Italy, leading to a reduced intake of Vitamin D. Magnesium supplementation is recommended when taking vitamin D supplements. Magnesium helps activate vitamin D, which in turn helps regulate calcium and phosphate homeostasis to influence the growth and maintenance of bones. All the enzymes that metabolize vitamin D seem to require magnesium, which acts as a cofactor in the enzymatic reactions in the liver and kidneys. [64]

Practical tips to avoid vitamin D deficiency during quarantine are: taking short walks, increasing sun exposure; consume foods rich in vitamin D; and/or drug supplementation

Preliminary observations support the hypothesis that vitamin D supplementation can reduce the risk of influenza and COVID-19. However, the incidence and death should be investigated in trials to determine the appropriate doses, serum 25(OH)D concentrations,

Effects of stress and anxiety on physical activity

The “GLOBAL ACTION PLAN ON PHYSICAL ACTIVITY 2018-2030” published by WHO indicated physical activity as mandatory for prevention of non-communicable disease. [65] Regular physical activity is associated with reduction in cardiovascular risk. [65-67]

The “2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease” recommended that “Adults should engage in at least 150 minutes per week of accumulated moderate-intensity or 75 minutes per week of vigorous-intensity aerobic physical activity (or an equivalent combination of moderate and vigorous activity) to reduce ASCVD risk” (Class I LOE B-NR) or “for adults unable to meet the minimum physical activity recommendations, engaging in some moderate- or vigorous-intensity physical activity,

304 even if less than this recommended amount, can be beneficial to reduce ASCVD risk”
305 (Class IIa LOE B-NR).[68] Prolonged TV viewing time is associated with an increased risk
306 of type 2 diabetes mellitus, CVD, and all-cause mortality [68]. Prolonged sedentary time is
307 independently associated with deleterious health outcomes, regardless of levels of physical
308 activity [67,68] During quarantine Governments prohibited the great majority of outdoor
309 exercise and social activities (e.g. going to the gym) resulting in a reduction of physical
310 activity. Inflammation is an underlying pathophysiological process in chronic diseases,
311 such as obesity, T2 diabetes, and cardiovascular disease. Regular physical activity reduces
312 inflammation, and oxidative stress and helps to maintain normal weight and to reduce
313 visceral fat accumulation. [65,66] Limited physical activity and inability to take a regular
314 walk out of one’s home as a consequence of collective quarantine, may be associated with
315 several metabolic effects that would increase the cardiovascular risk. It is also established
316 that many beneficial metabolic and cardiovascular adjustment in response to physical
317 exercise can be lost in just two weeks of inactivity, impairing aerobic capacity and/or
318 increasing blood pressure. Sudden exercise cessation has been associated with rapid onset
319 of insulin resistance in muscle tissue and decreased muscle glucose utilization that
320 worsened muscle performance. [69]

321 During quarantine, staying active and maintaining a physical exercise routine will be
322 essential for mental and physical health. [70,71,72] The WHO has just released guidance
323 intended for people in self-quarantine without any symptoms or diagnosis of acute
324 respiratory illness, containing practical advice on how to stay active and reduce sedentary
325 behavior while at home. They suggest following on-line exercise classes, and using video-
326 or app-guided aerobics training at home. Table 2 summarizes the WHO indication “Stay
327 physically active during self-quarantine”. [73] Little is known about the effects of home-
328 based physical activity on chronic disease. [67,74,75] The Web could prove useful, since
329 today there are many workout videos available that can assist people in exercising on their

own. However, trying to execute all steps (body pose alignments) in a workout accurately, in order to avoid long-term muscle and joint injuries is risk related. [74,75]

In addition, almost all modern Smartphone provide step-count and application for nutrition. Today, it is estimated that more than 5 billion people have mobile devices, and over half of these connections are smartphones. Therefore, there are higher potential apps to be developed for lifestyle behavior change in the community. Many people to control their diet and maintain their personal ideal weight use mobile applications related to nutrition. The increasing number of health and nutrition applications available on Google Play and the Apple App Store proves the awareness of community regarding adopting a healthy lifestyle. App programs may be more effective when social support is advocated and could be a useful instrument in order to reduce the negative impact of quarantine on lifestyle. [76,77]

Telemedicine is an important tool for patient home monitoring and is very useful for nutritional, motor, psychiatric and psychological support in quarantined patients.

Sex differences.

Sex differences are present in several diseases. Preliminary data suggest that SARS-CO-2 affected more men than women. One reason seems to be that men have higher rates of chronic disease, a risk factor for COVID-19. However, women are less physically active than men. Severe COVID-19 patients had significantly higher levels of Th1 cytokines (IL-6 and TNF- α) and higher incidence rate of ARDS, compared with non-severe cases, suggesting a more intense inflammatory response. [78] It is known that after ischemic damage, male gender has a significantly higher level of IL-6. [79] Similarly, exercise-induced muscle damage triggers inflammatory responses that result in elevations of inflammation markers such as C-reactive protein (CRP) and some inflammatory

interleukins (IL-1, IL-6, tumor necrosis factor (TNF- α)). [80] On contrary, regular physical activity seems to reduce IL-6 and C-reactive protein. Usually women are less physically active than men and the gap increase after menopause [66]. During and after menopause, most women tend to reduce their physical activity levels and together with the reduction in basal metabolic rate, these lead to a loss of skeletal muscle mass as well as loss of bone mineral density. [66,81] During perimenopause period, fat deposition shifts to favor the visceral depot that, in addition to the decreased protective effect of estrogen, contributes to endothelial dysfunction, inflammation, altered fatty acid metabolism, insulin resistance, all markers of CVDs. [66,67]. There is a direct relationship between time spent sitting, physical activity and the CVD risk in postmenopausal women, independent of leisure-time physical activity. Prolonged sitting time determines many detrimental adaptations, such as increased energy intake and reduction of skeletal muscle lipoprotein lipase activity that might explain its effect on cardiovascular risk factors [66] In addition, the described food craving has a higher prevalence in women than in men. [43,44]

Quarantine induced a reduction of physical activity and an increase in the sitting time leading to an increase of cardiovascular risk in women. It is mandatory to promote a call of action for women mostly after 40 years, for a healthy life style during quarantine to reduce, as consequence, cardiovascular risks.

Impact of quarantine on people with obesity

During the quarantine, patients suffering from obesity experienced immense stress which made them more vulnerable to over-eating and sedentary lifestyle, thus predisposing them to further weight gain. Moreover, the incoming economic downturn will also lead to more consumption of unhealthy foods as it is cheaper. This will lead to further increase in obesity prevalence especially in weaker sections of the society.[Recently, growing scientific evidence has reported an important role of obesity in Covid-19's prognosis [82]

Obesity is a leading risk factor of cardiovascular disease, diabetes, renal disease, and has a detrimental effect on lung function. A pro-inflammatory state coupled with malnutrition may lead to impaired immune response in patients suffering from obesity and increased susceptibility to all influenza viruses including COVID-19 [83,84] Obesity has some detrimental effects on immune system that include alterations in leukocyte development, phenotypes and activity. In particular, obesity has been shown to impair memory CD8+ T cell responses to influenza virus infection, resulting in increased mortality, viral titers in lung, and worsened lung pathology. [82] Two possible mechanisms have been hypothesized to explain the relationship between obesity and more serious disease in patients with Covid-19: lung function disorders and endothelial dysfunction. Both mechanisms have a common pathway: the increase in pro-inflammatory cytokines and the "cytokine storm". SARS-Cov-2 binds the trans-membrane angiotensin converting enzyme-II receptor inducing an acute endothelial injury. The storm of pro-inflammatory cytokines increases the expression of adhesion molecules which promotes endothelial activation and cascade activation of coagulation, leading to a worsening of the microcirculation system and tissue perfusion.[85] The "cytokine storm" can lead to acute respiratory distress syndrome or even multiple organ failure. It represents a phenomenon of immune hyper activation very similar to Cytokine-release syndrome (CRS).

Subjects presented with obesity have a pro-inflammatory status and the exposure to COVID-19 could further exacerbate inflammation exposing them to higher levels of circulating inflammatory molecules [82]

Obese subjects must be carefully informed about the risk of an unhealthy lifestyle during the quarantine due to the increasing risk of disease. These subjects need careful monitoring of their health and strong psychological support to reduce stress and anxiety.

404 Weight stigma is not uncommon in social media outlets such as Twitter, Facebook, and
405 Instagram and the recent COVID-19 quarantine has sparked social media posts that refer
406 to "quarantine-15". Social media posts that stigmatize obesity and mock or diminish real
407 struggles with weight and eating can be particularly harmful to individuals with obesity
408 who are actively trying to manage their weight. [86]

413 **Conclusions**

414 We need to be prepared to confront the likely increase in cardiovascular risk burden
415 following the pandemic. During quarantine we must promote healthy diet and physical
416 activity at home. After quarantine we need to re-evaluate the cardiovascular risk in
417 patients, assessing biometrical and metabolic parameters. Patients also need to be
418 evaluated by psychologist to early identify the persistence of anxiety and stress and/or
419 evolution to a post-traumatic syndrome. Global action supporting healthy Diet and
420 physical activity is mandatory to encourage people to return to a good lifestyle. This action
421 needs to be stronger for individuals of a low socio economic level that will suffer to a higher
422 degree from the inevitable restrictions and economic crisis following a vast and prolonged
423 quarantine.

424 **Acknowledgements**

425 We would like to thank Mrs Janet Ann Carter for reviewing the paper.
426 We thank everyone who is working to resolve the SARS - CoV - 2 pandemic.

427

References

1. Li G, Hu R, Gu X. A close-up on COVID-19 and cardiovascular diseases, *Nutrition, Metabolism & Cardiovascular Diseases*, <https://doi.org/10.1016/j.numecd.2020.04.00>
2. WHO announces COVID-19 outbreak a pandemic. 2020 at <http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>
3. Parmet WE, Sinha MS. Covid-19 — The Law and Limits of Quarantine. *N Engl J Med* 2020 DOI: 10.1056/NEJMp2004211
4. Mackowiak PA. The origin of quarantine. *Clinical Infectious Diseases* 2002; 35:1071–2.
5. WHO 2003, Update 58 – First Consultation on SARS epidemiology, travel recommendations for Hebei Province (China), situation in Singapore, http://www.who.int/csr/sars/archive/2003_05_17/en/
6. Giubilini A, Douglas T, Maslen H, Savulescu J. Quarantine, isolation and the duty of easy rescue in public health. *Dev World Bioeth.* 2018 Jun;18(2):182-189. doi: 10.1111/dewb.12165. Epub 2017 Sep 18.
7. Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *Lancet.* 2020;395(6):931-4
8. Mattioli AV, Ballerini Puviani M. Lifestyle at time of COVID-19, how could quarantine affect cardiovascular risk. *Am J lifestyle* 2020 in press. <https://doi.org/10.1177/1559827620918808>
9. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence *Lancet* 2020; 395: 912–20. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)

- 454 10. Nacoti M, Ciocca A, Giupponi A, Brambillasca P, Lussana F, Pisano M et al. At the
455 Epicenter of the Covid-19 Pandemic and Humanitarian Crises in Italy: Changing
456 Perspectives on Preparation and Mitigation *N Engl J Medicine* 2020 March 21, 2020
457 DOI: 10.1056/CAT.20.0080
- 458 11. DiGiovanni C, Conley J, Chiu D, Zaborski J. Factors influencing compliance with
459 quarantine in Toronto during the 2003 SARS outbreak. *Biosecur Bioterror* 2004; 2:
460 265–72.
- 461 12. Hawryluck L, Gold WL, Robinson S, Pogorski S, Galea S, Styra R. SARS control and
462 psychological effects of quarantine, Toronto, Canada. *Emerg Infect Dis* 2004; 10:
463 1206–12.
- 464 13. Jeong H, Yim HW, Song Y-J, Ki M, Min JA, Cho J, et al. Mental health status of people
465 isolated due to Middle East respiratory syndrome. *Epidemiol Health* 2016; 38:
466 e2016048
- 467 14. Lee S, Chan LY, Chau AM, Kwok KP, Kleinman A. The experience of SARS-related
468 stigma at Amoy Gardens. *Soc Sci Med* 2005; 61: 2038–46.
- 469 15. Hall RCW, Chapman MJ. The 1995 Kikwit Ebola outbreak: Lessons hospitals and
470 physicians can apply to future viral epidemics. *Gen. Hosp. Psychiatry* 2008, 30, 446–
471 452.
- 472 16. Rubin GJ, Potts HWW, Michie S. The impact of communications about swine flu
473 (influenza A H1N1v) on public responses to the outbreak: Results from 36 national
474 telephone surveys in the UK. *Health Technol. Assess.* 2010, 14, 183–266.
- 475 17. Kloner RA, Leor J. Natural disaster plus wake-up time: A deadly combination of
476 triggers. *Am Heart J* 1999;137:779-81.
- 477 18. Brotman DJ, Golden SH, Wittstein IS. The cardiovascular toll of stress. *Lancet* 2007;
478 370, 1089–1100.

19. Steptoe A, Kivimäki M. Stress and cardiovascular disease. *Nat. Rev. Cardiol.* 2012; 9, 360–370.
20. Huang QH, Takaki A, Arimura A. Central noradrenergic system modulates plasma interleukin-6 production by peripheral interleukin-1. *Am. J. Physiol.* 1997; 273, R731-738
21. Goebel MU, Mills PJ, Irwin MR, Ziegler MG. Interleukin-6 and tumor necrosis factor- α production after acute psychological stress, exercise, and infused isoproterenol: differential effects and pathways. *Psychosom. Med.* 2000; 62, 591–598 (2000).
22. Steptoe A, Shankar A, Demakakos P, Wardle J. Social isolation, loneliness, and all-cause mortality in older men and women. *Proc Natl Acad Sci U S A.* 2013;110:5797–801.
23. Yu B, Steptoe A, Chen LJ, Chen YH, Lin CH, Ku PW. Social Isolation, Loneliness, and All-Cause Mortality in Patients With Cardiovascular Disease: A 10-Year Follow-up Study. *Psychosom Med.* 2020;82(2):208–214. doi:10.1097/PSY.0000000000000777
24. Cacioppo JT, Cacioppo S, Capitanio JP, Cole SW. The neuroendocrinology of social isolation. *Annu Rev Psychol.* 2015;66:733–67.
25. Grippo AJ, Lamb DG, Carter CS, Porges SW. Social isolation disrupts autonomic regulation of the heart and influences negative affective behaviors. *Biol Psychiatry.* 2007;62:1162–70
26. Loucks EB, Berkman LF, Gruenewald TL, Seeman TE. Relation of social integration to inflammatory marker concentrations in men and women 70 to 79 years. *Am J Cardiol.* 2006;97:1010–6.
27. Elovainio M, Hakulinen C, Pulkki-Råback L, Virtanen M, Josefsson K, Jokela M, et al. Contribution of risk factors to excess mortality in isolated and lonely individuals: an analysis of data from the UK biobank cohort study. *Lancet Public Health.* 2017;2:e260–6.

28. Kutalek R, Wang S, Fallah M, Wesseh CS, Gilbert J. Ebola interventions: listen to communities. *Lancet* 2015; 3:e131. doi: 10.1016/S2214-109X(15)70010-0
29. ACAPS 2016. WASH in Guinea, Liberia, and Sierra Leone. The Impact of Ebola, Available at <https://www.acaps.org/special-report/wash-guinea-liberia-and-sierra-leone-impact-ebola>,
30. Sivasankaran S. The cardio-protective diet. *Indian J Med Res.* 2010;132(5):608–616.
31. Mattioli AV, Ballerini Puviani M, Nasi M, Farinetti A. COVID-19 pandemic: the effects of quarantine on cardiovascular risk. *Eur J Clin Nutr.* 2020 May 5:1-4. doi: 10.1038/s41430-020-0646-z.
32. Mattioli AV, Coppi F, Migaldi M, Scicchitano P, Ciccone MM, Farinetti A. Relationship between Mediterranean diet and asymptomatic peripheral arterial disease in a population of pre-menopausal women. *Nutr Metab Cardiovasc Dis.* 2017; 27 (11) 985-990. doi: 10.1016/j.numecd.2017.09.011
33. Leupker RV. Decline in incident coronary heart disease. Why are the rates falling? *Circulation.* 2008;117:592–3.
34. Ford AS, Ajani UA, Croft JB, Critchley JA, Labarthe DR, Kottke TE. Explaining the decrease in US deaths from Coronary disease: 1980-2000. *N Engl J Med* 2007; 356:2388–98.
35. Benzie IFF. Evolution of dietary antioxidants. *Com Biochem Physiol A Physiol.* 2003;136:113–26.
36. Kopp W. High-insulinogenic nutrition - an etiologic factor for obesity and the metabolic syndrome. *Metabolism.* 2003;52:840–4.
37. O'Keefe JH, Jr, Cordain L. Cardiovascular disease resulting from a diet and lifestyle at odds with our Paleolithic genome: How to become a 21st century Hunter-gatherer. *Mayo Clin Proc.* 2004;79:101–8

38. Johnson RK, Appel LJ, Brands M, Howard BV, Lefevre M, Lustig RH, Sacks F, Steffen LM, Wylie-Rosett J. American Heart Association Nutrition Committee of the Council on Nutrition, Physical Activity, and Metabolism and the Council on Epidemiology and Prevention. Dietary sugars intake and cardiovascular health: a scientific statement from the American Heart Association. *Circulation*. 2009 Sep 15;120(11):1011-20.
39. Vreman RA, Goodell AJ, Rodriguez LA, Porco TC, Lustig RH, Kahn JG. Health and economic benefits of reducing sugar intake in the USA, including effects via non-alcoholic fatty liver disease: a microsimulation model. *BMJ Open*. 2017 Aug 3;7(8):e013543. d
40. Schiffman SS, Graham BG, Sattely-Miller EA, Peterson-Dancy M. Elevated and sustained desire for sweet taste in African-Americans: a potential factor in the development of obesity. *Nutrition* 2000;16: 886 –93.
41. Bracale R, Vaccaro CM Changes in food choice following restrictive measures due to Covid-19 *Nutrition, Metabolism & Cardiovascular Diseases*, 2020, in press
42. Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Nutrition*. 2007;23(11- 12):887-894. <https://doi.org/10.1016/j.nut.2007.08.008>
43. Muscogiuri G, Barrea L, Savastano S, Colao A. Nutritional recommendations for CoVID-19 quarantine [published online ahead of print, 2020 Apr 14]. *Eur J Clin Nutr*. 2020;1 - 2. doi:10.1038/s41430-020-0635-2
44. Rodríguez-Martín BC, Meule A. Food craving: new contributions on its assessment, moderators, and consequences. *Front Psychol*. 2015;6:21. Published 2015 Jan 22. doi:10.3389/fpsyg.2015.00021
45. Walcott-McQuigg JA. The relationship between stress and weight control behavior in African-American women. *J Natl Med Assoc* 1995;87:427–32

46. Heraclides AM, Chandola T, Witte DR, Brunner EJ. Work stress, obesity and the risk of type 2 diabetes: gender-specific bidirectional effect in the Whitehall II study. *Obesity (Silver Spring)*. 2012;20(2):428–433. doi:10.1038/oby.2011.95
47. Wu Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med*. 2020. <https://doi.org/10.1001/jamainternmed.2020.0994>.
48. Laitinen J, Ek E, Sovio U. Stress-related eating and drinking behavior and body mass index and predictors of this behavior. *Prev Med (Baltim)*. 2002;34(1):29-39. <https://doi.org/10.1006/pmed.2001.0948>
49. Mattioli AV, Coppi F, Migaldi M, Farinetti A. Fruit and vegetables in hypertensive women with asymptomatic peripheral arterial disease. *Clinical Nutrition ESPEN* 2018; 27: 110-112
50. Badimon L, Vilahur G, Padro T. Nutraceuticals and atherosclerosis: human trials. *Cardiovasc Ther*. 2010;28(4):202 - 215. doi:10.1111/j.1755-5922.2010.00189.x
51. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* 2020.
52. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020
53. Panarese A, Shahini E. Letter: Covid-19, and vitamin D [published online ahead of print, 2020 Apr 12]. *Aliment Pharmacol Ther*. 2020;10.1111/apt.15752. doi:10.1111/apt.15752
54. Panarese A, Pesce F, Porcelli P, Riezzo G Iacovazzi PA, Leone CM et al. Chronic functional constipation is strongly linked to vitamin D deficiency. *World J Gastroenterol*. 2019; 25: 1729- 1740.

- 580 55. Greiller CL, Martineau AR. Modulation of the immune response to respiratory viruses
 581 by vitamin D. *Nutrients*. 2015; 7: 4240- 4270.
- 582 56. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P et al.
 583 Vitamin D supplementation to prevent acute respiratory tract infections: systematic
 584 review and meta-analysis of individual participant data. *BMJ*. 2017; 356: i6583.
 585 doi:10.1136/bmj.i6583.
- 586 57. Kara M, Ekiz T, Ricci V, Kara Ö, Chang KV, Özçakar L. 'Scientific Strabismus' or Two
 587 Related Pandemics: COVID-19 & Vitamin D Deficiency [published online ahead of
 588 print, 2020 May 12]. *Br J Nutr*. 2020;1 - 20. doi:10.1017/S0007114520001749
- 589 58. Alipio, Mark, Vitamin D Supplementation Could Possibly Improve Clinical Outcomes
 590 of Patients Infected with Coronavirus-2019 (COVID-19) (April 9, 2020). Available at
 591 SSRN: <https://ssrn.com/abstract=3571484> or
 592 <http://dx.doi.org/10.2139/ssrn.3571484>
- 593 59. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL et al.
 594 Evidence that Vitamin D Supplementation Could Reduce Risk of Influenza and
 595 COVID-19 Infections and Deaths. *Nutrients* 2020, 12, 988.
- 596 60. Beard JA, Bearden A, Striker R. Vitamin D and the anti-viral state. *J. Clin. Virol*. 2011,
 597 50, 194–200.
- 598 61. Hewison M. An update on vitamin D and human immunity. *Clin. Endocrinol*. 2012,
 599 76, 315–325.
- 600 62. Schwalfenberg GK. A review of the critical role of vitamin D in the functioning of the
 601 immune system and the clinical implications of vitamin D deficiency. *Mol. Nutr. Food*
 602 *Res*. 2011, 55, 96–108
- 603 63. Crowe F, Steur M, Allen N, Appleby P, Travis R, Key T. Plasma concentrations of 25-
 604 hydroxyvitamin D in meat eaters, fish eaters, vegetarians and vegans: Results from

the EPIC–Oxford study. *Public Health Nutrition*, 2011; 14(2), 340–346.

doi:10.1017/S1368980010002454

64. Uwitonze AM, Razzaque MS. Role of Magnesium in Vitamin D Activation and Function. *J. Am. Osteopath Assoc.* 2018, 118, 181–189.

65. WHO Global action plan on physical activity 2018–2030: more active people for a healthier world. Geneva: World Health Organization; 2018. Licence: CC BY-NC-SA 3.0 IGO.

66. Mattioli AV, Sciomer S, Moscucci F, Maiello M, Cugusi L, Gallina S et al. Cardiovascular prevention in women: a narrative review from the Italian Society of Cardiology working groups on 'Cardiovascular Prevention, Hypertension and peripheral circulation' and on 'Women Disease'. *J Cardiovasc Med* 2019; 20:575–583. doi:10.2459/JCM.0000000000000831.

67. Nasi M, Patrizi G, Pizzi C, Landolfo M, Boriani G, Dei Cas A et al. The role of physical activity in individuals with cardiovascular risk factors: An opinion paper from Italian Society of Cardiology-Emilia Romagna-Marche and SIC-Sport. *J Cardiovasc Med* 2019;20:631–639. doi: 10.2459/JCM.0000000000000855

68. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;000:e□□□–e□□□. DOI: 10.1161/CIR.0000000000000678

69. Womack VY, De Chavez PJ, Albrecht SS, Durant N, Loucks EB, Puterman E et al. A Longitudinal Relationship between Depressive Symptoms and Development of Metabolic Syndrome: The Coronary Artery Risk Development in Young Adults Study. *Psychosom. Med.* 2016, 78, 867–873

70. Thompson PD, Franklin BA, Balady GJ, Blair SN, Corrado D, Estes NA 3rd, et al. Exercise and acute cardiovascular events placing the risks into perspective: A scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism and the Council on Clinical Cardiology. *Circulation* 2007; 115: 2358–2368.
71. Ricci F, Izzicupo P, Moscucci F, Sciomer S, Maffei S, Di Baldassarre A, Mattioli AV, Gallina S. Recommendations for physical inactivity and sedentary behavior during COVID-19. *Front. Public Health* | doi: 10.3389/fpubh.2020.00199
72. Mattioli AV, Nasi M, Cocchi C, Farinetti A. COVID 19 outbreak: impact of the quarantine-induced stress on cardiovascular disease risk burden. *Future Cardiology* 2020 Published online: 30 April 2020. Doi: 10.2217/fca-2020-0055
73. WHO guideline “Stay physically active during self-quarantine” <http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-coronavirus-2019-ncov-technical-guidance/stay-physically-active-during-self-quarantine>
74. Nagarkoti A, Teotia R, Mahale AK, Das PK. Realtime Indoor Workout Analysis Using Machine Learning & Computer Vision. *Conf Proc IEEE Eng Med Biol Soc.* 2019 Jul;2019:1440-1443. doi: 10.1109/EMBC.2019.8856547.
75. Taylor JK, Buchan IE van der Veer SN. Assessing life-space mobility for a more holistic view on wellbeing in geriatric research and clinical practice. *Aging Clinical and Experimental Research* (2019) 31:439–445 <https://doi.org/10.1007/s40520-018-0999-5>
76. Paramastri R, Pratama SA, Ho DKN, Purnamsari SD, Mohamed AZ, Galvin CJ et al. Use of mobile applications to improve nutrition behaviour: A systematic review [published online ahead of print, 2020 Mar 19]. *Comput Methods Programs Biomed.* 2020;192:1054-59. doi:10.1016/j.cmpb.2020.105459

77. DiFilippo KN, Huang WH, Andreade JE. The use of mobile apps to improve nutrition outcomes: a systematic literature review, *J. Telemed. Telecare* 2015; 21: 243–253
78. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan [published online ahead of print, 2020 Apr 12]. *J Allergy Clin Immunol.* 2020; S0091-6749 (20) 30495-4. doi:10.1016/j.jaci.2020.04.006
79. Mattioli AV, Bonetti L, Zennaro M, Bertoncelli P, Mattioli G. Acute myocardial infarction in young patients: Nutritional status and biochemical factors. *Int J Cardiol* 2005; 101,(2), 185-190
80. Fatouros, IG, Jamurtas AZ. Insights into the molecular etiology of exercise-induced inflammation: Opportunities for optimizing performance. *J. Inflamm. Res.* 2016, 9, 175–186
81. Mattioli AV, Coppi F, Migaldi M, Farinetti A. Physical activity in premenopausal women with asymptomatic peripheral arterial disease. *J Cardiovasc Med* 2018 Nov;19(11):677-680. doi: 10.2459/JCM.0000000000000714
82. Muscogiuri G, Pugliese G, Barrea L, Savastano S, Colao A. Obesity: The "Achilles heel" for COVID-19? [published online ahead of print, 2020 Apr 27]. *Metabolism.* 2020;108:154251. doi:10.1016/j.metabol.2020.154251]
83. Green WD, Beck MA. Obesity impairs the adaptive immune response to influenza virus. *Ann Am Thorac Soc.* 2017;14:S406–9.
84. Bhasker AG, Greve JW. Are Patients Suffering from Severe Obesity Getting a Raw Deal During COVID-19 Pandemic? [published online ahead of print, 2020 May 12]. *Obes Surg.* 2020;1-2. doi:10.1007/s11695-020-04677-z
85. Tibiriçá E, De Lorenzo A. Increased severity of COVID-19 in people with obesity: are we overlooking plausible biological mechanisms? [published online ahead of print,

681 2020 May 13]. Obesity (Silver Spring). 2020;10.1002/oby.22887.
682 doi:10.1002/oby.22887]
683 86.Pearl RL. Weight Stigma and the "Quarantine-15" [published online ahead of print,
684 2020 Apr 23]. Obesity (Silver Spring). 2020;10.1002/oby.22850.
685 doi:10.1002/oby.22850]

686

687

688

689

690

691

692

693

Table 2

WHO guide for quarantine: “Stay physically active during self-quarantine”

(This guidance is intended for people in self-quarantine without any symptoms or diagnosis of acute respiratory illness. It should not replace medical guidance in case of any health condition.) Modify from WHO guide, freely available at

<http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/novel-coronavirus-2019-ncov-technical-guidance/stay-physically-active-during-self-quarantine>

- **Take short active breaks during the day.** Short bouts of physical activity add up to the weekly recommendations. You may use the suggested exercises below as inspiration to be active every day. Dancing, playing with children, and performing domestic chores such as cleaning and gardening are other means to stay active at home.
- **Follow an online exercise class.** Take advantage of the wealth of online exercise classes. Many of these are free and can be found on YouTube. If you have no experience, be cautious.
- **Walk.** Even in small spaces, walking around or walking on the spot, can help you remain active. If you have a call, stand or walk around your home while you speak, instead of sitting down. If you can go outside to walk or exercise, be sure to maintain at least a 1-meter distance from other people.
- **Stand up.** Reduce your sedentary time by standing up whenever possible. Ideally, aim to interrupt sitting and reclining time every 30 minutes. Consider setting up a standing desk by using a high table or stacking a pile of books or other materials, to

continue working while standing. During sedentary leisure time prioritize cognitively stimulating activities, such as reading, board games, and puzzles.

- **Relax.** Meditation and deep breaths can help you remain calm.
- **Eat healthily and stay hydrated.** WHO recommends drinking water instead of sugar-sweetened beverages. Limit or avoid alcoholic beverages for adults and strictly avoid these in young people. Ensure plenty of fruits and vegetables, and limit the intake of salt, sugar and fat.

Table 1

What is know

- Quarantine and isolation are effective measures to reduce diffusion of infection and to prevent pandemic. However these conditions can induce depression, anxiety, anger, and stress.
- Stress, depression and anxiety induce people to eat sugar-rich food and drink alcohol to feel better.
- During quarantine and isolation people reduce physical activity, and also reduce relaxing activities (i.e. yoga)
- Apps for Smartphone can help to control diet and maintain personal ideal weight.
- After quarantine, economic crisis could maintain or sometimes worsen unhealthy lifestyle, mainly in individuals of a low socio-economic level.