

This is a pre print version of the following article:

A stronger form of Neumann's BFC-theorem / Acciarri, C; Shumyatsky, P. - In: ISRAEL JOURNAL OF MATHEMATICS. - ISSN 0021-2172. - 242:(2021), pp. 269-278. [10.1007/s11856-021-2133-1]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

20/04/2024 12:24

(Article begins on next page)

A stronger form of Neumann's BFC-theorem

Cristina Acciarri and Pavel Shumyatsky

Dedicated to Andrea Lucchini on the occasion of his 60th birthday

ABSTRACT. Given a group G , we write x^G for the conjugacy class of G containing the element x . A famous theorem of B. H. Neumann states that if G is a group in which all conjugacy classes are finite with bounded size, then the derived group G' is finite. We establish the following result.

Let n be a positive integer and K a subgroup of a group G such that $|x^G| \leq n$ for each $x \in K$. Let $H = \langle K^G \rangle$ be the normal closure of K . Then the order of the derived group H' is finite and n -bounded.

Some corollaries of this result are also discussed.

1. Introduction

Given a group G and an element $x \in G$, we write x^G for the conjugacy class containing x . Of course, if the number of elements in x^G is finite, we have $|x^G| = [G : C_G(x)]$. A group is called a BFC-group if its conjugacy classes are finite and have bounded size. In 1954 B. H. Neumann discovered that in a BFC-group the derived group G' is finite [10]. It follows that if $|x^G| \leq n$ for each $x \in G$, then the order of G' is bounded by a number depending only on n . A first explicit bound for the order of G' was found by J. Wiegold [17], and the best known was obtained in [7] (see also [11] and [14]). The article [3] deals with groups G in which conjugacy classes containing commutators are bounded. In particular, [3] contains a proof of the following theorem.

2010 *Mathematics Subject Classification.* 20E45, 20F24.

This research was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and Fundação de Apoio à Pesquisa do Distrito Federal (FAPDF), Brazil.

Let n be a positive integer and G a group. If $|x^G| \leq n$ for any commutator x , then $|G''|$ is finite and n -bounded.

Here G'' denotes the second commutator subgroup of G . Throughout the article we use the expression “ (a, b, \dots) -bounded” to mean that a quantity is finite and bounded by a certain number depending only on the parameters a, b, \dots . The following extension of the aforementioned theorems to higher commutator subgroups was obtained in [1].

Let n be a positive integer and w a multilinear commutator word. Suppose that G is a group in which $|x^G| \leq n$ for each w -value $x \in G$. Then the verbal subgroup $w(G)$ has derived group of finite n -bounded order.

A related result for groups in which the conjugacy classes containing squares have finite bounded sizes was established in [4].

In the present paper we obtain a variation of different nature for the Neumann theorem.

THEOREM 1.1. *Let n be a positive integer, G a group having a subgroup K such that $|x^G| \leq n$ for each $x \in K$, and let $H = \langle K^G \rangle$. Then the order of the derived group H' is finite and n -bounded.*

Here, as usual, $\langle K^G \rangle$ denotes the normal closure of K in G .

Theorem 1.1 will be proved in the next section. Section 3 contains several easy but surprising corollaries of Theorem 1.1 for finite groups. In Section 4 we handle profinite groups with restricted centralizers. A group G is said to have restricted centralizers if for each g in G the centralizer $C_G(g)$ either is finite or has finite index in G . This notion was introduced by Shalev in [15] where he showed that a profinite group with restricted centralizers is virtually abelian. We say that a profinite group has a property virtually if it has an open subgroup with that property. The recent article [2] handles profinite groups with restricted centralizers of w -values for a multilinear commutator word w . The theorem proved in [2] says that if w is a multilinear commutator word and G is a profinite group in which the centralizer of any w -value is either finite or open, then the verbal subgroup $w(G)$ is virtually abelian. In Section 4 we establish

THEOREM 1.2. *Let p be a prime and G a profinite group in which the centralizer of each p -element is either finite or open. Then G has a normal abelian pro- p subgroup N such that G/N is virtually pro- p' .*

The proof of Theorem 1.2 is based on Theorem 1.1. It also uses the celebrated theorem of Zelmanov [18] which states that a torsion pro- p group is locally nilpotent.

2. Proof of the main result

The purpose of this section is to prove Theorem 1.1. We start with the following result which is somewhat similar to Dietzmann's lemma which says that if x is an element of finite order in a group G such that x^G is finite, then also the subgroup $\langle x^G \rangle$ is finite (see [13]).

LEMMA 2.1. *Let i, j be positive integers and G a group having a subgroup K such that $|x^G| \leq i$ for each $x \in K$. Suppose that $|K| \leq j$. Then $\langle K^G \rangle$ has finite (i, j) -bounded order.*

PROOF. Since $C_G(K) = \bigcap_{x \in K} C_G(x)$, we observe that the index of $C_G(K)$ in G is at most i^j . Let C be the maximal normal subgroup of G contained in $C_G(K)$. Of course, the index of C in G is at most $(i^j)!$. Set $H = \langle K^G \rangle$. It follows that $|H/Z(H)| \leq (i^j)!$ and Schur's theorem [12, 10.1.4] tells us that the order of H' is (i, j) -bounded. Passing to the quotient G/H' we can assume that H is abelian. Then H is a product of at most i^j conjugates of K and so the order of H is at most j^{i^j} , as desired. \square

Let G be a group generated by a set X such that $X = X^{-1}$. Given an element $g \in G$, we write $l_X(g)$ for the minimal number l with the property that g can be written as a product of l elements of X . The symbol $l_X(g)$ denotes the length of g with respect to X . The proof of the following result can be found in [3, Lemma 2.1].

LEMMA 2.2. *Let H be a group generated by a set $X = X^{-1}$ and let K be a subgroup of finite index m in H . Then each coset Kb contains an element g such that $l_X(g) \leq m - 1$.*

We now fix some notation and hypothesis that we will use within the current section.

HYPOTHESIS 2.3. *Let G be a group having a subgroup K such that $|x^G| \leq n$, for any $x \in K$, and let $H = \langle K^G \rangle$. Denote by X the set of all conjugates of elements of K , that is, $X = \bigcup_{g \in G} K^g$. Let m be the maximum of indices of $C_H(x)$ in H for $x \in K$. Select $a \in K$ such that $|a^H| = m$. Choose b_1, \dots, b_m in H such that $l_X(b_i) \leq m - 1$ and $a^H = \{a^{b_i}; i = 1, \dots, m\}$. (The existence of the elements b_i is guaranteed by Lemma 2.2.) Set $U = C_G(\langle b_1, \dots, b_m \rangle)$.*

Some of the arguments we use in the proof of the next lemmas are similar to those used in [3].

LEMMA 2.4. *Assume Hypothesis 2.3. Then for any $x \in X$ the subgroup $[H, x]$ has finite m -bounded order.*

PROOF. Take $x \in X$. Since the index of $C_H(x)$ in H is at most m , by Lemma 2.2, we can choose elements y_1, \dots, y_m in H such that $l_X(y_i) \leq m-1$ and the subgroup $[H, x]$ is generated by the commutators $[y_i, x]$, for $i = 1, \dots, m$. For any such i write $y_i = y_{i1} \dots y_{i(m-1)}$, with $y_{ij} \in X$. By using standard commutator identities we can rewrite $[y_i, x]$ as a product of conjugates in H of the commutators $[y_{ij}, x]$. Let $\{h_1, \dots, h_s\}$ be the conjugates in H of all elements from the set $\{x, y_{ij} \mid 1 \leq i, j \leq m\}$. Note that the number s here is m -bounded. This follows from the fact that $C_H(x)$ has index at most m in H for each $x \in X$. Put $T = \langle h_1, \dots, h_s \rangle$. Since $[H, x]$ is contained in T' , it is enough to show that T' has finite m -bounded order. Observe that the center $Z(T)$ has index at most m^s in T , since the index of $C_H(h_i)$ is at most m in H for any $i = 1, \dots, s$. Thus, by Schur's theorem [12, 10.1.4], we conclude that T' has finite m -bounded order, as desired. \square

Note that the index of U in G is n -bounded. Indeed, since $l_X(b_i) \leq m-1$ we can write $b_i = b_{i1} \dots b_{i(m-1)}$, where $b_{ij} \in X$ and $i = 1, \dots, m$. By the hypothesis the index of $C_G(b_{ij})$ in G is at most n for any such element b_{ij} . Thus, $[G : U] \leq n^{(m-1)m}$.

The next result is somewhat analogous to Lemma 4.5 of Wiegold in [17].

LEMMA 2.5. *Assume Hypothesis 2.3. Suppose that $u \in U$ and $ua \in X$. Then $[H, u] \leq [H, a]$.*

PROOF. Recall that $U = C_G(\langle b_1, \dots, b_m \rangle)$. For each $i = 1, \dots, m$ we have $(ua)^{b_i} = ua^{b_i}$, since u belongs to U . We know that $ua \in X$, so, taking into account the hypothesis on the order of the conjugacy class of ua in H , we deduce that $(ua)^H$ consists exactly of the elements ua^{b_i} , for $i = 1, \dots, m$. Thus, given an arbitrary element $h \in H$, there exists $b \in \{b_1, \dots, b_m\}$ such that $(ua)^h = ua^b$ and so $u^h a^h = ua^b$. It follows that $[u, h] = a^b a^{-h} \in [H, a]$, and the result holds. \square

We are now ready to embark on the proof of Theorem 1.1.

PROOF OF THEOREM 1.1. Recall that G is a group having a subgroup K such that $|x^G| \leq n$ for any x in K . Let $H = \langle K^G \rangle$. We need to show that $|H'|$ is finite and n -bounded.

As above, the symbol X denotes the set of all conjugates of elements of K . Thus $H = \langle X \rangle$. Let m be the maximum of indices of $C_H(x)$ in H , where $x \in K$. Note that $m \leq n$. Choose $a \in K$ such that $C_H(a)$ has index precisely m in H . By Lemma 2.2 we can choose b_1, \dots, b_m

in H such that $l_X(b_i) \leq m - 1$ and $a^H = \{a^{b_i}; i = 1, \dots, m\}$. Set $U = C_G(\langle b_1, \dots, b_m \rangle)$. Note that the index of U in G is n -bounded.

By the hypothesis a has at most n conjugates in G , say $\{a^{g_1}, \dots, a^{g_n}\}$, that are elements of X . Let T be the normal closure in G of the subgroup $[H, a]$, thus $T = [H, a^{g_1}] \cdots [H, a^{g_n}]$. By Lemma 2.4 each of these subgroups has n -bounded order. We conclude that T has finite n -bounded order.

Let $K_0 = K \cap U$. Note that for any $u \in K_0$ the product ua belongs to K . Therefore, by Lemma 2.5, for any u in K_0 , the subgroup $[H, u]$ is contained in $[H, a]$.

Since T has finite n -bounded order, it is sufficient to show that the derived group of the quotient H/T has finite n -bounded order. We pass now to the quotient G/T and for the sake of simplicity the images of G, H, U, K and K_0 will be denoted by the same symbols. Note that the subgroup K_0 becomes central in H modulo T . Next we consider the quotient $G/Z(H)$. Since the image of K_0 in $G/Z(H)$ is trivial, we deduce that the subgroup K has n -bounded order modulo $Z(H)$. Indeed, in $G/Z(H)$ the order of the image of K is at most the index of U in G . Now it follows from Lemma 2.1 that H has n -bounded order modulo $Z(H)$. Thus, by Schur's theorem [12, 10.1.4], we conclude that $|H'|$ is n -bounded, as desired. \square

3. Corollaries concerning finite groups

In this section we will record several easy corollaries of Theorem 1.1 concerning finite groups. If ϕ is an automorphism of a group G , then the centralizer $C_G(\phi)$ is the subgroup formed by the elements $x \in G$ such that $x^\phi = x$. In the case where $C_G(\phi) = 1$ the automorphism ϕ is called fixed-point-free. A famous result of Thompson [16] says that a finite group admitting a fixed-point-free automorphism of prime order is nilpotent. Higman proved that for each prime p there exists a number $h = h(p)$ such that whenever a nilpotent group G admits a fixed-point-free automorphism of order p , it follows that G is nilpotent of class at most h [8]. Therefore the nilpotency class of a finite group admitting a fixed-point-free automorphism of order p is at most h .

If ϕ is an automorphism of a finite group G such that $(|G|, |\phi|) = 1$, then for each normal ϕ -invariant subgroup N of G we have $C_{G/N}(\phi) = C_G(\phi)N/N$ (see for example [6, Theorem 6.2.2 (iv)]). Write $\gamma_i(G)$ for the i th term of the lower central series of a group G . We have

COROLLARY 3.1. *Let n be a positive integer and G a finite group admitting an automorphism ϕ of prime order p such that $(|G|, p) = 1$*

and $|x^G| \leq n$ for each $x \in C_G(\phi)$. Let $h = h(p)$. Then the order of the derived group of $\gamma_{h+1}(G)$ is n -bounded.

PROOF. Let H be the normal closure of $C_G(\phi)$. Theorem 1.1 tells us that the order of H' is n -bounded. Note that ϕ acts fixed-point-freely on the quotient G/H , whence G/H is nilpotent of class at most h . Therefore $\gamma_{h+1}(G) \leq H$ and so the derived group of $\gamma_{h+1}(G)$ is contained in H' . The result follows. \square

Observe that in the particular case where $p = 2$ Corollary 3.1 states that G'' , the second derived group of G , has n -bounded order. This is because $h(2) = 1$, that is, a finite group admitting a fixed-point-free automorphism of order two is necessarily abelian.

Recall that any finite soluble group G has a Sylow basis — a family of pairwise permutable Sylow p_i -subgroups P_i of G , exactly one for each prime divisor of the order of G , and any two Sylow bases are conjugate. The system normalizer (also known as the basis normalizer) of such a Sylow basis in G is the intersection $T = \bigcap_i N_G(P_i)$. If G is a finite soluble group and T is a system normalizer in G , then $G = \langle T^G \rangle$ (see [12, Theorem 9.2.8]). Therefore we deduce

COROLLARY 3.2. *Let n be a positive integer, G a finite soluble group having a system normalizer T such that $|x^G| \leq n$ for each $x \in T$. Then the order of the derived group G' is n -bounded.*

Given a finite group G , the soluble residual of G is defined as the smallest normal subgroup R with the property that the quotient G/R is soluble.

COROLLARY 3.3. *Let n be a positive integer and G a finite group in which $|x^G| \leq n$ for each 2-element $x \in G$. Then the order of the soluble residual of G is n -bounded.*

PROOF. If G has odd order, then by the Feit-Thompson Theorem [5] G is soluble and so the soluble residual of G is trivial. Therefore we assume that the order of G is even. Let S be a Sylow 2-subgroup of G , and let H be the normal closure of S . Theorem 1.1 tells us that the order of H' is n -bounded. The Feit-Thompson Theorem shows that G/H' is soluble. Hence, H' contains the soluble residual of G and the result follows. \square

4. Proof of Theorem 1.2

Recall that a group G is said to have restricted centralizers if for each g in G the centralizer $C_G(g)$ either is finite or has finite index in G . Of course, in the case where G is profinite this is equivalent to

saying that $C_G(g)$ either is finite or open. Shalev showed in [15] that a profinite group with restricted centralizers is virtually abelian. The goal of this section is to establish Theorem 1.2:

Let p be a prime and G a profinite group in which the centralizer of each p -element is either finite or open. Then G has a normal abelian pro- p subgroup N such that G/N is virtually pro- p' .

Throughout, by a subgroup of a profinite group we mean a closed subgroup. As usual, we say that a group has certain property locally if every finitely generated subgroup has that property. The next lemma is almost obvious. We include the proof for the reader's convenience. Note that the lemma is no longer true if we drop the assumption that G is residually finite (cf. the non-abelian semidirect product of the Prüfer group C_{2^∞} by the group of order 2).

LEMMA 4.1. *Let G be a locally nilpotent group containing an element with finite centralizer. Suppose that G is residually finite. Then G is finite.*

PROOF. Choose $x \in G$ such that $C_G(x)$ is finite. Let N be a normal subgroup of finite index such that $N \cap C_G(x) = 1$. Assume that $N \neq 1$ and let $1 \neq y \in N$. The subgroup $\langle x, y \rangle$ is nilpotent and so the center of $\langle x, y \rangle$ has nontrivial intersection with N . This is a contradiction since $N \cap C_G(x) = 1$. The result follows. \square

The next lemma is somewhat similar to Lemma 2.6 in Shalev [15].

LEMMA 4.2. *Let G be a profinite group having a subgroup K such that the conjugacy class x^G is finite for each $x \in K$. Then there is an integer n such that $|x^G| \leq n$ for each $x \in K$.*

PROOF. For each integer $i \geq 1$ set $S_i = \{x \in K; |x^G| \leq i\}$. The sets S_i are closed. Thus, we have at most countably many sets which cover the subgroup K . The Baire Category Theorem [9, Theorem 34] says that at least one of these sets has non-empty interior. It follows that there exists an open normal subgroup N in G , an element $a \in K$, and a positive integer m such that $|y^G| \leq m$ whenever $y \in a(N \cap K)$. Set $K_0 = N \cap K$. Since every element x of K_0 can be written as a product of a^{-1} and ax and since the centralizers of both a^{-1} and ax have indices at most m , we conclude that $|x^G| \leq m^2$ whenever $x \in K_0$. Let x_1, \dots, x_s be a full system of coset representatives for K_0 in K and set $m_j = |x_j^G|$ for $j = 1, \dots, s$. Let n be the product $m_1 \cdots m_s m^2$. We deduce that $|x^G| \leq n$ for each $x \in K$, as required. \square

We can now proceed with the proof of Theorem 1.2.

PROOF OF THEOREM 1.2. Let P be a Sylow p -subgroup of G . Suppose first that the conjugacy class x^G is finite whenever x is a p -element in G . By Lemma 4.2 there is an integer n such that $|x^G| \leq n$ for each $x \in P$. Let $H = \langle P^G \rangle$. Theorem 1.1 tells us that the derived group H' is finite. Choose an open normal subgroup L in G such that $L \cap H' = 1$. Then $L \cap H$ is a normal abelian pro- p subgroup while $G/(L \cap H)$ is virtually pro- p' . This proves the theorem in the case where the conjugacy classes of p -elements are finite.

Therefore we assume that P has an element whose centralizer in G is finite. Suppose that P is torsion. Zelmanov's theorem [18] says that P is locally nilpotent. In view of Lemma 4.1 the Sylow subgroup P is finite and so G is virtually pro- p' , as required.

Hence, we assume that P contains an element a of infinite order. This implies that $C_G(a)$ is open, because $\langle a \rangle$ is infinite and contained in $C_G(a)$. Let C be an open normal subgroup of G contained in $C_G(a)$. Since every element of C centralizes $\langle a \rangle$, it follows that the centralizer in G of every element from $P_0 = P \cap C$ is open. By Lemma 4.2 there is an integer m such that $|x^G| \leq m$ for each $x \in P_0$. Let $M = \langle P_0^G \rangle$. In virtue of Theorem 1.1 the derived group M' is finite. Choose an open normal subgroup J in G such that $J \cap M' = 1$. Then $J \cap M$ is a normal abelian pro- p subgroup while $G/(J \cap M)$ is virtually pro- p' . The proof is complete. \square

References

- [1] E. Detomi, M. Morigi, P. Shumyatsky, BFC-theorems for higher commutator subgroups, *Quarterly J. Math.* **70** (2019), no. 3, 849–858.
- [2] E. Detomi, M. Morigi, P. Shumyatsky, Profinite groups with restricted centralizers of commutators, *Proceedings of the Royal Society of Edinburgh, Section A: Mathematics*, doi:10.1017/prm.2019.17.
- [3] G. Dierings, P. Shumyatsky, Groups with boundedly finite conjugacy classes of commutators, *Quarterly J. Math.* **69** (2018), no. 3, 1047–1051.
- [4] G. Dierings, P. Shumyatsky, Groups in which squares have boundedly many conjugates, *J. Group Theory* **22** (2019), 133–136.
- [5] W. Feit and J. G. Thompson, Solvability of groups of odd order, *Pacific J. Math.* **13** (1963), 773–1029.
- [6] D. Gorenstein, *Finite Groups*, Chelsea Publishing Company, New York, 1980.
- [7] R. M. Guralnick, A. Maroti, Average dimension of fixed point spaces with applications, *J. Algebra* **226** (2011), 298–308.
- [8] G. Higman, Groups and rings having automorphisms without non-trivial fixed elements, *J. London Math. Soc.* **32** (1957), 321–334.
- [9] J. L. Kelley, *General topology*, Grad. Texts in Math., vol. 27, Springer, New York, 1975.
- [10] B. H. Neumann, Groups covered by permutable subsets, *J. London Math. Soc.* (3) **29** (1954), 236–248.

- [11] P. M. Neumann, M.R. Vaughan-Lee, An essay on BFC groups, Proc. Lond. Math. Soc. **35** (1977), 213–237.
- [12] D. J. S. Robinson, *A course in the theory of groups*, Second edition. Graduate Texts in Mathematics, 80. Springer-Verlag, New York, 1996.
- [13] D.J.S. Robinson, *Finiteness Conditions and Generalized Soluble Groups*, part 1, Springer-Verlag, 1972.
- [14] D. Segal, A. Shalev, On groups with bounded conjugacy classes, Quart. J. Math. Oxford **50** (1999), 505–516.
- [15] A. Shalev, Profinite groups with restricted centralizers. Proc. Amer. Math. Soc. **122** (1994), 1279–1284.
- [16] J. Thompson, Finite groups with fixed-point-free automorphisms of prime order, Proc. Nat. Acad. Sci. U.S.A., **45** (1959), 578–581.
- [17] J. Wiegold, Groups with boundedly finite classes of conjugate elements, Proc. Roy. Soc. London Ser. A **238** (1957), 389–401.
- [18] E. I. Zelmanov, On periodic compact groups. Israel J. Math. **77**, 83–95 (1992).

CRISTINA ACCIARRI: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA,
BRASILIA-DF, 70910-900 BRAZIL

Email address: acciarricristina@yahoo.it

PAVEL SHUMYATSKY: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRASILIA,
BRASILIA-DF, 70910-900 BRAZIL

Email address: pavel@unb.br