

This is the peer reviewed version of the following article:

Comment on "Western Dietary Pattern Antioxidant Intakes and Oxidative Stress: Importance during the SARS-CoV-2/COVID-19 Pandemic" / Mattioli, Anna V; Farinetti, Alberto. - In: ADVANCES IN NUTRITION. - ISSN 2161-8313. - 12:3(2021), pp. 1044-1045-1045. [10.1093/advances/nmab029]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

23/09/2024 17:18

(Article begins on next page)

study. Thus, in contrast to the authors' interpretation (1), we deemed the effect to be trivial not in relation to generalized, heuristic definitions, but in relation to expected effect sizes for apparently similar interventions. Moreover, to directly address the authors' query, the unstandardized meta-analytic effect of coaching on weight loss was -1.4 kg [95% credible interval (CrI): -3.0 , -0.1]. Considering only the highest-quality studies, the effect was -1.1 kg (95% CrI: -3.1 , 1.1). Although there is room for discussion about what constitutes meaningful weight loss in different contexts, we believe most would agree that these average effects are, indeed, trivial.

The root of the discrepant outcomes found in our study and in the compendium is methodological in nature. While we assessed the quality of studies using a clear systematic approach, and in accordance with GRADE recommendations, their compendium (3) did not use a systematic approach to assess either study quality or effect magnitude. The authors' conclusion that health coaching is beneficial for treating obesity is based on the observation that "A large portion of the studies showed a positive effect on weight reduction" (3). This is not a valid assertion, particularly considering the publication bias identified in our study. This highlights the need for exploring the potential impact of conflicts of interest on health coaching literature, in which objective research by those with no vested interest should be important.

To conclude, our findings represent a first attempt to systematically assess the health coaching literature. The main conclusions are that most of the studies present serious methodological flaws and divergent theoretical backgrounds, hampering the clinical use of this technique in an effective, uniform way. This is not to say that health coaching cannot be an effective and well-structured intervention. A few good examples on how this strategy can be a useful co-adjutant therapy in obesity management do exist (9, 10). Expanding the number of high-quality studies, with detailed information of their interventions and reporting on all aspects of their study design, is essential to pave the way for a more evidence-based use of health coaching in clinical practice. Simply incorporating any lifestyle behavioral intervention under the obscure umbrella of health coaching will not help build a scientifically oriented body of knowledge.

Sofia Mendes Sieczkowska
Alisson Padilha de Lima
Paul Alan Swinton
Eimear Dolan
Hamilton Roschel
Bruno Gualano

From the Applied Physiology & Nutrition Research Group, School of Physical Education and Sport, Laboratory of Assessment and Conditioning in Rheumatology; School of Medicine, University of Sao Paulo, Sao Paulo, Brazil (SMS, APL, ED, HR; BG, e-mail: gualano@usp.br); School of Physical Education, Faculty IELUSC, Joinville, Santa Catarina, Brazil (APL); and School of Health Sciences, Robert Gordon University, Aberdeen, United Kingdom (PAS).

The authors reported no funding received for this letter.
Author disclosures: The authors report no conflicts of interest.
All authors read and approved the final manuscript.

References

- Sforzo G, Moore M, Moore G, Harenberg S. Comment on "Health Coaching: 100 Strategies for Weight Loss: A Systematic Review and Meta-Analysis". *Adv Nutr* 2021;12(3):1042–3.
- Sieczkowska SM, de Lima AP, Swinton PA, Dolan E, Roschel H, Gualano B. Health coaching strategies for weight loss: a systematic review and meta-analysis. *Adv Nutr* 2020; nmaa159. <https://doi.org/10.1093/advances/nmaa159>
- Sforzo GA, Kaye MP, Harenberg S, Costello K, Cobus-Kuo L, Rauff E, Edman JS, Frates E, Moore M. Compendium of health and wellness coaching: 2019 addendum. *Am J Lifestyle Med* 2020;14(2):155–68.
- Janssen V, De Gucht V, Van Exel H, Maes S. A self-regulation lifestyle program for post-cardiac rehabilitation patients has long-term effects on exercise adherence. *J Behav Med*. 2014;37(2):308–21.
- Lin CH, Chiang SL, Heitkemper MML, Hung YJ, Lee MS, Tzeng WC, Chiang LC. Effects of telephone-based motivational interviewing in lifestyle modification program on reducing metabolic risks in middle-aged and older women with metabolic syndrome: a randomized controlled trial. *Int J Nurs Stud*. 2016;60:12–23.
- Miller WR. Motivational interviewing with problem drinkers. *Behav Psychother* 1983;11(2):147–72.
- Rubak S, Sandbæk A, Lauritzen T, Christensen B. Motivational interviewing: a systematic review and meta-analysis. *Br J Gen Pract* 2005;55(513):305–12.
- Armstrong MJ, Mottershead TA, Ronksley PE, Sigal RJ, Campbell TS, Hemmelgarn BR. Motivational interviewing to improve weight loss in overweight and/or obese patients: a systematic review and meta-analysis of randomized controlled trials. *Obes Rev* 2011;12(9):709–23.
- Kelly JT, Conley M, Hoffmann T, Craig JC, Tong A, Reidlinger DP, Reeves MM, Howard K, Krishnasamy R, Kurtkoti J, et al. A coaching program to improve dietary intake of patients with CKD: Entice-CKD. *Clin J Am Soc Nephrol* 2020;15(3):330–40.
- 40.40, Yun YH, Lim C III, Lee ES, Kim YT, Shin KH, Kim Y, Park KJ, Jeong S, Ryu KW, Han W, et al. Efficacy of health coaching and a web-based program on physical activity, weight, and distress management among cancer survivors: a multi-centered randomised controlled trial. *Psycho-oncology* 2020;29(7):1–10.

doi: <https://doi.org/10.1093/advances/nmab021>

Comment on "Western Dietary Pattern Antioxidant Intakes and Oxidative Stress: Importance during the SARS-CoV-2/COVID-19 Pandemic"

Dear Editor:

We have read with great interest the article "Western Dietary Pattern Antioxidant Intakes and Oxidative Stress: Importance during the SARS-CoV-2/COVID-19 Pandemic" by Trujillo-Mayol et al. (1) and we found it significant in the context of clinical prevention.

The relevant point explored by this review is the importance of balanced dietary habits including appropriate

amounts of antioxidants to maintain the immune system during the current severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)/coronavirus disease 2019 (COVID-19) pandemic. Viral infections are characterized by high oxidative stress and an appropriate intake of antioxidants would help to fight against the virus. The authors suggested that lifestyle factors could influence the impact of the disease.

With reference to the findings reported in the article, we would like to make the following contribution to the discussion. The review analyzes in depth the potential beneficial effects of a healthy diet, including a Mediterranean Diet, on viral infection. In recent articles we also supported the idea that a healthy lifestyle could contribute to a higher immune response to the virus and would therefore be beneficial in the fight against the pandemic (2, 3). It has been suggested that there would be a synergy among the antioxidant-rich foods that would foster favorable changes in inflammatory pathways (4). Humans do not consume 1 type of food but a wide variety of combinations of different foods forming a dietary pattern. Consequently, in analyzing eating habits it is mandatory to take the interactions between different foods and their components into consideration. Moreover, it is known that plant-derived phenolic compounds could differently influence the health of males and females (5). The recent COVID-19 pandemic strongly affected women owing to increased psychological distress leading to unhealthy lifestyles, and women were specifically more likely to develop food cravings (2, 6, 7). Food cravings are characterized by a high intake of fat- and sugar-rich foods and by a low intake of fruit and vegetables. In a previous article (2) we hypothesized that vitamin D prophylaxis can help reduce the severity of the disease caused by SARS-CoV-2, particularly in contexts where hypovitaminosis D is common, i.e., women currently living in Northern countries (8). Several factors may have contributed to the widespread infection in Italy, despite the strong adherence to the Mediterranean diet. Lifestyle and diet can influence our inflammatory response to the virus and therefore it is important that both are optimized. The review by Trujillo-Mayol et al. (1) emphasizes these aspects of prevention but does not emphasize the potential increased risk in women. Women's health is strongly influenced by lifestyle, which includes dietary pattern and nutritional status, physical activity, and socioeconomic stress. A healthy lifestyle is mandatory to prevent chronic diseases and to fight infections. Women are less likely to adopt a healthy lifestyle owing to the high social pressures that oblige women to play numerous roles in the family, in society, and at work, all of which are debilitating and time consuming (9, 10). We need educational programs and prevention measures specifically dedicated to women's health.

Anna V Mattioli
Alberto Farinetti

From the Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of

Modena and Reggio Emilia, Modena, Italy (AVM, e-mail: annavittoria.mattioli@unimore.it; AF).

Author disclosures: The authors report no conflicts of interest.
We thank Mrs. Janet Ann Carter for reviewing and editing the letter.

References

1. Trujillo-Mayol I, Guerra-Valle M, Casas-Forero N, Sobral MMC, Viegas O, Alarcón-Enos J, Ferreira IM, Pinho O. Western dietary pattern antioxidant intakes and oxidative stress: importance during the SARS-CoV-2/COVID-19 pandemic. *Adv Nutr* 2021;nmaa171. (Epub ahead of print; doi: 10.1093/advances/nmaa171).
2. Mattioli AV, Pinti M, Farinetti A, Nasi M. Obesity risk during collective quarantine for the COVID-19 epidemic. *Obes Med* 2020;20:100263.
3. Mattioli AV, Ballerini Puviani M, Nasi M, Farinetti A. COVID-19 pandemic: the effects of quarantine on cardiovascular risk. *Eur J Clin Nutr* 2020;74:852–5.
4. El-Missiry MA, Fekri A, Kesar LA, Othman AI. Polyphenols are potential nutritional adjuvants for targeting COVID-19. *Phytother Res* 2020 Dec 22 (Epub ahead of print; doi: 10.1002/ptr.6992).
5. Campesi I, Marino M, Cipolletti M, Romani A, Franconi F. Put “gender glasses” on the effects of phenolic compounds on cardiovascular function and diseases. *Eur J Nutr* 2018;57(8):2677–91.
6. Rodríguez-Martín BC, Meule A. Food craving: new contributions on its assessment, moderators, and consequences. *Front Psychol* 2015;6:21.
7. Mattioli AV, Sciomer S, Moscucci F, Maiello M, Cugusi L, Gallina S, Dei Cas A, Lombardi C, Pengo M, Parati G, et al. Cardiovascular prevention in women: a narrative review from the Italian Society of Cardiology working groups on ‘Cardiovascular Prevention, Hypertension and peripheral circulation’ and on ‘Women Disease’. *J Cardiovasc Med* 2019;20(9):575–83.
8. Schwalfenberg GK. A review of the critical role of vitamin D in the functioning of the immune system and the clinical implications of vitamin D deficiency. *Mol Nutr Food Res* 2011;55:96–108.
9. Sciomer S, Moscucci F, Maffei S, Gallina S, Mattioli AV. Prevention of cardiovascular risk factors in women: the lifestyle paradox and stereotypes we need to defeat. *Eur J Prev Cardiol* 2019;26(6):609–10.
10. Mattioli AV, Sciomer S, Maffei S, Gallina S. Lifestyle and stress management in women during COVID-19 pandemic: impact on cardiovascular risk burden. *Am J Lifestyle Med* 2020 Dec 10 (Epub ahead of print; doi: 10.1177/1559827620981014).

doi: <https://doi.org/10.1093/advances/nmab029>

Comment on "Western Dietary Pattern Antioxidant Intakes and Oxidative Stress: Importance during the SARS-CoV-2/COVID-19 Pandemic"

Dear Editor:

In an article published in *Advances in Nutrition*, Trujillo-Mayol et al. (1) present their argument for increasing dietary antioxidant intake to improve coronavirus disease