



UNIMORE
UNIVERSITÀ DEGLI STUDI DI
MODENA E REGGIO EMILIA



2017 SCIENTIFIC MEETING
ITALIAN ASSOCIATION FOR THE STUDY OF
TRACE ELEMENTS IN LIVING ORGANISMS - AISETOV

REGGIO NELL'EMILIA (ITALY), OCTOBER 20, 2017



Prampolini square and the Dome of Reggio Emilia

TRACE ELEMENTS IN HUMAN AND ANIMAL
HEALTH: FOCUS ON NEUROLOGICAL DISEASE

ABSTRACT BOOK

Organized by:

AISETOV and UNIVERSITY OF MODENA AND REGGIO EMILIA

P-03. Assessment of selenium levels and risk factors for stroke and other cardiovascular disease: a cross sectional study in a seleniferous area of Punjab, India

Rajinder Chawla¹, Silvia Cilloni², Rinchu Loomba¹, Chaudhary Rohit¹, Datt Chander¹,
Karaj S. Dhillon³, Tommaso Filippini², Marco Vinceti²

¹Christian Medical College & Hospital, Ludhiana (Pb), India;

²CREAGEN - Environmental, Genetic and Nutritional Epidemiology Research Center, University of Modena and Reggio Emilia, Italy;

³Punjab Agricultural University, Ludhiana (Pb), India.

Background and aims: Rural areas of Punjab in India have been found to have soil rich in selenium (Se); about 2160 hectare area is seleniferous and is populated by about 10,000 inhabitants. Selenium concentrations in these villages were reported to be as high as 65 times over non-seleniferous areas. The aim of this cross-sectional study was to evaluate selenium levels in blood, hair and nails in a group of subjects living in this area, and to evaluate the correlation between selenium exposure levels and a relevant cardiovascular risk factor and blood pressure.

Methods: In a random sample of rural residents in three districts of a seleniferous area of Punjab, we determined selenium concentration in hair, nail clippings and serum samples. Analyses were carried out using atomic absorption spectrophotometry at National Dairy Research Institute (NDRI), Karnal, India. Data analysis was performed using the STATA 15.0 software (STATA Corp. TX).

Results: A total of 680 human subjects were recruited in this study, with a male/female ratio of 0.65 and a median age of 43 (IQR 32-52). Medium selenium levels in blood, hair and nail were 86.7 µg/l (IQR 55.9-200.3), 20.7 µg/g (IQR 12.6-40.3) and 56.9 µg/g (IQR 42.8-83.9), respectively, with lower levels in women in all three kind of samples. Concerning systolic blood pressure, Pearson's correlation coefficients were 0.102 (95 % CI -0.025 to 0.226, p=0.116); 0.076 (95% CI -0.010 to 0.160, p=0.085); 0.072 (95% CI -0.015 to 0.157, p=0.104) with blood, hair and nail, respectively. For diastolic blood pressure, Pearson's correlation coefficients are 0.106 (95% CI -0.022 to 0.230, p=0.104), 0.036 (95% CI -0.050 to 0.122, p=0.409), 0.049 (95% CI -0.038 to 0.135, p=0.272), respectively.

Conclusions: Our findings indicate a positive correlation between selenium content in blood, hair and nails and increasing systolic and diastolic pressure levels, in line with previous epidemiologic findings, indicating a possible health concern for this highly exposed population. The possible relation between selenium over-exposure and onset of hypertension and other cardiovascular diseases deserves further investigation.

Printed in Reggio Emilia (Italy)

ISBN 978-88-943098-0-5



20 October 2017