

This is the peer reviewed version of the following article:

Intake of arsenic and mercury from fish and seafood in a Northern Italy community / Filippini, Tommaso; Malavolti, Marcella; Cilloni, Silvia; Wise, Lauren A.; Violi, Federica; Malagoli, Carlotta; Vescovi, Luciano; Vinceti, Marco. - In: FOOD AND CHEMICAL TOXICOLOGY. - ISSN 0278-6915. - 116:Pt B(2018), pp. 20-26. [10.1016/j.fct.2018.04.010]

Terms of use:


The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

09/04/2024 15:38

Accepted Manuscript

Intake of arsenic and mercury in a northern Italy community from fish and seafood

Tommaso Filippini, Marcella Malavolti, Silvia Cilloni, Lauren A. Wise, Federica Violi, Carlotta Malagoli, Luciano Vescovi, Marco Vinceti



PII: S0278-6915(18)30212-6

DOI: [10.1016/j.fct.2018.04.010](https://doi.org/10.1016/j.fct.2018.04.010)

Reference: FCT 9699

To appear in: *Food and Chemical Toxicology*

Received Date: 2 March 2018

Revised Date: 27 March 2018

Accepted Date: 4 April 2018

Please cite this article as: Filippini, T., Malavolti, M., Cilloni, S., Wise, L.A., Violi, F., Malagoli, C., Vescovi, L., Vinceti, M., Intake of arsenic and mercury in a northern Italy community from fish and seafood, *Food and Chemical Toxicology* (2018), doi: 10.1016/j.fct.2018.04.010.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Intake of arsenic and mercury in a Northern Italy community from fish and seafood**Authors:**

Tommaso Filippini^a, Marcella Malavolti^a, Silvia Cilloni^a, Lauren A. Wise^b, Federica Violi^a, Carlotta Malagoli^a, Luciano Vescovi^c, Marco Vinceti^{a,b}

Affiliations:

^aCREAGEN, Environmental, Genetic and Nutritional Epidemiology Research Center - Section of Public Health - Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, 287 Via Campi, 41125 Modena, Italy;

^bDepartment of Epidemiology, Boston University School of Public Health, Boston, MA, USA;

^cIREN, Reggio Emilia, Italy.

Corresponding author: Marco Vinceti, CREAGEN – Section of Public Health - Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Via Campi 287, 41125 Modena, Italy (Tel. +39 059 2055481; fax +39 059 2055483; E-mail marco.vinceti@unimore.it)

1.1 Introduction

During the last few decades, the consumption of seafood has increased considerably, mainly due to growing consciousness that seafood is an important source of protein, omega-3 polyunsaturated fatty acids (n-3 PUFAs), vitamins, and minerals (Domingo et al., 2007; Matos et al., 2015). The European Food Safety Authority (EFSA) has recently reviewed the role of seafood in European diets, assessing the beneficial effects of its consumption on health outcomes, and recommending fish consumption of 1-2 servings per week (European Food Safety Authority, 2014), similarly to World Health Organization and Food and Agriculture Organization (FAO/WHO, 2010b). However, fish and seafood also represent a source of environmental contaminants (Adel et al., 2016; Copat et al., 2014; Gonzalez-Alzaga et al., 2017; Vilavert et al., 2017). In fact, urbanization, globalization and human activity have adversely influenced water quality during the last decades, due to increased pollution in global aquatic ecosystem and the decreased capacity of freshwater ecosystems to dilute anthropogenic contaminants (Alava et al., 2017; Delpla et al., 2009; Duh et al., 2008; Pal et al., 2014; WHO, 2017). Therefore, fish and seafood are foods with the highest contribution to total dietary uptake of chemical contaminants by humans (Bae et al., 2017; Bosch et al., 2016; Cano-Sancho et al., 2015b; Copat et al., 2013; Perello et al., 2014; Taylor et al., 2016). In particular, fish and seafood are a relevant source of heavy metals and metalloids, which can spread through the food chain and accumulate in human body reaching levels which could represent a potential risk to human health (Arslan et al., 2017; Chiocchetti et al., 2017; Copat et al., 2013; Kaya and Turkoglu, 2017; Vinceti et al., 2013). Arsenic (As) and mercury (Hg) are two of the heavy metals of major environmental and toxicological concern (ATSDR, 2012a, b; Donaldson et al., 2010; Papadopoulos et al., 2015).

In this study, we used validated semi-quantitative food frequency questionnaires to assess the dietary habits of a representative sample of individuals aged 18-87 years from Northern Italy.

We estimated weekly dietary intakes of Arsenic (As) and Mercury (Hg), and we compared them with safety standards set by the European Food Safety Authority.

2.1 Methods

2.1.1 Food collection and analysis

We determined the arsenic and mercury content of foods by characterizing the usual diet of a sample of the population of the Emilia-Romagna region, Northern Italy. As previously described, we selected the most frequently consumed foods in a random sample of this population (Bottecchi and Vinceti, 2012; Cilloni et al., 2017), and we bought samples of these food items in markets and community canteens during the period from October 2016 to February 2017, as previously described (Filippini et al., 2018). To summarize, we strove to avoid cross-contamination with metals from food containers using plastic tubes or jars as well plastic cutlery or stainless-steel knife during food collection and handling. We then cut with a clean stainless-steel knife solid food samples collecting specimens in six different points from the plate. These samples were homogenised separately in a food blender, equipped with a stainless-steel blade, for subsequent analyses. We placed portions of the samples (0.5-1.0 g) in quartz containers previously washed with MilliQ water (MilliQPlus, Millipore, MA, USA) and HNO_3 . Food samples were liquid-ashed (5 ml HNO_3 + 5 ml H_2O_2) in a microwave digestion system (Discover SP-D, CEM Corporation, NC, USA) and then stored in plastic tubes and diluted to 50 ml with deionised water before analyses. We performed trace element determination using an inductively coupled plasma-mass spectrometer (Agilent 7500ce, Agilent Technologies, CA, USA). We implemented quality controls including both blank solution (MilliQ water, Millipore, MA, USA), and using a control solution of tap-water additionally enriched with 22 ppb of each element under investigation. Limit of quantification was 0.05 $\mu\text{g/kg}$ for arsenic and 0.02 $\mu\text{g/kg}$ for mercury, corresponding limit of

detection (LOD) was 0.012 µg/kg for arsenic and 0.005 µg/kg for mercury. Values below the LOD were set equal to LOD/2. We performed all the analyses in duplicate.

We then reported the concentrations of investigated trace elements according to the food consumption patterns and food categories typical of this Italian population, as assessed through a food frequency questionnaire (FFQ), as explained in detail (Filippini et al., 2017; Turrini et al., 2001). Our final list of food categories included cereals, meat, milk and dairy products, eggs, fish and seafood, vegetables, legumes, potatoes, fresh fruits, dry fruits, sweets and beverages. We further divided fish and seafood according to detailed food consumption patterns (Pala et al., 2003) into seafood (i.e. crustaceans, cephalopods, bivalves) and fish (i.e. preserved fish, tinned fish, and fresh fish). In particular group of fresh fish was further divided in codfish, sole or flounder, sardine, trout, swordfish, and other fish (including tuna and salmon).

2.1.2. Study population and estimation of daily element intake

We carried out the evaluation of dietary habits of a representative sample of a Northern Italy population, as previously described (Malagoli et al., 2015; Vinceti et al., 2011). Briefly, we identified a list of 2825 potentially eligible subjects from the population database of the Emilia-Romagna Region residents, using the National Health Service directory (namely from the provinces of Bologna, Ferrara, Modena, Parma, and Reggio Emilia). The eligible subjects were selected among all residents in the study provinces having the same sex, year of birth (± 5 years) and province of residence of 572 melanoma patients diagnosed during 2005-2006 in the same provinces. A total of 747 (26.4%) controls agreed to participate in the study, after giving their informed written consent, and returned the lifestyle and FFQ which we had mailed to them. Twenty-eight individuals were later excluded from subsequent analysis because of incomplete data or extreme values (ratio of total energy intake:calculated basal metabolic rate lower than the

0.5th percentile or higher than the 99.5th percentile) derived from the FFQ. The final population sample comprised 719 subjects, 319 males and 400 females, aged from 18 to 87 years (median 57, interquartile range (IQR) years in all subjects, with corresponding figures 60 (IQR: 50-69) and 53 (IQR: 41-63) for males and females, respectively. Median energy intake was 1906 kcal/day (IQR: 1538-2364) in all subjects, with 2024 kcal/day (IQR: 1649-2462) and 1800 kcal/day (IQR: 1455-2296) in male and female participants, respectively.

In this population, we assessed food and nutrient intake using a validated semi-quantitative FFQ specifically developed for the Central-Northern Italy population (Pala et al., 2003; Pasanisi et al., 2002) within the EPIC (European Prospective Investigation into Cancer and Nutrition) project. The FFQ was designed to estimate frequency and amount of consumption of 188 food items over the previous year, using also photos of serving sizes to help proper completion by participants. Dietary intake of individual nutrients and contaminants were calculated using *ad hoc* software (Malagoli et al., 2015; Malavolti et al., 2013).

We combined data on the estimated trace elements in foods and the EPIC FFQ to compute total weekly trace element intake using the equation presented below.

$$\text{Weekly dietary exposure} \left(\frac{\mu\text{g}}{\text{kg bw/week}} \right) = \sum \frac{\text{element food content} \left(\frac{\mu\text{g}}{\text{kg}} \right) \times \text{food intake} \left(\frac{\text{g}}{\text{day}} \right) \times 7}{\text{body weight (kg)} \times 1000}$$

We multiplied the element content measured in food ($\mu\text{g}/\text{kg}$) with the intake as estimated with the FFQ (g/day) and we divided by the body weight (kg) of each participant. Accordingly, we estimated weekly dietary As and Hg intake for the total diet and for each food category by reporting median and interquartile ranges of intake. Finally, we estimated inorganic As, inorganic Hg and methylmercury (methyl-Hg) intake according to the conservative EFSA approach (European Food Safety Authority, 2009, 2012). In particular, for the Hg species, we assumed that 100% of Hg in fish is in the organic form (i.e. methyl-Hg). However, in order to ensure that dietary exposure to

inorganic was not underestimated, 20% of total mercury from fish was assumed to be inorganic (See Supplemental Table S1 for details).

3.1 Results

Table 1 shows the distribution of As and Hg in our study population, taking into account the 890 food items we assessed and their percentage contribution to each food category. Levels were below the LOD in 4 and 90 samples for arsenic and mercury, respectively.

We report in table 2 the average As and Hg intake according to food category consumption for all participants and by sex.

Table 3 summarizes the estimated levels of weekly dietary intake of total and inorganic As, respectively of 5.34 $\mu\text{g/kg}$ of body weight (bw) per week (IQR: 2.75-8.84) and 0.58 $\mu\text{g/kg}$ bw/week (IQR:0.44-0.75), and total and inorganic Hg, respectively of 0.39 $\mu\text{g/kg}$ bw/week (IQR: 0.25-0.59) and 0.14 $\mu\text{g/kg}$ bw/week (IQR: 0.10-0.19). For both elements, the highest contribution to total dietary intake was driven by fish and seafood intake. Concerning inorganic trace element intake, aside from fish and seafood, major contributors to inorganic As intake were cereals and beverages, while for inorganic Hg, they were cereals and vegetables. On the converse, estimated intake of methyl-Hg corresponded to total contribution of Hg by fish and seafood, namely 0.32 (IQR: 0.18-0.51) $\mu\text{g/kg}$ bw/week, on the basis of the most conservative value of 100% as suggested in the EFSA exposure assessment (European Food Safety Authority, 2012). Sex-specific estimates showed slightly higher values of total As among females, due to higher contribution from fish and seafood, while results for inorganic As were substantially comparable (Tables 4-5). Regarding total and inorganic Hg, dietary exposure was higher in females, due to a higher contribution from fish and vegetable intake. Methyl-Hg intake was 0.287 $\mu\text{g/kg}$ bw/week (IQR: 0.159-0.457) and 0.345 $\mu\text{g/kg}$ bw/week (IQR: 0.192-0.556) in males and females, respectively.

Since fish and seafood showed the highest contamination levels and contributed to the highest amount of exposure in the study population, we further investigated their concentrations (white bars) in different types of fish (Figure 1). Interestingly, the highest As values were found in sardine, sole/flounder and cephalopods (i.e. octopus, cuttlefish and squid). On the converse, for Hg, the highest concentrations were found in typical big and predatory fishes, i.e. sword fish, and both fresh and tinned tuna. Supplemental Figures S1-S3 show the concentration of each element and relative contribution to estimated dietary intake, respectively, for cereals, vegetables, and beverages. Among cereals, rice showed the highest concentration and contribution (60%) to total As intake, followed by bread and pasta (around 17% each). Bread demonstrated the highest contribution to total Hg intake (55%), followed by crispbread/rusks (20%) and rice (16%). With regard to vegetables, the highest concentrations for both total As and total Hg were found in mushrooms, followed by leafy vegetables. Similarly, leafy vegetables (42%), root vegetables (13%) and mushrooms (12%) made large contributions to total As, while leafy vegetables (35%), mushrooms (25%) and tomatoes (21%) made large contributions to total Hg. Finally, among beverages, the major contributors to As and Hg were coffee and wine, accounting for 46% and 35% for total As and 90% and 6% for total Hg, respectively, within this category.

4.1 Discussion

In this study, we assessed total As and Hg intake within a Northern Italian community by measuring trace element content in foods and assessing usual dietary intake of foods containing these elements. While our findings are new for the northern part of Italy due to the absence of previous studies carried out in this region, they agree with previous studies from other European countries (Cirillo et al., 2010; Marti-Cid et al., 2008; Martorell et al., 2011; Miklavcic et al., 2013; Perello et al., 2015), especially for fish and seafood, which showed the highest levels of

contamination (Brambilla et al., 2013; Calatayud et al., 2012; Cano-Sancho et al., 2015a). Therefore, our findings indicate that, in industrialized nations, fish and seafood are likely to be the most important contributors to total As and Hg intake.

Both As and Hg are environmental contaminants that have both natural and anthropogenic origin. For both elements, their toxicity is strongly related not only to the environmental levels but also to their species, either elemental or inorganic or organic (ATSDR, 2012a, b). As is most toxic in its inorganic form (Sattar et al., 2016). In our population, the estimated weekly intake of inorganic arsenic of 0.58 $\mu\text{g/kg bw/ week}$ (IQR: 0.44-0.75) is consistent with previous findings (Barone et al., 2015; Copat et al., 2014; Storelli and Barone, 2013), and it is within the range values of benchmark dose suggested by EFSA (European Food Safety Authority, 2009).

For Hg, its organic compounds, including the most abundant methyl-Hg form, are the most common species entering the food chain, and they show the highest toxicity to humans (Bjorklund et al., 2017; European Food Safety Authority, 2012). Based on our estimates, weekly intake of inorganic Hg and methyl-Hg are 0.14 (IQR: 0.10-0.19) and 0.32 (IQR: 0.18-0.51), respectively, below the tolerable weekly intake (TWI) of 4 $\mu\text{g/kg bw}$ and 1.6 $\mu\text{g/kg bw}$ established by both EFSA and the Joint FAO/WHO Expert Committee on Food Additives (European Food Safety Authority, 2012; FAO/WHO, 2010a).

Since large, predator fish showed the highest Hg concentrations (Burger and Gochfeld, 2005; Donald et al., 2015), as expected, subjects consuming a large portion of their diet from these type of fishes, such as swordfish or tuna, were at higher risk of a relevant exposure (Adel et al., 2016; Hightower and Moore, 2003). Interestingly, we did not find appreciable differences between tinned and fresh tuna in Hg levels, while for As, similarly with previous findings, slightly lower concentrations were detected in tinned tuna as compared with fresh tuna (Nunez et al., 2018).

Sex-related differences in dietary As and Hg exposure have already been described (Llobet et al., 2003; Marzec and Schlegel-Zawadzka, 2004). Our analysis showed a higher intake of total As in females, due to the higher contribution of fish category in females, while intake of inorganic As was substantially similar between males and females. Consistent with previous findings (Liu et al., 2018), we found higher levels of total intake of Hg, inorganic Hg and methyl-Hg in females, entirely driven by higher fish and seafood consumption.

Previous studies suggested that only a fraction of total heavy metal intake is absorbed after ingestion (Calatayud et al., 2012; Cano-Sancho et al., 2015a; Maulvault et al., 2011).

Study strengths include collection of data on a large number of foods, which were divided in categories based on food consumption patterns derived from a validated dietary assessment in the same population (Filippini et al., 2018; Malagoli et al., 2015; Pala et al., 2003; Pasanisi et al., 2002; Turrini et al., 2017; Vinceti et al., 2011). We could therefore obtain both a weighted average of As and Hg food category content and an estimate of overall and food-specific intake in our study population, allowing comparability of our results with other studies. Moreover, use of local groceries and supermarkets for purchasing foodstuffs provided a close-to-real estimation of dietary intake in our Northern Italy community. Finally, we further divided food categories by sub-categories reported in the FFQ, thereby allowing estimation of As and Hg intake in individual species of fish and seafood.

Study limitations include misclassification of elemental species. While speciation analysis is of increasing scientific interest (Piras et al., 2015; Ruiz-de-Cenzano et al., 2017; Ruzik, 2012; Solovyev et al., 2017) due to the heterogeneous and sometimes opposite health effects of the various species (Karri et al., 2016; Michalke et al., 2017; Vinceti et al., 2012), we attempted to estimate intake of both inorganic and organic compounds. Unfortunately, we were unable to directly determine these single element species in foods, thus hampering their direct estimation in

food samples. We therefore used an indirect and conservative approach to assess elements fraction from total concentration according to the EFSA methodology (European Food Safety Authority, 2009, 2012), which may have induced misclassification (Bradley et al., 2017).

Since food collection was carried out mainly during winter and autumn, we could only partially account for possible seasonal variation of heavy metal content in food (Turrini et al., 2017; Vincevica-Gaile and Klavins, 2013). However, conflicting results have been reported about the existence of seasonal effects for several elements, specifically for As and Hg (Biswas et al., 2014; Copat et al., 2012; Nacano et al., 2014; Nevarez et al., 2015; Piras et al., 2015), and they have not been shown to occur for other elements (Agudo et al., 2002; Halkjaer et al., 2009; Olsen et al., 2009; Welch et al., 2009).

Finally, we assessed dietary intake in an adult population only, and our estimates may not apply to adolescents and children, due to differences in dietary habits, heavy metal metabolism, and toxic effects.

5.1 Conclusions

In a population-based study from Northern Italy, we provided detailed information about the foods most likely to contain appreciable amounts of As and Hg. In agreement with previous studies of individuals from industrialized nations, we found that fish and seafood contributed the highest levels of As and Hg to the human diet. Weekly dietary intake of As and Hg in our population did not appear to exceed the recommended safety limits set by EFSA.

Declaration of interest

All authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by a grant from the Italian Association for Cancer Research (AIRC) to Dr.

Marco Vinceti.

ACCEPTED MANUSCRIPT

Table 1. Distribution of arsenic (As) and mercury (Hg) concentrations in foods composing the usual diet of the study population.

Food (N)	As ($\mu\text{g/kg}$)			Hg ($\mu\text{g/kg}$)		
	50 th	IQR	<LOD n (%)	50 th	IQR	<LOD n (%)
Cereals (126)	6.01	(3.63 - 9.90)	1 (0.8)	0.78	(0.16 - 3.51)	17 (13.7)
Meat (86)	3.97	(2.73 - 8.23)	0 (0.0)	0.65	(0.18 - 1.80)	12 (14.0)
Milk and dairy products (72)	3.92	(1.43 - 8.64)	0 (0.0)	0.37	(0.14 - 0.92)	6 (8.3)
Eggs (9)	2.58	(1.45 - 8.29)	1 (11.1)	0.99	(0.12 - 2.79)	0 (0.0)
Fish and seafood (62)	1346.95	(686.1 - 3831.32)	0 (0.0)	55.03	(25.27 - 142.26)	0 (0.0)
Vegetables (193)	2.51	(0.94 - 6.59)	1 (0.5)	0.63	(0.18 - 2.06)	14 (7.3)
Legumes (42)	4.32	(3.74 - 8.24)	0 (0.0)	0.66	(0.14 - 3.65)	3 (7.1)
Potatoes (14)	2.61	(1.08 - 4.68)	0 (0.0)	0.36	(0.15 - 0.91)	1 (7.1)
Fresh fruits (65)	1.20	(0.62 - 3.33)	0 (0.0)	0.28	(0.06 - 0.60)	7 (10.8)
Dry fruits (39)	10.82	(3.81 - 30.67)	0 (0.0)	0.82	(0.33 - 3.71)	5 (12.8)
Sweets (64)	5.76	(2.50 - 8.90)	0 (0.0)	0.27	(0.01 - 0.99)	15 (23.4)
Oils and fats (22)	2.59	(1.76 - 10.82)	1 (4.5)	0.03	(0.00 - 0.19)	10 (45.5)
Beverages (96)	2.70	(0.89 - 7.77)	0 (0.0)	0.01	(0.00 - 0.04)	0 (0.0)

Table 2. Median (interquartile range - IQR) of daily dietary intake (g/day) for main food categories in the study population.

Food	Total (n=719)		Males (N=319)		Females (N=400)	
	50 th	IQR	50 th	IQR	50 th	IQR
Cereals	175.4	(120.6 - 245.5)	194.7	(129.5 - 267.4)	165.9	(113.9 - 221.5)
Meat	119.3	(79.5 - 164.3)	136.6	(92.1 - 174.3)	107.5	(71.8 - 150.6)
Milk and dairy products	185.0	(85.2 - 301.7)	161.3	(63.9 - 286.1)	209.8	(120.7 - 309.6)
Eggs	13.5	(7.3 - 21.2)	12.4	(7.0 - 20.0)	14.0	(7.6 - 21.4)
Fish and seafood	28.7	(15.9 - 45.7)	29.1	(16.1 - 48.2)	28.6	(15.7 - 44.8)
Vegetables	140.8	(96.3 - 210.8)	137.2	(99.2 - 204.3)	143.6	(92.2 - 212.6)
Legumes	13.6	(6.3 - 25.3)	14.2	(6.2 - 25.7)	13.5	(6.4 - 23.9)
Potatoes	18.0	(10.7 - 32.0)	18.1	(10.7 - 30.8)	17.9	(10.1 - 32.3)
Fresh fruits	302.0	(206.1 - 436.9)	284.0	(202.8 - 426.9)	318.7	(208.8 - 452.1)
Dry fruits	0.3	(0.2 - 1.6)	0.3	(0.2 - 1.8)	0.3	(0.2 - 1.6)
Sweets	68.7	(38.6 - 112.5)	65.5	(32.6 - 108.8)	74.2	(43.0 - 114.0)
Oils and fats	24.7	(18.4 - 33.4)	25.6	(18.9 - 34.5)	24.1	(17.9 - 32.7)
Beverages	290.7	(159.0 - 483.1)	360.7	(185.4 - 574.6)	239.9	(141.2 - 413.5)

Table 3. Weekly dietary intake of estimated total and inorganic arsenic (As) and mercury (Hg) in the study population. Values in $\mu\text{g}/\text{kg}$ of body weight per week. Median and interquartile range (IQR) are reported. Range of benchmark dose lower confidence limit (BMDL) values for 1% excess risk of cancers of the lung, skin and bladder, and skin lesions (BMDL₀₁) of 2.1-24 $\mu\text{g}/\text{kg}$ bw/week corresponding to 0.3-8 $\mu\text{g}/\text{kg}$ bw/day. Tolerable weekly intake (TWI) for inorganic mercury 4 $\mu\text{g}/\text{kg}$ bw/week.

Food	Total As		Inorg As	
	50 th	IQR	50 th	IQR
Cereals	0.195	(0.120 - 0.289)	0.195	(0.120 - 0.289)
Meat	0.046	(0.031 - 0.064)	0.023	(0.016 - 0.032)
Milk and dairy products	0.036	(0.021 - 0.053)	0.018	(0.011 - 0.027)
Eggs	0.003	(0.002 - 0.005)	0.002	(0.001 - 0.003)
Fish and seafood	4.686	(2.163 - 8.128)	0.094	(0.043 - 0.163)
Vegetables	0.036	(0.023 - 0.053)	0.018	(0.011 - 0.027)
Legumes	0.006	(0.003 - 0.011)	0.003	(0.001 - 0.005)
Potatoes	0.005	(0.003 - 0.008)	0.002	(0.001 - 0.004)
Fresh fruits	0.03	(0.018 - 0.044)	0.015	(0.009 - 0.022)
Dry fruits	0	(0.000 - 0.002)	0	(0.000 - 0.001)
Sweets	0.032	(0.017 - 0.057)	0.016	(0.009 - 0.028)
Oils and fats	0.007	(0.004 - 0.012)	0.003	(0.002 - 0.006)
Beverages	0.121	(0.071 - 0.187)	0.121	(0.071 - 0.187)
Total	5.337	(2.749 - 8.835)	0.578	(0.435 - 0.749)

	Total Hg		Inorg Hg	
	50 th	IQR	50 th	IQR
Cereals	0.024	(0.015 - 0.035)	0.024	(0.015 - 0.035)
Meat	0.005	(0.003 - 0.008)	0.005	(0.003 - 0.008)
Milk and dairy products	0.004	(0.002 - 0.006)	0.004	(0.002 - 0.006)
Eggs	0.001	(0.001 - 0.002)	0.001	(0.001 - 0.002)
Fish and seafood	0.319	(0.178 - 0.511)	0.064	(0.036 - 0.102)
Vegetables	0.017	(0.011 - 0.026)	0.017	(0.011 - 0.026)
Legumes	0.001	(0.000 - 0.002)	0.001	(0.000 - 0.002)
Potatoes	0.001	(0.000 - 0.001)	0.001	(0.000 - 0.001)
Fresh fruits	0.006	(0.004 - 0.01)	0.006	(0.004 - 0.01)
Dry fruits	0	-	0	-
Sweets	0.002	(0.001 - 0.004)	0.002	(0.001 - 0.004)
Oils and fats	0	-	0	-
Beverages	0.003	(0.002 - 0.004)	0.003	(0.002 - 0.004)
Total	0.391	(0.248 - 0.588)	0.139	(0.102 - 0.188)

Table 4. Weekly dietary intake of estimated total and inorganic arsenic (As) and mercury (Hg) in male study population. Values in $\mu\text{g/kg}$ of body weight (bw) per week. Median and interquartile range (IQR) are reported. Range of benchmark dose lower confidence limit values for 1% excess risk of cancers of the lung, skin and bladder, and skin lesions of 2.1-24 $\mu\text{g/kg}$ bw/week corresponding to 0.3-8 $\mu\text{g/kg}$ bw/day. Tolerable weekly intake for inorganic Hg=4 $\mu\text{g/kg}$ bw/week.

Food	Total As		Inorganic As	
	50 th	IQR	50 th	IQR
Cereals	0.191	(0.120 - 0.290)	0.191	(0.120 - 0.290)
Meat	0.047	(0.032 - 0.064)	0.023	(0.016 - 0.032)
Milk and dairy products	0.030	(0.017 - 0.046)	0.015	(0.008 - 0.023)
Eggs	0.003	(0.002 - 0.005)	0.001	(0.001 - 0.002)
Fish and seafood	4.279	(1.852 - 7.684)	0.086	(0.037 - 0.154)
Vegetables	0.030	(0.020 - 0.046)	0.015	(0.010 - 0.023)
Legumes	0.005	(0.002 - 0.010)	0.003	(0.001 - 0.005)
Potatoes	0.004	(0.002 - 0.007)	0.002	(0.001 - 0.004)
Fresh fruits	0.025	(0.016 - 0.037)	0.013	(0.008 - 0.019)
Dry fruits	0	(0.000 - 0.002)	0	(0.000 - 0.001)
Sweets	0.027	(0.012 - 0.049)	0.013	(0.006 - 0.024)
Oils and fats	0.007	(0.004 - 0.011)	0.003	(0.002 - 0.006)
Beverages	0.127	(0.077 - 0.199)	0.127	(0.077 - 0.199)
Total	4.793	(2.345 - 8.364)	0.571	(0.423 - 0.725)


	Total Hg		Inorganic Hg	
	50 th	IQR	50 th	IQR
Cereals	0.023	(0.013 - 0.033)	0.023	(0.013 - 0.033)
Meat	0.005	(0.003 - 0.007)	0.005	(0.003 - 0.007)
Milk and dairy products	0.003	(0.002 - 0.005)	0.003	(0.002 - 0.005)
Eggs	0.001	(0.001 - 0.002)	0.001	(0.001 - 0.002)
Fish and seafood	0.287	(0.158 - 0.457)	0.057	(0.032 - 0.091)
Vegetables	0.015	(0.010 - 0.024)	0.015	(0.010 - 0.024)
Legumes	0.001	(0.000 - 0.002)	0.001	(0.000 - 0.002)
Potatoes	0.001	(0.000 - 0.001)	0.001	(0.000 - 0.001)
Fresh fruits	0.006	(0.003 - 0.008)	0.006	(0.003 - 0.008)
Dry fruits	0	-	0	-
Sweets	0.002	(0.001 - 0.004)	0.002	(0.001 - 0.004)
Oils and fats	0	-	0	-
Beverages	0.003	(0.001 - 0.004)	0.003	(0.001 - 0.004)
Total	0.352	(0.222 - 0.54)	0.128	(0.096 - 0.167)

Table 5. Weekly dietary intake of estimated total and inorganic arsenic (As) and mercury (Hg) in female study population. Values in $\mu\text{g}/\text{kg}$ of body weight (bw) per week. Median and interquartile range (IQR) are reported. Range of benchmark dose lower confidence limit values for 1% excess risk of cancers of the lung, skin and bladder, and skin lesions of 2.1-24 $\mu\text{g}/\text{kg}$ bw/week corresponding to 0.3-8 $\mu\text{g}/\text{kg}$ bw/day. Tolerable weekly intake for inorganic Hg=4 $\mu\text{g}/\text{kg}$ bw/week.

Food	Total As		Inorganic As	
	50 th	IQR	50 th	IQR
Cereals	0.201	(0.120 - 0.288)	0.201	(0.120 - 0.288)
Meat	0.046	(0.031 - 0.064)	0.023	(0.015 - 0.032)
Milk and dairy products	0.043	(0.027 - 0.058)	0.022	(0.013 - 0.029)
Eggs	0.004	(0.002 - 0.006)	0.002	(0.001 - 0.003)
Fish and seafood	5.088	(2.422 - 8.531)	0.102	(0.048 - 0.171)
Vegetables	0.041	(0.026 - 0.061)	0.020	(0.013 - 0.031)
Legumes	0.006	(0.003 - 0.011)	0.003	(0.002 - 0.006)
Potatoes	0.005	(0.003 - 0.008)	0.003	(0.001 - 0.004)
Fresh fruits	0.035	(0.021 - 0.049)	0.017	(0.011 - 0.025)
Dry fruits	0	(0.000 - 0.002)	0	(0.000 - 0.001)
Sweets	0.036	(0.020 - 0.062)	0.018	(0.010 - 0.031)
Oils and fats	0.007	(0.004 - 0.012)	0.003	(0.002 - 0.006)
Beverages	0.116	(0.067 - 0.180)	0.116	(0.067 - 0.180)
Total	5.761	(2.957 - 9.159)	0.583	(0.454 - 0.781)

	Total Hg		Inorganic Hg	
	50 th	IQR	50 th	IQR
Cereals	0.025	(0.017 - 0.036)	0.025	(0.017 - 0.036)
Meat	0.005	(0.003 - 0.008)	0.005	(0.003 - 0.008)
Milk and dairy products	0.004	(0.003 - 0.006)	0.004	(0.003 - 0.006)
Eggs	0.001	(0.001 - 0.002)	0.001	(0.001 - 0.002)
Fish and seafood	0.345	(0.192 - 0.556)	0.069	(0.038 - 0.111)
Vegetables	0.019	(0.013 - 0.029)	0.019	(0.013 - 0.029)
Legumes	0.001	(0.000 - 0.002)	0.001	(0.000 - 0.002)
Potatoes	0.001	(0.000 - 0.001)	0.001	(0.000 - 0.001)
Fresh fruits	0.008	(0.005 - 0.011)	0.008	(0.005 - 0.011)
Dry fruits	0	-	0	-
Sweets	0.003	(0.001 - 0.005)	0.003	(0.001 - 0.005)
Oils and fats	0	-	0	-
Beverages	0.003	(0.002 - 0.005)	0.003	(0.002 - 0.005)
Total	0.423	(0.264 - 0.621)	0.149	(0.111 - 0.200)

Figure 1. Trace elements content (white bars) according to fish and seafood category and % contribution to trace element daily intake (gray bars) in study population.



References

- Adel, M., Oliveri Conti, G., Dadar, M., Mahjoub, M., Copat, C., Ferrante, M., 2016. Heavy metal concentrations in edible muscle of whitecheek shark, *Carcharhinus dussumieri* (*elasmobranchii, chondrichthyes*) from the Persian Gulf: A food safety issue. *Food Chem Toxicol* 97, 135-140.
- Agudo, A., Slimani, N., Ocke, M.C., Naska, A., Miller, A.B., Kroke, A., Bamia, C., Karalis, D., Vineis, P., Palli, D., Bueno-de-Mesquita, H.B., Peeters, P.H., Engeset, D., Hjartaker, A., Navarro, C., Martinez Garcia, C., Wallstrom, P., Zhang, J.X., Welch, A.A., Spencer, E., Stripp, C., Overvad, K., Clavel-Chapelon, F., Casagrande, C., Riboli, E., 2002. Consumption of vegetables, fruit and other plant foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Public Health Nutr* 5, 1179-1196.
- Alava, J.J., Cheung, W.W.L., Ross, P.S., Sumaila, U.R., 2017. Climate change-contaminant interactions in marine food webs: Toward a conceptual framework. *Glob Chang Biol* 23, 3984-4001.
- Arslan, B., Djamgoz, M.B.A., Akun, E., 2017. Arsenic: A review on exposure pathways, accumulation, mobility and transmission into the human food chain. *Rev Environ Contam Toxicol* 243, 27-51.
- ATSDR, 2012a. Toxicological profile for arsenic, in: U.S. Department of Health and Human Services - Public Health Service (Ed.). Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- ATSDR, 2012b. Toxicological profile for mercury, in: U.S. Department of Health and Human Services - Public Health Service (Ed.). Agency for Toxic Substances and Disease Registry, Atlanta, GA.
- Bae, H.S., Kang, I.G., Lee, S.G., Eom, S.Y., Kim, Y.D., Oh, S.Y., Kwon, H.J., Park, K.S., Kim, H., Choi, B.S., Yu, I.J., Park, J.D., 2017. Arsenic exposure and seafood intake in Korean adults. *Hum Exp Toxicol* 36, 451-460.
- Barone, G., Storelli, A., Garofalo, R., Busco, V.P., Quaglia, N.C., Centrone, G., Storelli, M.M., 2015. Assessment of mercury and cadmium via seafood consumption in Italy: estimated dietary intake (EWI) and target hazard quotient (THQ). *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 32, 1277-1286.
- Biswas, A., Deb, D., Ghose, A., Santra, S.C., Guha Mazumder, D.N., 2014. Seasonal perspective of dietary arsenic consumption and urine arsenic in an endemic population. *Environ Monit Assess* 186, 4543-4551.
- Bjorklund, G., Dadar, M., Mutter, J., Aaseth, J., 2017. The toxicology of mercury: Current research and emerging trends. *Environ Res* 159, 545-554.
- Bosch, A.C., O'Neill, B., Sigge, G.O., Kerwath, S.E., Hoffman, L.C., 2016. Heavy metals in marine fish meat and consumer health: a review. *J Sci Food Agric* 96, 32-48.
- Bottecchi, I., Vinceti, M., 2012. [Trace elements in foods. Analysis of diet of Reggio Emilia and Modena], Department of Public Health. University of Modena and Reggio Emilia, p. 124.
- Bradley, M.A., Barst, B.D., Basu, N., 2017. A Review of Mercury Bioavailability in Humans and Fish. *Int J Environ Res Public Health* 14.
- Brambilla, G., Abete, M.C., Binato, G., Chiaravalle, E., Cossu, M., Dellatte, E., Miniero, R., Orletti, R., Piras, P., Roncarati, A., Ubaldi, A., Chessa, G., 2013. Mercury occurrence in Italian seafood from the Mediterranean Sea and possible intake scenarios of the Italian coastal population. *Regul Toxicol Pharmacol* 65, 269-277.
- Burger, J., Gochfeld, M., 2005. Heavy metals in commercial fish in New Jersey. *Environ Res* 99, 403-412.

- Calatayud, M., Devesa, V., Virseda, J.R., Barbera, R., Montoro, R., Velez, D., 2012. Mercury and selenium in fish and shellfish: occurrence, bioaccessibility and uptake by Caco-2 cells. *Food Chem Toxicol* 50, 2696-2702.
- Cano-Sancho, G., Perello, G., Maulvault, A.L., Marques, A., Nadal, M., Domingo, J.L., 2015a. Oral bioaccessibility of arsenic, mercury and methylmercury in marine species commercialized in Catalonia (Spain) and health risks for the consumers. *Food Chem Toxicol* 86, 34-40.
- Cano-Sancho, G., Sioen, I., Vandermeersch, G., Jacobs, S., Robbens, J., Nadal, M., Domingo, J.L., 2015b. Integrated risk index for seafood contaminants (IRISC): Pilot study in five European countries. *Environ Res* 143, 109-115.
- Chiocchetti, G., Jadan-Piedra, C., Velez, D., Devesa, V., 2017. Metal(loid) contamination in seafood products. *Crit Rev Food Sci Nutr* 57, 3715-3728.
- Cilloni, S., Malavolti, M., Malagoli, C., Violi, F., Filippini, T., Vescovi, L., Vinceti, M., 2017. Selenium, zinc and copper content of foods consumed in an Italian community. *J Trace Elem Med Biol* 41S, 59.
- Cirillo, T., Fasano, E., Viscardi, V., Arnese, A., Amodio-Cocchieri, R., 2010. Survey of lead, cadmium, mercury and arsenic in seafood purchased in Campania, Italy. *Food Addit Contam Part B Surveill* 3, 30-38.
- Copat, C., Arena, G., Fiore, M., Ledda, C., Fallico, R., Sciacca, S., Ferrante, M., 2013. Heavy metals concentrations in fish and shellfish from eastern Mediterranean Sea: Consumption advisories. *Food Chem Toxicol* 53, 33-37.
- Copat, C., Brundo, M.V., Arena, G., Grasso, A., Oliveri Conti, G., Ledda, C., Fallico, R., Sciacca, S., Ferrante, M., 2012. Seasonal variation of bioaccumulation in *Engraulis encrasicolus* (Linnaeus, 1758) and related biomarkers of exposure. *Ecotoxicol Environ Saf* 86, 31-37.
- Copat, C., Vinceti, M., D'Agati, M.G., Arena, G., Mauceri, V., Grasso, A., Fallico, R., Sciacca, S., Ferrante, M., 2014. Mercury and selenium intake by seafood from the Ionian Sea: A risk evaluation. *Ecotoxicol Environ Saf* 100, 87-92.
- Delpla, I., Jung, A.V., Baures, E., Clement, M., Thomas, O., 2009. Impacts of climate change on surface water quality in relation to drinking water production. *Environ Int* 35, 1225-1233.
- Domingo, J.L., Bocio, A., Falco, G., Llobet, J.M., 2007. Benefits and risks of fish consumption Part I. A quantitative analysis of the intake of omega-3 fatty acids and chemical contaminants. *Toxicology* 230, 219-226.
- Donald, D.B., Wissel, B., Anas, M.U., 2015. Species-specific mercury bioaccumulation in a diverse fish community. *Environ Toxicol Chem* 34, 2846-2855.
- Donaldson, S.G., Van Oostdam, J., Tikhonov, C., Feeley, M., Armstrong, B., Ayotte, P., Boucher, O., Bowers, W., Chan, L., Dallaire, F., Dallaire, R., Dewailly, E., Edwards, J., Egeland, G.M., Fontaine, J., Furgal, C., Leech, T., Loring, E., Muckle, G., Nancarrow, T., Pereg, D., Plusquellec, P., Potyrala, M., Receveur, O., Shearer, R.G., 2010. Environmental contaminants and human health in the Canadian Arctic. *Sci Total Environ* 408, 5165-5234.
- Duh, J.D., Shandas, V., Chang, H., George, L.A., 2008. Rates of urbanisation and the resiliency of air and water quality. *Sci Total Environ* 400, 238-256.
- European Food Safety Authority, 2009. Arsenic in Food. *EFSA Journal* 7, 1351.
- European Food Safety Authority, 2012. Mercury and methylmercury in food. *EFSA Journal* 10, 2985.
- European Food Safety Authority, 2014. Scientific Opinion on health benefits of seafood (fish and shellfish) consumption in relation to health risks associated with exposure to methylmercury. *EFSA Journal* 12, 3761-n/a.
- FAO/WHO, 2010a. Evaluation of certain food additives, Geneva, 8–17 June 2010. Seventy-third Meeting.


- FAO/WHO, 2010b. Joint FAO/WHO expert consultation on the risks and benefits of fish consumption, 25-29 January 2010, Rome, Italy.
- Filippini, T., Cilloni, S., Malavolti, M., Violi, F., Malagoli, C., Tesauro, M., Bottecchi, I., Ferrari, A., Vescovi, L., Vinceti, M., 2018. Dietary intake of cadmium, chromium, copper, manganese, selenium and zinc in a Northern Italy community. *J Trace Elem Med Biol* 2018.
- Filippini, T., Ferrari, A., Michalke, B., Grill, P., Vescovi, L., Salvia, C., Malagoli, C., Malavolti, M., Sieri, S., Krogh, V., Bargellini, A., Martino, A., Ferrante, M., Vinceti, M., 2017. Toenail selenium as an indicator of environmental exposure: A cross-sectional study. *Mol Med Rep* 15, 3405-3412.
- Gonzalez-Alzaga, B., Lacasana, M., Hernandez, A.F., Arrebola, J.P., Lopez-Flores, I., Artacho-Cordon, F., Bonde, J.P., Olea, N., Aguilar-Garduno, C., 2017. Serum concentrations of organochlorine compounds and predictors of exposure in children living in agricultural communities from South-Eastern Spain. *Environ Pollut*.
- Halkjaer, J., Olsen, A., Bjerregaard, L.J., Deharveng, G., Tjonneland, A., Welch, A.A., Crowe, F.L., Wirfalt, E., Hellstrom, V., Niravong, M., Touvier, M., Linseisen, J., Steffen, A., Ocke, M.C., Peeters, P.H., Chirlaque, M.D., Larranaga, N., Ferrari, P., Contiero, P., Frasca, G., Engeset, D., Lund, E., Misirli, G., Kostis, M., Riboli, E., Slimani, N., Bingham, S., 2009. Intake of total, animal and plant proteins, and their food sources in 10 countries in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* 63 Suppl 4, S16-36.
- Hightower, J.M., Moore, D., 2003. Mercury levels in high-end consumers of fish. *Environ Health Perspect* 111, 604-608.
- Karri, V., Schuhmacher, M., Kumar, V., 2016. Heavy metals (Pb, Cd, As and MeHg) as risk factors for cognitive dysfunction: A general review of metal mixture mechanism in brain. *Environ Toxicol Pharmacol* 48, 203-213.
- Kaya, G., Turkoglu, S., 2017. Bioaccumulation of heavy metals in various tissues of some fish species and green tiger shrimp (*Penaeus semisulcatus*) from Iskenderun Bay, Turkey, and risk assessment for human health. *Biol Trace Elem Res* 180, 314-326.
- Liu, Y., Buchanan, S., Anderson, H.A., Xiao, Z., Persky, V., Turyk, M.E., 2018. Association of methylmercury intake from seafood consumption and blood mercury level among the Asian and Non-Asian populations in the United States. *Environ Res* 160, 212-222.
- Llobet, J.M., Falco, G., Casas, C., Teixido, A., Domingo, J.L., 2003. Concentrations of arsenic, cadmium, mercury, and lead in common foods and estimated daily intake by children, adolescents, adults, and seniors of Catalonia, Spain. *J Agric Food Chem* 51, 838-842.
- Malagoli, C., Malavolti, M., Agnoli, C., Crespi, C.M., Fiorentini, C., Farnetani, F., Longo, C., Ricci, C., Albertini, G., Lanzoni, A., Veneziano, L., Virgili, A., Pagliarello, C., Santini, M., Fanti, P.A., Dika, E., Sieri, S., Krogh, V., Pellacani, G., Vinceti, M., 2015. Diet quality and risk of melanoma in an Italian population. *J Nutr* 145, 1800-1807.
- Malavolti, M., Malagoli, C., Fiorentini, C., Longo, C., Farnetani, F., Ricci, C., Albertini, G., Lanzoni, A., Reggiani, C., Virgili, A., Pagliarello, C., Santini, M., Fanti, P.A., Dika, E., Sieri, S., Krogh, V., Pellacani, G., Vinceti, M., 2013. Association between dietary vitamin C and risk of cutaneous melanoma in a population of Northern Italy. *Int J Vitam Nutr Res* 83, 291-298.
- Marti-Cid, R., Llobet, J.M., Castell, V., Domingo, J.L., 2008. Dietary intake of arsenic, cadmium, mercury, and lead by the population of Catalonia, Spain. *Biol Trace Elem Res* 125, 120-132.
- Martorell, I., Perello, G., Marti-Cid, R., Llobet, J.M., Castell, V., Domingo, J.L., 2011. Human exposure to arsenic, cadmium, mercury, and lead from foods in Catalonia, Spain: temporal trend. *Biol Trace Elem Res* 142, 309-322.
- Marzec, Z., Schlegel-Zawadzka, M., 2004. Exposure to cadmium, lead and mercury in the adult population from Eastern Poland, 1990-2002. *Food Addit Contam* 21, 963-970.

- Matos, J., Lourenco, H.M., Brito, P., Maulvault, A.L., Martins, L.L., Afonso, C., 2015. Influence of bioaccessibility of total mercury, methyl-mercury and selenium on the risk/benefit associated to the consumption of raw and cooked blue shark (*Prionace glauca*). *Environ Res* 143, 123-129.
- Maulvault, A.L., Machado, R., Afonso, C., Lourenco, H.M., Nunes, M.L., Coelho, I., Langerholc, T., Marques, A., 2011. Bioaccessibility of Hg, Cd and As in cooked black scabbard fish and edible crab. *Food Chem Toxicol* 49, 2808-2815.
- Michalke, B., Willkommen, D., Drobyshchev, E., Solovyev, N., 2017. The importance of speciation analysis in neurodegeneration research. *Trends Analyt Chem* (in press).
- Miklavcic, A., Casetta, A., Snoj Tratnik, J., Mazej, D., Krsnik, M., Mariuz, M., Sofianou, K., Spiric, Z., Barbone, F., Horvat, M., 2013. Mercury, arsenic and selenium exposure levels in relation to fish consumption in the Mediterranean area. *Environ Res* 120, 7-17.
- Nacano, L.R., de Freitas, R., Barbosa, F., Jr., 2014. Evaluation of seasonal dietary exposure to arsenic, cadmium and lead in schoolchildren through the analysis of meals served by public schools of Ribeirao Preto, Brazil. *J Toxicol Environ Health A* 77, 367-374.
- Nevarez, M., Leal, L.O., Moreno, M., 2015. Estimation of seasonal risk caused by the intake of lead, mercury and cadmium through freshwater fish consumption from urban water reservoirs in arid areas of northern Mexico. *Int J Environ Res Public Health* 12, 1803-1816.
- Nunez, R., Garcia, M.A., Alonso, J., Melgar, M.J., 2018. Arsenic, cadmium and lead in fresh and processed tuna marketed in Galicia (NW Spain): Risk assessment of dietary exposure. *Sci Total Environ* 627, 322-331.
- Olsen, A., Halkjaer, J., van Gils, C.H., Buijsse, B., Verhagen, H., Jenab, M., Boutron-Ruault, M.C., Ericson, U., Ocke, M.C., Peeters, P.H., Touvier, M., Niravong, M., Waaseth, M., Skeie, G., Khaw, K.T., Travis, R., Ferrari, P., Sanchez, M.J., Agudo, A., Overvad, K., Linseisen, J., Weikert, C., Sacerdote, C., Evangelista, A., Zylis, D., Tsiotas, K., Manjer, J., van Guelpen, B., Riboli, E., Slimani, N., Bingham, S., 2009. Dietary intake of the water-soluble vitamins B1, B2, B6, B12 and C in 10 countries in the European Prospective Investigation into Cancer and Nutrition. *Eur J Clin Nutr* 63 Suppl 4, S122-149.
- Pal, A., He, Y., Jekel, M., Reinhard, M., Gin, K.Y., 2014. Emerging contaminants of public health significance as water quality indicator compounds in the urban water cycle. *Environ Int* 71, 46-62.
- Pala, V., Sieri, S., Palli, D., Salvini, S., Berrino, F., Bellegotti, M., Frasca, G., Tumino, R., Sacerdote, C., Fiorini, L., Celentano, E., Galasso, R., Krogh, V., 2003. Diet in the Italian EPIC cohorts: presentation of data and methodological issues. *Tumori* 89, 594-607.
- Papadopoulos, A., Sioen, I., Cubadda, F., Ozer, H., Basegmez, H.I., Turrini, A., Lopez Esteban, M.T., Fernandez San Juan, P.M., Sokolic-Mihalak, D., Jurkovic, M., De Henauw, S., Aureli, F., Vin, K., Sirot, V., 2015. TDS exposure project: application of the analytic hierarchy process for the prioritization of substances to be analyzed in a total diet study. *Food Chem Toxicol* 76, 46-53.
- Pasanisi, P., Berrino, F., Bellati, C., Sieri, S., Krogh, V., 2002. Validity of the Italian EPIC questionnaire to assess past diet. *IARC Sci Publ* 156, 41-44.
- Perello, G., Llobet, J.M., Gomez-Catalan, J., Castell, V., Centrich, F., Nadal, M., Domingo, J.L., 2014. Human health risks derived from dietary exposure to toxic metals in Catalonia, Spain: temporal trend. *Biol Trace Elem Res* 162, 26-37.
- Perello, G., Vicente, E., Castell, V., Llobet, J.M., Nadal, M., Domingo, J.L., 2015. Dietary intake of trace elements by the population of Catalonia (Spain): results from a total diet study. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 32, 748-755.
- Piras, P., Orletti, R., Chessa, G., Carloni, C., Griffoni, F., Palombo, P., Velieri, F., 2015. Arsenic speciation in fish products and seafood as a prerequisite for proper risk assessment. *Ital J Food Saf* 4, 4577.

- Ruiz-de-Cenzano, M., Rochina-Marco, A., Cervera, M.L., de la Guardia, M., 2017. Evaluation of the content of antimony, arsenic, bismuth, selenium, tellurium and their inorganic forms in commercially baby foods. *Biol Trace Elem Res* 180, 355-365.
- Ruzik, L., 2012. Speciation of challenging elements in food by atomic spectrometry. *Talanta* 93, 18-31.
- Sattar, A., Xie, S., Hafeez, M.A., Wang, X., Hussain, H.I., Iqbal, Z., Pan, Y., Iqbal, M., Shabbir, M.A., Yuan, Z., 2016. Metabolism and toxicity of arsenicals in mammals. *Environ Toxicol Pharmacol* 48, 214-224.
- Solovyev, N., Vinceti, M., Grill, P., Mandrioli, J., Michalke, B., 2017. Redox speciation of iron, manganese, and copper in cerebrospinal fluid by strong cation exchange chromatography - sector field inductively coupled plasma mass spectrometry. *Anal Chim Acta* 973, 25-33.
- Storelli, M.M., Barone, G., 2013. Toxic metals (Hg, Pb, and Cd) in commercially important demersal fish from Mediterranean sea: contamination levels and dietary exposure assessment. *J Food Sci* 78, T362-366.
- Taylor, C.M., Golding, J., Emond, A.M., 2016. Blood mercury levels and fish consumption in pregnancy: Risks and benefits for birth outcomes in a prospective observational birth cohort. *Int J Hyg Environ Health* 219, 513-520.
- Turrini, A., Lombardi-Boccia, G., Aureli, F., Cubadda, F., D'Addezio, L., D'Amato, M., D'Evoli, L., Darnerud, P., Devlin, N., Dias, M.G., Jurkovic, M., Kelleher, C., Le Donne, C., Lopez Esteban, M., Lucarini, M., Martinez Burgos, M.A., Martinez-Victoria, E., McNulty, B., Mistura, L., Nugent, A., Oktay Basegmez, H.I., Oliveira, L., Ozer, H., Perello, G., Pite, M., Presser, K., Sokolic, D., Vasco, E., Volatier, J.L., 2017. A conceptual framework for the collection of food products in a Total Diet Study. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess*, 1-20.
- Turrini, A., Saba, A., Perrone, D., Cialfa, E., D'Amicis, A., 2001. Food consumption patterns in Italy: the INN-CA Study 1994-1996. *Eur J Clin Nutr* 55, 571-588.
- Vilavert, L., Borrell, F., Nadal, M., Jacobs, S., Minnens, F., Verbeke, W., Marques, A., Domingo, J.L., 2017. Health risk/benefit information for consumers of fish and shellfish: FishChoice, a new online tool. *Food Chem Toxicol* 104, 79-84.
- Vinceti, M., Bottecchi, I., Fan, A., Finkelstein, Y., Mandrioli, J., 2012. Are environmental exposures to selenium, heavy metals, and pesticides risk factors for amyotrophic lateral sclerosis? *Rev Environ Health* 27, 19-41.
- Vinceti, M., Crespi, C.M., Bonvicini, F., Malagoli, C., Ferrante, M., Marmiroli, S., Stranges, S., 2013. The need for a reassessment of the safe upper limit of selenium in drinking water. *Sci Total Environ* 443, 633-642.
- Vinceti, M., Malagoli, C., Fiorentini, C., Longo, C., Crespi, C.M., Albertini, G., Ricci, C., Lanzoni, A., Reggiani, M., Virgili, A., Osti, F., Lombardi, M., Santini, M., Fanti, P.A., Dika, E., Sieri, S., Krogh, V., Seidenari, S., Pellacani, G., 2011. Inverse association between dietary vitamin D and risk of cutaneous melanoma in a northern Italy population. *Nutr Cancer* 63, 506-513.
- Vincevica-Gaile, Z., Klavins, M., 2013. Concentration of elements in food: How can it reflect impact of environmental and other influencing factors? *Environmental and Climate Technologies* 12, 15-19.
- Welch, A.A., Fransen, H., Jenab, M., Boutron-Ruault, M.C., Tumino, R., Agnoli, C., Ericson, U., Johansson, I., Ferrari, P., Engeset, D., Lund, E., Lentjes, M., Key, T., Touvier, M., Niravong, M., Larranaga, N., Rodriguez, L., Ocke, M.C., Peeters, P.H., Tjonneland, A., Bjerregaard, L., Vasilopoulou, E., Dilis, V., Linseisen, J., Nothlings, U., Riboli, E., Slimani, N., Bingham, S., 2009. Variation in intakes of calcium, phosphorus, magnesium, iron and potassium in 10 countries in the European Prospective Investigation into Cancer and Nutrition study. *Eur J Clin Nutr* 63 Suppl 4, S101-121.

WHO, 2017. Guidelines for drinking-water quality: Fourth Edition, Geneva.

ACCEPTED MANUSCRIPT



Highlights:

- We assessed arsenic and mercury contamination levels in foods composing dietary habits of Northern Italy community
- Sardine and sole species showed higher concentration of arsenic, while swordfish and tuna of mercury
- Estimated dietary intake of both trace elements were below the EFSA recommended tolerable intake