

This is the peer reviewed version of the following article:

The natural pseudo-distance as a quotient pseudo-metric, and applications / Cagliari, F.; Di Fabio, B.; Landi, Claudia. - In: FORUM MATHEMATICUM. - ISSN 0933-7741. - STAMPA. - 27:3(2015), pp. 1729-1742. [10.1515/forum-2012-0152]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

21/12/2025 06:11

THE NATURAL PSEUDO-DISTANCE AS A QUOTIENT PSEUDO-METRIC, AND APPLICATIONS

F. CAGLIARI, B. DI FABIO, AND C. LANDI

ABSTRACT. The natural pseudo-distance is a similarity measure conceived for the purpose of comparing shapes. In this paper we revisit this pseudo-distance from the point of view of quotients. In particular, we show that the natural pseudo-distance coincides with the quotient pseudo-metric on the space of continuous functions on a compact manifold, endowed with the uniform convergence metric, modulo self-homeomorphisms of the manifold. As applications of this result, the natural pseudo-distance is shown to be actually a metric on a number of function subspaces such as the space of topological embeddings, of isometries, and of simple Morse functions on surfaces.

1. INTRODUCTION

The natural pseudo-distance was introduced by Frosini and Mulazzani in [13], and further studied in [8, 9, 10, 12], as a measure of similarity that behaves nicely when invariance to deformations or different poses of the compared objects is a key requirement. These contributions stemmed from the actual need for these sort of similarity measures in pattern recognition to cope with the matching of natural or articulated objects. In general, such measures are reckoned to be beneficial for the organization of the huge collections of digital models produced nowadays through massive data acquisitions and shape modeling. In recent years, the development and study of topology-invariant metrics with stability properties has widely increased, as the numerous studies on similarity of non-rigid shapes testify (cf., e.g., [2, 6]).

In many practical applications, data are endowed with measurements on their points. For instance, in applications to shape comparison, recognition, and retrieval, the measurements can correspond to perceptions (e.g., height, depth, curvature, or color). Therefore, it is natural to model data as manifolds endowed with \mathbb{R}^n -valued functions defined on them [1].

The natural pseudo-distance is usually defined on the space $\mathcal{C}(M, \mathbb{R}^n)$ of \mathbb{R}^n -valued continuous functions with domain a compact manifold M in the following way. Assuming \mathbb{R}^n endowed with the usual maximum norm: $\|(x_1, x_2, \dots, x_n)\|_\infty = \max_{1 \leq i \leq n} |x_i|$, for $f, g \in \mathcal{C}(M, \mathbb{R}^n)$,

$$\delta(f, g) = \inf_{h \in \mathcal{H}(M)} \max_{p \in M} \|f(p) - g \circ h(p)\|_\infty,$$

$\mathcal{H}(M)$ being the set of self-homeomorphisms of M .

Starting from an idea presented in [4], the aim of this paper is to put the natural pseudo-distance in context with the classical notion of quotient pseudo-metric. We think that this link between the natural pseudo-distance and quotients is not only interesting per se, but can also enable to derive new results. For example, a result that, for the case of curves, was obtained in [5] using a constructive technique, is proved here for surfaces by indirect arguments. Indeed, as a further contribution of this paper, we show that for simple Morse functions on a surface, the natural

2010 *Mathematics Subject Classification.* Primary 54C35, 57S25, 18A32; secondary 68U05, 54E35.
Key words and phrases. Function space, Morse function, homeomorphism group action, isometry, shape comparison.

pseudo-distance is actually a distance, i.e. it distinguishes non-isometric surfaces (Subsection 4.3). Besides this result, we also prove that the natural pseudo-distance turns to a distance when defined on subspaces of $\mathcal{C}(M, \mathbb{R}^n)$ such as the spaces of embeddings and immersions (Subsection 4.1), and on quotient spaces induced by compact subgroups of $\mathcal{H}(M)$ (Subsection 4.2).

In perspective, we hope that other well-known properties of quotient pseudo-metrics will turn useful for the study of the natural pseudo-distance.

2. QUOTIENTS OF PSEUDO-METRIC SPACES

In this section we review the notion of a quotient pseudo-metric. More details can be found in [3] where the quotient pseudo-metric is called the quotient semi-metric.

Let us consider the category **PMet** of pseudo-metric spaces and non-expansive maps, i.e. functions between pseudo-metric spaces that do not increase distances: A map $h : (Z, d) \rightarrow (Z', d')$ is *non-expansive* if, for all $z_1, z_2 \in Z$,

$$d'(h(z_1), h(z_2)) \leq d(z_1, z_2).$$

A map h between pseudo-metric spaces is an isometry if and only if it is non-expansive, it is a bijection, and its inverse is also non-expansive. In particular, since non-expansive maps are always continuous, any isometry in **PMet** is a homeomorphism.

Non-expansive maps are the suitable maps between pseudo-metric spaces to pass to quotients. Indeed, if (Z, d) is a pseudo-metric space and \sim is an equivalence relation on Z , the quotient set Z/\sim can be endowed with the following pseudo-distance: Given two equivalence classes $[z]$ and $[y]$, the *quotient pseudo-metric* is defined by

$$d_\sim([z], [y]) = \inf \left\{ \sum_{i=1}^n d(z_i, y_i) \right\}$$

where the infimum is taken over all finite sequences (z_1, z_2, \dots, z_n) and (y_1, y_2, \dots, y_n) with $[z_1] = [z], [y_{i-1}] = [z_i], \dots, [y_n] = [y], i = 2, \dots, n$ (see [3, Def. 3.1.12]).

The quotient pseudo-distance d_\sim is characterized by the following universal property. If $h : (Z, d) \rightarrow (Z', d')$ is a non-expansive map between pseudo-metric spaces such that $h(z) = h(y)$ whenever $z \sim y$, then the induced quotient map $h_\sim : (Z/\sim, d_\sim) \rightarrow (Z', d')$ is non-expansive, that is the following diagram commutes in **PMet**:

$$\begin{array}{ccc} Z & \xrightarrow{\pi} & Z/\sim \\ & \searrow h & \downarrow h_\sim \\ & & Z' \end{array}$$

In general the quotient topology induced by \sim is different from the topology induced by the quotient pseudo-distance d_\sim . However, when the equivalence classes are the orbits of the action of a group of isometries the following result holds.

Theorem 2.1. *If (Z, d) is a pseudo-metric space endowed with an equivalence relation where the equivalence classes are the orbits of the action of a group of isometries on (Z, d) , then*

- (i) $d_\sim([z], [y]) = \inf \{d(z', y') : z' \sim z; y' \sim y\}$.
- (ii) *The topology induced by the quotient pseudo-metric coincides with the quotient topology.*
- (iii) *d_\sim is a metric if and only if the topology it induces is T_0 .*
- (iv) *d_\sim is a metric if and only if the orbits of the action are closed.*

Proof. Statements (i) and (ii) have been proved in [14, Thm. 4], while statement (iii) can be found in [20, p. 85].

It remains to verify (iv). From (iii), $(Z/\sim, d_\sim)$ is a metric space if and only if the topology it generates is T_0 . Let us observe that a pseudo-metric space is T_0 if and only if is T_1 . Therefore $(Z/\sim, d_\sim)$ is a metric space if and only if its points are closed, i.e. if and only if the equivalence classes $[z] \in Z/\sim$ are closed. Moreover, since from (ii) the projection $\pi : (Z, d) \rightarrow (Z/\sim, d_\sim)$ is a topological quotient, $[z] \in Z/\sim$ is closed if and only if $\pi^{-1}([z])$, that is the orbit of z in Z induced by the action, is closed. \square

3. THE NATURAL PSEUDO-DISTANCE AS A QUOTIENT PSEUDO-METRIC

In this section we show that the natural pseudo-distance is a quotient pseudo-metric. In order to see this, we endow the space of continuous functions $\mathcal{C}(M, \mathbb{R}^n)$ with the uniform convergence metric $d : d(f, g) = \max_{p \in M} \|f(p) - g(p)\|_\infty$, which induces the compact open topology. In this way, $\mathcal{C}(M, \mathbb{R}^n)$ belongs to **PMet**.

Next we quotient $\mathcal{C}(M, \mathbb{R}^n)$ by the following equivalence relation: denoting by $\mathcal{H}(M)$ the set of self-homeomorphisms of M , for $f, f' \in \mathcal{C}(M, \mathbb{R}^n)$, $f \sim f'$ if and only if there exists $h \in \mathcal{H}(M)$ such that $f' = f \circ h$. In other words, the equivalence classes of \sim coincide with the orbits induced by the action of $\mathcal{H}(M)$ on $\mathcal{C}(M, \mathbb{R}^n)$ given by:

$$\begin{aligned} \circ : \mathcal{H}(M) \times \mathcal{C}(M, \mathbb{R}^n) &\longrightarrow \mathcal{C}(M, \mathbb{R}^n) \\ (h, f) &\mapsto f \circ h \end{aligned}$$

By definition, the natural pseudo-distance is trivial on the orbits of this action: $\delta(f, f \circ h) = 0$ for every $f \in \mathcal{C}(M, \mathbb{R}^n)$ and every $h \in \mathcal{H}(M)$. Therefore it is natural to identify functions in the same orbit and consider the natural pseudo-distance on $\mathcal{C}(M, \mathbb{R}^n)/\sim$ rather than on $\mathcal{C}(M, \mathbb{R}^n)$:

Definition 3.1. The natural pseudo-distance $\delta : \mathcal{C}(M, \mathbb{R}^n)/\sim \times \mathcal{C}(M, \mathbb{R}^n)/\sim \rightarrow \mathbb{R}$ is defined by setting

$$\delta([f], [g]) = \inf_{h \in \mathcal{H}(M)} \max_{p \in M} \|f(p) - g \circ h(p)\|_\infty.$$

Clearly this definition does not depend on the choice of the representatives f, g .

Let us observe that δ is not in general a distance, even when we define it on $\mathcal{C}(M, \mathbb{R}^n)/\sim$ rather than on $\mathcal{C}(M, \mathbb{R}^n)$. Indeed, there may exist functions $f, g \in \mathcal{C}(M, \mathbb{R}^n)$ such that $\delta([f], [g]) = 0$, but with no $h \in \mathcal{H}(M)$ for which $f = g \circ h$. Some examples of this fact can be found in [5, Sec. 2].

Now, as a corollary of Theorem 2.1, we get the result below.

Corollary 3.2. *The following statements hold:*

- (i) *The natural pseudo-distance δ is the quotient pseudo-metric induced by the action of $\mathcal{H}(M)$ on $(\mathcal{C}(M, \mathbb{R}^n), d)$: $\delta = d_\sim$.*
- (ii) *The topology induced on $\mathcal{C}(M, \mathbb{R}^n)/\sim$ by δ coincides with the quotient topology.*
- (iii) *$(\mathcal{C}(M, \mathbb{R}^n)/\sim, \delta)$ is a metric space if and only if the topology induced by δ is T_0 .*
- (iv) *$(\mathcal{C}(M, \mathbb{R}^n)/\sim, \delta)$ is a metric space if and only if each orbit induced by the action of $\mathcal{H}(M)$ on $\mathcal{C}(M, \mathbb{R}^n)$ is closed.*

Proof. We observe that

$$\begin{aligned} \delta([f], [g]) &= \inf \left\{ \max_{p \in M} \|f'(p) - g'(p)\|_\infty : f' \in [f], g' \in [g] \right\} \\ &= \inf \{d(f', g') : f' \sim f, g' \sim g\}. \end{aligned}$$

Furthermore, any self-homeomorphism of M induces an isometry on $(\mathcal{C}(M, \mathbb{R}^n), d)$: for every $h \in \mathcal{H}(M)$, $\max_{p \in M} \|f(p) - g(p)\|_\infty = \max_{p \in M} \|f \circ h(p) - g \circ h(p)\|_\infty$. Hence, it is sufficient to apply Theorem 2.1(i) to obtain that $\delta = d_\sim$ and Theorem 2.1(ii – iv) to obtain the other three claims. \square

We end the section by considering the case when the natural pseudo-distance is defined using only homeomorphisms in a subgroup $\mathcal{K}(M)$ of $\mathcal{H}(M)$.

Definition 3.3. Let $\mathcal{K}(M)$ be a subgroup of $\mathcal{H}(M)$, and let us consider the action of $\mathcal{K}(M)$ on $\mathcal{C}(M, \mathbb{R}^n)$: $f \sim_{\mathcal{K}} f'$ if and only if $f' = f \circ k$, for some $k \in \mathcal{K}(M)$. We define $\delta_{\mathcal{K}} : \mathcal{C}(M, \mathbb{R}^n)/_{\sim_{\mathcal{K}}} \times \mathcal{C}(M, \mathbb{R}^n)/_{\sim_{\mathcal{K}}} \rightarrow \mathbb{R}$ by

$$\delta_{\mathcal{K}}([f]_{\mathcal{K}}, [g]_{\mathcal{K}}) = \inf_{k \in \mathcal{K}(M)} \max_{p \in M} \|f(p) - g \circ k(p)\|_\infty.$$

Proposition 3.4. *The following statements hold:*

- (1) $\delta_{\mathcal{K}}$ is a quotient pseudo-metric induced by the action of $\mathcal{K}(M)$ on $(\mathcal{C}(M, \mathbb{R}^n), d)$.
- (2) The topology induced on $\mathcal{C}(M, \mathbb{R}^n)/_{\sim_{\mathcal{K}}}$ by $\delta_{\mathcal{K}}$ coincides with the quotient topology.
- (3) If $\mathcal{K}(M)$ is compact in $\mathcal{H}(M)$ with the compact open topology, then $(\mathcal{C}(M, \mathbb{R}^n), \delta_{\mathcal{K}})$ is a metric space.

Proof. The proofs of (1) and (2) follow immediately from Theorem 2.1(i – ii). As for (3), by Theorem 2.1(iv) it is sufficient to show that each orbit induced by $\mathcal{K}(M)$ on $\mathcal{C}(M, \mathbb{R}^n)$ is closed. Let $[\bar{f}]_{\mathcal{K}} \in (\mathcal{C}(M, \mathbb{R}^n)/_{\sim_{\mathcal{K}}}, \delta_{\mathcal{K}})$, and let (f_i) be a sequence such that $d(f_i, \bar{f}) \xrightarrow{i \rightarrow \infty} 0$ for some $\bar{f} \in \mathcal{C}(M, \mathbb{R}^n)$, and $f_i \in [\bar{f}]_{\mathcal{K}}$ for every i . Since $f_i = \bar{f} \circ k_i$, with $k_i \in \mathcal{K}(M)$, for every i , and $\mathcal{K}(M)$ is compact, there exists a subsequence (k_{i_j}) of (k_i) converging to a certain $k \in \mathcal{K}(M)$. Then we can take the subsequence (f_{i_j}) of (f_i) , with $f_{i_j} = \bar{f} \circ k_{i_j}$ for every j . Since composition is continuous with the compact open topology [11, Thm. 2.2], $\bar{f} \circ k_{i_j}$ converges to $\bar{f} \circ k$, and hence, $f = \bar{f} \circ k$. This proves that $f \in [\bar{f}]_{\mathcal{K}}$, i.e. that the orbit $\pi^{-1}([\bar{f}]_{\mathcal{K}})$ is closed, being π a topological quotient from (2). \square

4. APPLICATIONS

This section concerns some applications of Corollary 3.2 and Proposition 3.4 to subspaces of $\mathcal{C}(M, \mathbb{R}^n)$ under the action of $\mathcal{H}(M)$ or its subgroups $\mathcal{K}(M)$. In particular, in Subsection 4.1 we show that the natural pseudo-distance is a distance when induced by the action of $\mathcal{H}(M)$ on the space $\mathcal{E}(M, \mathbb{R}^n)$ of topological embeddings, and of the group $\mathcal{D}(M)$ of diffeomorphisms on the space $\mathcal{I}(M, \mathbb{R}^n)$ of immersions; Subsection 4.2 provides some examples of compact subgroups $\mathcal{K}(M)$ whose action on $\mathcal{C}(M, \mathbb{R}^n)$ induces a distance $\delta_{\mathcal{K}}$; Subsection 4.3 is devoted to prove that the natural pseudo-distance is a metric when we consider the space of simple Morse functions on surfaces under the action of C^2 -diffeomorphisms.

4.1. Embeddings in \mathbb{R}^n under the action of $\mathcal{H}(M)$. Let us consider the space $(\mathcal{E}(M, \mathbb{R}^n), d)$ of topological embeddings of M in \mathbb{R}^n (i.e. of homeomorphisms onto their image) endowed with the uniform convergence distance d . If K is a compact subset of M , and U is an open subset of \mathbb{R}^n , then $V(K, U) = \{f \in \mathcal{E}(M, \mathbb{R}^n), f(K) \subset U\}$ is an open set in the compact open topology induced by d .

Let us pass to the quotient by considering the action of $\mathcal{H}(M)$ on $\mathcal{E}(M, \mathbb{R}^n)$, and take \mathbb{R}^n endowed with the maximum norm. The following result, whose proof is inspired by [18, Lemma 13.9], holds.

Proposition 4.1. $(\mathcal{E}(M, \mathbb{R}^n)/_{\sim}, \delta)$ is a metric space.

Proof. Since $(\mathcal{E}(M, \mathbb{R}^n)/\sim, \delta)$ is pseudo-metric, from Corollary 3.2(iii), it is sufficient to verify that it is T_0 . Let $[f], [g] \in \mathcal{E}(M, \mathbb{R}^n)/\sim$, $[f] \neq [g]$. We want to show that there exists an open set containing $[f]$ and not $[g]$ or viceversa.

Let us observe that $[f] \neq [g]$ implies $f(M) \neq g(M)$. Indeed, if $f(M) = g(M)$, then $g = f \circ (f^{-1} \circ g)$, with $f^{-1} \circ g \in \mathcal{H}(M)$, i.e. $[f] = [g]$. Therefore, since $[f] \neq [g]$, without loss of generality, we can assume that there exists $y \in f(M) \setminus g(M)$. Since $g(M)$ is compact as the image of a compact space through a continuous function, an open set $U \subset \mathbb{R}^n$ can be found such that $g(M) \subset U$, $y \notin U$. Then, considering the open subset $V(M, U)$ of $\mathcal{E}(M, \mathbb{R}^n)$, we have $g \in V(M, U)$, while $f \notin V(M, U)$. Now, let us observe that $\pi(V(M, U))$ is open if and only if $\pi^{-1} \circ \pi(V(M, U))$ is open. Given that $\pi^{-1} \circ \pi(V(M, U)) = V(M, U)$ which is open, then $\pi(V(M, U))$ is open. Clearly, $[f]$ cannot belong to $\pi(V(M, U))$, while $[g]$ does. Hence, $\mathcal{E}(M, \mathbb{R}^n)/\sim$ is T_0 . \square

The above result can be generalized by considering the action of the group $\mathcal{D}(M)$ of diffeomorphisms of class \mathcal{C}^1 of M on the space $(\mathcal{I}(M, \mathbb{R}^n), d)$ of \mathcal{C}^1 immersions of M in \mathbb{R}^n (i.e. of diffeomorphisms whose differential is injective) without points of self-tangency. Let us recall that an immersion $f : M \rightarrow \mathbb{R}^n$ has no points of self-tangency if whenever p, p' are distinct points of M with $f(p) = f(p')$, then $\text{im } df_p \neq \text{im } df_{p'}$.

To prove that $\mathcal{I}(M, \mathbb{R}^n)/\sim_{\mathcal{D}}$ is T_0 , let us consider $f, g \in \mathcal{I}(M, \mathbb{R}^n)$ with $[f]_{\mathcal{D}} \neq [g]_{\mathcal{D}}$ and show that this implies $f(M) \neq g(M)$. The rest of the proof can be obtained using the same arguments as in the case of embeddings.

Let us assume $f(M) = g(M)$ and prove the existence of $h \in \mathcal{D}(M)$ such that $f = g \circ h$, i.e. $[f]_{\mathcal{D}} = [g]_{\mathcal{D}}$. Let $\mathcal{P}_f, \mathcal{P}_g \subset M$ be the sets of preimages of multiple points of f and g . Since $f, g \in \mathcal{I}(M, \mathbb{R}^n)$, $f|_{M \setminus \mathcal{P}_f}, g|_{M \setminus \mathcal{P}_g}$ are \mathcal{C}^1 -diffeomorphisms between $M \setminus \mathcal{P}_f$ and $f(M \setminus \mathcal{P}_f)$, and between $M \setminus \mathcal{P}_g$ and $g(M \setminus \mathcal{P}_g)$, respectively.

Moreover, since $f(M) = g(M)$, for any $p \in M$, the set $g^{-1}(f(p))$ is not empty. Seeing that in particular $f(\mathcal{P}_f) = g(\mathcal{P}_g)$, if $p \in M \setminus \mathcal{P}_f$, the set $g^{-1}(f(p))$ contains only one point p' and we can define $h(p) = p'$. If $p \in \mathcal{P}_f$, then $g^{-1}(f(p)) \subset \mathcal{P}_g$. In this case, there is just one point $p' \in g^{-1}(f(p))$ verifying $\text{im } dg_{p'} = \text{im } df_p$ because multiple points of g are without self-tangency. Thus, we can define $h(p) = p'$. Because of its definition, the function h verifies the equality $g \circ h = f$. Let us show that $h \in \mathcal{D}(M)$. Recalling that $f(M) = g(M)$, the definition of h implies that h is injective and surjective. Furthermore, for each point $p \in M$, there exist an open neighborhood $U(p)$ of p in M such that $f|_{U(p)}$ is a \mathcal{C}^1 -diffeomorphism, a point $p' \in M$ for which $g(p') = f(p)$, and an open neighborhood $U'(p')$ of p' in M such that $g|_{U'(p')} \circ f|_{U(p)}$ is a \mathcal{C}^1 -diffeomorphism and $g(U'(p')) = f(U(p))$. Hence, $h|_{U(p)}$ equals the \mathcal{C}^1 -diffeomorphism $g|_{U'(p')}^{-1} \circ f|_{U(p)}$. This proves that $[f]_{\mathcal{D}} = [g]_{\mathcal{D}}$.

4.2. $\mathcal{C}(M, \mathbb{R}^n)$ under the action of compact groups. As an application of Proposition 3.4(3), let us consider the space $(\mathcal{C}(M, \mathbb{R}^n), d)$ with M a submanifold of \mathbb{R}^n verifying one of the following properties:

- M is of revolution;
- M is invariant with respect to a rotation of $\frac{2\pi}{n}$.

In the previous cases or in each combination of them we can consider respectively $\mathcal{K}(M) \cong S^1$, $\mathcal{K}(M) \cong \mathbb{Z}_n$ or the corresponding product of these compact groups. Then the orbits induced by the action of $\mathcal{K}(M)$ on $\mathcal{C}(M, \mathbb{R}^n)$ are closed, so that $\delta_{\mathcal{K}}$ is a metric on $\mathcal{C}(M, \mathbb{R}^n)/\sim_{\mathcal{K}}$.

More in general, we can consider the action of the group $\mathfrak{I}(M)$ of isometries on M (i.e. of distance preserving self-homeomorphisms of M). Indeed, as stated in [15,

Thm. 1.2], in the case M is a compact manifold, the group $\mathfrak{I}(M)$ results to be compact in the compact open topology. Consequently, because of Proposition 3.4(3), we obtain that $(\mathcal{C}(M, \mathbb{R}^n)/_{\sim_{\mathfrak{J}}}, \delta_{\mathfrak{J}})$ is always a metric space.

4.3. The Morse functions space under the action of the group of diffeomorphisms. In what follows, M will denote a 2-dimensional smooth compact connected manifold without boundary, and $(\mathcal{M}_0(M, \mathbb{R}), d)$ the space of simple Morse functions on M endowed with the uniform convergence metric d . Let us recall the following facts: $f : M \rightarrow \mathbb{R}$ is a Morse function if it is of class C^2 and all its critical points are non-degenerate (i.e. the Hessian matrix at each critical point is non-singular); the number of negative eigenvalues of the Hessian matrix at a critical point is called the index of f at the critical point; as a consequence of the compactness of M and the property of being non-degenerate, Morse functions' critical points are isolated [19]. Moreover, the Morse function f is said to be simple if each of its critical values corresponds to a different critical point. Accordingly, it makes sense to use the terminology index of a critical value c to indicate the index of f at the only critical point whose value is c .

Given $f \in \mathcal{M}_0(M, \mathbb{R})$, we denote by $K(f)$ the set of its critical points, and by f^a the set $f^{-1}((-\infty, a])$, $a \in \mathbb{R}$.

We want to show that, under the action of the group $\mathcal{D}(M)$ of C^2 -diffeomorphisms on M , the natural pseudo-distance $\delta_{\mathcal{D}}$ turns out to be a distance:

Theorem 4.2. $(\mathcal{M}_0(M, \mathbb{R})/_{\sim_{\mathcal{D}}}, \delta_{\mathcal{D}})$ is a metric space.

In virtue of Corollary 3.2(iv), the proof of Theorem 4.2 will be provided showing that any orbit in $\mathcal{M}_0(M, \mathbb{R})$ induced by the action of the group $\mathcal{D}(M)$ is closed. To be more precise, we will prove that, if (f_i) is a converging sequence of simple Morse functions with $d(f_i, f) \xrightarrow{i \rightarrow \infty} 0$ for some $f \in \mathcal{M}_0(M, \mathbb{R})$, and $f_i \in [\bar{f}]_{\mathcal{D}} = \{f' \in \mathcal{M}_0(M, \mathbb{R}) : f' = \bar{f} \circ h, h \in \mathcal{D}(M)\}$ for every i , then $f \in [\bar{f}]_{\mathcal{D}}$. All these notations will be maintained throughout the section.

The main tool we will use is a result by Kudryavsteva [16, Lemma 1], rewritten here in Lemma 4.11. It states that two Morse functions sharing the same collection of critical points, the same graph in the sense of Definition 4.9, and the same values at critical points, belong to the same equivalence class under the action of $\mathcal{D}(M)$.

The proof of Theorem 4.2 is by steps. Firstly, we prove that f and \bar{f} share the same set of critical values with the same indices (Proposition 4.5); secondly, we show that each converging sequence of critical points of (f_i) corresponding to a certain critical value converges to the critical point of f corresponding to the same critical value (Corollary 4.7); thirdly, we demonstrate the existence of a function $f' \in [\bar{f}]_{\mathcal{D}}$ with the same collection of critical points, the same values at critical points as f (Proposition 4.8), and the same graph as the one of f (Remark 4.10). In this way, applying Lemma 4.11 to f and f' , Theorem 4.2 is proved.

The following two lemmas will be used to prove that f and the functions in $[\bar{f}]_{\mathcal{D}}$ have the same critical values with the same indices (Proposition 4.5).

Lemma 4.3. ([17, Lemma 4.1]) *Let $X_1, X_2, X_3, X'_1, X'_2, X'_3$ be topological spaces such that $X_1 \subseteq X_2 \subseteq X_3 \subseteq X'_1 \subseteq X'_2 \subseteq X'_3$. Let $H_k(X_3, X_1) = 0$, $H_k(X'_3, X'_1) = 0$ for every $k \in \mathbb{Z}$. Then the homomorphism induced by inclusion $H_k(X'_1, X_1) \rightarrow H_k(X'_2, X_2)$ is injective for every $k \in \mathbb{Z}$.*

Lemma 4.4. ([7, Thm. A.3]) *Let $g \in \mathcal{M}_0(M, \mathbb{R})$, and let c be a critical value of index k of g . Then there exists a real number $\eta(g, c) > 0$ such that each $g' \in \mathcal{M}_0(M, \mathbb{R})$ verifying $d(g, g') \leq \eta(g, c)$ admits at least one critical value c' of index k for which $|c - c'| \leq d(g, g')$.*

Proposition 4.5. *The functions f , \bar{f} , and all f_i have the same critical values with the same indices.*

Proof. Since $f_i \in [\bar{f}]_{\mathcal{D}}$, i.e. $f_i = \bar{f} \circ h_i$ for some $h_i \in \mathcal{D}(M)$, and critical values are preserved under diffeomorphisms, f_i and \bar{f} share the same set of critical values with the same indices for every i .

Let us prove the claim for f_i and f . By Lemma 4.4, if c is a critical value of index k of f , then there exists a real number $\eta(f, c) > 0$ such that each f_i verifying $d(f_i, f) \leq \eta(f, c)$ admits at least one critical value c' of index k , with $|c - c'| \leq d(f_i, f)$. Let us underline that c' does not depend on the index i as seen at the beginning of the proof. Letting i tend to infinity, we obtain $c = c'$. This proves that the set of critical values of f is contained in the set of critical values of f_i for every i .

To show that this inclusion cannot be proper, let us assume, by contradiction, that there exists $c \in \mathbb{R}$ that is a critical value for some, and hence all, f_i , and it is regular for f . Since $f \in \mathcal{M}_0(M, \mathbb{R})$, there exists a real number $\eta(f, c) > 0$ such that $[c - \eta(f, c), c + \eta(f, c)]$ does not contain any critical value of f . If we consider i large enough that $d(f_i, f) \leq \eta(f, c)$, Lemma 4.4 implies the existence of at least one critical value of f distant less than $\eta(f, c)$ from c . This gives a contradiction. \square

The result below shows that there exists a subsequence of critical points of (f_i) corresponding to a certain critical value which converges to the critical point of f corresponding to the same critical value.

Proposition 4.6. *Let c be a critical value of \bar{f} , and hence of f and all f_i . Let $q_i \in K(f_i) \cap f_i^{-1}(c)$ for every i . Then $q \in K(f) \cap f^{-1}(c)$ if and only if there exists a subsequence of (q_i) converging to q .*

Proof. Let us begin by proving that, if $q \in K(f) \cap f^{-1}(c)$, then there exists a subsequence of (q_i) converging to q .

Let us assume by contradiction that no subsequences of (q_i) converging to q exist. Then a sufficiently small neighborhood $U \subset M$ of q can be found such that U contains neither any other critical point of f besides q , nor critical points of f_i for any i . Moreover, since $f \in \mathcal{M}_0(M, \mathbb{R})$ and we are assuming $f(q) = c$, there exists a real number $\eta(f, c) > 0$ such that $[c - 3 \cdot \eta(f, c), c + 3 \cdot \eta(f, c)]$ does not contain any critical value of f besides c .

Let us consider an index \bar{i} so large that $\eta' = d(f_{\bar{i}}, f) < \eta(f, c)$, and write $f^{c-3 \cdot \eta'} \cap U = X_1$, $f_{\bar{i}}^{c-2 \cdot \eta'} \cap U = X_2$, $f^{c-\eta'} \cap U = X_3$, $f^{c+\eta'} \cap U = X'_1$, $f_{\bar{i}}^{c+2 \cdot \eta'} \cap U = X'_2$, $f^{c+3 \cdot \eta'} \cap U = X'_3$. Since $X_1 \subseteq X_2 \subseteq X_3 \subseteq X'_1 \subseteq X'_2 \subseteq X'_3$, and both $H_k(X_3, X_1)$ and $H_k(X'_3, X'_1)$ are trivial for every $k \in \mathbb{Z}$, we can apply Lemma 4.3 to obtain that the homomorphism $H_k(X'_1, X_1) \rightarrow H_k(X'_2, X_2)$ induced by inclusion is injective for every $k \in \mathbb{Z}$. But assuming that c is a critical value of index \bar{k} of f , $H_{\bar{k}}(X'_1, X_1)$ is not trivial because the critical point $q \in f^{-1}([c - 3 \cdot \eta', c + \eta']) \cap U$, while $H_{\bar{k}}(X'_2, X_2) = 0$ because $f_{\bar{i}}^{-1}([c - 2 \cdot \eta', c + 2 \cdot \eta']) \cap U$ does not contain any critical point of $f_{\bar{i}}$. This implies a contradiction.

Let us prove now that if there exists a subsequence of (q_i) converging to q , then $q \in K(f) \cap f^{-1}(c)$.

Let us denote again by (q_i) the subsequence converging to q . The fact that $d(f_i, f) \xrightarrow[i \rightarrow \infty]{} 0$, with $f_i(q_i) = c$ for every i , immediately implies that $f(q) = c$. By contradiction, let us assume that q is a regular value of f . Since f is a simple Morse function, an arbitrarily small neighborhood $U \subset M$ of q can be found such that U does not contain critical points of f . Because of the convergence of (q_i) , U contains the critical points q_i of f_i for every $i > \hat{i}$, for a certain index \hat{i} . Moreover, since $f_i \in \mathcal{M}_0(M, \mathbb{R})$ for every i , it is not restrictive to assume that U does not

contain any other critical point of f_i besides q_i for every i large enough. Hence, under the assumption $f_i(q_i) = c$, a real number $\eta(f_i, c) > 0$ can be chosen such that $[c - 3 \cdot \eta(f_i, c), c + 3 \cdot \eta(f_i, c)]$ does not contain any critical value of f_i besides c .

Fixed an index \bar{i} large enough, we obtain again a contradiction using the same arguments as in the first part of the proof with the roles of f and $f_{\bar{i}}$ exchanged. \square

Corollary 4.7. *Let c be a critical value of \bar{f} , and hence of f and all f_i . Let $q_i \in K(f_i) \cap f_i^{-1}(c)$ for every i . Every converging subsequence of (q_i) converges to $q \in K(f) \cap f^{-1}(c)$.*

Proof. It is sufficient to observe that, if two different converging subsequences of (q_i) converged to two different points q and q' , respectively, by Proposition 4.6 we would have that both q and q' are critical points of f , with $f(q) = f(q')$, against the assumption that f is simple. \square

The following result shows the existence of a function in $[\bar{f}]_{\mathcal{D}}$ having $K(f)$ as the set of its critical points, as well as the same values at critical points as f .

Proposition 4.8. *There exists $f' \in [\bar{f}]_{\mathcal{D}}$ such that f and f' have the same collection of critical points, the same index and the same value at each of them.*

Proof. Let $K(f_i) = \{q_i^1, \dots, q_i^n\}$ and $K(f) = \{q^1, \dots, q^n\}$, and assume $f_i(q_i^j) = f(q^j)$ for every $j = 1, \dots, n$. We apply iteratively Corollary 4.7 to extract a subsequence of (f_i) , say again (f_i) , such that, for every $j = 1, \dots, n$, the sequences of critical points (q_i^j) converge to $q^j \in K(f)$. Moreover, for every $j = 1, \dots, n$, let (U^j, ψ^j) be a local chart centered at q^j , and fix an index i large enough that $q_i^j \in U^j$.

Let $h : M \rightarrow M$ be a \mathcal{C}^2 -diffeomorphism such that

$$h(p) = \begin{cases} p, & p \in M \setminus \bigcup_{j=1}^n U^j \\ (\psi^j)^{-1} \circ h_i^j \circ \psi^j(p), & p \in U^j, j = 1, \dots, n \end{cases}$$

where, denoting by D^2 the unit 2-disk in \mathbb{R}^2 , $h_i^j : D^2 \rightarrow D^2$ is a diffeomorphism which takes $\psi^j(q^j)$ to $\psi^j(q_i^j)$, and is the identity in a neighborhood of ∂D^2 . Then we can define the function $f' : M \rightarrow \mathbb{R}$ as $f' = f_i \circ h$.

We observe that f' is a simple Morse function because obtained from f_i by composition with a diffeomorphism. Hence $f' \in [\bar{f}]_{\mathcal{D}}$. By Proposition 4.5, f' and f have the same critical values with the same indices. Furthermore, by construction, $h(q^j) = q_i^j$, and therefore, $f'(q^j) = f_i(h(q^j)) = f_i(q_i^j) = f(q^j)$. Since diffeomorphisms take critical points to critical points, $q^j \in K(f')$. This shows that $K(f') \subset K(f)$. On the other side, $K(f')$ and $K(f)$ have the same cardinality, so $K(f') = K(f)$. In conclusion, f and f' have the same set of critical points $\{q^1, \dots, q^n\}$, and the same values at them since $f'(q^j) = f(q^j)$ for every $j = 1, \dots, n$. \square

Let us recall the following concept introduced in [16].

Definition 4.9. Let $g : M \rightarrow \mathbb{R}$ be a Morse function with r saddle points p_1, \dots, p_r . The graph G_g associated with g is the graph obtained from $g^{-1}\{g(p_1), \dots, g(p_r)\}$ by removing all connected components containing no saddle critical points.

The graph G_g in Definition 4.9 has r vertices (which are the saddle points p_1, \dots, p_r); the degree of each vertex is equal to 4 and hence the graph has $2r$ edges.

Remark 4.10. The functions f and f' considered in Proposition 4.8 have the same associated graph in virtue on the fact that they are both simple Morse functions.

To prove Theorem 4.2, we use the following Lemma 4.11.

Lemma 4.11. ([16, Lemma 1]) *Let $g, g' : M \rightarrow \mathbb{R}$ be Morse functions with the same collection of critical points, the same graph $G_g = G_{g'}$, and the same values at critical points. Then*

$$g = g' \circ h$$

for some $h \in \mathcal{D}(M)$ homotopic to id_M .

Proof of Theorem 4.2. From Proposition 4.8 and Remark 4.10, we can apply Lemma 4.11 to f and f' . This proves that $[f]_{\mathcal{D}} = [f']_{\mathcal{D}}$, and therefore that the orbits induced by $\mathcal{D}(M)$ on $\mathcal{M}_0(M, \mathbb{R})$ are closed. Eventually, applying Corollary 3.2(iv), the claim follows. \square

REFERENCES

1. S. Biasotti, L. De Floriani, B. Falcidieno, P. Frosini, D. Giorgi, C. Landi, L. Papaleo, and M. Spagnuolo, *Describing shapes by geometrical-topological properties of real functions*, ACM Comput. Surv. **40** (2008), no. 4, 1–87.
2. A.M. Bronstein, M.M. Bronstein, and R. Kimmel, *Topology-invariant similarity of nonrigid shapes*, Int. J. Comput. Vision **81** (2009), no. 3, 281–301.
3. D. Burago, Y. Burago, and S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics, American Mathematical Society, 2001.
4. Francesca Cagliari, *Natural pseudodistance for shape comparison and quotients of metric spaces*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia **57** (2010), 63–67.
5. A. Cerri and B. Di Fabio, *On certain optimal diffeomorphisms between closed curves*, Forum Mathematicum (to appear).
6. F. Chazal, D. Cohen-Steiner, L. J. Guibas, F. Mémoli, and S. Y. Oudot, *Gromov-Hausdorff stable signatures for shapes using persistence*, Computer Graphics Forum (proc. SGP 2009) (2009), 1393–1403.
7. B. Di Fabio and C. Landi, *Reeb graphs of curves are stable under function perturbations*, Mathematical Methods in the Applied Sciences **35** (2012), no. 12, 1456–1471.
8. P. Donatini and P. Frosini, *Natural pseudodistances between closed manifolds*, Forum Mathematicum **16** (2004), no. 5, 695–715.
9. ———, *Natural pseudodistances between closed surfaces*, Journal of the European Mathematical Society **9** (2007), no. 2, 231–253.
10. ———, *Natural pseudo-distances between closed curves*, Forum Mathematicum **21** (2009), no. 6, 981–999.
11. James Dugundji, *Topology*, Series in advanced mathematics, Allyn and Bacon, 1966.
12. P. Frosini and C. Landi, *No embedding of the automorphisms of a topological space into a compact metric space endows them with a composition that passes to the limit*, Appl. Math. Lett. **24** (2011), no. 10, 1654–1657.
13. P. Frosini and M. Mulazzani, *Size homotopy groups for computation of natural size distances*, Bulletin of the Belgian Mathematical Society **6** (1999), no. 3, 455–464.
14. C. J. Himmelberg, *Pseudo-metrizability of quotient spaces*, Fundamenta Mathematicae **63** (1968), no. 1, pp. 1–6.
15. Shoshichi Kobayashi, *Transformation groups in differential geometry*, Springer-Verlag, Berlin, New York, 1972 (English).
16. E.A. Kudryavtseva, *Uniform Morse lemma and isotopy criterion for Morse functions on surfaces*, Moscow University Mathematics Bulletin **64** (2009), 150–158.
17. A. Marino and G. Prodi, *Metodi perturbativi nella teoria di Morse*, Boll. Un. Mat. Ital. (4) **11** (1975), no. 3, suppl., 1–32.
18. P.W. Michor, *Manifolds of differentiable mappings*, Shiva, Orpington, 1980.
19. J. Milnor, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, no. 51, Princeton University Press, Princeton, N.J., 1963.
20. Stephen Willard, *General Topology*, Addison-Wesley Series in Mathematics, Addison-Wesley, 1970.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, ITALIA
E-mail address: `francesca.cagliari@unibo.it`

ARCES, UNIVERSITÀ DI BOLOGNA, ITALIA
E-mail address: `barbara.difabio@unibo.it`

DIPARTIMENTO DI SCIENZE E METODI DELL'INGEGNERIA, UNIVERSITÀ DI MODENA E REGGIO
EMILIA, ITALIA
E-mail address: `clandi@unimore.it`