

This is the peer reviewed version of the following article:

Aluminum and tin: Food contamination and dietary intake in an Italian population / Filippini, Tommaso; Tancredi, Stefano; Malagoli, Carlotta; Cilloni, Silvia; Malavolti, Marcella; Violi, Federica; Vescovi, Luciano; Bargellini, Annalisa; Vinceti, Marco. - In: JOURNAL OF TRACE ELEMENTS IN MEDICINE AND BIOLOGY. - ISSN 0946-672X. - 52:(2019), pp. 293-301. [10.1016/j.jtemb.2019.01.012]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.


08/09/2024 09:08

(Article begins on next page)

Accepted Manuscript

Title: Aluminum and tin: Food contamination and dietary intake in an Italian population

Authors: Tommaso Filippini, Stefano Tancredi, Carlotta Malagoli, Silvia Cilloni, Marcella Malavolti, Federica Violi, Luciano Vescovi, Annalisa Bargellini, Marco Vinceti



PII: S0946-672X(18)30687-4
DOI: <https://doi.org/10.1016/j.jtemb.2019.01.012>
Reference: JTEMB 26287

To appear in:

Received date: 8 November 2018
Revised date: 20 January 2019
Accepted date: 22 January 2019

Please cite this article as: Filippini T, Tancredi S, Malagoli C, Cilloni S, Malavolti M, Violi F, Vescovi L, Bargellini A, Vinceti M, Aluminum and tin: Food contamination and dietary intake in an Italian population, *Journal of Trace Elements in Medicine and Biology* (2019), <https://doi.org/10.1016/j.jtemb.2019.01.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Aluminum and tin: Food contamination and dietary intake in an Italian population

Authors: Tommaso Filippini^a, Stefano Tancredi^a, Carlotta Malagoli^a, Silvia Cilloni^a, Marcella Malavolti^a, Federica Violi^a, Luciano Vescovi^a, Annalisa Bargellini^a, Marco Vinceti^{a,b}

Affiliations

^aEnvironmental, Genetic and Nutritional Epidemiology Research Center (CREAGEN), Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, ^bDepartment of Epidemiology, Boston University School of Public Health, Boston, USA.

Corresponding author

Tommaso Filippini, CREAGEN – Section of Public Health - Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Via Campi 287, 41125 Modena, Italy (Tel. +39 059 2055716; fax +39 059 2055483; E-mail tommaso.filippini@unimore.it)

Highlights

- We estimated dietary intake of aluminum and tin in an adult Italian population sample.
- Legumes, sweets, and cereals show the highest contents of aluminum, whereas sweets, meat, and fish contain large amounts of tin.
- The estimated median aluminum and tin intake are 4.1 mg/day and 66.8 µg/day, respectively.
- Major intake contributors are beverages, leafy vegetables, and cereals for aluminum, and tomatoes, fruit and processed meat for tin.

Abstract

Aluminum and tin are ubiquitous in the environment. In normal biological systems, however, they are present only in trace amounts and have no recognized biological

functions in humans. High exposure to these metals can result in adverse health effects such as neurodegenerative diseases. In non-occupationally exposed subjects, diet is the primary source of exposure. In this study, we aimed at estimating dietary aluminum and tin intake in an Italian adult population.

We measured aluminum and tin concentrations through inductively-coupled plasma mass spectrometry in 908 food samples. We also estimated dietary intake of these two metals, by using a validated semi-quantitative food-frequency questionnaire administered to 719 subjects (319 men and 400 women) recruited from the general population of the Emilia Romagna region, Northern Italy.

We found the highest aluminum levels in legumes, sweets, and cereals, while the highest tin levels were in sweets, meat and seafood. The estimated median daily dietary intake of aluminum was 4.1 mg/day (Interquartile range - IQR: 3.3-5.2), with a major contribution from beverages (28.6%), cereals (16.9%), and leafy vegetables (15.2%). As for tin, we estimated a median intake of 66.8 $\mu\text{g/day}$ (IQR: 46.7-93.7), with a major contribution from vegetables (mainly tomatoes) (24.9%), fruit (15.5%), aged cheese (12.2%) and processed meat (10.4%).

This study provides an updated estimate of the dietary intake of aluminum and tin in a Northern-Italy adult population, based on data from a validated food-frequency questionnaire. The intake determined for this population does not exceed the established thresholds of tolerable intake.

Keywords: aluminum, tin, dietary intake, food contamination, food safety.

Introduction

Aluminum and tin are ubiquitous in the environment. Both metals have no known biological function in humans, while a markedly increased exposure can result in toxic effects involving several systems [1, 2]. Non-occupational exposure to aluminum has been linked to adverse effects, particularly anemia, bone disease, and dialysis encephalopathy [3, 4]. Moreover, overexposure to such trace elements including high chronic aluminum exposure has been related to an increased risk of neurodegenerative diseases, including Alzheimer's Dementia, Parkinson's disease, and amyotrophic lateral sclerosis [5-14]. With reference to tin, its different chemical species play a major role in the assessment of adverse health effects [15]. High exposure to inorganic compounds mainly leads to gastrointestinal problems [15-17]. On the other hand, exposure to organotin compounds, such as trimethyltin and triethyltin, has been related to gastrointestinal, metabolic and neurological adverse effects [16, 18-20].

In non-occupationally exposed individuals, the primary sources of exposure to these metals is diet. In industrialized societies, more specifically, the use of food packaging materials or food additives enhancing food properties has markedly increased exposure to these metals [21-24]. In particular, aluminum compounds may be added during food industrial transformation to produce convenience foods. Aluminum containers are thus widely used to cook, freeze or wrap foods through aluminum foil [25-28], while food additives containing aluminum are used as coloring and/or anticaking agents [29-31].

Similarly, the main use of tin is for the manufacture of cans and containers, typically as tinfoil [16, 21, 32]. Another source of exposure to tin, chiefly for organic compounds, is its use as a stabilizer for polyvinyl chloride [33, 34]. As a consequence, direct contact of tin with foods for general populations may result in migration from the inside wall of cans and other packaging materials to edible contents.

For these reasons, dietary intake, particularly through convenience foods produced through food industrialized transformation and/or packaging processes, is generally the most important source of exposure to these metals [16, 29, 35, 36]. In this study, we sought to assess dietary aluminum and tin intake in order to provide an updated estimate of their exposure in the general adult population of the Emilia Romagna Region, Northern Italy.

Methods

Study population and assessment of dietary habits

We assessed the dietary habits of a sample population from Northern Italy. Detailed information of participant identification and recruitment has been previously reported [37, 38]. In brief, we identified 2,825 potentially eligible subjects from the population database of the Emilia-Romagna Region residents. These were from the Bologna, Ferrara, Modena, Parma, and Reggio Emilia provinces. A list was compiled by accessing the National Health Service directory. Of the eligible subjects, 747 (26.4%) agreed to study participation. After providing written informed consent, they returned a lifestyle and food frequency questionnaire (FFQ), which we had mailed to them. We excluded twenty-eight individuals from the final analysis because of incomplete data or extreme values (ratio of total energy intake: basal metabolic rate lower than the 0.5th percentile or higher than the 99.5th percentile) derived from the FFQ. The final sample comprised 719 participants aged 18-87 years (mean 55 years), including 319 men (mean age 59 years) and 400 women (mean age 52 years) [39, 40]. With regard to the evaluation of dietary habits, we assessed food and nutrient intake using a validated semi-quantitative FFQ specifically developed for the Central-Northern Italy population [41, 42] within the EPIC (European Prospective Investigation into Cancer and Nutrition) project. The FFQ was designed to estimate frequency and amount of consumption of 188 food items over the previous year, also using photos of serving sizes to ensure proper completion by participants. The median energy intake was 1906 kcal/day (interquartile range-IQR: 1538-2364) in all subjects, 2024 kcal/day (IQR: 1649-2462) in men and 1800 kcal/day (IQR: 1455-2296) in women, respectively.

Food collection and analysis

We determined aluminum and tin contents of food by characterizing the usual diet of a sample from the Emilia-Romagna population. As previously described in detail, we selected food consumed by a random sample of this population. Furthermore, we bought samples of these food items in markets and community canteens between October 2016 and February 2017 [43]. We strove to avoid cross-contamination with metals and other trace elements from food containers, by using plastic tubes or jars as well as plastic cutlery or stainless-steel knives during food collection and handling. With a clean stainless-steel knife, we then cut solid food samples by collecting specimens from the plate in six different points. For the purpose of subsequent analysis, these samples were homogenised separately in a food blender equipped with a stainless-steel blade. We placed portions of the samples (0.5 g) in quartz containers previously washed with MilliQ water (MilliQPlus,

Millipore, MA, USA) and HNO₃. Food samples were liquid-ashed with 10 ml solution (5 ml HNO₃ + 5 ml H₂O) in a microwave digestion system (Discover SP-D, CEM Corporation, NC, USA). We then stored them in plastic tubes and diluted them to 50 ml with deionised water before analysis. We performed trace element determination using an inductively coupled plasma-mass spectrometer (Agilent 7500ce, Agilent Technologies, CA, USA). We performed all analyses in duplicate and implemented quality controls as previously described [44, 45]. Limits of quantification amounted to 0.5 µg/kg for aluminum and 0.02 µg/kg for tin, corresponding to limits of detection (LOD) of 0.17 µg/kg and 0.007 µg/kg, respectively. Values below the LOD were set equal to LOD/2.

We reported aluminum and tin concentration according to food consumption patterns and food categories typical of this Italian population, as assessed through the FFQ [46, 47]. Our final list of main food categories included the following items: cereals and cereal products (pasta and other grains: durum wheat pasta, whole grain pasta, filled pasta, egg pasta, and pasta containing other grains such as barley and spelt; rice: both white (Carnaroli, Arborio) and brown; bread: white bread; whole-wheat bread; multigrain bread; crackers, crispbread and salty snacks, including maltose crackers, whole-wheat crackers and taralli), meat and meat products (red meat: beef, calf, pork, horse and game, including lean and fatty cuts; white meat: chicken, turkey, rabbit; processed meat: several types of salt-cured meat, especially those traditional of Emilia-Romagna including ham, sausage, salami, mortadella, bacon, etc.; offal: liver, heart, spleen); milk and dairy products, e.g. milk and yogurt (both whole milk, reduced-fat or skim milk, and fruit yogurts), fresh cheese including cream cheese, ricotta, stracchino, mozzarella, etc., aged cheese such as Parmigiano-Reggiano, Grana Padano, and Emmental, etc.); eggs, fish and seafood (preserved fish such as smoked salmon, and tinned tuna or mackerel; non-piscivorous fish such as sole and plaice; piscivorous fish including tuna, swordfish and salmon; crustaceans and molluscs such as shrimp, octopus, cuttlefish, mussels and clams); vegetables (leafy vegetables: various types of salad, spinach, chard, etc.; other vegetables: sweet, pepper, fennel, celery, artichoke, etc.; tomatoes; root vegetables: carrot, turnip; cabbage: green, Savoy, red, and Brussels sprouts; mushrooms, both cultivated and wild; onion and garlic); legumes (kidney beans, peas, lentils, etc.); potatoes (mainly regular but also sweet potatoes); fresh fruit (citrus fruit: orange, mandarin, grapefruit; other fruit: apple, pear, apricot, plum, banana, etc.); dried fruit and seeds (raisins, prunes, walnuts, hazelnuts, etc.); sweets including chocolate and cakes (sugar, non-chocolate confectionery with honey, jam and candies; chocolate and chocolate-based candy bars; ice cream: chocolate, non-chocolate and fruit ice cream; cakes, pies and

pastries; biscuits and dried cakes); oils and fats (vegetables oils and non-olive oil including margarine; olive oil; butter and other animal fats), and beverages (coffee and tea: caffeinated and decaffeinated coffee made using moka pots as well as espresso machines, black and green tea; wine: several types of red and white wine mainly from Emilia-Romagna but also from other Italian regions; aperitif wines and beers; spirits and liqueurs such as rum, whiskey and limoncello; fruit juices: orange juice and non-citrus juices; soft drinks: cola-type drinks, soda-type drinks, orangeade, etc.).

2.3 Estimate of metal dietary intake

We combined data from the measurement of metals in foods and the dietary habits assessed by the FFQ, in order to compute total daily metal intake by using the equation presented below.

$$\text{Daily dietary exposure } \left(\frac{\mu\text{g}}{\text{day}} \right) = \sum \frac{\text{element food content } \left(\frac{\mu\text{g}}{\text{kg}} \right) \times \text{food intake } \left(\frac{\text{g}}{\text{day}} \right)}{1000}$$

We multiplied contents measured in food ($\mu\text{g}/\text{kg}$) with intake as estimated through the FFQ (g/day). Accordingly, we estimated daily dietary aluminum and tin intake for the diet as a whole and for each food category, by reporting median and interquartile ranges (IQR). For the estimate of dietary intake by body weight (bw), we divided by participant weight (kg). Compared to other studies, during conversion from $\mu\text{g}/\text{day}$ to $\mu\text{g}/\text{kg}$ of bw per day or week, we considered a body weight of 60 kg for the purpose of our calculations, in accordance with guidelines from international agencies [29, 36, 48].

Results

Table 1 reports details about dietary habits in the study population. In general, we found a comparable intake of main food categories in both sexes, with the partial exception of a higher intake of cereal products, meat and beverages (mainly due to wine intake) in men. Conversely, women showed higher consumption of milk and yogurt, coffee and tea and slightly higher consumption of citrus fruits.

The analysis of aluminum and tin concentration in food samples is presented in Table 2 according to main food categories and subgroups. All values were above the LOD for aluminum. As far as tin is concerned, in contrast, 61 specimens (6.7% of total sampled foods) showed concentrations below the LOD (Supplemental Table S1). Considering the main food categories, the highest aluminum contents were found in legumes ahead of

sweets (mainly in chocolate-based products) and cereals, mainly due to higher contents in bread, crackers and crispbread. Results from the subgroup analysis showed high aluminum concentrations in processed meat, crustaceans/molluscs, leafy vegetables, dried fruit and coffee/tea, compared with the median value of the main food categories. As for tin, likewise, the highest contents were found in sweets and meat, ahead of dried fruit and fish, mainly preserved and canned samples. The subgroup analysis demonstrated high tin contents in bread, mushrooms and leafy vegetables.

The dietary intake of the two metals is presented in Table 3. The median total aluminum intake was 4.1 mg/day (IQR: 3.3-5.2 mg/day), corresponding to 407.43 (IQR: 319.75-524.05) $\mu\text{g}/\text{kg}$ of bw/week or 58.20 (IQR: 45.68-74.86) $\mu\text{g}/\text{kg}$ bw/day. The food categories which mainly contributed to total intake were vegetables (25.2%), mainly leafy vegetables (15.2%) and tomatoes (3.8%), followed by beverages (28.6%), mainly coffee and tea (24.9%), and finally cereal products (16.9%) (Figure 1). The median tin intake was 66.80 $\mu\text{g}/\text{day}$ (IQR: 46.72-93.66 $\mu\text{g}/\text{day}$), corresponding to 6.70 (IQR: 4.85-9.25) $\mu\text{g}/\text{kg}$ bw/week or 0.96 (IQR: 0.69-1.32) $\mu\text{g}/\text{kg}$ bw/day. The main contribution to total tin intake was made by vegetables (30.3%) – particularly tomatoes (24.9%) – and fresh fruit (15.5%), with a substantial contribution from aged cheese (12.2%), processed meat (10.4%) and other animal fats (4.1%) as well (Figure 1). Estimates stratified by sex generally showed comparable results, except for the slightly higher intake of tin in men related to a higher intake of vegetables, mainly tomatoes (Supplemental Tables S2-S3 and Supplemental Figures S1-S2).

Finally in Table 4, we compared the estimated aluminum and tin intake in our population with dietary intake and tolerable upper intake levels reported by international agencies for European [29, 36] and worldwide populations [24, 49, 50]. While median and upper (95th percentile) values of aluminum intake fall in the range reported by other European and non-European populations, the corresponding figures for tin are below the range of intakes reported in other populations. Finally, Table 4 demonstrated that for both metals, the tolerable levels established by EFSA and WHO are not exceeded even considering the upper estimated intake of the two metals.

Discussion

In the context of a Northern Italian population, we assessed dietary intake of aluminum and tin by determining their concentration in a large number of food samples. Aluminum intake fell within the range of values reported by EFSA for European populations [29], consistent with other studies [26, 51-54]. However, other studies have

yielded higher [55-61] or lower values [25, 27, 62-66]. Variations in dietary exposure may be accounted for by differences in soil composition in the regions where food is produced or in individual dietary patterns. The population investigated in our study was characterized by generally high adherence to the Mediterranean diet [38], as compared with other populations with different dietary habits [67]. This may have influenced amount and frequency of consumption of the kind of foods that we observed to be major contributors to aluminum and tin intake. In particular, cereals and vegetables were found to be major contributors to aluminum intake, consistent with other studies on European populations [56, 61, 65, 68, 69]. We found the highest levels of aluminum in legumes, which were shown to naturally accumulate large amount of the metal [56, 70, 71]. We also found high levels of aluminum in cereal products and sweets, especially in bread products (including crackers and rusks), cakes and biscuits. Such high levels might be due both to the use of food additives, e.g. leavening agents such as acid sodium aluminum phosphate in cereal products, or anti-caking agents such as sodium aluminum silicate in cake mixes and dried products [31, 56, 72], and to generally very high aluminum contents in cocoa powder and chocolate [70].

As expected for vegetables, they showed fairly marked variation in aluminum contents. Thus, higher levels were observed in leafy vegetables [57, 69, 70] and tomatoes, especially in canned samples. This may be due to migration from containers and the low pH of foodstuffs as well as higher temperatures during storage time [73-76]. Furthermore, beverages (chiefly coffee and tea) were an important contributor to aluminum intake, consistent with previous studies [60, 69]. Consumption of coffee and tea may influence overall aluminum intake considerably, due to substantial variation in consumption and metal content [77, 78]. Interestingly, the analysis of aluminum distribution in different parts of tea plants also showed higher accumulation in mature leaves. On the other hand, lower values were detected in roots and young leaves [79], thus explaining the high amount of aluminum in tea samples. Consistent with other studies, we found generally high aluminum contents in aged cheese, arguably related to aluminum additives especially in (semi-)hard and industrially processed samples [31, 53, 56, 80].

Low levels of dietary tin intake were generally reported for our population, compared with others [36, 59, 61, 63, 65, 66, 81-83]. Overall, tin intake levels were similar to or slightly higher than values reported for few other countries [32, 84, 85]. Our findings indicated that vegetables and fruits were the main contributors to tin intake. This could be due to contamination from can containers [16, 23], as shown by the high levels found in canned tomatoes. Both the high content and the high consumption of this food item

resulted in a major contribution to dietary intake, tomatoes being the highest source among all vegetables. Similarly to aluminum, the migration of this metal from packaging materials could have been favored by the acidity of this type of food [86, 87].

By contrast, the contribution of tinned fish in our population was not relevant compared with that observed in other populations. This may be due to varying concentrations of tin as well as lower daily intake of this type of preserved foodstuff [32, 88]. The relevant role of processed meat and aged cheese to overall tin intake was due to high metal contents in these foods, probably correlated with industrial processing, use of higher amounts of food additives and/or migration from packaging materials [16, 36].

Overall, the risk assessment of both aluminum and tin intake in our study population indicated that safety levels were not exceeded. This view is based on the upper levels established by FAO/WHO and EFSA, considering neither median nor upper levels of estimated intake for both metals.

A strength of our study lies in the collection and analysis of a large number of food samples. In order to identify food relevant to the dietary habits of our population, we organized sample collection according to main categories, as in previous studies on dietary intake of other trace elements [44, 45]. We also followed guidelines from total diet studies, thus improving the validity of our findings and comparability with other studies [89]. Moreover, we aimed at mimicking a close-to-real mode of food collection in the general population through the purchase of food samples in both supermarkets and groceries of two provinces of Emilia Romagna (i.e. Modena and Reggio Emilia). However, we could not assess other kinds of possible contamination between local and imported foodstuff, since we did not evaluate the geographical origin of products.

Another strength of the study is the estimation of dietary habits in a large population sample through a detailed and validated food frequency questionnaire specifically developed for a Northern Italian population [41, 42].

A first limitation is that we assessed dietary habits in an adult population. Therefore, our findings may not be applicable to other population subgroups, such as children and adolescents.

We also carried out no speciation analysis for the two elements, hampering intake evaluation for different species characterized by heterogeneous or sometimes opposite health effects. With reference to tin, however, a study carried out using duplicate diet samples demonstrated relatively low levels of organotin compounds in sampled foods, and estimated daily intake was lower than the established tolerable daily intake [90].

In addition, we did not assess metal bioavailability. The average fraction of aluminum absorbed from a given dose is estimated to be low, subject though it is to wide variation (from less than 1% to 24% in humans). This depends on a number of factors: aluminum intake, intraluminal presence of substances with structural similarity (as for essential metals such as calcium and iron), or with binding potential such as that of citrate, as well as intraluminal pH, iron status and calcium status [72, 91-93]. As regards tin, metallic ions are practically insoluble and do not break through gastrointestinal barriers. As a result, they are mainly absorbed as organotin compounds [90, 94], for the bioavailability of which pH is considered an important factor [21]. In this regard, a bioconcentration study indicated that, as pH increases, the uptake of organotin compounds increases [95].

Furthermore, we did evaluate the health status of the study subjects, including possible malabsorption disease affecting metal uptake. With regard to overall aluminum intake, finally, we did not evaluate the influence of other dietary sources of the metal such as drinking water [96] and aluminum-containing medications, above all antacid drugs likely to increase daily aluminum intake by up to several grams [97, 98]. Moreover, our assessment was limited to the dietary intake of the two metals, while evaluating overall exposure in humans requires that additional sources be considered, such as inhalation and dermal contact. However, with the exception of rare occupational exposure or severe environmental pollution, these sources generally play a negligible role in determining overall exposure, compared with diet [21, 22].

Conclusions

In the Italian area investigated in the present study, legumes, sweets, and cereals presented the highest aluminum contents, whereas the highest tin levels were found in sweets, meat, fish and seafood. Overall, aluminum and tin intake was lower compared with the tolerable intake, as established by international agencies.

Founding

This work was supported by the Italian Association for Cancer Research (AIRC) and by the National Health Service - Local Health Authority of Reggio Emilia.

Declarations of interest: none.

References

- [1] R.A. Yokel, Aluminum, in: E. Merian, M. Anke, M. Ihnat, M. Stoeppler (Eds.), Elements and Their Compounds in the Environment, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2004, pp. 635-658.
- [2] J.P. Anger, Tin, in: E. Merian, M. Anke, M. Ihnat, M. Stoeppler (Eds.), Elements and Their Compounds in the Environment, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2004, pp. 1113-1124.
- [3] D. Krewski, R.A. Yokel, E. Nieboer, D. Borchelt, J. Cohen, J. Harry, S. Kacew, J. Lindsay, A.M. Mahfouz, V. Rondeau, Human health risk assessment for aluminium, aluminium oxide, and aluminium hydroxide, *J Toxicol Environ Health B Crit Rev* 10 Suppl 1 (2007) 1-269.
- [4] P. Nayak, Aluminum: impacts and disease, *Environ Res* 89(2) (2002) 101-15.
- [5] B. Bocca, G. Forte, R. Oggiano, S. Clemente, Y. Asara, A. Peruzzu, C. Farace, S. Pala, A.G. Fois, P. Pirina, R. Madeddu, Level of neurotoxic metals in amyotrophic lateral sclerosis: A population-based case-control study, *J Neurol Sci* 359(1-2) (2015) 11-7.
- [6] F. Cavaleri, Review of Amyotrophic Lateral Sclerosis, Parkinson's and Alzheimer's diseases helps further define pathology of the novel paradigm for Alzheimer's with heavy metals as primary disease cause, *Med Hypotheses* 85(6) (2015) 779-90.
- [7] C.E. Cicero, G. Mostile, R. Vasta, V. Rapisarda, S.S. Signorelli, M. Ferrante, M. Zappia, A. Nicoletti, Metals and neurodegenerative diseases. A systematic review, *Environ Res* 159 (2017) 82-94.
- [8] M.T. Colomina, F. Peris-Sampedro, Aluminum and Alzheimer's Disease, *Adv Neurobiol* 18 (2017) 183-197.
- [9] E.M. Garzillo, M. Lamberti, G. Genovese, P. Pedata, D. Feola, N. Sannolo, L. Daniele, F. Trojsi, M.R. Monsurro, N. Miraglia, Blood lead, manganese, and aluminum levels in a regional Italian cohort of ALS patients: does aluminum have an influence?, *J Occup Environ Med* 56(10) (2014) 1062-6.
- [10] L. Levesque, C.A. Mizzen, D.R. McLachlan, P.E. Fraser, Ligand specific effects on aluminum incorporation and toxicity in neurons and astrocytes, *Brain Res* 877(2) (2000) 191-202.
- [11] M. Mold, D. Umar, A. King, C. Exley, Aluminium in brain tissue in autism, *J Trace Elem Med Biol* 46 (2018) 76-82.

- [12] P.M. Roos, O. Vesterberg, T. Syversen, T.P. Flaten, M. Nordberg, Metal concentrations in cerebrospinal fluid and blood plasma from patients with amyotrophic lateral sclerosis, *Biol Trace Elem Res* 151(2) (2013) 159-70.
- [13] M. Vinceti, M. Fiore, C. Signorelli, A. Odone, M. Tesaro, M. Consonni, E. Arcolin, C. Malagoli, J. Mandrioli, S. Marmiroli, S. Sciacca, M. Ferrante, Environmental risk factors for amyotrophic lateral sclerosis: methodological issues in epidemiologic studies, *Ann Ig* 24(5) (2012) 407-15.
- [14] J.R. Walton, Chronic aluminum intake causes Alzheimer's disease: applying Sir Austin Bradford Hill's causality criteria, *J Alzheimers Dis* 40(4) (2014) 765-838.
- [15] J.P. Anger, *Tin, Elements and Their Compounds in the Environment* (2008).
- [16] S. Blunden, T. Wallace, Tin in canned food: a review and understanding of occurrence and effect, *Food Chem Toxicol* 41(12) (2003) 1651-62.
- [17] P.J. Boogaard, M. Boisset, S. Blunden, S. Davies, T.J. Ong, J.P. Taverne, Comparative assessment of gastrointestinal irritant potency in man of tin(II) chloride and tin migrated from packaging, *Food Chem Toxicol* 41(12) (2003) 1663-70.
- [18] J.J. Heindel, B. Blumberg, M. Cave, R. Machtinger, A. Mantovani, M.A. Mendez, A. Nadal, P. Palanza, G. Panzica, R. Sargis, L.N. Vandenberg, F. Vom Saal, Metabolism disrupting chemicals and metabolic disorders, *Reprod Toxicol* 68 (2017) 3-33.
- [19] P. Rantakokko, K.M. Main, C. Wohlfart-Veje, H. Kiviranta, R. Airaksinen, T. Vartiainen, N.E. Skakkebaek, J. Toppari, H.E. Virtanen, Association of placenta organotin concentrations with congenital cryptorchidism and reproductive hormone levels in 280 newborn boys from Denmark and Finland, *Hum Reprod* 28(6) (2013) 1647-60.
- [20] P. Rantakokko, K.M. Main, C. Wohlfart-Veje, H. Kiviranta, R. Airaksinen, T. Vartiainen, N.E. Skakkebaek, J. Toppari, H.E. Virtanen, Association of placenta organotin concentrations with growth and ponderal index in 110 newborn boys from Finland during the first 18 months of life: a cohort study, *Environ Health* 13(1) (2014) 45.
- [21] ATSDR, Toxicological profile for tin, in: U.S. Department of Health and Human Services - Public Health Service (Ed.) Agency for Toxic Substances and Disease Registry, Atlanta, GA, 2005.
- [22] ATSDR, Toxicological profile for aluminum, in: U.S. Department of Health and Human Services - Public Health Service (Ed.) Agency for Toxic Substances and Disease Registry, Atlanta, GA, 2008.
- [23] C. Reilly, *The packaging metals: Aluminium and Tin, Metal Contamination of Food: Its Significance for Food Quality and Human Health*, Third Edition, Blackwell Publishing Ltd 2007.

- [24] WHO, Trace elements in human health and nutrition, World Health Organization, Geneva, 1996.
- [25] V. Fekete, S. Vandevijvere, F. Bolle, J. Van Loco, Estimation of dietary aluminum exposure of the Belgian adult population: evaluation of contribution of food and kitchenware, *Food Chem Toxicol* 55 (2013) 602-8.
- [26] L. Gramiccioni, G. Ingrao, M.R. Milana, P. Santaroni, G. Tomassi, Aluminium levels in Italian diets and in selected foods from aluminium utensils, *Food Addit Contam* 13(7) (1996) 767-74.
- [27] J.P. Muller, A. Steinegger, C. Schlatter, Contribution of aluminum from packaging materials and cooking utensils to the daily aluminum intake, *Z Lebensm Unters Forsch* 197(4) (1993) 332-41.
- [28] R. Ranau, J. Oehlenschläger, H. Steinhart, Aluminium levels of fish fillets baked and grilled in aluminium foil, *Food Chemistry* 73(1) (2001) 1-6.
- [29] European Food Safety Authority, Scientific Opinion of the Panel on Food Additives, Flavourings, Processing Aids and Food Contact Materials on a request from European Commission on Safety of aluminium from dietary intake, *EFSA Journal* 754 (2008) 1-34.
- [30] M. Ogimoto, K. Suzuki, N. Haneishi, Y. Kikuchi, M. Takanashi, N. Tomioka, Y. Uematsu, K. Monma, Aluminium content of foods originating from aluminium-containing food additives, *Food Addit Contam Part B Surveill* 9(3) (2016) 185-90.
- [31] S.M. Saiyed, R.A. Yokel, Aluminium content of some foods and food products in the USA, with aluminium food additives, *Food Addit Contam* 22(3) (2005) 234-44.
- [32] S. Shimbo, N. Matsuda-Inoguchi, T. Watanabe, K. Sakurai, C. Date, A. Nishimura, H. Nakatsuka, H. Saito, K. Arisawa, M. Ikeda, Dietary intake of tin in Japan, and the effects on intake of canned food and beverage consumption, *Food Addit Contam* 24(5) (2007) 535-45.
- [33] W.A. Adams, Y. Xu, J.C. Little, A.F. Fristachi, G.E. Rice, C.A. Impellitteri, Predicting the migration rate of dialkyl organotins from PVC pipe into water, *Environ Sci Technol* 45(16) (2011) 6902-7.
- [34] A.I. Sadiki, D.T. Williams, A study on organotin levels in Canadian drinking water distributed through PVC pipes, *Chemosphere* 38(7) (1999) 1541-8.
- [35] R.A. Yokel, Aluminum in Food – The Nature and Contribution of Food Additives, in: Y. El-Samragy (Ed.), *Food Additive*, IntechOpen2012.

[36] European Food Safety Authority, Opinion of the Scientific Panel on Dietetic Products, Nutrition and Allergies on a request from the Commission related to the Tolerable Upper Intake Level of Tin, *EFSA Journal* 254 (2005) 1-25.

[37] M. Malavolti, C. Malagoli, C.M. Crespi, F. Brighenti, C. Agnoli, S. Sieri, V. Krogh, C. Fiorentini, F. Farnetani, C. Longo, C. Ricci, G. Albertini, A. Lanzoni, L. Veneziano, A. Virgili, C. Pagliarello, C. Feliciani, P.A. Fanti, E. Dika, G. Pellacani, M. Vinceti, Glycaemic index, glycaemic load and risk of cutaneous melanoma in a population-based, case-control study, *Br J Nutr* 117(3) (2017) 432-438.

[38] C. Malagoli, M. Malavolti, C. Agnoli, C.M. Crespi, C. Fiorentini, F. Farnetani, C. Longo, C. Ricci, G. Albertini, A. Lanzoni, L. Veneziano, A. Virgili, C. Pagliarello, M. Santini, P.A. Fanti, E. Dika, S. Sieri, V. Krogh, G. Pellacani, M. Vinceti, Diet quality and risk of melanoma in an Italian population, *J Nutr* 145(8) (2015) 1800-7.

[39] M. Malavolti, C. Malagoli, C. Fiorentini, C. Longo, F. Farnetani, C. Ricci, G. Albertini, A. Lanzoni, C. Reggiani, A. Virgili, C. Pagliarello, M. Santini, P.A. Fanti, E. Dika, S. Sieri, V. Krogh, G. Pellacani, M. Vinceti, Association between dietary vitamin C and risk of cutaneous melanoma in a population of Northern Italy, *Int J Vitam Nutr Res* 83(5) (2013) 291-8.

[40] M. Vinceti, C. Malagoli, C. Fiorentini, C. Longo, C.M. Crespi, G. Albertini, C. Ricci, A. Lanzoni, M. Reggiani, A. Virgili, F. Osti, M. Lombardi, M. Santini, P.A. Fanti, E. Dika, S. Sieri, V. Krogh, S. Seidenari, G. Pellacani, Inverse association between dietary vitamin D and risk of cutaneous melanoma in a Northern Italy population, *Nutr Cancer* 63(4) (2011) 506-13.

[41] V. Pala, S. Sieri, D. Palli, S. Salvini, F. Berrino, M. Bellegotti, G. Frasca, R. Tumino, C. Sacerdote, L. Fiorini, E. Celentano, R. Galasso, V. Krogh, Diet in the Italian EPIC cohorts: presentation of data and methodological issues, *Tumori* 89(6) (2003) 594-607.

[42] P. Pasanisi, F. Berrino, C. Bellati, S. Sieri, V. Krogh, Validity of the Italian EPIC questionnaire to assess past diet, *IARC Sci Publ* 156 (2002) 41-4.

[43] T. Filippini, M. Malavolti, S. Cilloni, L.A. Wise, F. Violi, C. Malagoli, L. Vescovi, M. Vinceti, Intake of arsenic and mercury from fish and seafood in a Northern Italy community, *Food Chem Toxicol* 116(Pt B) (2018) 20-26.

[44] T. Filippini, B. Michalke, C. Salvia, C. Malagoli, M. Malavolti, P. Grill, L. Vescovi, S. Sieri, V. Krogh, M. Vinceti, Dietary determinants of serum selenium species in Italian populations, *J Trace Elem Med Biol* 41S (2017) 39.

- [45] T. Filippini, S. Cilloni, M. Malavolti, F. Violi, C. Malagoli, M. Tesauro, I. Bottecchi, A. Ferrari, L. Vescovi, M. Vinceti, Dietary intake of cadmium, chromium, copper, manganese, selenium and zinc in a Northern Italy community, *J Trace Elem Med Biol* 50 (2018) 508-517.
- [46] T. Filippini, A. Ferrari, B. Michalke, P. Grill, L. Vescovi, C. Salvia, C. Malagoli, M. Malavolti, S. Sieri, V. Krogh, A. Bargellini, A. Martino, M. Ferrante, M. Vinceti, Toenail selenium as an indicator of environmental exposure: A cross-sectional study, *Mol Med Rep* 15(5) (2017) 3405-3412.
- [47] A. Turrini, A. Saba, D. Perrone, E. Cialfa, A. D'Amicis, Food consumption patterns in Italy: the INN-CA Study 1994-1996, *Eur J Clin Nutr* 55(7) (2001) 571-88.
- [48] WHO, Evaluation of certain food additives and contaminants. Eightieth report of the Joint FAO/WHO Expert Committee on Food Additives, *World Health Organ Tech Rep Ser* (995) (2016) 1-114.
- [49] WHO, Evaluation of certain food additives and contaminants. Seventy-seven report of the Joint FAO/WHO Expert Committee on Food Additives (JECFA), WHO, Geneva, 2007.
- [50] WHO, Safety evaluation of certain food additives and contaminants, in: *W.F.a.s. 65* (Ed.) WHO, Geneva, 2012.
- [51] N. Arnich, V. Sirot, G. Riviere, J. Jean, L. Noel, T. Guerin, J.C. Leblanc, Dietary exposure to trace elements and health risk assessment in the 2nd French Total Diet Study, *Food Chem Toxicol* 50(7) (2012) 2432-49.
- [52] G.H. Biego, M. Joyeux, P. Hartemann, G. Debry, Daily intake of essential minerals and metallic micropollutants from foods in France, *Sci Total Environ* 217(1-2) (1998) 27-36.
- [53] J.A. Pennington, S.A. Schoen, Estimates of dietary exposure to aluminium, *Food Addit Contam* 12(1) (1995) 119-28.
- [54] M. Rose, M. Baxter, N. Brereton, C. Baskaran, Dietary exposure to metals and other elements in the 2006 UK Total Diet Study and some trends over the last 30 years, *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 27(10) (2010) 1380-404.
- [55] N.N. Aung, J. Yoshinaga, J.I. Takahashi, Dietary intake of toxic and essential trace elements by the children and parents living in Tokyo Metropolitan Area, Japan, *Food Addit Contam* 23(9) (2006) 883-94.
- [56] S.M. Bratakos, A.E. Lazou, M.S. Bratakos, E.S. Lazos, Aluminium in food and daily dietary intake estimate in Greece, *Food Addit Contam Part B Surveill* 5(1) (2012) 33-44.

- [57] D. Gonzalez-Weller, A.J. Gutierrez, C. Rubio, C. Revert, A. Hardisson, Dietary intake of aluminum in a Spanish population (Canary Islands), *J Agric Food Chem* 58(19) (2010) 10452-7.
- [58] J.L. Greger, Dietary and other sources of aluminium intake, *Ciba Found Symp* 169 (1992) 26-35; discussion 35-49.
- [59] G.V. Iyenga, W.R. Wolfe, J.T. Tanner, E.R. Morris, Content of minor and trace elements, and organic nutrients in representative mixed total diet composites from the USA, *Sci Total Environ* 256(2-3) (2000) 215-26.
- [60] L. Jorhem, G. Haegglund, Aluminium in foodstuffs and diets in Sweden, *Z Lebensm Unters Forsch* 194(1) (1992) 38-42.
- [61] G. Ysart, P. Miller, H. Crews, P. Robb, M. Baxter, C. De L'Argy, S. Lofthouse, C. Sargent, N. Harrison, Dietary exposure estimates of 30 elements from the UK Total Diet Study, *Food Addit Contam* 16(9) (1999) 391-403.
- [62] G. Ellen, E. Egmond, J.W. Van Loon, E.T. Sahertian, K. Tolsma, Dietary intakes of some essential and non-essential trace elements, nitrate, nitrite and N-nitrosamines, by Dutch adults: estimated via a 24-hour duplicate portion study, *Food Addit Contam* 7(2) (1990) 207-21.
- [63] L. Noel, J.C. Leblanc, T. Guerin, Determination of several elements in duplicate meals from catering establishments using closed vessel microwave digestion with inductively coupled plasma mass spectrometry detection: estimation of daily dietary intake, *Food Addit Contam* 20(1) (2003) 44-56.
- [64] E.E. Santos, D.C. Lauria, C.L. Porto da Silveira, Assessment of daily intake of trace elements due to consumption of foodstuffs by adult inhabitants of Rio de Janeiro city, *Sci Total Environ* 327(1-3) (2004) 69-79.
- [65] G. Turconi, C. Minoia, A. Ronchi, C. Roggi, Dietary exposure estimates of twenty-one trace elements from a Total Diet Study carried out in Pavia, Northern Italy, *Br J Nutr* 101(8) (2009) 1200-8.
- [66] G. Ysart, P. Miller, M. Croasdale, H. Crews, P. Robb, M. Baxter, C. de L'Argy, N. Harrison, 1997 UK Total Diet Study--dietary exposures to aluminium, arsenic, cadmium, chromium, copper, lead, mercury, nickel, selenium, tin and zinc, *Food Addit Contam* 17(9) (2000) 775-86.
- [67] C. Vilarnau, D.M. Stracker, A. Funtikov, R. da Silva, R. Estruch, A. Bach-Faig, Worldwide adherence to Mediterranean Diet between 1960 and 2011, *Eur J Clin Nutr* (2018).

- [68] J.C. Leblanc, T. Guerin, L. Noel, G. Calamassi-Tran, J.L. Volatier, P. Verger, Dietary exposure estimates of 18 elements from the 1st French Total Diet Study, *Food Addit Contam* 22(7) (2005) 624-41.
- [69] J.A. Pennington, Aluminium content of foods and diets, *Food Addit Contam* 5(2) (1988) 161-232.
- [70] M. Müller, M. Anke, H. Illing-Günther, Aluminium in foodstuffs, *Food Chemistry* 61(4) (1998) 419-428.
- [71] G.S. Xu, R.P. Jin, Z.W. Zhang, W.Q. Zhang, D.L. Ren, J. Chen, G.W. Huang, Preliminary study on aluminum content of foods and aluminum intake of residents in Tianjin, *Biomed Environ Sci* 6(3) (1993) 319-25.
- [72] R.A. Yokel, C.L. Hicks, R.L. Florence, Aluminum bioavailability from basic sodium aluminum phosphate, an approved food additive emulsifying agent, incorporated in cheese, *Food Chem Toxicol* 46(6) (2008) 2261-6.
- [73] D.H. Calloway, J.J. McMullen, Fecal excretion of iron and tin by men fed stored canned foods, *Am J Clin Nutr* 18(1) (1966) 1-6.
- [74] J.M. Duggan, J.E. Dickeson, P.F. Tynan, A. Houghton, J.E. Flynn, Aluminium beverage cans as a dietary source of aluminium, *Med J Aust* 156(9) (1992) 604-5.
- [75] T. Inoue, H. Ishiwata, K. Yoshihira, Aluminum levels in food-simulating solvents and various foods cooked in aluminum pans, *J Agric Food Chem* 36(3) (1988) 599-601.
- [76] H. Liukkonen-Lilja, S. Piepponen, Leaching of aluminium from aluminium dishes and packages, *Food Addit Contam* 9(3) (1992) 213-23.
- [77] R.A. Yokel, R.L. Florence, Aluminum bioavailability from tea infusion, *Food Chem Toxicol* 46(12) (2008) 3659-63.
- [78] R.F. Milani, L.K. Silvestre, M.A. Morgano, S. Cadore, Investigation of twelve trace elements in herbal tea commercialized in Brazil, *Journal of Trace Elements in Medicine and Biology* 52 (2019) 111-117.
- [79] K.F. Fung, H.P. Carr, B.H. Poon, M.H. Wong, A comparison of aluminum levels in tea products from Hong Kong markets and in varieties of tea plants from Hong Kong and India, *Chemosphere* 75(7) (2009) 955-62.
- [80] T. Stahl, H. Taschan, H. Brunn, Aluminium content of selected foods and food products, *Environ Sci Eur* 23(1) (2011) 37.
- [81] S. Marin, O. Pardo, R. Baguena, G. Font, V. Yusa, Dietary exposure to trace elements and health risk assessment in the region of Valencia, Spain: a total diet study, *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 34(2) (2017) 228-240.

- [82] J.C. Sherlock, G.A. Smart, Tin in foods and the diet, *Food Addit Contam* 1(3) (1984) 277-82.
- [83] W. van Dokkum, R.H. de Vos, T. Muys, J.A. Wesstra, Minerals and trace elements in total diets in The Netherlands, *Br J Nutr* 61(1) (1989) 7-15.
- [84] A. Bocio, M. Nadal, J.L. Domingo, Human exposure to metals through the diet in Tarragona, Spain: temporal trend, *Biol Trace Elem Res* 104(3) (2005) 193-201.
- [85] J.M. Llobet, S. Granero, M. Schuhmacher, J. Corbella, J.L. Domingo, Biological monitoring of environmental pollution and human exposure to metals in Tarragona, Spain. IV. Estimation of the dietary intake, *Trace elements and electrolytes* 15(3) (1998) 136-141.
- [86] A. Buculei, G. Gutt, S. Amariei, D. Adriana, G. Constantinescu, Study regarding the tin and iron migration from metallic cans into foodstuff during storage, *Journal of Agroalimentary Processes and Technologies* 18(4) (2012) 299-303.
- [87] F.M. Makki, P. Ziarati, Determination of histamine and heavy metal concentrations in tomato pastes and fresh tomato (*Solanum lycopersicum*) in Iran, *Biosci Biotech Res Asia* 11(2) (2014) 537-544.
- [88] S. Mol, Determination of trace metals in canned anchovies and canned rainbow trouts, *Food Chem Toxicol* 49(2) (2011) 348-51.
- [89] A. Turrini, G. Lombardi-Boccia, F. Aureli, F. Cubadda, L. D'Addezio, M. D'Amato, L. D'Evoli, P. Darnerud, N. Devlin, M.G. Dias, M. Jurkovic, C. Kelleher, C. Le Donne, M. Lopez Esteban, M. Lucarini, M.A. Martinez Burgos, E. Martinez-Victoria, B. McNulty, L. Mistura, A. Nugent, H.I. Oktay Basegmez, L. Oliveira, H. Ozer, G. Perello, M. Pite, K. Presser, D. Sokolic, E. Vasco, J.L. Volatier, A conceptual framework for the collection of food products in a Total Diet Study, *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* (2017) 1-20.
- [90] A.C.A. Sousa, S.D. Coelho, M.R. Pastorinho, L. Taborda-Barata, A.J.A. Nogueira, T. Isobe, T. Kunisue, S. Takahashi, S. Tanabe, Levels of TBT and other selected organotin compounds in duplicate diet samples, *Sci Total Environ* 574 (2017) 19-23.
- [91] G. Berthon, Aluminium speciation in relation to aluminium bioavailability, metabolism and toxicity, *Coordination Chemistry Reviews* 228(2) (2002) 319-341.
- [92] R.A. Yokel, R.L. Florence, Aluminum bioavailability from the approved food additive leavening agent acidic sodium aluminum phosphate, incorporated into a baked good, is lower than from water, *Toxicology* 227(1-2) (2006) 86-93.
- [93] G.B. van der Voet, Intestinal Absorption of Aluminium, *Ciba Foundation Symposium 169 - Aluminium in Biology and Medicine* (2007).

[94] H. Rudel, Case study: bioavailability of tin and tin compounds, *Ecotoxicol Environ Saf* 56(1) (2003) 180-9.

[95] P.W. Looser, S. Bertschi, K. Fent, Bioconcentration and bioavailability of organotin compounds: Influence of pH and humic substances, *Applied Organometallic Chemistry* 12(8-9) (1998) 601-611.

[96] M. Ferrante, S.S. Signorelli, S.L. Ferlito, A. Grasso, A. Dimartino, C. Copat, Groundwater-based water wells characterization from Guinea Bissau (Western Africa): A risk evaluation for the local population, *Sci Total Environ* 619-620 (2018) 916-926.

[97] A. Lione, Aluminum toxicology and the aluminum-containing medications, *Pharmacol Ther* 29(2) (1985) 255-85.

[98] C.M. Reinke, J. Breitzkreutz, H. Leuenberger, Aluminium in over-the-counter drugs: risks outweigh benefits?, *Drug Saf* 26(14) (2003) 1011-25.

ACCEPTED MANUSCRIPT

Table 1. Food intake (g/day) according to different categories for the whole study population and by sex.

Foods	All (N=719)	Men (N=319)	Women (N=400)
	Mean (St. Dev.)	Mean (St. Dev.)	Mean (St. Dev.)
Cereals and cereal products	188.5 (99.3)	206.4 (104.6)	174.2 (92.5)
Pasta, other grain	57.1 (40.7)	70.1 (45.3)	46.8 (33.2)
Rice	5.6 (7.6)	6.2 (8.7)	5.0 (6.7)
Bread	78.3 (72.0)	83.9 (73.4)	73.9 (70.6)
Crackers, crispbread, salty snacks	47.5 (34.2)	46.2 (35.7)	48.5 (33.0)
Meat and meat products	128.4 (70.9)	142.4 (73.5)	117.2 (66.8)
Red meat	67.4 (45.1)	76.8 (49.6)	59.9 (39.7)
White meat	29.7 (26.2)	31.2 (25.6)	28.4 (26.6)
Processed meat	29.5 (24.9)	32.3 (27.2)	27.3 (22.7)
Offal	1.8 (4.6)	2.1 (5.2)	1.6 (4.0)
Milk and dairy products	229.4 (216.1)	201.8 (191.7)	251.5 (231.6)
Milk and yogurt	188.9 (209.3)	159.4 (188.3)	212.5 (222.0)
Cheese	40.5 (33.9)	42.4 (35.5)	39.0 (32.5)
Fresh cheese	14.6 (19.0)	11.9 (15.5)	16.8 (21.1)
Aged cheese	25.9 (24.0)	30.5 (27.6)	22.2 (20.1)
Eggs	15.1 (11.4)	14.6 (11.2)	15.4 (11.5)
Fish and seafood	35.1 (28.1)	35.5 (26.8)	34.9 (29.0)
Fish	27.9 (23.1)	28.5 (22.6)	27.5 (23.6)
Preserved and tinned fish	9.3 (10.7)	10.2 (9.7)	8.5 (11.4)
Non-piscivorous fish	10.4 (12.3)	10.8 (13.1)	10.2 (11.6)
Piscivorous fish	8.2 (11.7)	7.5 (10.7)	8.7 (12.4)
Crustaceans and molluscs	7.2 (10.1)	7.0 (9.4)	7.4 (10.6)
All vegetables	160.7 (94.7)	156.6 (88.8)	163.9 (99.2)
Leafy vegetables	31.6 (25.6)	29.4 (22.8)	33.3 (27.5)
Other vegetables	27.3 (20.0)	24.0 (17.0)	29.9 (21.8)
Tomatoes	63.4 (50.8)	67.0 (52.0)	60.5 (49.7)
Root vegetables	14.2 (19.7)	11.0 (16.7)	16.8 (21.4)
Cabbage	4.2 (7.1)	3.8 (8.0)	4.4 (6.2)
Mushrooms	2.5 (4.0)	2.5 (4.2)	2.5 (3.8)
Onion and garlic	17.5 (21.9)	18.9 (22.0)	16.5 (21.8)
Legumes	18.7 (18.6)	19.5 (19.3)	18.1 (18.1)
Potatoes	24.5 (24.2)	25.5 (26.4)	23.7 (22.3)
Fresh fruit	279.4 (165.7)	270.3 (162.8)	286.7 (167.8)
Citrus fruit	216.3 (135.6)	207.0 (132.9)	223.8 (137.5)
All other fruit	63.1 (50.0)	63.3 (49.5)	62.9 (50.5)
Dried fruit, nuts and seeds	1.7 (3.0)	1.8 (2.9)	1.6 (3.0)
Dried fruit	0.4 (1.1)	0.5 (1.2)	0.3 (1.0)
Nuts and seeds	1.3 (2.5)	1.3 (2.3)	1.2 (2.8)
Sweets, chocolate, cakes, etc.	86.5 (73.8)	82.9 (77.3)	89.4 (70.8)
Sugar, non-chocolate confectionery	18.5 (22.5)	17.3 (18.2)	19.4 (25.4)
Chocolate, candy bars, etc.	5.3 (8.7)	4.9 (8.5)	5.6 (8.9)
Ice cream	13.9 (15.6)	13.6 (16.9)	14.2 (14.5)
Cakes, pies and pastries	35.1 (54.3)	34.0 (61.6)	36.0 (47.7)
Biscuits, dried cakes	13.7 (17.1)	13.1 (17.2)	14.2 (17.1)
Oils and fats	27.2 (13.5)	27.9 (12.7)	26.7 (14.1)
Vegetable fats and oils (non-olive)	2.4 (5.7)	2.6 (6.1)	2.2 (5.4)
Olive oil	22.0 (12.6)	22.0 (12.1)	22.1 (12.9)
Butter and other animal fats	2.8 (3.5)	3.3 (4.1)	2.4 (2.9)
Beverages	429.5 (341.3)	479.5 (322.5)	389.7 (350.8)
Coffee and tea	148.4 (160.3)	120.6 (109.3)	170.6 (188.7)
Wines	126.0 (164.1)	191.4 (190.0)	73.9 (116.3)
Red wine	74.5 (119.0)	116.6 (144.1)	41.0 (80.0)
White wine	51.5 (102.0)	74.8 (123.8)	32.9 (75.8)
Aperitif wines and beers	41.4 (117.5)	49.9 (123.8)	34.6 (126.0)
Spirits and liqueurs	3.1 (11.0)	5.3 (14.9)	1.3 (5.7)
Fruit juices	66.4 (135.0)	67.6 (126.7)	65.5 (141.4)
Soft drinks	44.2 (118.2)	44.7 (100.7)	43.8 (130.6)

Table 2. Aluminum and tin concentrations ($\mu\text{g}/\text{kg}$) in food samples by food category. Median (50^{th}) and interquartile range (IQR) are reported.

Foods	N	Aluminum ($\mu\text{g}/\text{kg}$)		Tin ($\mu\text{g}/\text{kg}$)	
		50^{th}	(IQR)	50^{th}	(IQR)
Cereals and cereal products	112	2470.36	(1466.80-5685.81)	3.60	(0.97-8.58)
Pasta, other grain	41	1602.21	(960.83-2498.67)	2.97	(0.78-5.53)
Rice	8	276.12	(120.03-439.64)	0.52	(0.13-4.09)
Bread	42	3555.10	(2257.65-7756.52)	6.81	(3.60-11.38)
Crackers, crispbread, salty snacks	21	3515.11	(2607.98-5691.34)	2.66	(1.21-4.91)
Meat and meat products	86	584.59	(367.16-1181.77)	5.73	(3.00-12.37)
Red meat	28	478.27	(339.87-801.01)	5.90	(3.14-8.48)
White meat	12	386.29	(272.67-493.60)	3.08	(2.32-6.44)
Processed meat	36	1227.92	(750.19-2004.49)	8.54	(4.35-17.99)
Offal	10	334.31	(80.83-494.37)	4.49	(1.91-10.22)
Milk and dairy products	72	442.54	(218.02-1117.32)	3.28	(1.53-6.11)
Milk and yogurt	13	56.84	(29.30-63.00)	0.37	(0.17-0.88)
Cheese	59	647.04	(322.31-1622.85)	3.74	(2.39-7.56)
Fresh cheese	17	794.34	(312.66-1215.75)	3.00	(1.40-4.89)
Aged cheese	42	613.66	(336.53-1622.85)	5.05	(2.58-8.09)
Eggs	9	127.59	(77.54-168.41)	0.45	(0.26-0.97)
Fish and seafood	62	973.05	(432.43-2948.74)	4.31	(2.26-14.55)
Fish	41	530.75	(312.13-1181.82)	4.88	(2.62-16.64)
Preserved and tinned fish	9	589.84	(274.58-1088.94)	10.41	(3.82-15.56)
Non-piscivorous fish	15	568.86	(269.82-1584.56)	3.96	(1.51-11.35)
Piscivorous fish	17	478.79	(393.96-1012.41)	6.74	(2.37-23.37)
Crustaceans and molluscs	21	4258.89	(1633.63-5995.71)	3.50	(1.83-8.92)
All vegetables	201	858.22	(283.84-2732.59)	3.79	(1.26-13.01)
Leafy vegetables	40	9039.40	(1921.10-40667.56)	7.36	(4.14-18.92)
Other vegetables	63	815.34	(286.17-2121.31)	3.22	(1.14-17.70)
Tomatoes	19	356.47	(155.76-956.53)	2.94	(0.97-12.12)
Root vegetables	14	1358.58	(283.84-2428.15)	2.28	(1.12-5.34)
Cabbage	28	666.84	(290.25-1432.59)	4.03	(2.07-8.65)
Mushrooms	5	1855.89	(1484.48-2339.60)	17.09	(17.02-49.23)
Onion and garlic	32	242.67	(146.24-674.40)	1.68	(0.77-3.31)
Legumes	43	7370.23	(1231.42-15515.08)	1.65	(0.00-4.99)
Potatoes	14	471.93	(336.16-1225.13)	2.19	(1.32-4.89)
Fresh fruit	60	353.20	(177.85-706.13)	1.58	(1.04-2.52)
Citrus fruit	12	102.51	(60.22-187.47)	1.72	(1.30-5.46)
All other fruit	48	437.92	(235.44-846.96)	1.58	(0.92-2.39)
Dried fruit, nuts and seeds	45	1303.11	(571.52-3226.65)	4.53	(2.37-8.21)
Dried fruit	8	8318.11	(2737.64-26133.22)	5.43	(2.73-34.94)
Nuts and seeds	37	845.03	(367.23-2535.28)	3.65	(2.31-8.10)
Sweets, chocolate, cakes, etc.	79	4387.24	(1349.90-7949.51)	6.01	(3.14-10.21)
Sugar, non-chocolate confectionery	8	3555.18	(644.19-7552.50)	9.73	(5.70-14.62)
Chocolate, candy bars, etc.	21	14050.13	(7830.60-26330.64)	8.50	(4.89-13.25)
Ice cream	5	368.01	(90.60-627.10)	2.29	(1.84-2.40)
Cakes, pies and pastries	30	2339.53	(866.36-5138.26)	4.20	(1.78-8.33)
Biscuits, dried cakes	15	4436.53	(2584.85-5335.13)	7.19	(4.79-13.85)
Oils and fats	23	308.76	(167.71-449.64)	2.04	(1.00-39.95)
Vegetable fats and oils (non-olive)	12	284.61	(185.04-496.17)	6.62	(1.21-20.89)
Olive oil	4	238.54	(48.57-589.19)	1.63	(0.99-22.43)
Butter and other animal fats	7	314.81	(193.24-449.64)	1.90	(0.14-5.30)
Beverages	102	364.73	(91.96-885.03)	1.09	(0.45-2.84)
Coffee and tea	8	2411.92	(597.90-8213.29)	1.81	(0.16-7.49)
Wines	50	564.51	(324.03-978.96)	1.17	(0.60-1.99)
Red wine	27	546.80	(324.03-843.16)	1.72	(1.16-3.30)
White wine	23	807.22	(227.28-1019.64)	0.63	(0.45-1.11)
Aperitif wines and beers	8	102.75	(30.95-523.99)	0.95	(0.56-1.91)
Spirits and liqueurs	21	73.14	(33.44-167.52)	3.58	(0.60-6.23)
Fruit juices	8	71.78	(42.07-99.46)	0.39	(0.08-0.71)
Soft drinks	7	282.42	(21.93-1122.58)	0.21	(0.11-3.38)

Table 3. Estimated aluminum and tin intake ($\mu\text{g}/\text{day}$) by food category. Values of median (50th), interquartile range (IQR), and 95th percentile are reported.

Foods	Aluminum ($\mu\text{g}/\text{day}$)			Tin ($\mu\text{g}/\text{day}$)		
	50 th	(IQR)	95 th	50 th	(IQR)	95 th
<i>Total intake ($\mu\text{g}/\text{day}$)</i>	<i>4103.98</i>	<i>(3309.68-5162.74)</i>	<i>7226.66</i>	<i>66.80</i>	<i>(46.72-93.66)</i>	<i>147.67</i>
Cereals and cereal products	693.05	(430.42-1031.53)	1592.24	2.46	(1.60-3.46)	5.26
Pasta, other grain	194.72	(111.94-302.00)	525.48	1.19	(0.68-1.84)	3.20
Rice	1.14	(0.36-1.99)	6.31	0.02	(0.01-0.04)	0.11
Bread	337.62	(111.98-611.09)	1157.47	0.77	(0.25-1.39)	2.62
Crackers, crispbread, salty snacks	88.53	(51.95-141.85)	263.03	0.19	(0.10-0.34)	0.65
Meat and meat products	90.95	(60.16-127.91)	224.74	7.67	(4.36-12.65)	24.88
Red meat	31.77	(17.70-47.52)	80.68	0.52	(0.29-0.81)	1.40
White meat	10.00	(4.93-18.51)	35.90	0.13	(0.06-0.23)	0.43
Processed meat	40.18	(21.39-67.65)	135.31	6.96	(3.71-11.72)	23.44
Offal	0.00	(0.00-0.57)	2.84	0.00	(0.00-0.02)	0.09
Milk and dairy products	52.96	(31.81-78.90)	146.82	8.29	(4.08-14.78)	28.24
Milk and yogurt	9.37	(2.50-16.12)	34.03	0.09	(0.03-0.15)	0.33
Cheese	40.93	(22.06-67.41)	129.52	8.14	(4.03-14.72)	28.06
Fresh cheese	9.79	(3.26-19.57)	51.32	0.03	(0.01-0.07)	0.18
Aged cheese	26.31	(13.02-47.79)	91.55	8.06	(3.99-14.65)	28.06
Eggs	1.77	(0.96-2.78)	4.36	0.01	(0.00-0.01)	0.02
Fish and seafood	68.38	(31.53-170.46)	409.74	0.75	(0.40-1.22)	2.41
Fish	23.83	(11.54-36.94)	80.34	0.71	(0.35-1.15)	2.35
Preserved and tinned fish	5.63	(2.61-11.27)	22.96	0.48	(0.20-1.01)	2.03
Non-piscivorous fish	8.47	(2.03-18.44)	46.71	0.05	(0.01-0.12)	0.34
Piscivorous fish	3.60	(0.39-10.16)	27.94	0.06	(0.01-0.20)	0.58
Crustaceans and molluscs	39.08	(5.58-130.51)	376.91	0.02	(0.00-0.07)	0.16
All vegetables	1032.95	(671.34-1598.96)	2831.48	20.26	(9.29-38.41)	81.36
Leafy vegetables	624.49	(337.32-1100.17)	2061.89	0.32	(0.17-0.56)	1.05
Other vegetables	95.48	(52.80-163.30)	294.07	1.95	(1.08-3.34)	6.00
Tomatoes	156.48	(60.98-317.06)	670.26	16.66	(6.07-35.89)	77.42
Root vegetables	26.85	(6.32-72.65)	202.94	0.04	(0.01-0.11)	0.29
Cabbage	2.64	(0.00-8.06)	25.64	0.01	(0.00-0.04)	0.12
Mushrooms	2.32	(0.58-7.75)	16.66	0.04	(0.01-0.12)	0.26
Onion and garlic	4.80	(2.15-9.59)	28.15	0.03	(0.01-0.06)	0.18
Legumes	139.59	(64.66-259.68)	550.15	0.21	(0.10-0.40)	0.84
Potatoes	13.66	(8.12-24.29)	53.13	0.19	(0.11-0.33)	0.73
Fresh fruit	145.18	(92.04-210.48)	348.22	10.32	(5.17-14.88)	27.02
Citrus fruit	135.74	(83.75-196.77)	322.08	0.70	(0.43-1.02)	1.67
All other fruit	9.13	(4.36-13.08)	24.43	9.54	(4.56-13.68)	25.54
Dried fruit, nuts and seeds	2.44	(1.44-12.22)	56.06	0.003	(0.002-0.014)	0.07
Dried fruit	1.48	(0.00-1.48)	35.47	0.002	(0.000-0.002)	0.04
Nuts and seeds	0.96	(0.96-6.22)	33.99	0.001	(0.001-0.009)	0.05
Sweets, chocolate, cakes, etc.	283.39	(147.38-480.23)	957.61	0.54	(0.30-0.91)	1.77
Sugar, non-chocolate confectionery	52.96	(17.65-117.31)	235.89	0.13	(0.04-0.29)	0.57
Chocolate, candy bars, etc.	34.04	(0.00-114.14)	450.56	0.02	(0.00-0.06)	0.25
Ice cream	4.24	(1.02-8.08)	17.50	0.03	(0.01-0.05)	0.11
Cakes, pies and pastries	45.95	(10.60-168.05)	385.58	0.11	(0.03-0.42)	0.96
Biscuits, dried cakes	29.28	(0.00-111.55)	224.02	0.06	(0.00-0.21)	0.43
Oils and fats	8.10	(6.00-10.97)	17.92	4.41	(1.37-10.26)	25.44
Vegetable fats and oils (non-olive)	0.18	(0.09-0.41)	5.60	0.25	(0.13-0.51)	15.96
Olive oil	6.41	(4.46-9.02)	14.70	0.24	(0.16-0.33)	0.54
Butter and other animal fats	0.57	(0.17-1.30)	2.78	2.77	(0.31-7.08)	14.78
Beverages	1175.13	(770.85-1676.82)	2825.79	1.12	(0.95-3.44)	4.64
Coffee and tea	1020.53	(656.61-1533.78)	2628.13	0.51	(0.26-0.73)	1.24
Wines	40.77	(1.53-189.86)	380.32	0.22	(0.00-1.40)	3.78
Red wine	12.07	(0.00-86.27)	253.12	0.18	(0.00-1.28)	3.75
White wine	1.62	(0.00-37.05)	192.57	0.00	(0.00-0.04)	0.21
Aperitif wines and beers	1.17	(0.00-3.32)	23.52	0.02	(0.00-0.04)	0.30
Spirits and liqueurs	0.00	(0.00-0.13)	3.08	0.00	(0.00-0.00)	0.06
Fruit juices	2.09	(0.00-9.70)	33.98	0.01	(0.00-0.04)	0.13
Soft drinks	0.00	(0.00-20.88)	125.40	0.00	(0.00-0.04)	0.27

Table 4. Dietary weekly intake (DWI) and tolerable weekly intake (TWI) of aluminum and tin in the study population compared with data reported by international agencies. Values reported in mg/kg of body weight (bw) per week, based on a body weight of 60 kg^a.

Metal	This study		EFSA ^b		WHO ^c	
	50 th DWI	95 th DWI	DWI	TWI	DWI	TWI
Aluminum	0.41	0.78	0.20-1.50	1.00	0.20-2.26	2.00
Tin	0.007	0.015	0.18-0.63	ND ^d	0.06-0.35	2.00

^aDI: dietary intake; EFSA: European Food Safety Authority; WDI: Weekly dietary intake; TWI: tolerable weekly intake; WHO: World Health Organization.

^bData from ranges of average dietary intake obtained from EFSA Scientific Opinions [29, 36].

^cData from ranges of average dietary intake obtained from FAO/WHO [24, 49, 50].

^dTolerable upper intake level was not derived due to insufficient data, but regulatory limits of tin contents in canned food (200 mg/kg) and beverages (100 mg/kg) have been established [36].

Figure 1. Aluminum (A) and tin (B) food content ($\mu\text{g}/\text{kg}$), and contribution to dietary intake (% - percentage) by food category in the whole population.

