

ADOPTED: 25 November 2021

doi: 10.2903/j.efsa.2022.7024

## Pest categorisation of *Maconellicoccus hirsutus*

EFSA Panel on Plant Health (PLH),

Claude Bragard, Paula Baptista, Elisavet Chatzivassiliou, Francesco Di Serio, Paolo Gonthier, Josep Anton Jaques Miret, Annemarie Fejer Justesen, Christer Sven Magnusson, Panagiotis Milonas, Juan A Navas-Cortes, Stephen Parnell, Roel Potting, Philippe Lucien Reignault, Emilio Stefani, Hans-Hermann Thulke, Wopke Van der Werf, Antonio Vicent Civera, Jonathan Yuen, Lucia Zappalà, Jean-Claude Gregoire, Chris Malumphy, Spyridon Antonatos, Virag Kertesz, Andrea Maiorano, Dimitrios Papachristos and Alan MacLeod

### Abstract

The EFSA Panel on Plant Health performed a pest categorisation of *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), the pink hibiscus mealybug, for the EU. *M. hirsutus* is native to Southern Asia and has established in many countries in tropical and subtropical regions throughout the world. Within the EU, the pest has been reported from Cyprus and Greece (Rhodes). *M. hirsutus* is not listed in Annex II of Commission Implementing Regulation (EU) 2019/2072. It is highly polyphagous, feeding on plants assigned to 229 genera in 78 plant families, and shows some preference for hosts in the families Malvaceae, Fabaceae and Moraceae. Economically important crops in the EU such as cotton (*Gossypium* spp.), citrus (*Citrus* spp.), ornamentals (*Hibiscus* spp.), grapes (*Vitis vinifera*), soybean (*Glycine max*), avocado (*Persea americana*) and mulberry trees (*Morus alba*) may be significantly affected by *M. hirsutus*. The lower and upper developmental temperature threshold of *M. hirsutus* on *Hibiscus rosa-sinensis* are 14.5 and 35.0°C, respectively, with optimal female development estimated to be at 29.0°C. There are about 10 generations a year in the subtropics but as many as 15 may occur under optimal conditions. Plants for planting, fruits, vegetables and cut flowers provide potential pathways for entry into the EU. Climatic conditions in EU member states around the Mediterranean Sea and host plant availability in those areas are conducive for establishment. The introduction of *M. hirsutus* is expected to have an economic impact in the EU through damage to various ornamental plants, as already observed in Cyprus and Greece, and reduction in yield and quality of many significant crops. Phytosanitary measures are available to reduce the likelihood of entry and further spread. Some uncertainties include the area of establishment, whether it could become a greenhouse pest, impact, and the influence of natural enemies. *M. hirsutus* meets the criteria that are within the remit of EFSA to assess for it to be regarded as a potential Union quarantine pest.

© 2022 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

**Keywords:** pink hibiscus mealybug, Hemiptera, pest risk, plant health, plant pest, Pseudococcidae, quarantine

**Requestor:** European Commission

**Question number:** EFSA-Q-2021-00490

**Correspondence:** alpha@efsa.europa.eu

**Panel members:** Claude Bragard, Paula Baptista, Elisavet Chatzivassiliou, Francesco Di Serio, Paolo Gonthier, Josep Anton Jaques Miret, Annemarie Fejer Justesen, Alan MacLeod, Christer Sven Magnusson, Panagiotis Milonas, Juan A Navas-Cortes, Stephen Parnell, Roel Potting, Philippe L Reignault, Emilio Stefani, Hans-Hermann Thulke, Wopke Van der Werf, Antonio Vicent, Jonathan Yuen and Lucia Zappalà.

**Declarations of interest:** The declarations of interest of all scientific experts active in EFSA's work are available at <https://ess.efsa.europa.eu/doi/doiweb/doisearch>.

**Acknowledgments:** EFSA wishes to acknowledge the contribution of Caterina Campese and Oresteia Sfyra to this opinion.

**Suggested citation:** EFSA PLH Panel (EFSA Panel on Plant Health), Bragard C, Baptista P, Chatzivassiliou E, Di Serio F, Gonthier P, Jaques Miret JA, Justesen AF, Magnusson CS, Milonas P, Navas-Cortes JA, Parnell S, Potting R, Reignault PL, Stefani E, Thulke H-H, Van der Werf W, Vicent Civera A, Yuen J, Zappalà L, Gregoire J-C, Malumphy C, Antonatos S, Kertesz V, Maiorano A, Papachristos D and MacLeod A, 2022. Scientific Opinion on the pest categorisation of *Maconellicoccus hirsutus*. EFSA Journal 2022;20(1):7024, 45 pp. <https://doi.org/10.2903/j.efsa.2022.7024>

**ISSN:** 1831-4732

© 2022 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs](https://creativecommons.org/licenses/by/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.

Reproduction of the images listed below is prohibited and permission must be sought directly from the copyright holder:

Figure 1: © Courtesy of Cris Malumphy



The EFSA Journal is a publication of the European Food Safety Authority, a European agency funded by the European Union.



## Table of contents

Abstract.....	1
1. Introduction.....	4
1.1. Background and Terms of Reference as provided by the requestor.....	4
1.1.1. Background.....	4
1.1.2. Terms of reference.....	4
1.2. Interpretation of the Terms of Reference.....	4
1.3. Additional information.....	5
2. Data and methodologies.....	5
2.1. Data.....	5
2.1.1. Information on pest status from NPPOs.....	5
2.1.2. Literature search.....	5
2.1.3. Database search.....	5
2.2. Methodologies.....	5
3. Pest categorisation.....	6
3.1. Identity and biology of the pest.....	6
3.1.1. Identity and taxonomy.....	6
3.1.2. Biology of the pest.....	7
3.1.3. Host range/species affected.....	9
3.1.4. Intraspecific diversity.....	9
3.1.5. Detection and identification of the pest.....	9
3.2. Pest distribution.....	10
3.2.1. Pest distribution outside the EU.....	10
3.2.2. Pest distribution in the EU.....	11
3.3. Regulatory status.....	11
3.3.1. Commission Implementing Regulation 2019/2072.....	11
3.3.2. Hosts or species affected that are prohibited from entering the Union from third countries.....	11
3.4. Entry, establishment and spread in the EU.....	13
3.4.1. Entry.....	13
3.4.2. Establishment.....	14
3.4.2.1. EU distribution of main host plants.....	14
3.4.2.2. Climatic conditions affecting establishment.....	15
3.4.3. Spread.....	15
3.5. Impacts.....	16
3.6. Available measures and their limitations.....	16
3.6.1. Identification of potential additional measures.....	16
3.6.1.1. Additional potential risk reduction options.....	16
3.6.1.2. Additional supporting measures.....	18
3.6.1.3. Biological or technical factors limiting the effectiveness of measures.....	19
3.7. Uncertainty.....	19
4. Conclusions.....	19
References.....	20
Abbreviations.....	22
Glossary.....	23
Appendix A – <i>Maconellicoccus hirsutus</i> host plants/species affected.....	24
Appendix B – Distribution of <i>Maconellicoccus hirsutus</i> .....	37
Appendix C – Import data.....	41

## 1. Introduction

### 1.1. Background and Terms of Reference as provided by the requestor

#### 1.1.1. Background

The new Plant Health Regulation (EU) 2016/2031, on the protective measures against pests of plants, is applying from 14 December 2019. Conditions are laid down in this legislation in order for pests to qualify for listing as Union quarantine pests, protected zone quarantine pests or Union regulated non-quarantine pests. The lists of the EU regulated pests together with the associated import or internal movement requirements of commodities are included in Commission Implementing Regulation (EU) 2019/2072. Additionally, as stipulated in the Commission Implementing Regulation 2018/2019, certain commodities are provisionally prohibited to enter in the EU (high risk plants, HRP). EFSA is performing the risk assessment of the dossiers submitted by exporting to the EU countries of the HRP commodities, as stipulated in Commission Implementing Regulation 2018/2018. Furthermore, EFSA has evaluated a number of requests from exporting to the EU countries for derogations from specific EU import requirements.

In line with the principles of the new plant health law, the European Commission with the Member States are discussing monthly the reports of the interceptions and the outbreaks of pests notified by the Member States. Notifications of an imminent danger from pests that may fulfil the conditions for inclusion in the list of the Union quarantine pest are included. Furthermore, EFSA has been performing horizon scanning of media and literature.

As a follow-up of the above-mentioned activities (reporting of interceptions and outbreaks, HRP, derogation requests and horizon scanning), a number of pests of concern have been identified. EFSA is requested to provide scientific opinions for these pests, in view of their potential inclusion by the risk manager in the lists of Commission Implementing Regulation (EU) 2019/2072 and the inclusion of specific import requirements for relevant host commodities, when deemed necessary by the risk manager.

#### 1.1.2. Terms of reference

EFSA is requested, pursuant to Article 29(1) of Regulation (EC) No 178/2002, to provide scientific opinions in the field of plant health.

EFSA is requested to deliver 53 pest categorisations for the pests listed in Annex 1A, 1B, 1D and 1E (for more details see mandate M-2021-00027 on the [Open.EFSA portal](#)). Additionally, EFSA is requested to perform pest categorisations for the pests so far not regulated in the EU, identified as pests potentially associated with a commodity in the commodity risk assessments of the HRP dossiers (Annex 1C; for more details see mandate M-2021-00027 on the [Open.EFSA portal](#)). Such pest categorisations are needed in the case where there are not available risk assessments for the EU.

When the pests of Annex 1A are qualifying as potential Union quarantine pests, EFSA should proceed to phase 2 risk assessment. The opinions should address entry pathways, spread, establishment, impact and include a risk reduction options analysis.

Additionally, EFSA is requested to develop further the quantitative methodology currently followed for risk assessment, in order to have the possibility to deliver an express risk assessment methodology. Such methodological development should take into account the EFSA Plant Health Panel Guidance on quantitative pest risk assessment and the experience obtained during its implementation for the Union candidate priority pests and for the likelihood of pest freedom at entry for the commodity risk assessment of High Risk Plants.

### 1.2. Interpretation of the Terms of Reference

*Maconellicoccus hirsutus* is one of a number of pests listed in Annex 1A to the Terms of Reference (ToR) (Section 1.1.2) to be subject to pest categorisation to determine whether it fulfils the criteria of a potential Union quarantine pest for the area of the EU excluding Ceuta, Melilla and the outermost regions of Member States referred to in Article 355(1) of the Treaty on the Functioning of the European Union (TFEU), other than Madeira and the Azores, and so inform European Commission decision making as to its appropriateness for potential inclusion in the lists of pests of Commission Implementing Regulation (EU) 2019/2072. If a pest fulfils the criteria to be potentially listed as a Union quarantine pest, risk reduction options will be identified.

### 1.3. Additional information

This pest categorisation was initiated following the commodity risk assessment of *Ficus carica* plants from Israel performed by EFSA (EFSA PLH Panel, 2021), in which *M. hirsutus* was identified as a relevant non-regulated EU pest which could potentially enter the EU on *F. carica*.

## 2. Data and methodologies

### 2.1. Data

#### 2.1.1. Information on pest status from NPPOs

In the context of the commodity risk assessment of *Ficus carica* plants from Israel (EFSA PLH Panel, 2021), EFSA consulted (in April-May 2020) the NPPOs where the pest is present, in order to have an updated information on the pest status. For the information on pest status in Cyprus and Greece, please see Section 3.2.2.

#### 2.1.2. Literature search

A literature search on *M. hirsutus* was conducted at the beginning of the categorisation in the ISI Web of Science bibliographic database, using the scientific name of the pest as search term. Papers relevant for the pest categorisation were reviewed, and further references and information were obtained from experts, as well as from citations within the references and grey literature.

#### 2.1.3. Database search

Pest information, on host(s) and distribution, was retrieved from the European and Mediterranean Plant Protection Organization (EPPO) Global Database (EPPO, online), the CABI databases and scientific literature databases as referred above in Section 2.1.2.

Data about the import of commodity types that could potentially provide a pathway for the pest to enter the EU and about the area of hosts grown in the EU were obtained from EUROSTAT (Statistical Office of the European Communities).

The Europhyt and TRACES databases were consulted for pest-specific notifications on interceptions and outbreaks. Europhyt is a web-based network run by the Directorate General for Health and Food Safety (DG SANTÉ) of the European Commission as a subproject of PHYSAN (Phyto-Sanitary Controls) specifically concerned with plant health information. TRACES is the European Commission's multilingual online platform for sanitary and phytosanitary certification required for the importation of animals, animal products, food and feed of non-animal origin and plants into the EU, and the intra-EU trade and EU exports of animals and certain animal products. Up until May 2020, the Europhyt database managed notifications of interceptions of plants or plant products that do not comply with EU legislation, as well as notifications of plant pests detected in the territory of the Member States and the phytosanitary measures taken to eradicate or avoid their spread. The recording of interceptions switched from Europhyt to TRACES in May 2020.

### 2.2. Methodologies

The Panel performed the pest categorisation for *M. hirsutus*, following guiding principles and steps presented in the EFSA guidance on quantitative pest risk assessment (EFSA PLH Panel, 2018), the EFSA guidance on the use of the weight of evidence approach in scientific assessments (EFSA Scientific Committee, 2017) and the International Standards for Phytosanitary Measures No. 11 (FAO, 2013).

The criteria to be considered when categorising a pest as a potential Union quarantine pest (QP) is given in Regulation (EU) 2016/2031 Article 3 and Annex I, Section 1 to this Regulation. Table 1 presents the Regulation (EU) 2016/2031 pest categorisation criteria on which the Panel bases its conclusions. In judging whether a criterion is met the Panel uses its best professional judgement (EFSA Scientific Committee, 2017) by integrating a range of evidence from a variety of sources (as presented above in Section 2.1) to reach an informed conclusion as to whether or not a criterion is satisfied.

The Panel's conclusions are formulated respecting its remit and particularly with regard to the principle of separation between risk assessment and risk management (EFSA founding regulation (EU) No 178/2002); therefore, instead of determining whether the pest is likely to have an unacceptable impact, deemed to be a risk management decision, the Panel will present a summary of the observed impacts in the areas where the pest occurs, and make a judgement about potential likely impacts in the EU. While the Panel may quote impacts reported from areas where the pest occurs in monetary terms, the Panel will seek to express potential EU impacts in terms of yield and quality losses and not in monetary terms, in agreement with the EFSA guidance on quantitative pest risk assessment (EFSA PLH Panel, 2018). Article 3 (d) of Regulation (EU) 2016/2031 refers to unacceptable social impact as a criterion for quarantine pest status. Assessing social impact is outside the remit of the Panel.

**Table 1:** Pest categorisation criteria under evaluation, as defined in Regulation (EU) 2016/2031 on protective measures against pests of plants (the number of the relevant sections of the pest categorisation is shown in brackets in the first column)

Criterion of pest categorisation	Criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest (article 3)
<b>Identity of the pest (Section 3.1)</b>	Is the identity of the pest established, or has it been shown to produce consistent symptoms and to be transmissible?
<b>Absence/presence of the pest in the EU territory (Section 3.2)</b>	Is the pest present in the EU territory? If present, is the pest widely distributed within the EU? Describe the pest distribution briefly
<b>Regulatory status (Section 3.3)</b>	If the pest is present in the EU but not widely distributed in the risk assessment area, it should be under official control or expected to be under official control in the near future
<b>Pest potential for entry, establishment and spread in the EU territory (Section 3.4)</b>	Is the pest able to enter into, become established in, and spread within, the EU territory? If yes, briefly list the pathways
<b>Potential for consequences in the EU territory (Section 3.5)</b>	Would the pests' introduction have an economic or environmental impact on the EU territory?
<b>Available measures (Section 3.6)</b>	Are there measures available to prevent the entry into the EU such that the likelihood of introduction becomes mitigated?
<b>Conclusion of pest categorisation (Section 4)</b>	A statement as to whether (1) all criteria assessed by EFSA above for consideration as a potential quarantine pest were met and (2) if not, which one(s) were not met

### 3. Pest categorisation

#### 3.1. Identity and biology of the pest

##### 3.1.1. Identity and taxonomy

*Is the identity of the pest established, or has it been shown to produce consistent symptoms and/or to be transmissible?*

**Yes**, the identity of the pest is established and *Maconellicoccus hirsutus* (Green) is the accepted name.

The pink hibiscus mealybug, also known as the hibiscus mealybug, *Maconellicoccus hirsutus* (Green, 1908) is an insect within the order Hemiptera, family Pseudococcidae. This species was initially described by Green in 1908 as *Phenacoccus hirsutus* from specimens collected on an undetermined shrub attended by ants in India (García Morales et al., 2016). Indeed, this species is likely to be native to southern Asia (Williams, 2004). Former scientific names include *Maconellicoccus pasaniae*, *Maconellicoccus perforatus*, *Paracoccus pasaniae*, *Phenacoccus glomeratus*, *Phenacoccus hirsutus*, *Phenacoccus quaternus*, *Pseudococcus hibisci* and *Spilococcus perforatus* (CABI, 2021). The genus *Maconellicoccus* includes eight described species (Williams, 1996; CABI, 2021). Detailed morphological descriptions, illustrations and keys to the eight species of the genus *Maconellicoccus* can be found in



Williams (1996), Meyerdirk et al. (2001) and EPPO (2006). The EPPO code (Griessinger & Roy, 2015; EPPO, 2019) for this species is PHENHI (EPPO, 2021).

### 3.1.2. Biology of the pest

Adult females of *M. hirsutus* in Jordan appear in early February and show their highest abundance in mid-July (Al-Fwaeer et al., 2014). *M. hirsutus* reproduces parthenogenetically or sexually (Williams, 1996). Reproduction is mostly parthenogenetic in Egypt and the State of Bihar, India (Hall, 1921; Singh and Ghosh, 1970), while it is sexual in the Indian state of West Bengal (Ghose, 1971) and probably in the Caribbean (Williams, 1996). According to Bartlett (1978) and Mani (1989), an adult female lays 150–600 eggs over a period of about 1 week on the host plants. The eggs are laid in an ovisac, consisting of a mass of sticky wax filaments. Oviposition occurs mainly in the outer parts of the host, such as the growing points, buds and fruits, but in case of cold weather conditions the females search for shelter to oviposit (Meyerdirk et al., 2001). The lower and upper developmental temperature threshold of *M. hirsutus* on *Hibiscus rosa-sinensis* are 14.5 and 35.0°C, respectively. The optimal developmental temperature for females was estimated to be 29.0°C (Chong et al., 2008). In warm, but unspecified conditions, it takes 5 weeks for a generation to be completed (Bartlett, 1978). Chong et al. (2008) stated that the generation time is 41 days at 25°C and 82 days at 20°C. In countries with a cool winter the species overwinters as eggs (Bartlett, 1978) or other stages in protected parts of the host plant or as eggs in the soil (Pollard, 1995). There are about 10 generations a year in the subtropics (Meyerdirk et al., 2001). However, under optimum conditions, there may be as many as 15 generations per year (Pollard, 1995).

There are three immature instars in the female and four in the male (EPPO, 2005). First instar nymphs are known as crawlers and are mobile. The crawlers prefer the apical and tender regions of the host. However, large populations of nymphs may also settle on the older plant parts including stems, leaves, petioles, roots, tubers and pods (Ghose, 1972). After locating a suitable feeding site on a host plant, nymphs settle to feed and develop. Later instars turn grey–pink and start to secrete white wax that covers their bodies (Chong et al., 2015). In heavy infestations white masses of wax concealing the insect may occur in axils and on twigs and stems (EPPO, 2006) (Figure 1). Female adults live for 19–28 days (Chong et al., 2008; Sahito et al., 2012; Negrini et al., 2017). Males have one pair of wings, but they are weak flyers, only live a day or two, and are not commonly observed (Chong et al., 2015).

Key features of the biology of each life stage are summarised in Table 2.



**Figure 1:** *Maconellicoccus hirsutus*: (A) adult female; (B) adult female covered in waxy filaments; (C) large infestation on hibiscus; (D) ovisacs in the crevices of *Annona* fruit; (E) distorted growth characteristic of plants infested by *M. hirsutus*; (F) hibiscus plant in Rhodes, severely damaged by *M. hirsutus* © Chris Malumphy

**Table 2:** Important features of the life history strategy of *Maconellicoccus hirsutus*

Life stage	Phenology and relation to host	Other relevant information
<b>Egg</b>	Adult female lays 150–600 eggs in a sticky waxy ovisac. Oviposition occurs mainly on the outer areas of the host, including the buds and fruit	The eggs hatch in 6–9 days at temperatures between 25 and 35°C but it requires 16 days at 20°C. The lower and upper threshold for the eggs and the optimal developmental temperature were estimated at 14.5, 39.8 and 33.4°C, respectively (Chong et al., 2008)
<b>First instar nymph</b>	First instar nymphs are known as crawlers. They prefer the apical and tender parts of the host. However, large populations of nymphs may also settle on the older plant parts	The crawlers disperse by walking to other parts of the host plant. They may also be transported by water, wind or animals



Life stage	Phenology and relation to host	Other relevant information
	including stems, leaves, petioles, roots, tubers, and pods. After locating a suitable host plant, nymphs settle on the host to feed and develop	
<b>Later instar nymphs</b>	Later instars start to secrete white wax that covers their bodies. There are three immature instars in the female and four in the male	White masses of wax concealing the insect may occur in axils and on twigs and stems. The nymphal development is affected by both temperature and host plant. At 25°C, the female nymphs need 23 and 26.6 days on <i>H. rosa-sinensis</i> and <i>Morus alba</i> , respectively, to complete their development (Chong et al., 2008; Sahito et al., 2012). On <i>H. rosa-sinensis</i> and at 27°C nymphal development was reported to last either 17.5 or 20.6 days (Chong et al., 2008; Negrini et al., 2017). Whereas at 30 and 20°C the female nymphal stages last 26.6 and 50.1 days, respectively. The lower and upper threshold and the optimal developmental temperature for female nymphs were estimated at 15.1, 35.0 and 28.8°C, respectively (Chong et al., 2008)
<b>Adult</b>	Males have one pair of wings, but they are weak flyers. Female adults live for 19–28 days (Chong et al., 2008; Sahito et al., 2012; Negrini et al., 2017) while males only 1 or 2 days and are not commonly observed (Chong et al., 2015)	<i>M. hirsutus</i> reproduces parthenogenetically or sexually. The lower and upper developmental temperature threshold on <i>H. rosa-sinensis</i> were 14.5 and 35°C, respectively. The optimal developmental temperature for females was estimated to be 29°C

### 3.1.3. Host range/species affected

There is a long list of host plants of *M. hirsutus* worldwide. The host range of *M. hirsutus* is broad with more than 229 plant genera from 78 plant families (García Morales et al., 2016). Appendix A provides the full list of plant species reported to be *M. hirsutus* hosts. Economically important crops in the EU such as cotton (*Gossypium* spp.), citrus (*Citrus* spp.), ornamentals (*Hibiscus* spp.), grapes (*Vitis vinifera*), soybean (*Glycine max*), avocado (*Persea americana*) and mulberry trees (*Morus alba*) may be significantly affected by *M. hirsutus*. *M. hirsutus* has also been recorded on several rosaceous crops that are important in the EU, including apple (*Malus domestica*), apricot (*Prunus armeniaca*), peach (*Prunus persica*), pear (*Pyrus communis*) and plum (*Prunus domestica*), but there appears to be no economic impact recorded on these hosts.

### 3.1.4. Intraspecific diversity

No intraspecific diversity is reported for this species.

### 3.1.5. Detection and identification of the pest

Are detection and identification methods available for the pest?

**Yes.** There are methods available for detection, and morphological and molecular identification of *M. hirsutus*.

## Detection

Careful visual examination of plants is an effective way for the detection of the insect. The white waxy covering of mealybug instars and white waxy filaments in the egg mass allow detection (Meyerdirk et al., 2001). The mealybugs themselves are in general visible, although they are hidden in the swollen growth. Male adults can also be caught using sticky cards baited with a sex pheromone

which contains esters of lavandulyl and maconellyl and allow detection in areas of low density of the pest (Francis et al., 2007).

## Symptoms

The main symptoms of *M. hirsutus* infestation are (Dufour and Léon, 1997; Sagarra and Peterkin, 1999; Kairo et al., 2000; Alleyne, 2004; Chong et al., 2015):

- large quantities of honeydew
- black sooty mould
- leaf curling
- shoot and leave malformation
- fruit malformation
- bunchy top appearance
- premature senescence of flowers and foliage
- heavy infestation may cause a complete defoliation of the plant, leading to their death

## Identification

The identification of *M. hirsutus* requires microscopic examination of slide-mounted adult females and verification of the presence of key morphological characteristics as given in Meyerdirk et al. (2001) and Williams (1996). Moreover, a key is available (EPPO, 2006) to distinguish *M. hirsutus* from other species of the genus. Molecular techniques for species identification have also been developed (Malausa et al., 2011; Abd-Rabou et al., 2012).

**Description** (detailed morphological descriptions are available from Meyerdirk et al. (2001) and EPPO (2006))


The main morphological characteristics of *M. hirsutus* are:

- The eggs are 0.3 mm long and initially orange, turning pink before hatching (Chong et al., 2015).
- Crawlers 0.37 mm long (Aristizábal et al., 2012), pink and oval with antennae; they lack the waxy body coating (CABI, 2021).
- Second instars average length 0.70 mm, third instars 1.1 mm and male fourth instar 1.1 mm (Aristizábal et al., 2012). Immature females and newly matured females have greyish-pink bodies dusted with mealy white wax (CABI, 2021).
- Mature adult females are wingless, elongate oval, slightly flattened in profile, 2.5–4 mm long, and their ovisacs cover most of the body. Body is greyish pink or occasionally purple, and covered with a thin white cotton like wax forming a protective ovisac for her eggs. The entire colony tends to become covered by white, waxy ovisac material (EPPO, 2005, 2006; Chong et al., 2015).
- On microscopic examination of slide-mounted females, the combination of nine-segmented antennae, anal lobe bars, numerous large dorsal oral rim ducts on all parts of the body, and long, flagellate dorsal setae make the species fairly easy to recognize in parts of the world where other *Maconellicoccus* species do not occur. Males have one pair of very simple wings, long antennae, white wax filaments projecting posteriorly and lack mouthparts CABI (2021).

## 3.2. Pest distribution

### 3.2.1. Pest distribution outside the EU

*M. hirsutus* has established in many tropical and subtropical regions throughout the world in the past 100 years (Culik et al., 2013). It has a wide distribution which includes many countries in Africa, South Asia, Australia, Central America, South America, Caribbean and the southern part of North America (EPPO, 2021) (Figure 2). For a detailed list of countries where *M. hirsutus* is present, see Appendix B.



**Figure 2:** Global distribution of *Maconellicoccus hirsutus* (Source: EPPO Global Database accessed on 15/10/2021)

### 3.2.2. Pest distribution in the EU

*Is the pest present in the EU territory? If present, is the pest widely distributed within the EU?*

*M. hirsutus* has a restricted distribution in the EU. It is present in Greece and Cyprus.

The pest is widespread only in the island of Rhodes in southern eastern part of Greece. The pest is present, widespread and under official control in Cyprus (EPPO GD, online).

According to Miller et al. (2014), between 1995 and 2012 the species had been intercepted in USA ports in commodities originating from France and Italy. However, there are no records of the species from France and Italy. This has probably resulted from produce being imported to Europe from areas where the mealybug occurs and re-exported to the USA.

## 3.3. Regulatory status

### 3.3.1. Commission Implementing Regulation 2019/2072

*M. hirsutus* is not listed in Annex II of Commission Implementing Regulation (EU) 2019/2072.

### 3.3.2. Hosts or species affected that are prohibited from entering the Union from third countries

According to the Commission Implementing Regulation (EU) 2019/2072, Annex VI, introduction of several *M. hirsutus* hosts in the Union from certain third countries is prohibited (Table 3).

**Table 3:** List of plants, plant products and other objects that are *Maconellicoccus hirsutus* hosts whose introduction into the Union from certain third countries is prohibited (Source Commission Implementing Regulation (EU) 2019/2072, Annex VI)

<b>List of plants, plant products and other objects whose introduction into the Union from certain third countries is prohibited</b>			
	<b>Description</b>	<b>CN Code</b>	<b>Third country, group of third countries or specific area of third country</b>
8.	Plants for planting of <i>Chaenomeles</i> Ldl., <i>Crateagus</i> L., <i>Cydonia</i> Mill., <i>Malus</i> Mill., <i>Prunus</i> L., <i>Pyrus</i> L. and <i>Rosa</i> L., other than dormant plants free from leaves, flowers and fruits	ex 0602 10 90 ex 0602 20 20 ex 0602 20 80 ex 0602 40 00 ex 0602 90 41 ex 0602 90 45 ex 0602 90 46 ex 0602 90 47 ex 0602 90 48 ex 0602 90 50 ex 0602 90 70 ex 0602 90 91 ex 0602 90 99	Third countries other than: Albania, Andorra, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canary Islands, Faeroe Islands, Georgia, Iceland, Liechtenstein, Moldova, Monaco, Montenegro, North Macedonia, Norway, Russia (only the following parts: Central Federal District (Tsentralny federalny okrug), Northwestern Federal District (Severo-Zapadny federalny okrug), Southern Federal District (Yuzhny federalny okrug), North Caucasian Federal District (Severo-Kavkazsky federalny okrug) and Volga Federal District (Privolzhsky federalny okrug)), San Marino, Serbia, Switzerland, Turkey and Ukraine
9.	Plants for planting of <i>Cydonia</i> Mill., <i>Malus</i> Mill., <i>Prunus</i> L. and <i>Pyrus</i> L. and their hybrids, and <i>Fragaria</i> L., other than seeds	ex 0602 10 90 ex 0602 20 20 ex 0602 90 30 ex 0602 90 41 ex 0602 90 45 ex 0602 90 46 ex 0602 90 48 ex 0602 90 50 ex 0602 90 70 ex 0602 90 91 ex 0602 90 99	Third countries, other than: Albania, Algeria, Andorra, Armenia, Australia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canada, Canary Islands, Egypt, Faeroe Islands, Georgia, Iceland, Israel, Jordan, Lebanon, Libya, Liechtenstein, Moldova, Monaco, Montenegro, Morocco, New Zealand, North Macedonia, Norway, Russia (only the following parts: Central Federal District (Tsentralny federalny okrug), Northwestern Federal District (Severo-Zapadny federalny okrug), Southern Federal District (Yuzhny federalny okrug), North Caucasian Federal District (Severo-Kavkazsky federalny okrug) and Volga Federal District (Privolzhsky federalny okrug)), San Marino, Serbia, Switzerland, Syria, Tunisia, Turkey, Ukraine, and United States other than Hawaii
10.	Plants of <i>Vitis</i> L., other than fruits	0602 10 10 0602 20 10 ex 0604 20 90 ex 1404 90 00	Third countries other than Switzerland
11.	Plants of <i>Citrus</i> L., <i>Fortunella</i> Swingle, <i>Poncirus</i> Raf., and their hybrids, other than fruits and seed	ex 0602 10 90 ex 0602 20 20 0602 20 30 ex 0602 20 80 ex 0602 90 45 ex 0602 90 46 ex 0602 90 47 ex 0602 90 50 ex 0602 90 70 ex 0602 90 91 ex 0602 90 99 ex 0604 20 90 ex 1404 90 00	
18.	Plants for planting of Solanaceae other than seeds and the plants covered by entries 15, 16 or 17	ex 0602 90 30 ex 0602 90 45 ex 0602 90 46 ex 0602 90 48 ex 0602 90 50 ex 0602 90 70 ex 0602 90 91 ex 0602 90 99	Third countries other than: Albania, Algeria, Andorra, Armenia, Azerbaijan, Belarus, Bosnia and Herzegovina, Canary Islands, Egypt, Faeroe Islands, Georgia, Iceland, Israel, Jordan, Lebanon, Libya, Liechtenstein, Moldova, Monaco, Montenegro, Morocco, North Macedonia, Norway, Russia (only the following parts: Central Federal District (Tsentralny federalny okrug), Northwestern Federal District (Severo-Zapadny federalny okrug), Southern Federal

**List of plants, plant products and other objects whose introduction into the Union from certain third countries is prohibited**

	Description	CN Code	Third country, group of third countries or specific area of third country
			District (Yuzhny federalny okrug), North Caucasian Federal District (Severo-Kavkazsky federalny okrug) and Volga Federal District (Privolzhsky federalny okrug)), San Marino, Serbia, Switzerland, Syria, Tunisia, Turkey and Ukraine
20.	Growing medium as such, other than soil, consisting in whole or in part of solid organic substances, other than that composed entirely of peat or fibre of <i>Cocos nucifera</i> L., previously not used for growing of plants or for any agricultural purposes	ex 2530 10 00 ex 2530 90 00 ex 2703 00 00 ex 3101 00 00 ex 3824 99 93	Third countries other than Switzerland

### 3.4. Entry, establishment and spread in the EU

#### 3.4.1. Entry

*Is the pest able to enter into the EU territory? If yes, identify and list the pathways.*

*Comment on plants for planting as a pathway.*

**Yes.** The pest has already entered the EU territory. The main pathways are plants for planting, fruits, vegetables and cut flowers.

Plants for planting, fruits, vegetables and cut flowers are the main pathways for entry of *M. hirsutus* (EPPO, 2005; Culik et al., 2013). It can also be associated with soil, which could however be considered as a closed pathway (Table 4).

**Table 4:** Potential pathways for *Maconellicoccus hirsutus* into the EU 27

Pathways description (e.g. host/intended use/source)	Life stage	Relevant mitigations [e.g. prohibitions (Annex VI), special requirements (Annex VII) or phytosanitary certificates (Annex XI) within Implementing Regulation 2019/2072]
Plants for planting	Eggs, nymphs and adults	Plants for planting that are hosts of <i>M. hirsutus</i> , and are prohibited to import from third countries (Regulation 2019/2072, Annex VI), are listed in Table 3.  The growing medium attached to or associated with plants, intended to sustain the vitality of the plants, are regulated in Regulation 2019/2072, Annex VII.  Plants for planting from third countries require a phytosanitary certificate (Regulation 2019/2072, Annex XI, Part A)
Fruits, vegetables and cut flowers	Eggs, nymphs and adults	Fruits, vegetables and cut flowers from third countries require a phytosanitary certificate to import into the EU (2019/2072, Annex XI, Part A). However, no requirements are specified for <i>M. hirsutus</i> .  According to Regulation 2019/2072, Annex XI, Part C there is a list of plants which a phytosanitary certificate is not required for their introduction into the Union territory.



Pathways description (e.g. host/intended use/source)	Life stage	Relevant mitigations [e.g. prohibitions (Annex VI), special requirements (Annex VII) or phytosanitary certificates (Annex XI) within Implementing Regulation 2019/2072]
		<i>M. hirsutus</i> infests fruits that are included in that list ( <i>Ananas comosus</i> and <i>Musa</i> spp.)
Soil	Eggs	Import of soil from third countries is prohibited (Regulation 2019/2072, Annex VI)

The import of some host plants of *M. hirsutus* for planting from third countries is not allowed (Regulation 2019/2072, Annex VI), while there are many other hosts that can be imported to the EU with a phytosanitary certificate.

Vegetables, cut flowers and most fruits that are imported into the EU must have a phytosanitary certificate. However, pineapple (*Ananas comosus*) and banana (*Musa* spp.), which are hosts for *M. hirsutus*, are exempt by Regulation 2019/2072, Annex XI, Part C.

EU legislation (2019/2072) prohibits the import of soil from third countries so that pathway can be considered as closed.

Annual imports of *M. hirsutus* hosts from countries where the pest is known to occur are provided in Appendix C.

Notifications of interceptions of harmful organisms began to be compiled in Europhyt in May 1994 and in TRACES in May 2020. As at 16/9/2021 (search date) there were two records of interceptions of *M. hirsutus* in the Europhyt and TRACES databases:

- in 2008 on *Colocasia* sp. plants for planting imported from India
- in 2018 on *Annona squamosa* fruits imported from Brazil

In the UK, a former member of the EU, there were more than 240 interceptions of *M. hirsutus* between 1994 and 2021, mostly on *Annona squamosa* fruits from India. *M. hirsutus* was also found on *Annona* fruits from Egypt, Indonesia, Kenya, Pakistan, Saint Lucia and Vietnam, and a range of fresh fruits and vegetables imported from Asia, Africa, and the Caribbean (Fera unpublished records). No action was taken against these findings.

### 3.4.2. Establishment

*Is the pest able to become established in the EU territory?*

**Yes**, in the EU countries of southern Europe the climate is suitable and there are many available hosts that can support establishment. Given that *M. hirsutus* occurs in Greece and has a wide distribution in Cyprus, it must have been able to transfer following entry.

#### 3.4.2.1. EU distribution of main host plants

*M. hirsutus* is a polyphagous pest. The main hosts of the pest cultivated in the EU 27 between 2016 and 2020 are shown in Table 5. Among others, citrus, cotton, soybeans, grapes, pome fruits and stone fruits are highly economically important crops in the EU.


**Table 5:** Crop area of *Maconellicoccus hirsutus* hosts in EU 27 in 1,000 ha (Eurostat accessed on 21/09/2021)

Crop	2016	2017	2018	2019	2020
Citrus	519.01	502.84	508.99	512.53	487.08
Cotton	301.34	326.12	345.64	361.78	349.94
Soybeans	831.18	962.39	955.40	907.91	939.86
Grapes	3,136.04	3,133.21	3,135.02	3,158.32	3,160.27
Cucumbers	32.33	31.81	32.65	33.69	33.15
Bananas	20.30	18.91	17.94	18.19	19.61
Pome fruits	No data	627.98	629.42	610.11	589.85

Crop	2016	2017	2018	2019	2020
Stone fruits	No data	625.46	621.32	612.33	No data
Avocados	12.24	12.72	13.22	15.52	17.27

### 3.4.2.2. Climatic conditions affecting establishment

*M. hirsutus* occurs mainly in tropical and subtropical regions in Asia, Africa, Australia, and America. Moreover, it has also been recorded in Greece, Cyprus and Turkey, countries with a Mediterranean climate. According to the global Köppen-Geiger climate zones (Kottek et al., 2006), *M. hirsutus* is present in countries with climate zones Aw (Equatorial savannah with dry winter), Am (Equatorial monsoon), Af (Equatorial rainforest, fully humid), BWh (Desert climate, hot desert), Bsh (Steppe climate, hot steppe) and Csa (warm temperate climate with dry hot summer). The lower and upper developmental temperature threshold of *M. hirsutus* on *H. rosa-sinensis* is 14.5 and 35°C, respectively (Chong et al., 2008), temperatures that are relatively high. Figure 3 shows the World distribution of Köppen–Geiger climate types that occur in the EU and which occur in countries where *M. hirsutus* has been reported. Southern EU countries provide suitable climatic conditions that would support the establishment of *M. hirsutus*. There is uncertainty as to whether *M. hirsutus* could establish in the EU countries of central Europe. It is unlikely that the insect could establish in the northern EU, and if it did, the populations are likely to be small and have no impact. Countries and areas of the EU most suitable include Cyprus, Greece, Malta, Portugal, Spain, coastal areas of southern France, including Corsica, as well as southern Italy, including Sardinia and Sicily. There is a possibility that *M. hirsutus* could occur in glasshouses and on indoor plantings in cooler areas.



**Figure 3:** World distribution of Köppen–Geiger climate types that occur in the EU and which occur in areas where *Maconellicoccus hirsutus* has been reported

### 3.4.3. Spread

*Describe how the pest would be able to spread within the EU territory following establishment?*

First instar nymphs are spread by crawling, wind, rainfall and on humans and animals. Overwintering eggs may be moved in soil. All stages may be moved over long distances in trade.

*Comment on plants for planting as a mechanism of spread.*

Plants for planting are one of the main pathways of spread of the pest over long distances.

First instar nymphs are active and spread by crawling, wind and rainfall. The sticky egg masses and mobile crawlers may also be carried to new areas on humans and other animals (Sagarra and Peterkin, 1999; EPPO, 2005; Culik et al., 2013). Moreover, overwintering eggs can be found in soil (Pollard, 1995) and spread through the soil attached to plants for planting and machinery.

The introduction of this pest to new territories over long distance is possible through the movement of infested plants for planting (e.g. fruit tree and ornamental nursery seedlings), and trade of infested fruit, vegetables, cut flowers or other plant products (Meyerdirk et al., 2001; CABI, 2021).

Plants for planting, fruits, vegetables and cut flowers are the main pathways of spread of the pest over long distances.

### 3.5. Impacts

*Would the pests' introduction have an economic or environmental impact on the EU territory?*

**Yes**, if *M. hirsutus* established more widely in the EU, it would most probably have an economic impact on the host species of the pest.

The pest may seriously affect the commercial value of various ornamental plants and potentially have a high economic impact on crop production in the EU. *M. hirsutus* egests large quantities of honeydew, and as a result black sooty mould develops on the plants, which reduces the aesthetic value, normal growth and reproduction (Kairo et al., 2000; Chong et al., 2015). *M. hirsutus* also injects toxic saliva into the plant during feeding, which results in leaf curling, fruit malformation, bunchy top appearance (Figure 1E) and premature senescence of flowers and foliage (Dufour and Léon, 1997; Chong et al., 2015). Heavy infestations may cause a complete defoliation of the plant, leading to its death (Figure 1F) (Dufour and Léon, 1997; Sagarra and Peterkin, 1999). These impacts have been documented in city parks and gardens in Cyprus (Ülgentürk et al., 2015) and Greece (Milonas and Partsinevelos, 2017).

The potential annual economic impact of *M. hirsutus* to avocado (*Persea americana*), citrus (*Citrus* spp.), cotton (*Gossypium hirsutum*), peanut (*Arachis hypogaea*), soybean (*Glycine max*), nursery and vegetable crops was estimated at US\$163 million in Florida or US\$1.6 billion for the entire United States (Ranjan, 2006). In Egypt, *M. hirsutus* was reported to cause damage to *Albizia lebbek*, mulberry, *Hibiscus* spp., and cotton. In Africa, it was considered as a possible pest of cocoa. In India, Bangladesh and Pakistan it is a pest of cotton, mulberry and several fibre crop species. In India, it has also been considered to be a severe pest of grapes (Muralidharan and Badaya, 2000; Culik et al., 2013). When *M. hirsutus* was introduced in the Caribbean islands it became a very serious problem. Grenada reported economic losses of \$3.5 to \$10 million for the season 1996–1997 and Trinidad and Tobago estimated potential losses exceeding \$125 million/year, if infestations continued to escalate (Meyerdirk et al., 2001). However, in many countries *M. hirsutus* is restricted to *Hibiscus* species and is not a serious pest, possibly because natural enemies effectively reduce its populations (Meyerdirk et al., 2001).

### 3.6. Available measures and their limitations

*Are there measures available to prevent pest entry, establishment, spread or impacts such that the risk becomes mitigated?*

**Yes**. Although the existing phytosanitary measures identified in Section 3.3.2 do not specifically target *M. hirsutus*, they mitigate the likelihood of its entry into and spread within the EU (see also Section 3.6.1).

#### 3.6.1. Identification of potential additional measures

Phytosanitary measures (prohibitions) are currently applied to some host plants for planting (see Section 3.3.2).

Additional potential risk reduction options and supporting measures are shown in Sections 3.6.1.1 and 3.6.1.2.

##### 3.6.1.1. Additional potential risk reduction options

Potential additional control measures are listed in Table 6.

**Table 6:** Selected control measures (a full list is available in EFSA PLH Panel, 2018) for pest entry/establishment/spread/impact in relation to currently unregulated hosts and pathways. Control measures are measures that have a direct effect on pest abundance

<b>Control measure/risk reduction option</b> <b>(Blue underline = Zenodo doc)</b>	<b>RRO summary</b>	<b>Risk element targeted (entry/establishment/spread/impact)</b>
<a href="#"><u>Growing plants in isolation</u></a>	Description of possible exclusion conditions that could be implemented to isolate the crop from pests and if applicable relevant vectors. E.g. a dedicated structure such as glass or plastic greenhouses. Used to mitigate likelihood of infestation by specified pest in vicinity of growing site. Plants could be grown in glass or plastic structures	Entry (reduce contamination/infestation)/spread
<a href="#"><u>Chemical treatments on crops including reproductive material</u></a>	Used to mitigate likelihood of infestation of pests susceptible to chemical treatments. Pesticide application for the control of <i>M. hirsutus</i> has been considered to be impractical (Culik et al., 2013). Some neonicotinoid and pyrethroid insecticides (e.g. imidacloprid, thiamethoxam, bifenthrin) as well as their mixture have provided encouraging results regarding the control of the pest (Castle and Prabhaker, 2011; Fatima et al., 2016). However, the use of some neonicotinoids for outdoor use in EU has been banned. Moreover, the natural wax coating covering the various stages of the insect protects it from pesticides (Meyerdirk et al., 2001)	Entry/establishment/impact
<a href="#"><u>Chemical treatments on consignments or during processing</u></a>	Use of chemical compounds that may be applied to plants or to plant products after harvest, during process or packaging operations and storage. The treatments addressed in this information sheet are: a) fumigation; b) spraying/dipping pesticides; c) surface disinfectants; d) process additives; e) protective compounds  Used to mitigate likelihood of infestation of pests susceptible to chemical treatments. Eggs, nymphs and adults of <i>M. hirsutus</i> were susceptible to methyl bromide fumigations. A dose of 48 mg/litre methyl bromide at 21–26°C produced 100% mortality of all life stages (Zettler et al., 2002)	Entry/spread
<a href="#"><u>Physical treatments on consignments or during processing</u></a>	This information sheet deals with the following categories of physical treatments: irradiation/ionisation; mechanical cleaning (brushing, washing); sorting and grading, and; removal of plant parts (e.g. debarking wood). This information sheet does not address: heat and cold treatment (information sheet 1.14); roguing and pruning (information sheet 1.12). Used to mitigate likelihood of infestation of pests susceptible to physical treatments Washing, brushing and other mechanical cleaning methods can be used to reduce the prevalence of the pest in the consignments to be exported or to be planted	Entry/spread
<a href="#"><u>Cleaning and disinfection of facilities, tools and machinery</u></a>	The physical and chemical cleaning and disinfection of facilities, tools, machinery, transport means, facilities and other accessories (e.g. boxes, pots, pallets, palox, supports, hand tools). The measures addressed in this information sheet are: washing, sweeping and fumigation.	Entry/spread

<b>Control measure/risk reduction option</b> <b>(Blue underline = Zenodo doc)</b>	<b>RRO summary</b>	<b>Risk element targeted (entry/ establishment/ spread/ impact)</b>
	Used to mitigate likelihood of entry or spread of soil borne pests	
Limits on soil	Used to mitigate likelihood of entry or spread of <i>M. hirsutus</i> eggs in soil	Entry/spread
<b><u>Soil treatment</u></b>	The control of soil organisms by chemical and physical methods listed below: a) Fumigation; b) Heating; c) Solarisation; d) Flooding; e) Soil suppression; f) Augmentative Biological control; g) Biofumigation Used to mitigate likelihood of presence of eggs in the soil	Entry/establishment/ impact
<b><u>Heat and cold treatments</u></b>	Controlled temperature treatments aimed to kill or inactivate pests without causing any unacceptable prejudice to the treated material itself. The measures addressed in this information sheet are: autoclaving; steam; hot water; hot air; cold treatment Used to mitigate likelihood of infestation of pests susceptible to physical treatments. Hot water immersion treatment of fruits has been reported as an effective measure for disinfestation of fresh fruits. Effective temperature time combinations for control of <i>M. hirsutus</i> on fruits were 55 min at 47°C, 23 min at 48°C and 13 min at 49°C (Hara and Jacobsen, 2005)	Entry/spread
<b><u>Controlled atmosphere</u></b>	Treatment of plants by storage in a modified atmosphere (including modified humidity, O <sub>2</sub> , CO <sub>2</sub> , temperature, pressure). Used to mitigate likelihood of infestation of pests susceptible to modified atmosphere (usually applied during transport) hence to mitigate entry. Controlled atmosphere storage can be used in commodities such as fresh and dried fruits, flowers and vegetables	Entry/spread (via commodity)
Post-entry quarantine and other restrictions of movement in the importing country	This information sheet covers post-entry quarantine (PEQ) of relevant commodities; temporal, spatial and end-use restrictions in the importing country for import of relevant commodities; Prohibition of import of relevant commodities into the domestic country. 'Relevant commodities' are plants, plant parts and other materials that may carry pests, either as infection, infestation, or contamination. Plants in PEQ are held in conditions that prevent the escape of pests; they can be carefully inspected and tested to verify they are of sufficient plant health status to be released, or may be treated, re-exported or destroyed. Tests on plants are likely to include laboratory diagnostic assays and bioassays on indicator hosts to check whether the plant material is infected with particular pathogens	Establishment/spread

### 3.6.1.2. Additional supporting measures

Potential additional supporting measures are listed in Table 7.



**Table 7:** Selected supporting measures (a full list is available in EFSA PLH Panel, 2018) in relation to currently unregulated hosts and pathways. Supporting measures are organisational measures or procedures supporting the choice of appropriate risk reduction options that do not directly affect pest abundance

Supporting measure	RRO summary	Risk element targeted (entry/establishment/spread/impact)
<a href="#">Inspection and trapping</a>	<p>Inspection is defined as the official visual examination of plants, plant products or other regulated articles to determine if pests are present or to determine compliance with phytosanitary regulations (ISPM 5). The effectiveness of sampling and subsequent inspection to detect pests may be enhanced by including trapping and luring techniques.</p> <p>Used to mitigate likelihood of infestation by specified pest at origin. Any shipments of fresh plant material from an infested country to another that is not infested should be examined thoroughly to detect <i>M. hirsutus</i> (CABI, 2021)</p>	Establishment/spread
Phytosanitary certificate and plant passport	<p>An official paper document or its official electronic equivalent, consistent with the model certificates of the IPPC, attesting that a consignment meets phytosanitary import requirements (ISPM 5)</p> <p>a) export certificate (import) b) plant passport (EU internal trade)</p> <p>Used to attest which of the above requirements have been applied</p>	Entry/spread

### 3.6.1.3. Biological or technical factors limiting the effectiveness of measures

- *M. hirsutus* hide in cracks and crevices on the plant bark and in the calyx of *fruits*, making its detection, especially in early infestations and low population, difficult.
- The high number of host plants and the wide distribution of *M. hirsutus* makes the inspections of all consignments imported from countries where the pest occurs difficult.
- The natural wax coating covering the various stages of *M. hirsutus* protects it from treatments with contact insecticides.

## 3.7. Uncertainty

- Uncertainty exists regarding the suitability of the climate of EU countries in central Europe for the establishment of *M. hirsutus*. However, its establishment in the southern EU countries is very likely since it has already been detected in Cyprus and Greece (Rhodes).
- In many countries where climate is suitable, *M. hirsutus* is not a serious pest, largely due to natural enemies (Kairo et al., 2000), thus there is uncertainty on the magnitude of impact. For example, it is not known if, and how quickly, natural enemies such as the parasitoid *Anagyrus kamali*, will follow the spread of *M. hirsutus* in the EU.
- The presence of *M. hirsutus* in France and Italy, implied by some interceptions in the USA, is uncertain (Miller et al., 2014). It is likely that the interceptions recorded in the US are on produce imported into the EU from other countries and reexported (see Section 3.2.2).

## 4. Conclusions

The criteria assessed by EFSA for consideration of *M. hirsutus* as a potential EU quarantine pest are met (Table 8).

**Table 8:** The Panel's conclusions on the pest categorisation criteria defined in Regulation (EU) 2016/2031 on protective measures against pests of plants (the number of the relevant sections of the pest categorisation is shown in brackets in the first column)

Criterion of pest categorisation	Panel's conclusions against criterion in Regulation (EU) 2016/2031 regarding Union quarantine pest	Key uncertainties
<b>Identity of the pest (3.1)</b>	The identity of the pest is established. Taxonomic keys based on morphology of female adults exist	None
<b>Absence/presence of the pest in the EU (3.2)</b>	The pest has a restricted distribution in the EU territory (Rhodes Island in Greece and Cyprus)	None
<b>Regulatory status (3.3)</b>	<i>Maconellicoccus hirsutus</i> is not regulated as a quarantine pest in the EU; the Cypriot NPPO is taking official action	None
<b>Pest potential for entry, establishment and spread in the EU (3.4)</b>	<i>Maconellicoccus hirsutus</i> is able to enter into, become established, and spread within the EU territory. The main pathways are: <ul style="list-style-type: none"> <li>– plants for planting (regulated, some prohibited, some permitted)</li> <li>– fruits, vegetables and cut flowers (regulated, except bananas and pineapple)</li> </ul>	None
<b>Potential for consequences in the EU (3.5)</b>	The pests' introduction could reduce the aesthetic value of various ornamental plants and the production of many crops	In many countries <i>M. hirsutus</i> is not a serious pest, possibly due to the climate being less favourable, and natural enemies reducing its population levels
<b>Available measures (3.6)</b>	There are measures available to prevent the entry, establishment and spread of <i>M. hirsutus</i> within the EU. Risk reduction options include the inspections and physical treatments on consignments of fresh plant material from infested countries and the production of plants for import into the EU in pest free areas (this could be difficult due to wide distribution of the pest)	Eradication and containment actions taken in the Caribbean (for example, restricting the movement of host plant material) were unsuccessful. There is uncertainty regarding how effective risk reduction measures would be in the EU
<b>Conclusion (4)</b>	The criteria assessed by EFSA for consideration as a potential quarantine pest are met	
<b>Aspects of assessment to focus on/scenarios to address in future if appropriate</b>	Establishment, impact, and natural enemies	

## References

- Abd-Rabou S, Shalaby H, Germain JF, Ris N, Kreiter P and Malausa T, 2012. Identification of mealybug pest species (Hemiptera: Pseudococcidae) in Egypt and France, using a DNA barcoding approach. *Bulletin of Entomological Research*, 24, 1–9.
- Al-Fwaeer M, Abu-Obeid I, Al-Zyoud F, Abo-Alosh A and Halaybeh M, 2014. Population Dynamics of the hibiscus mealybug *Maconellicoccus hirsutus* Green (Hom., Pseudococcidae) and its parasitoid on guava trees in Madaba-Jordan. *International Journal of Agriculture and Forestry*, 4, 171–177.
- Alleyne JC, 2004. Controlling a dangerous pest-pink hibiscus mealybug. *Newsletter of the University of Florida - IFAS Extension - Pinellas County Extension*, 38, 2–3.
- Aristizábal LF, Mannion C, Bergh C and Arthurs S, 2012. Life history of pink hibiscus mealybug, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) on three *Hibiscus rosa-sinensis* cultivars. *Florida Entomologist*, 95, 89–94.

- Bartlett BR, 1978. Pseudococcidae. In: Clausen CP (ed.). Introduced parasites and predators of arthropod pests and weeds: a world review. Agriculture Handbook no. 480. USDA, Washington, US. pp. 137–170.
- CABI (Centre for Agriculture and Biosciences International), 2021. Available online: [www.cabi.org](http://www.cabi.org) [Accessed: 15 September 2021]
- Castle SJ and Prabhaker N, 2011. Field evaluation of two systemic neonicotinoid insecticides against pink hibiscus mealybug (*Maconellicoccus hirsutus* (Green)) on mulberry trees. *Journal of Pest Science*, 84, 363–371.
- Chong JH, Roda AL and Mannion CM, 2008. Life history of the mealybug, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), at constant temperatures. *Environmental Entomology*, 37, 323–332.
- Chong JH, Aristizabal LF and Arthurs SP, 2015. Biology and management of *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) on ornamental plants. *Journal of Integrated Pest Management*, 6, 5.
- Culik MP, Fornazier MJ, dos Santos Martins D, Zanuncio JS, Ventura JA, Peronti ALBG and Zanuncio JC, 2013. The invasive mealybug *Maconellicoccus hirsutus*: lessons for its current range expansion in South America and invasive pest management in general. *Journal of Pest Science*, 86, 387–398.
- Dufour BP and Léon J, 1997. Informe de mission de informacion sobre el control de la cochinilla rosada del hibisco (*Maconellicoccus hirsutus* Green) en la region del Caribe. IICA - El Salvador. 22 pp.
- EFSA PLH Panel (EFSA Panel on Plant Health), Jeger M, Bragard C, Caffier D, Candresse T, Chatzivassiliou E, Dehnen-Schmutz K, Gregoire J-C, Jaques Miret JA, MacLeod A, Navajas Navarro M, Niere B, Parnell S, Potting R, Rafoss T, Rossi V, Urek G, Van Bruggen A, Van Der Werf W, West J, Winter S, Hart A, Schans J, Schrader G, Suffert M, Kertesz V, Kozelska S, Mannino MR, Mosbach-Schulz O, Pautasso M, Stancanelli G, Tramontini S, Vos S and Gilioli G, 2018. Guidance on quantitative pest risk assessment. *EFSA Journal* 2018;16(8):5350, 86 pp. Available online: <https://doi.org/10.2903/j.efsa.2018.5350>
- EFSA PLH Panel (EFSA Panel on Plant Health), Bragard C, Dehnen-Schmutz K, Di Serio F, Jacques M-A, Jaques Miret JA, Justesen AF, MacLeod A, Magnusson CS, Milonas P, Navas-Cortes JA, Parnell S, Potting R, Reignault PL, Thulke H-H, van der Werf W, Civera AV, Yuen J, Zappalá L, Battisti A, Mas H, Rigling D, Mosbach-Schulz O, and Gonthier P, 2021. Scientific Opinion on the commodity risk assessment of *Ficus carica* plants from Israel. *EFSA Journal* 2021;19(1):6353, 249 pp. Available online: <https://doi.org/10.2903/j.efsa.2021.6353>
- EFSA Scientific Committee, Hardy A, Benford D, Halldorsson T, Jeger MJ, Knutsen HK, More S, Naegeli H, Noteborn H, Ockleford C, Ricci A, Rycken G, Schlatter JR, Silano V, Solecki R, Turck D, Benfenati E, Chaudhry QM, Craig P, Frampton G, Greiner M, Hart A, Hogstrand C, Lambre C, Luttik R, Makowski D, Siani A, Wahlstroem H, Aguilera J, Dorne J-L, Fernandez Dumont A, Hempten M, Valtueña Martínez S, Martino L, Smeraldi C, Terron A, Georgiadis N and Younes M, 2017. Scientific Opinion on the guidance on the use of the weight of evidence approach in scientific assessments. *EFSA Journal* 2017;15(8):4971, 69 pp. <https://doi.org/10.2903/j.efsa.2017.4971>
- EPPO (European and Mediterranean Plant Protection Organization), 2005. OEPP/EPPO, data sheets on quarantine pests *Maconellicoccus hirsutus*. *Bulletin OEPP/EPPO Bulletin*, 35, 413–415.
- EPPO (European and Mediterranean Plant Protection Organization), 2006. OEPP/EPPO, diagnostics *Maconellicoccus hirsutus*. *Bulletin OEPP/EPPO Bulletin*, 36, 167–169.
- EPPO (European and Mediterranean Plant Protection Organization), 2019. EPPO codes. Available online: [https://www.eppo.int/RESOURCES/eppo\\_databases/eppo\\_codes](https://www.eppo.int/RESOURCES/eppo_databases/eppo_codes)
- EPPO (European and Mediterranean Plant Protection Organization), 2021. EPPO Global Database. Available online: <https://gd.eppo.int> [Accessed: 20 September 2021]
- EPPO (European and Mediterranean Plant Protection Organization), online. EPPO Global Database. Available online: <https://gd.eppo.int> [Accessed: 15/10/2021]
- FAO (Food and Agriculture Organization of the United Nations), 2013. ISPM (International Standards for Phytosanitary Measures) 11—Pest risk analysis for quarantine pests. FAO, Rome. 36 pp. Available online: [https://www.ippc.int/sites/default/files/documents/20140512/isp11\\_2013\\_en\\_2014-04-30\\_201405121523-494.65%20KB.pdf](https://www.ippc.int/sites/default/files/documents/20140512/isp11_2013_en_2014-04-30_201405121523-494.65%20KB.pdf)
- FAO (Food and Agriculture Organization of the United Nations), 2018. International Standards for Phytosanitary Measures. ISPM 5 Glossary of phytosanitary terms. Revised version adopted CPM 13, April 2018. FAO, Rome. Available online: <https://www.ippc.int/en/publications/621/>
- Fatima S, Hussain M, Shafiqat S, Faheem Malik M, Abbas Z, Noureen N and Ane N, 2016. Laboratory evaluation of different insecticides against hibiscus mealybug, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae). *Scientifica*, 2016, 9312013.
- Francis A, Bloem KA, Roda AL, Lapointe SL, Zhang A and Onokpise O, 2007. Development of trapping methods with a synthetic sex pheromone of the pink hibiscus mealybug, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae). *Florida Entomologist*, 90, 440–446.
- García Morales G, Denno BD, Miller DR, Miller GL, Ben-Dov Y and Hardy NB, 2016. ScaleNet: a literature-based model of scale insect biology and systematics. Database. <https://doi.org/10.1093/database/bav118>. Available online: <http://scalenet.info> [Accessed: 20 September 2021].
- Ghose SK, 1971. Morphology of various instars of both sexes of the mealy-bug, *Maconellicoccus hirsutus*. *Indian Journal of Agricultural Sciences*, 41, 602–611.
- Ghose SK, 1972. Biology of the mealybug, *Maconellicoccus hirsutus* (Green) (Pseudococcidae, Hemiptera). *Indian Agriculturalist*, 16, 323–332.

- Griessinger D and Roy A-S, 2015. EPPO codes: a brief description. Available online: [https://www.eppo.int/media/uploaded\\_images/RESOURCES/eppo\\_databases/A4\\_EPPO\\_Codes\\_2018.pdf](https://www.eppo.int/media/uploaded_images/RESOURCES/eppo_databases/A4_EPPO_Codes_2018.pdf)
- Hall WJ, 1921. The hibiscus mealy bug (*Phenacoccus hirsutus*). Bulletin Ministry of Agriculture Egypt Technical and Scientific Service Entomological Section, 17, 1–28.
- Hara AH and Jacobsen CM, 2005. Hot water immersion for surface disinfestations of *Maconellicoccus hirsutus* (Homoptera: Pseudococcidae). Journal of Economic Entomology, 98, 284–288.
- Kairo MTK, Pollard GV, Peterkin DD and Lopez VF, 2000. Biological control of the hibiscus mealybug, *Maconellicoccus hirsutus* Green (Hemiptera: Pseudococcidae) in the Caribbean. Integrated Pest Management Reviews, 5, 241–254.
- Kottek M, Grieser J, Beck C, Rudolf B and Rubel F, 2006. World map of Köppen-Geiger climate classification updated. Meteorologische Zeitschrift, 15, 259–263.
- Malausca T, Fenis A, Warot S, Germain JF, Ris N, Prado E, Botton M, Vanlerberghe-Masutti F, Sforza R, Cruaud C, Couloux A and Kreiter P, 2011. DNA markers to disentangle complexes of cryptic taxa in mealybugs (Hemiptera: Pseudococcidae). Journal of Applied Entomology, 135, 142–155.
- Mani M, 1989. A review of the pink mealybug – *Maconellicoccus hirsutus*. Insect Science and Its Application, 10, 157–167.
- Meyerdirk DE, Warkentin R, Attavian B, Gersabeck E, Francis A, Adams J and Francis G, 2001. Biological control of pink hibiscus mealybug project manual, USDA, Washington.
- Miller D, Rung A, Parikh G, Venable G, Redford AJ, Evans GA and Gill RJ, 2014. Scale insects. Edition 2. USDA APHIS Identification Technology Program (ITP), Fort Collins, CO. Available online: <http://idtools.org/id/scales/> [Accessed 1 October 2021]
- Milonas PG and Partsinevelos GK, 2017. The pink hibiscus mealybug *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae) in Greece. Hellenic Plant Protection Journal, 10, 80–83.
- Muralidharan CM and Badaya SN, 2000. Mealy bug (*Maconellicoccus hirsutus*) (Pseudococcidae: Hemiptera) outbreak on herbaceous cotton (*Gossypium herbaceum*) in Wagad cotton belt of Kachchh. Indian Journal of Agricultural Sciences, 70, 405–706.
- Negrini M, de Moraes EGF and Zanuncio JC, 2017. Biology of *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) on *Hibiscus rosa-sinensis*. Revista Agro@ambiente On-line, 11, 336–346.
- Pollard GV, 1995. Pink or hibiscus mealybug in the Caribbean. CARAPHIN News, 12, 1–2.
- Ranjan R, 2006. Economic impacts of pink hibiscus mealybug in Florida and the United States. Stochastic Environmental Research and Risk Assessment, 20, 353–362.
- Sagarra LA and Peterkin DD, 1999. Invasion of the Caribbean by the hibiscus mealybug, *Maconellicoccus hirsutus* Green (Homoptera: Pseudococcidae). Phytoprotection, 80, 103–113.
- Sahito HA, Soomro RB, Tapur MA, Memon SA and Dhiloo KH, 2012. Biology of mulberry mealybug, *Maconellicoccus hirsutus* (Green) in laboratory conditions. Basic Research Journal of Agricultural Science and Review, 1, 11–18.
- Singh MP and Ghosh SN, 1970. Studies on *Maconellicoccus hirsutus* causing 'bunchy top' in mesta. Indian Journal of Science and Industry, A4, 99–105.
- Ülgentürk S, Kaydan MB and Hocaali Şişman S, 2015. New scale insect (Hemiptera: Coccoomorpha) records for the Turkish Republic of Northern Cyprus. Türkiye Entomoloji Bülteni, 5, 59–68.
- Williams DJ, 1996. A brief account of the hibiscus mealybug *Maconellicoccus hirsutus*, a pest of agriculture and horticulture, with descriptions of two related species from southern Asia. Bulletin of Entomological Research, 86, 617–628.
- Williams DJ, 2004. Mealybugs of Southern Asia. The Natural History Museum, Kuala Lumpur: Southdene SDN. BHD. 896 pp.
- Zettler JL, Follett PA and Gill RF, 2002. Susceptibility of *Maconellicoccus hirsutus* (Homoptera: Pseudococcidae) to Methyl Bromide. Journal of Economic Entomology, 95, 1169–1173.

## Abbreviations

EPPO	European and Mediterranean Plant Protection Organization
FAO	Food and Agriculture Organization
IPPC	International Plant Protection Convention
ISPM	International Standards for Phytosanitary Measures
MS	Member State
PLH	EFSA Panel on Plant Health
PZ	Protected Zone
TFEU	Treaty on the Functioning of the European Union
ToR	Terms of Reference

## Glossary

Containment (of a pest)	Application of phytosanitary measures in and around an infested area to prevent spread of a pest (FAO, 2018)
Control (of a pest)	Suppression, containment or eradication of a pest population (FAO, 2018)
Entry (of a pest)	Movement of a pest into an area where it is not yet present, or present but not widely distributed and being officially controlled (FAO, 2018)
Eradication (of a pest)	Application of phytosanitary measures to eliminate a pest from an area (FAO, 2018)
Establishment (of a pest)	Perpetuation, for the foreseeable future, of a pest within an area after entry (FAO, 2018)
Greenhouse	A walk-in, static, closed place of crop production with a usually translucent outer shell, which allows controlled exchange of material and energy with the surroundings and prevents release of plant protection products (PPPs) into the environment.
Impact (of a pest)	The impact of the pest on the crop output and quality and on the environment in the occupied spatial units
Introduction (of a pest)	The entry of a pest resulting in its establishment (FAO, 2018)
Pathway	Any means that allows the entry or spread of a pest (FAO, 2018)
Phytosanitary measures	Any legislation, regulation or official procedure having the purpose to prevent the introduction or spread of quarantine pests, or to limit the economic impact of regulated non-quarantine pests (FAO, 2018)
Quarantine pest	A pest of potential economic importance to the area endangered thereby and not yet present there, or present but not widely distributed and being officially controlled (FAO, 2018)
Risk reduction option (RRO)	A measure acting on pest introduction and/or pest spread and/or the magnitude of the biological impact of the pest should the pest be present. A RRO may become a phytosanitary measure, action or procedure according to the decision of the risk manager
Spread (of a pest)	Expansion of the geographical distribution of a pest within an area (FAO, 2018)



## Appendix A – *Maconellicoccus hirsutus* host plants/species affected

Source: EPPO Global Database (EPPO, online), García Morales et al. (2016) and other references.

Host status	Host name	Plant family	Common name	Reference
<b>Cultivated hosts</b>	<i>Abutilon indicum</i>	Malvaceae	Country mallow	CABI (2021)
	<i>Acalypha hispida</i>	Euphorbiaceae	Copperleaf	CABI (2021)
	<i>Aegle marmelos</i>	Rutaceae	Indian bael	Chong et al. (2015)
	<i>Aglaonema</i>	Araceae	Aglaonema	Chong et al. (2015)
	<i>Albizia niopoides</i>	Fabaceae	Guanacaste, monkey's earring	García Morales et al. (2016)
	<i>Albizia saman</i>	Fabaceae	Crow bean tree, monkey pod	García Morales et al. (2016)
	<i>Allamanda</i>	Apocynaceae		CABI (2021)
	<i>Allamanda cathartica</i>	Apocynaceae	Yellow allamanda	CABI (2021)
	<i>Alocasia cucullata</i>	Araceae	Chinese taro	Chong et al. (2015)
	<i>Alpinia</i>	Zingiberaceae	Alpinia (ginger and galangal)	Chong et al. (2015)
	<i>Alpinia purpurata</i>	Zingiberaceae	Red ginger	CABI (2021)
	<i>Althaea</i>	Malvaceae	Marshmallow	Chong et al. (2015)
	<i>Amaranthus</i>	Amatanthaceae	Amaranth	Chong et al. (2015)
	<i>Abelmoschus esculentus</i>	Malvaceae	Gumbo, lady's fingers, okra	EPPO GD (2021)
	<i>Abelmoschus manihot</i>	Malvaceae	Sunset musk mallow, sunset hibiscus, hibiscus manihot	García Morales et al. (2016)
	<i>Ananas comosus</i>	Bromeliaceae	Pineapple	EPPO GD (2021)
	<i>Annona</i>	Annonaceae		CABI (2021)
	<i>Annona cherimola</i>	Annonaceae	Cherimoya, custard apple, graveola	EPPO GD (2021)
	<i>Annona muricata</i>	Annonaceae	Prickly custard apple	EPPO GD (2021)
	<i>Annona reticulata</i>	Annonaceae	Bullock's heart	CABI (2021)
	<i>Annona squamosa</i>	Annonaceae	Cachiman, Cuban sugar apple, sugar apple, sweetsop	EPPO GD (2021)
	<i>Anthurium andraeanum</i>	Araceae	Flamingo flower, flamingo lily, oilcloth flower, tail flower	EPPO GD (2021)
	<i>Arachis hypogaea</i>	Fabaceae	Groundnut, monkeynut, peanut	EPPO GD (2021)
	<i>Aralia</i>	Araliaceae		CABI (2021)
	<i>Artocarpus</i>	Moraceae	Breadfruit trees	CABI (2021)
	<i>Artocarpus altilis</i>	Moraceae	Breadfruit	CABI (2021)
	<i>Asparagus</i>	Asparagaceae		CABI (2021)
	<i>Asparagus densiflorus</i>	Liliaceae	Sprenger's asparagus fern	Chong et al. (2015)
	<i>Asparagus officinalis</i>	Asparagaceae	Asparagus, garden asparagus, wild asparagus	EPPO GD (2021)
	<i>Asparagus setaceus</i>	Liliaceae	Asparagus fern	CABI (2021)
	<i>Averrhoa carambola</i>	Oxalidaceae	Caramba, carambola, Chinese gooseberry, country gooseberry, star fruit	EPPO GD (2021)
	<i>Azadirachta indica</i>	Meliaceae	Neem tree	CABI (2021)
	<i>Basella alba</i>	Basellaceae	Malabar spinach	García Morales et al. (2016)
	<i>Bauhinia</i>	Fabaceae	Camel's foot	CABI (2021)
	<i>Bauhinia variegata</i>	Fabaceae	Mountain ebony	CABI (2021)
	<i>Begonia</i>	Begoniaceae	Begonia	Chong et al. (2015)
	<i>Beta</i>	Chenopodiaceae		CABI (2021)

Host status	Host name	Plant family	Common name	Reference
	<i>Beta vulgaris</i>	Amaranthaceae	Beet	EPPO GD (2021)
	<i>Bignonia</i>	Bignoniaceae		CABI (2021)
	<i>Boehmeria</i>	Urticaceae		CABI (2021)
	<i>Boehmeria nivea</i>	Urticaceae	China grass, false nettle, ramie	EPPO GD (2021)
	<i>Bougainvillea</i>	Nyctaginaceae		CABI (2021)
	<i>Bougainvillea</i>	Nyctaginaceae	Bougainvillea	Chong et al. (2015)
	<i>Bougainvillea spectabilis</i>	Nyctaginaceae	Great bougainvillea	Chong et al. (2015)
	<i>Brassica oleracea</i>	Brassicaceae	Cabbage, cauliflower	CABI (2021)
	<i>Caesalpinia coriaria</i>	Fabaceae	Divi-divi	Chong et al. (2015)
	<i>Caesalpinia pulcherrima</i>	Fabaceae	Pride-of-Barbados	Chong et al. (2015)
	<i>Cajanus cajan</i>	Fabaceae	Bengal pea, cajan pea, Congo pea, dal, pigeon pea, red gram	EPPO GD (2021)
	<i>Calliandra</i>	Fabaceae	stick pea	Chong et al. (2015)
	<i>Callistemon</i>	Myrtaceae	Bottlebrush	Chong et al. (2015)
	<i>Calostemma</i>	Amatanthaceae	Wilcannia lily	Chong et al. (2015)
	<i>Camaesyce (Euphorbia) hypericifolia</i>	Euphorbiaceae	Graceful sandmat	Chong et al. (2015)
	<i>Campsis (Tecoma) grandiflora</i>	Bignoniaceae	Chinese trumpet vine	Chong et al. (2015)
	<i>Cananga odorata</i>	Annonaceae	Ilang-ilang (kenanga)	Chong et al. (2015)
	<i>Capsicum</i>	Solanaceae		García Morales et al. (2016)
	<i>Capsicum annuum</i>	Solanaceae	Bell pepper, chilli, paprika, red pepper, sweet pepper	EPPO GD (2021)
	<i>Capsicum frutescens</i>	Solanaceae	Bird chilli, bird pepper, cayenne pepper, chilli pepper, hot pepper	EPPO GD (2021)
	<i>Carica papaya</i>	Caricaceae	Papaw, papaya, pawpaw, tree melon	EPPO GD (2021)
	<i>Carissa macrocarpa</i>	Apocynaceae	Amatungulu (num-num)	Chong et al. (2015)
	<i>Cassia</i>	Fabaceae	Cassia	Chong et al. (2015)
	<i>Cassia javanica</i>	Fabaceae	APPLE BLOSSOM (JAVA CASSIA)	Chong et al. (2015)
	<i>Casuarina</i>	Casuarinaceae	BEEFWOOD	CABI (2021)
	<i>Catharanthus roseus</i>	Apocynaceae	Madagascar periwinkle	Chong et al. (2015)
	<i>Ceiba pentandra</i>	Bombacaceae	Kapok	CABI (2021)
	<i>Celosia argentea</i>	Amatanthaceae	Cock's comb	Chong et al. (2015)
	<i>Centipede tongavine</i>	Araceae		Chong et al. (2015)
	<i>Ceratonia</i>	Fabaceae		CABI (2021)
	<i>Ceratonia siliqua</i>	Fabaceae	Carob, carob tree, locust bean, locust tree, St John's bread	EPPO GD (2021)
	<i>Cestrum nocturnum</i>	Solanaceae	Night jessamine	Chong et al. (2015)
	<i>Chrysanthemum</i>	Asteraceae	Daisy	CABI (2021)
	<i>Chrysanthemum coronarium</i>	Asteraceae	Garland chrysanthemum	CABI (2021)
	<i>Chrysothemis pulchella</i>	Gesneriaceae	Squarestem	Chong et al. (2015)

Host status	Host name	Plant family	Common name	Reference
	<i>Cissus verticillata</i>	Vitaceae	Possum grape vine	García Morales et al. (2016)
	<i>Citrus</i>	Rutaceae		EPPO GD (2021)
	<i>Citrus aurantiifolia</i>	Rutaceae	Lime	CABI (2021)
	<i>Citrus aurantium</i>	Rutaceae	Bigarade, bitter orange, seville orange, sour orange	EPPO GD (2021)
	<i>Citrus maxima</i>	Rutaceae	Pummelo	CABI (2021)
	<i>Citrus medica</i>	Rutaceae	Citron	Chong et al. (2015)
	<i>Citrus nobilis</i>	Rutaceae	Tangor	Chong et al. (2015)
	<i>Citrus paradisi</i>	Rutaceae	Grapefruit, pomelo	EPPO GD (2021)
	<i>Citrus reticulata</i>	Rutaceae	Clementine, mandarin, tangerine	EPPO GD (2021)
	<i>Citrus sinensis</i>	Rutaceae	sweet orange	EPPO GD (2021)
	<i>Clerodendrum aculeatum</i>	Verbenaceae	Haggarbush	Chong et al. (2015)
	<i>Clerodendrum infortunatum</i>	Lamiaceae		CABI (2021)
	<i>Codiaeum</i>	Euphorbiaceae	Codiaeum	Chong et al. (2015)
	<i>Codiaeum variegatum</i>	Euphorbiaceae	Garden croton	CABI (2021)
	<i>Coffea</i>	Rubiaceae	Coffee	CABI (2021)
	<i>Coffea arabica</i>	Rubiaceae	Arabian coffee	EPPO GD (2021)
	<i>Coffea canephora</i>	Rubiaceae	Congo coffee, robusta coffee	EPPO GD (2021)
	<i>Colubrina arborescens</i>	Rhamnaceae	Greenheart	Chong et al. (2015)
	<i>Cordyline terminalis</i>	Liliaceae	Ti plant, palm lily	Chong et al. (2015)
	<i>Couroupita guianensis</i>	Lecythidaceae	Cannonball tree	Chong et al. (2015)
	<i>Crataegus</i>	Rosaceae	Hawthorn	Chong et al. (2015)
	<i>Crescentia cujete</i>	Bignoniaceae	Calabash tree	Chong et al. (2015)
	<i>Crotalaria</i>	Fabaceae		CABI (2021)
	<i>Croton</i>	Euphorbiaceae	Croton	Chong et al. (2015)
	<i>Cucumis sativus</i>	Cucurbitaceae	Cucumber, gherkin	EPPO GD (2021)
	<i>Cucurbita</i>	Cucurbitaceae	Pumpkin	CABI (2021)
	<i>Cucurbita maxima</i>	Cucurbitaceae	Giant pumpkin, marrow	EPPO GD (2021)
	<i>Cucurbita moschata</i>	Cucurbitaceae	Pumpkin	CABI (2021)
	<i>Cucurbita pepo</i>	Cucurbitaceae	Edible gourd, garden marrow, pumpkin, summer squash	EPPO GD (2021)
	<i>Cydonia oblonga</i>	Rosaceae	Quince	CABI (2021)
	<i>Dahlia</i>	Asteraceae		CABI (2021)
	<i>Delonix regia</i>	Fabaceae	Flamboyant	CABI (2021)
	<i>Dendrobium</i>	Orchidaceae	Dendrobium orchid	Chong et al. (2015)
	<i>Dieffenbachia</i>	Araceae	Dieffenbachia	Chong et al. (2015)
	<i>Dimocarpus longan</i>	Sapindaceae	Longan	Chong et al. (2015)
	<i>Diospyros kaki</i>	Ebenaceae	Chinese date plum, Chinese persimmon, Japanese persimmon, kaki, persimmon	EPPO GD (2021)
	<i>Dodonaea viscosa</i>	Sapindaceae	Switch sorrel	CABI (2021)
	<i>Dovyalis (Aberia)</i>	Flacourtiaceae	Ceylon goose berry	Chong et al. (2015)
	<i>Dracaena</i>	Liliaceae	Dracaena (dragon tree)	Chong et al. (2015)
	<i>Duranta</i>	Verbenaceae		CABI (2021)

Host status	Host name	Plant family	Common name	Reference
	<i>Duranta erecta</i>	Verbenaceae	Golden dewdrops	Chong et al. (2015)
	<i>Elaeagnus</i>	Elaeagnaceae	Elaeagnus (oleaster)	Chong et al. (2015)
	<i>Epipremnum pinnatum</i>	Araceae	Centipede tonga vine	Chong et al. (2015)
	<i>Eranthemum pulchellum</i>	Acanthaceae	Blue-sage	Chong et al. (2015)
	<i>Eriobotrya japonica</i>	Rosaceae	Loquat	Chong et al. (2015)
	<i>Eryngium foetidum</i>	Apiaceae	Culantro, shadow beni, Mexican coriander	García Morales et al. (2016)
	<i>Erythrina</i>	Fabaceae		CABI (2021)
	<i>Erythrina corallodendron</i>	Fabaceae	Coral erythrina	Chong et al. (2015)
	<i>Erythrina crista-galli</i>	Fabaceae	Cry baby tree	Chong et al. (2015)
	<i>Erythrina</i> spp.	Fabaceae		CABI (2021)
	<i>Erythrina subumbrans</i>	Fabaceae	December tree	CABI (2021)
	<i>Erythrina variegata</i>	Fabaceae	Flame tree, Indian coral tree, mountain ebony, tiger's claw	EPPO GD (2021)
	<i>Euphorbia</i>	Euphorbiaceae	Spurge	Chong et al. (2015)
	<i>Euphorbia pulcherrima</i>	Euphorbiaceae	Christmas flower, Christmas star, common poinsettia, lobster plant, Mexican flame-leaf, painted leaf, poinsettia	EPPO GD (2021)
	<i>Ficus</i>	Moraceae		CABI (2021)
	<i>Ficus benghalensis</i>	Moraceae	Banyan	CABI (2021)
	<i>Ficus benjamina</i>	Moraceae	Benjamin's fig, Java fig, small-leaved rubber plant, tropical laurel, weeping fig, Benjamin tree	EPPO GD (2021)
	<i>Ficus carica</i>	Moraceae	Common fig, edible fig	EPPO GD (2021)
	<i>Ficus elastica</i>	Moraceae	Rubber plant	CABI (2021)
	<i>Ficus laurifolia</i>	Moraceae		CABI (2021)
	<i>Ficus obtusifolia</i>	Moraceae		CABI (2021)
	<i>Ficus pertusa</i>	Moraceae		CABI (2021)
	<i>Ficus platyphylla</i>	Moraceae		CABI (2021)
	<i>Ficus pumila</i>	Moraceae	Creeping fig	CABI (2021)
	<i>Ficus racemosa</i>	Moraceae	Cluster tree	CABI (2021)
	<i>Ficus religiosa</i>	Moraceae	Sacred fig tree	CABI (2021)
	<i>Ficus semicordata</i>	Moraceae		CABI (2021)
	<i>Flacourtis indica</i>	Flacourtiaceae	Governor's plum	Chong et al. (2015)
	<i>Gerbera</i>	Asteraceae	Gerbera	Chong et al. (2015)
	<i>Glebionis coronaria</i>	Asteraceae	Garland chrysanthemum, chrysanthemum greens, edible chrysanthemum	García Morales et al. (2016)
	<i>Gliricidia sepium</i>	Fabaceae	Gliricidia	CABI (2021)
	<i>Glycine max</i>	Fabaceae	Soybean	EPPO GD (2021)
	<i>Glycosmis pentaphylla</i>	Rutaceae	Orange berry, gin berry	García Morales et al. (2016)
	<i>Cocos nucifera</i>	Arecaceae	Common coconut palm	EPPO GD (2021)
	<i>Colocasia</i>	Araceae		CABI (2021)
	<i>Colocasia esculenta</i>	Araceae	Chinese potato, cocoyam, dasheen, eddoe, Egyptian colocasia, elephant's-ear, kalo, taro, wild taro, yam	EPPO GD (2021)

Host status	Host name	Plant family	Common name	Reference
	<i>Gossypium</i>	Malvaceae	Cotton	CABI (2021)
	<i>Gossypium arboreum</i>	Malvaceae	Cotton, tree	CABI (2021)
	<i>Gossypium herbaceum</i>	Malvaceae	Short staple cotton	CABI (2021)
	<i>Gossypium hirsutum</i>	Malvaceae	American upland cotton, upland cotton	EPPO GD (2021)
	<i>Grevillea</i>	Proteaceae		CABI (2021)
	<i>Grevillea robusta</i>	Proteaceae	Silk oak	Chong et al. (2015)
	<i>Hamelia</i>	Rubiaceae	Hamelia (firebush)	Chong et al. (2015)
	<i>Helianthus annuus</i>	Asteraceae	Common sunflower, sunflower	EPPO GD (2021)
	<i>Hevea</i>	Euphorbiaceae		García Morales et al. (2016)
	<i>Hevea brasiliensis</i>	Euphorbiaceae	Brazilian rubber tree, para rubber, para rubber tree	EPPO GD (2021)
	<i>Hibiscus boryanus</i>	Malvaceae		García Morales et al. (2016)
	<i>Hibiscus</i>	Malvaceae	Rose mallows	CABI (2021)
	<i>Hibiscus acetosella</i>	Malvaceae	African rosemallow	Chong et al. (2015)
	<i>Hibiscus cannabinus</i>	Malvaceae	Bombay hemp, Deccan hemp, kenaf	EPPO GD (2021)
	<i>Hibiscus elatus</i>	Malvaceae	Blue mahoe	CABI (2021)
	<i>Hibiscus manihot</i>	Malvaceae	Bele	CABI (2021)
	<i>Hibiscus mutabilis</i>	Malvaceae	Cotton rose	CABI (2021)
	<i>Hibiscus rosa-sinensis</i>	Malvaceae	China rose, Chinese hibiscus, Chinese rose, Hawaiian hibiscus, rose mallow, rose of China, shoe-black plant, shoe-flower	EPPO GD (2021)
	<i>Hibiscus sabdariffa</i>	Malvaceae	Jamaica sorrel, red sorrel, roselle, tropical cranberry	EPPO GD (2021)
	<i>Hibiscus schizopetalus</i>	Malvaceae	Fringed hibiscus	CABI (2021)
	<i>Hibiscus surattensis</i>	Malvaceae		CABI (2021)
	<i>Hibiscus syriacus</i>	Malvaceae	Shrubby althaea	CABI (2021)
	<i>Hibiscus tiliaceus</i>	Malvaceae	Coast hibiscus, hau tree, linden hibiscus, mahoe, mahoe tree, wild cotton tree	EPPO GD (2021)
	<i>Holmskioldia sanguinea</i>	Verbenaceae	Chinese hatplant	Chong et al. (2015)
	<i>Jacaranda</i>	Bignoniaceae		CABI (2021)
	<i>Jacaranda mimusifolia</i>	Bignoniaceae	Black poui	Chong et al. (2015)
	<i>Jasminum</i>	Oleaceae	Jasmine	CABI (2021)
	<i>Jasminum sambac</i>	Oleaceae	Arabian jasmine	CABI (2021)
	<i>Kalanchoe</i>	Crassulaceae	Widow's-thrill	Chong et al. (2015)
	<i>Kigelia</i>	Bignoniaceae	Sausage tree	Chong et al. (2015)
	<i>Lactuca sativa</i>	Asteraceae	Garden lettuce, lettuce	EPPO GD (2021)
	<i>Lagerstroemia speciosa</i>	Lythraceae	Pride of India	Chong et al. (2015)
	<i>Lantana</i>	Verbenaceae	Lantana	Chong et al. (2015)
	<i>Lantana camara</i>	Verbenaceae	Lantana	CABI (2021)



Host status	Host name	Plant family	Common name	Reference
	<i>Leonotis</i>	Lamiaceae	Lion's ear	Chong et al. (2015)
	<i>Manihot esculenta</i>	Euphorbiaceae	Cassava, manioc, tapioca	EPPO GD (2021)
	<i>Mangifera</i>	Anacardiaceae		CABI (2021)
	<i>Mangifera indica</i>	Anacardiaceae	Mango	EPPO GD (2021)
	<i>Manilkara zapota</i>	Sapotaceae	Bully tree, chapoti, chicle, chiku, marmalade plum, noseberry, sapodilla, sapodilla plum, sapota	EPPO GD (2021)
	<i>Malpighia glabra</i>	Malpighiaceae	Barbados cherry	EPPO GD (2021)
	<i>Malus domestica</i>	Rosaceae	Apple	EPPO GD (2021)
	<i>Malus sylvestris</i>	Rosaceae	Crab apple, wild apple, wild crab	EPPO GD (2021)
	<i>Malvaviscus arboreus</i>	Malvaceae	Wax mallow	CABI (2021)
	<i>Medicago sativa</i>	Fabaceae	Lucerne	CABI (2021)
	<i>Melia azedarach</i>	Meliaceae	Chinaberry tree	Chong et al. (2015)
	<i>Melicocca bijugatus</i>	Sapindaceae	Spanish lime	Chong et al. (2015)
	<i>Mimosa</i>	Fabaceae	Sensitive plants	CABI (2021)
	<i>Mimosa caesalpinifolia</i>	Fabaceae		EPPO GD (2021)
	<i>Mimosa diplotricha</i>	Fabaceae	Creeping-sensitive plant	CABI (2021)
	<i>Mimosa hostilis</i>	Fabaceae		EPPO GD (2021)
	<i>Mimosa pigra</i>	Fabaceae	Giant sensitive plant	CABI (2021)
	<i>Mimosa pudica</i>	Fabaceae	Sensitive plant	CABI (2021)
	<i>Morus</i>	Moraceae	Mulberry tree	CABI (2021)
	<i>Morus alba</i>	Moraceae	Silkworm mulberry, white mulberry	EPPO GD (2021)
	<i>Morus nigra L.</i>	Moraceae	Black mulberry	Chong et al. (2015)
	<i>Murraya exotica</i>	Rutaceae	Chinese box, orange jessamine	Chong et al. (2015)
	<i>Murraya koenigii</i>	Rutaceae	Curry leaf, karapincha	EPPO GD (2021)
	<i>Murraya paniculata</i>	Rutaceae	Orange jasmine, orange jessamine, china box, mock orange	García Morales et al. (2016)
	<i>Musa</i>	Musaceae	Banana	CABI (2021)
	<i>Musa paradisiaca</i>	Musaceae	Plantain	CABI (2021)
	<i>Mussaenda</i>	Rubiaceae		CABI (2021)
	<i>Myrtus communis</i>	Myrtaceae	Myrtle	CABI (2021)
	<i>Nephrolepis biserrata</i>	Dryopteridaceae	Giant swordfern	Chong et al. (2015)
	<i>Nephrolepis exaltata</i>	Dryopteridaceae	Boston swordfern	Chong et al. (2015)
	<i>Nerium oleander</i>	Apocynaceae	Common oleander, oleander, rose bay	EPPO GD (2021)
	<i>Pachystachys lutea</i>	Acanthaceae	Pachystachys, lollipop-plant	Chong et al. (2015)
	<i>Passiflora</i>	Passifloraceae	Passionflower	CABI (2021)
	<i>Passiflora caerulea</i>	Passifloraceae	Bluecrown passionflower	Chong et al. (2015)
	<i>Passiflora edulis</i>	Passifloraceae	Passionfruit	CABI (2021)
	<i>Passiflora quadrangularis</i>	Passifloraceae	Giant granadilla	Chong et al. (2015)
	<i>Pavonia</i>	Malvaceae	Swampmallow	Chong et al. (2015)
	<i>Peperomia pellucid</i>	Piperaceae	Man-to-Man	Chong et al. (2015)
	<i>Pereskia bleo</i>	Cactaceae	Rose cactus	Chong et al. (2015)
	<i>Persea americana</i>	Lauraceae	Alligator pear, avocado, avocado pear, holly ghost pear	EPPO GD (2021)

Host status	Host name	Plant family	Common name	Reference
	<i>Petrea volubilis</i>	Verbenaceae	Queen's-wreath	Chong et al. (2015)
	<i>Phaseolus vulgaris</i>	Fabaceae	Bush bean, climbing French bean, climbing kidney bean, field bean, flageolet bean, French bean, garden bean, green bean, haricot bean, kidney bean, pop bean, snap bean, string bean	EPPO GD (2021)
	<i>Philodendron</i>	Araceae	Philodendron	Chong et al. (2015)
	<i>Phoenix dactylifera</i>	Arecaceae	Common date palm, date palm	EPPO GD (2021)
	<i>Phoenix sylvestris</i>	Arecaceae	East Indian wine palm, silver date palm, wild date palm	EPPO GD (2021)
	<i>Phyllanthus acidus</i>	Euphorbiaceae	Tahitian gooseberry tree	Chong et al. (2015)
	<i>Phyllanthus elsiae</i>	Euphorbiaceae		CABI (2021)
	<i>Phyllanthus niruri</i>	Euphorbiaceae	Seed-under-the-leaf	CABI (2021)
	<i>Plumbago auriculata</i>	Plumbaginaceae	Cape leadwort	Chong et al. (2015)
	<i>Portulaca grandiflora</i>	Portulacaceae	Rose moss	CABI (2021)
	<i>Portulaca oleracea</i>	Portulacaceae	Common purslane, duckweed, little hogweed, pursley	García Morales et al. (2016)
	<i>Portulaca pilosa</i>	Portulacaceae	Kiss-me-quick, rimson-flowered purslane, hairy pigweed, pink purslane, shaggy portulaca	García Morales et al. (2016)
	<i>Prunus armeniaca</i>	Rosaceae	Apricot	EPPO GD (2021)
	<i>Prunus domestica</i>	Rosaceae	European plum, garden plum, plum	EPPO GD (2021)
	<i>Prunus persica</i>	Rosaceae	Peach	EPPO GD (2021)
	<i>Prunus salicina</i>	Rosaceae	Japanese plum	CABI (2021)
	<i>Psidium</i>	Myrtaceae	Guava	CABI (2021)
	<i>Psidium guajava</i>	Myrtaceae	Common guava, guava, yellow guava	EPPO GD (2021)
	<i>Punica granatum</i>	Lythraceae	Pomegranate	EPPO GD (2021)
	<i>Pyrus communis</i>	Rosaceae	Common pear, pear	EPPO GD (2021)
	<i>Quercus</i>	Fagaceae	Oak	Chong et al. (2015)
	<i>Rhododendron</i>	Ericaceae	Azalea	CABI (2021)
	<i>Ricinus communis</i>	Euphorbiaceae	Castor-oil plant, castor bean	EPPO GD (2021)
	<i>Rivina humilis</i>	Phytolacaceae	Rougeplant	Chong et al. (2015)
	<i>Robinia pseudoacacia</i>	Fabaceae	Black locust	CABI (2021)
	<i>Rosa</i>	Rosaceae	Rose	Chong et al. (2015)
	<i>Russelia equisetiformis</i>	Scrophulariaceae	Fountainbush	Chong et al. (2015)
	<i>Saccharum officinarum</i>	Poaceae	Sugarcane	CABI (2021)
	<i>Salix</i>	Salicaceae	Willows	CABI (2021)
	<i>Schefflera</i>	Araliaceae	Schefflera	Chong et al. (2015)
	<i>Senna</i>	Fabaceae	Senna	Chong et al. (2015)
	<i>Senna siamea</i>	Fabaceae	Yellow cassia	CABI (2021)
	<i>Solanum aethiopicum</i>	Solanaceae	African scarlet eggplant	CABI (2021)
	<i>Solanum bicolor</i>	Solanaceae		Chong et al. (2015)
	<i>Solanum lycopersicum</i>	Solanaceae	Tomato	EPPO GD (2021)

Host status	Host name	Plant family	Common name	Reference
	<i>Solanum melongena</i>	Solanaceae	Aubergine, eggplant	EPPO GD (2021)
	<i>Spondias dulcis</i>	Anacardiaceae	Otaheite apple	CABI (2021)
	<i>Spondias purpurea</i>	Anacardiaceae	Red mombin, purple mombin	CABI (2021)
	<i>Stachytarpheta jamaicensis</i>	Verbenaceae	Light-blue snakeweed	Chong et al. (2015)
	<i>Syngonium podophyllum</i>	Araceae	American evergreen	Chong et al. (2015)
	<i>Syzygium cumini</i>	Myrtaceae	Black plum	CABI (2021)
	<i>Syzygium malaccense</i>	Myrtaceae	Malaysian apple	Chong et al. (2015)
	<i>Tabebuia</i>	Bignoniaceae	Trumpet-tree	Chong et al. (2015)
	<i>Tabebuia heterophylla</i>	Bignoniaceae	Pink trumpet tree	CABI (2021)
	<i>Tamarindus indica</i>	Fabaceae	Tamarind	Chong et al. (2015)
	<i>Tamarix</i>	Tamaricaceae	Tamarisk	CABI (2021)
	<i>Tecoma capensis</i>	Bignoniaceae	Cape honeysuckle	Chong et al. (2015)
	<i>Tecoma stans</i>	Bignoniaceae	Yellow trumpetbush	Chong et al. (2015)
	<i>Terminalia catappa</i>	Combretaceae	Singapore almond	CABI (2021)
	<i>Theobroma bicolor</i>	Malvaceae	Bacao, Nicaraguan cocoa	EPPO GD (2021)
	<i>Theobroma cacao</i>	Malvaceae	Cacao, cocoa, common cacao, common cocoa	EPPO GD (2021)
	<i>Theobroma grandiflorum</i>	Malvaceae	Cupuassu	EPPO GD (2021)
	<i>Thunbergia erecta</i>	Acanthaceae	Bush clockvine	Chong et al. (2015)
	<i>Vinca minor</i>	Apocynaceae	Common periwinkle, vinca	Chong et al. (2015)
	<i>Vitis</i>	Vitaceae	Rape	CABI (2021)
	<i>Vitis vinifera</i>	Vitaceae	Common grapevine, grapevine, European grape	EPPO GD (2021)
	<i>Zea mays</i>	Poaceae	Maize	CABI (2021)
	<i>Ziziphus</i>	Rhamnaceae		CABI (2021)
	<i>Ziziphus</i>	Rhamnaceae	Jujube	Chong et al. (2015)
	<i>Ziziphus jujuba</i>	Rhamnaceae	Common jujube	CABI (2021)
	<i>Ziziphus mauritiana</i>	Rhamnaceae	Indian jujube	EPPO GD (2021)
	<i>Ziziphus mucronata</i>	Rhamnaceae		CABI (2021)
	<i>Ziziphus spina-christi</i>	Rhamnaceae	Christ's thorn jujube	CABI (2021)
<b>Wild weed hosts</b>	<i>Abutilon fruticosum</i>	Malvaceae	Texas Indian mallow, pelotazo, sweet Indian mallow	García Morales et al. (2016)
	<i>Acacia</i>	Fabaceae	Wattles	CABI (2021)
	<i>Acacia acatensis</i>	Fabaceae		CABI (2021)
	<i>Acacia cochliacantha</i>	Fabaceae		CABI (2021)
	<i>Acacia farnesiana</i>	Fabaceae	Huisache	CABI (2021)
	<i>Acacia hindsii</i>	Fabaceae		CABI (2021)
	<i>Acacia nilotica</i>	Fabaceae	Gum arabic tree	CABI (2021)
	<i>Acalypha</i>	Euphorbiaceae	Copperleaf	CABI (2021)
	<i>Acalypha indica</i>	Euphorbiaceae	Indian acalypha, Indian mercury, Indian copperleaf, Indian nettle, Three-seeded mercury	García Morales et al. (2016)
	<i>Acalypha wilkesiana</i>	Euphorbiaceae	Copperleaf and Jacob's coat	García Morales et al. (2016)
	<i>Acanthus ilicifolius</i>	Acanthaceae	Copperleaf	CABI (2021)
	<i>Acharia</i>	Limacodidae		CABI (2021)

Host status	Host name	Plant family	Common name	Reference
	<i>Achyranthes aspera</i>	Amaranthaceae	Devil's horsewhip	CABI (2021)
	<i>Acokanthera</i>	Apocynaceae		García Morales et al. (2016)
	<i>Aegiphila martinicensis</i>	Lamiaceae		CABI (2021)
	<i>Albizia</i>	Fabaceae		CABI (2021)
	<i>Albizia lebbbeck</i>	Fabaceae	Indian siris	CABI (2021)
	<i>Angelica</i>	Apiaceae		CABI (2021)
	<i>Anthurium</i>	Araceae		CABI (2021)
	<i>Bauhinia forficata</i>	Fabaceae	Brazilian orchid tree	García Morales et al. (2016)
	<i>Bauhinia racemosa</i>	Fabaceae		García Morales et al. (2016)
	<i>Bauhinia vahlii</i>	Fabaceae		García Morales et al. (2016)
	<i>Biancaea decapetala</i>	Fabaceae	Shoofly, Mauritius, Mysore thorn	García Morales et al. (2016)
	<i>Bidens pilosa</i>	Asteraceae	Beggar tick, bur marigold, butterfly needles	García Morales et al. (2016)
	<i>Blighia sapida</i>	Sapindaceae	Akee	Chong et al. (2015)
	<i>Byttneria aculeata</i>	Malvaceae		CABI (2021)
	<i>Calathea warszewiczii</i>	Marantaceae		García Morales et al. (2016)
	<i>Calophyllum</i>	Calophyllaceae		García Morales et al. (2016)
	<i>Carissa bispinosa</i>	Apocynaceae		García Morales et al. (2016)
	<i>Cassia glauca</i>	Fabaceae		García Morales et al. (2016)
	<i>Cassia renigera</i>	Fabaceae		García Morales et al. (2016)
	<i>Cedrela odorata</i>	Meliaceae	Spanish cedar	CABI (2021)
	<i>Centrolobium paraense</i>	Fabaceae		EPPO GD (2021)
	<i>Chenopodium album</i>	Amaranthaceae	Goosefoot, green pigweed, lamb's quarters, wild spinach, fat-hen, white goosefoot, pigweed	EPPO GD (2021)
	<i>Clitoria ternatea</i>	Fabaceae	Butterfly-pea	CABI (2021)
	<i>Coccoloba uvifera</i>	Polygonaceae	Jamaica kino, platter leaf, sea grape, common sea grape	EPPO GD (2021)
	<i>Combretum</i>	Combretaceae		García Morales et al. (2016)
	<i>Corchorus</i>	Tiliaceae	Jutes	CABI (2021)
	<i>Corchorus capsularis</i>	Tiliaceae	White jute	CABI (2021)
	<i>Corchorus olitorius</i>	Tiliaceae	Jute	CABI (2021)
	<i>Cordia curassavica</i>	Boraginaceae	Black sage or wild sage	García Morales et al. (2016)
	<i>Cordia dichotoma</i>	Boraginaceae	Indian cherry	CABI (2021)
	<i>Cordyline fruticosa</i>	Asparagaceae	Bongbush, cabbage palm, kiwi, palm lily, ti-palm	García Morales et al. (2016)

Host status	Host name	Plant family	Common name	Reference
	<i>Cosmos</i>	Asteraceae		EPPO GD (2021)
	<i>Crotalaria micans</i>	Fabaceae		CABI (2021)
	<i>Croton flavens</i>	Euphorbiaceae		García Morales et al. (2016)
	<i>Cyperus</i>	Cyperaceae		García Morales et al. (2016)
	<i>Dalbergia</i>	Fabaceae	Rosewoods	CABI (2021)
	<i>Datura</i>	Solanaceae	Jimsonweed (angel trumpet)	Chong et al. (2015)
	<i>Daucus carota</i>	Apiaceae	Queen Anne's lace	Chong et al. (2015)
	<i>Desmanthus virgatus</i>	Fabaceae	False tamarind	CABI (2021)
	<i>Dioscorea</i>	Dioscoreaceae		García Morales et al. (2016)
	<i>Emilia</i>	Asteraceae		García Morales et al. (2016)
	<i>Enterolobium</i>	Fabaceae		CABI (2021)
	<i>Enterolobium cyclocarpum</i>	Fabaceae	Ear pod tree	CABI (2021)
	<i>Epipremnum aureum</i>	Araceae	Golden pothos, Ceylon creeper, Hunter's robe, ivy arum	García Morales et al. (2016)
	<i>Erythrina resinifera</i>	Fabaceae		García Morales et al. (2016)
	<i>Erythrina speciosa</i>	Fabaceae		García Morales et al. (2016)
	<i>Erythrina vespertilio</i>	Fabaceae		García Morales et al. (2016)
	<i>Eugenia uniflora</i>	Myrtaceae	Surinam cherry	CABI (2021)
	<i>Euphorbia atoto</i>			CABI (2021)
	<i>Euphorbia hypericifolia</i>	Euphorbiaceae	Graceful spurge, golden spurge, and chickenweed	García Morales et al. (2016)
	<i>Ficus amplissima</i>	Moraceae	Indian Bat tree, Indian Bat fig, Pimpri	García Morales et al. (2016)
	<i>Ficus lacor</i>	Moraceae		García Morales et al. (2016)
	<i>Flacourtia indica</i>	Flacourtiaceae	Governor's plum	García Morales et al. (2016)
	<i>Gliricidia</i>	Fabaceae		CABI (2021)
	<i>Gliricidia maculata</i>	Fabaceae		CABI (2021)
	<i>Grevillea robusta</i>	Proteaceae	Australian silky oak, silk oak, silk-bark oak, silky oak	EPPO GD (2021)
	<i>Grewia</i>	Tiliaceae		CABI (2021)
	<i>Guazuma ulmifolia</i>	Sterculiaceae	Bastard cedar	CABI (2021)
	<i>Gymnanthemum urticifolium</i>	Asteraceae		García Morales et al. (2016)
	<i>Haldina cordifolia</i>	Rubiaceae	Heart-leaf adina	García Morales et al. (2016)
	<i>Heliconia</i>	Heliconiaceae		EPPO GD (2021)
	<i>Hoya carnososa</i>	Asclepiadaceae	Wax plant	CABI (2021)
	<i>Inga</i>	Fabaceae		García Morales et al. (2016)
	<i>Inga edulis</i>	Fabaceae	Food inga, icecream bean, St John's bread	EPPO GD (2021)



Host status	Host name	Plant family	Common name	Reference
	<i>Inga ingoides</i>	Fabaceae		CABI (2021)
	<i>Inga vera</i>	Fabaceae		CABI (2021)
	<i>Ipomoea</i>	Convolvulaceae	Morning glory	CABI (2021)
	<i>Ipomoea batatas</i>	Convolvulaceae	Sweet potato	EPPO GD (2021)
	<i>Ixora</i>	Rubiaceae		EPPO GD (2021)
	<i>Ixora chinensis</i>	Rubiaceae	Flame of the woods, jungle flame, jungle geranium	EPPO GD (2021)
	<i>Jatropha curcas</i>	Euphorbiaceae	Barbados nut, purging nut, physic nut	EPPO GD (2021)
	<i>Laportea aestuans</i>	Urticaceae	West Indian woodnettle	García Morales et al. (2016)
	<i>Lawsonia</i>	Lythraceae		García Morales et al. (2016)
	<i>Lawsonia inermis</i>	Lythraceae	Egyptian privet	CABI (2021)
	<i>Leonotis nepetifolia</i>	Lamiaceae	Christmas candlestick	García Morales et al. (2016)
	<i>Leucaena</i>	Fabaceae		CABI (2021)
	<i>Leucaena leucocephala</i>	Fabaceae	Leucaena	CABI (2021)
	<i>Lithocarpus</i>	Fagaceae	Stone oak	Chong et al. (2015)
	<i>Macaranga</i>	Euphorbiaceae		CABI (2021)
	<i>Malachra alceifolia</i>	Malvaceae		CABI (2021)
	<i>Malpighia</i>	Malpighiaceae		CABI (2021)
	<i>Malpighia emarginata</i>	Malpighiaceae		CABI (2021)
	<i>Malvaviscus conzattii</i>	Malvaceae		CABI (2021)
	<i>Miconia cornifolia</i>	Melastomataceae		García Morales et al. (2016)
	<i>Mikania cordata</i>	Asteraceae		García Morales et al. (2016)
	<i>Mimosa tenuiflora</i>	Fabaceae		García Morales et al. (2016)
	<i>Momordica charantia</i>	Cucurbitaceae	Bitter gourd	CABI (2021)
	<i>Montanoa grandiflora</i>	Asteraceae		CABI (2021)
	<i>Mussaenda erythrophylla</i>	Rubiaceae	Ashanti blood, red flag bush, red flag mussaenda	EPPO GD (2021)
	<i>Nephelium lappaceum</i>	Sapindaceae	Rambutan	EPPO GD (2021)
	<i>Nerium indicum</i>	Apocynaceae		García Morales et al. (2016)
	<i>Opuntia</i>	Cactaceae		EPPO GD (2021)
	<i>Paritium</i>	Malvaceae		García Morales et al. (2016)
	<i>Parkinsonia aculeata</i>	Fabaceae	Mexican palo-verde	CABI (2021)
	<i>Parthenium hysterophorus</i>	Asteraceae	Parthenium weed	CABI (2021)
	<i>Persea</i>	Lauraceae		CABI (2021)
	<i>Petiveria alliacea</i>	Phytolaccaceae		García Morales et al. (2016)
	<i>Phyllanthus amarus</i>	Euphorbiaceae	Gale of the wind, carry me seed, seed on the leaf	García Morales et al. (2016)
	<i>Phyllanthus urinaria</i>	Euphorbiaceae	Leafflower	García Morales et al. (2016)

Host status	Host name	Plant family	Common name	Reference
	<i>Piper tuberculatum</i>	Piperaceae		García Morales et al. (2016)
	<i>Pithecellobium</i>	Fabaceae		CABI (2021)
	<i>Pithecellobium caribaeum</i>	Fabaceae		García Morales et al. (2016)
	<i>Plerandra elegantissima</i>	Araliaceae	False aralia	
	<i>Prosopis</i>	Fabaceae		CABI (2021)
	<i>Prosopis cineraria</i>	Fabaceae	Screw-bean	CABI (2021)
	<i>Prosopis laevigata</i>	Fabaceae		CABI (2021)
	<i>Quisqualis</i>	Combretaceae		CABI (2021)
	<i>Rosa obtusifolia</i>	Rosaceae		García Morales et al. (2016)
	<i>Samanea saman</i>	Fabaceae	Rain tree	CABI (2021)
	<i>Schefflera actinophylla</i>	Araliaceae	Octopus tree, Queensland umbrella tree, star leaf, umbrella tree	EPPO GD (2021)
	<i>Schefflera pueckleri</i>	Araliaceae	Mallet flower	García Morales et al. (2016)
	<i>Schinus molle</i>	Anacardiaceae	False pepper tree	García Morales et al. (2016)
	<i>Schinus terebinthifolia</i>	Anacardiaceae	Brazilian pepper tree	García Morales et al. (2016)
	<i>Scoparia dulcis</i>	Plantaginaceae	Licorice weed, goat weed, scoparia-weed, sweet-broom	García Morales et al. (2016)
	<i>Senna italica</i>	Fabaceae	Senegal senna	García Morales et al. (2016)
	<i>Senna obtusifolia</i>	Fabaceae	Sicklepod	García Morales et al. (2016)
	<i>Senna polyphylla</i>	Fabaceae		García Morales et al. (2016)
	<i>Senna sulfurea</i>	Fabaceae		García Morales et al. (2016)
	<i>Senna surattensis</i>	Fabaceae	Golden senna, foetid cassia, glaucous cassia, glossy shower	García Morales et al. (2016)
	<i>Sesbania sesban</i>	Fabaceae	Sesban, common sesban, Egyptian pea, Egyptian rattle pod	García Morales et al. (2016)
	<i>Sida acuta</i>	Malvaceae	Sida	CABI (2021)
	<i>Solandra</i>	Solanaceae		CABI (2021)
	<i>Solanum americanum</i>	Solanaceae	Eastern black nightshade, glossy nightshade, West Indian nightshade, American black nightshade	EPPO GD (2021)
	<i>Solanum donianum</i>	Solanaceae		García Morales et al. (2016)
	<i>Solanum umbellatum</i>	Solanaceae		CABI (2021)
	<i>Spondias</i>	Anacardiaceae	Purple mombin	CABI (2021)
	<i>Spondias mombin</i>	Anacardiaceae	Golden apple, hog-plum tree, yellow mombin	EPPO GD (2021)
	<i>Spondias tuberosa</i>	Anacardiaceae	Imbu	EPPO GD (2021)
	<i>Synedrella nodiflora</i>	Asteraceae	Synedrella	García Morales et al. (2016)

Host status	Host name	Plant family	Common name	Reference
	<i>Syzygium aqueum</i>	Myrtaceae	Water apple	García Morales et al. (2016)
	<i>Syzygium aromaticum</i>	Myrtaceae	Clove	CABI (2021)
	<i>Tabernaemontana divaricata</i>	Apocynaceae		García Morales et al. (2016)
	<i>Talinum paniculatum</i>	Talinaceae	Fame flower, Jewels-of-Opar, pink baby-breath	EPPO GD (2021)
	<i>Talipariti elatum</i>	Malvaceae	Blue mahoe	García Morales et al. (2016)
	<i>Tamarindus</i>	Fabaceae		CABI (2021)
	<i>Tectona grandis</i>	Lamiaceae	Common teak	EPPO GD (2021)
	<i>Templetonia</i>	Fabaceae		García Morales et al. (2016)
	<i>Tephrosia</i>	Fabaceae	Hoary-pea	CABI (2021)
	<i>Teramnus labialis</i>	Fabaceae	Blue wiss	CABI (2021)
	<i>Terminalia</i>	Combretaceae		García Morales et al. (2016)
	<i>Terminalia mantaly</i>	Combretaceae		CABI (2021)
	<i>Terminalia neotaliala</i>	Combretaceae	Madagascar almond tree	García Morales et al. (2016)
	<i>Tetracera</i>	Dilleniaceae		CABI (2021)
	<i>Theobroma speciosum</i>	Malvaceae		EPPO GD (2021)
	<i>Thespesia</i>	Malvaceae		CABI (2021)
	<i>Thespesia lampas</i>	Malvaceae		CABI (2021)
	<i>Thespesia populnea</i>	Malvaceae	Portia tree	CABI (2021)
	<i>Tithonia diversifolia</i>	Asteraceae	Mexican sunflower	CABI (2021)
	<i>Tradescantia</i>	Commelinaceae		García Morales et al. (2016)
	<i>Trema micrantha</i>	Cannabaceae	Jamaican nettle tree, capulin	García Morales et al. (2016)
	<i>Vachellia nilotica</i>	Fabaceae	Gum arabic tree, babul, thorn mimosa, Egyptian acacia, thorny acacia	García Morales et al. (2016)
	<i>Verbesina fastigiata</i>	Asteraceae		CABI (2021)
	<i>Viburnum odoratissimum</i>	Caprifoliaceae	Sweet viburnum	García Morales et al. (2016)
	<i>Vigna mungo</i>	Fabaceae	Black gram	García Morales et al. (2016)
	<i>Vigna unguiculata</i>	Fabaceae	Cowpea	García Morales et al. (2016)
	<i>Volkameria aculeata</i>	Lamiaceae		García Morales et al. (2016)
	<i>Xanthosoma</i>	Araceae	Cocoyam	CABI (2021)
	<i>Zinnia</i>	Asteraceae		CABI (2021)

## Appendix B – Distribution of *Maconellicoccus hirsutus*

Distribution records based on EPPO Global Database (EPPO, online) and García Morales et al. (2016).

Region	Country	Sub-national (e.g. State)	Status	Reference
North America	Mexico		Present, restricted distribution	EPPO GD (2021)
	USA		Present, restricted distribution	EPPO GD (2021)
	USA	Alabama	Present, no details	EPPO GD (2021)
	USA	California	Present, restricted distribution	EPPO GD (2021)
	USA	Florida	Present, few occurrences	EPPO GD (2021)
	USA	Georgia	Present, few occurrences	EPPO GD (2021)
	USA	Louisiana	Present, no details	EPPO GD (2021)
	USA	New York	Present, no details	EPPO GD (2021)
	USA	North Carolina	Present, no details	EPPO GD (2021)
	USA	Oklahoma	Present, no details	EPPO GD (2021)
	USA	South Carolina	Present, no details	EPPO GD (2021)
	USA	Tennessee	Present, no details	EPPO GD (2021)
	USA	Texas	Present, no details	García Morales et al. (2016)
Central America	Belize		Present, no details	EPPO GD (2021)
	Costa Rica		Present, restricted distribution	EPPO GD (2021)
	Guatemala		Absent, unreliable record	EPPO GD (2021)
	Nicaragua		Present, restricted distribution	EPPO GD (2021)
Caribbean	Anguilla		Absent, unreliable record	EPPO GD (2021)
	Antigua and Barbuda		Present, no details	EPPO GD (2021)
	Aruba		Present, no details	EPPO GD (2021)
	Bahamas		Present, no details	EPPO GD (2021)
	Barbados		Present, no details	EPPO GD (2021)
	Cayman Islands		Present, no details	EPPO GD (2021)
	Cuba		Present, no details	García Morales et al. (2016)
	Dominica		Present, no details	EPPO GD (2021)
	Dominican Republic		Absent, unreliable record	EPPO GD (2021)
	Grenada		Present, restricted distribution	EPPO GD (2021)
	Guadeloupe		Present, no details	EPPO GD (2021)
	Haiti		Present, widespread	EPPO GD (2021)
	Jamaica		Present, few occurrences	EPPO GD (2021)
	Martinique		Present, no details	EPPO GD (2021)
	Montserrat		Present, no details	EPPO GD (2021)
	Netherlands Antilles		Present, no details	EPPO GD (2021)
	Puerto Rico		Present, no details	EPPO GD (2021)
	Saint Lucia		Present, no details	EPPO GD (2021)
	St Kitts-Nevis		Present, no details	EPPO GD (2021)
	St Vincent and the Grenadines		Present, no details	EPPO GD (2021)
Saint Barthelemy		Present, no details	García Morales et al. (2016)	
Saint Martin		Present, no details	García Morales et al. (2016)	

Region	Country	Sub-national (e.g. State)	Status	Reference
	Trinidad and Tobago		Present, no details	EPPO GD (2021)
	Virgin Islands (British)		Present, no details	EPPO GD (2021)
	Virgin Islands (US)		Present, no details	EPPO GD (2021)
South America	Brazil		Present, restricted distribution	EPPO GD (2021)
	Brazil	Alagoas	Present, no details	EPPO GD (2021)
	Brazil	Bahia	Present, restricted distribution	EPPO GD (2021)
	Brazil	Espirito Santo	Present, restricted distribution	EPPO GD (2021)
	Brazil	Maranhao	Present, no details	EPPO GD (2021)
	Brazil	Mato Grosso	Present, no details	EPPO GD (2021)
	Brazil	Para	Present, no details	EPPO GD (2021)
	Brazil	Pernambuco	Present, no details	EPPO GD (2021)
	Brazil	Rio Grande do Sul	Present, no details	EPPO GD (2021)
	Brazil	Roraima	Present, restricted distribution	EPPO GD (2021)
	Brazil	Santa Catarina	Present, no details	EPPO GD (2021)
	Brazil	Sao Paulo	Present, no details	EPPO GD (2021)
	Colombia		Present, restricted distribution	EPPO GD (2021)
	French Guiana		Present, no details	EPPO GD (2021)
	Guyana		Present, widespread	EPPO GD (2021)
	Suriname		Present, restricted distribution	EPPO GD (2021)
Venezuela		Present, no details	EPPO GD (2021)	
EU (27)	Cyprus		Present, widespread	EPPO GD (2021)
	Greece		Present, restricted distribution	EPPO GD (2021)
Africa	Algeria		Absent, invalid record	EPPO GD (2021)
	Benin		Present, no details	EPPO GD (2021)
	Burkina Faso		Present, no details	EPPO GD (2021)
	Cameroon		Present, no details	EPPO GD (2021)
	Central African Republic		Present, no details	EPPO GD (2021)
	Chad		Present, no details	EPPO GD (2021)
	Congo		Present, no details	EPPO GD (2021)
	Congo, Democratic republic of the		Present, no details	EPPO GD (2021)
	Cote d'Ivoire		Present, no details	EPPO GD (2021)
	Egypt		Present, no details	EPPO GD (2021)
	Gabon		Present, no details	EPPO GD (2021)
	Gambia		Present, no details	EPPO GD (2021)
	Kenya		Present, no details	EPPO GD (2021)
	Liberia		Present, no details	EPPO GD (2021)
	Niger		Present, no details	EPPO GD (2021)
	Nigeria		Present, no details	EPPO GD (2021)
	Reunion		Present, no details	EPPO GD (2021)
	Senegal		Present, no details	EPPO GD (2021)
	Seychelles		Present, no details	EPPO GD (2021)
	Socotra Island		Present, no details	García Morales et al. (2016)
	Somalia		Present, no details	EPPO GD (2021)
	Sudan		Present, no details	EPPO GD (2021)



Region	Country	Sub-national (e.g. State)	Status	Reference
	Tanzania		Present, no details	EPPO GD (2021)
	Tunisia		Present, restricted distribution	EPPO GD (2021)
	Zambia		Absent, invalid record	EPPO GD (2021)
	Zaire		Present, no details	García Morales et al. (2016)
	Zanzibar		Present, no details	García Morales et al. (2016)
Asia	Bali		Present, no details	García Morales et al. (2016)
	Bangladesh		Present, no details	EPPO GD (2021)
	Brunei Darussalam		Present, no details	EPPO GD (2021)
	Cambodia		Present, no details	EPPO GD (2021)
	China		Present, restricted distribution	EPPO GD (2021)
	China	Aomen (Macau)	Present, no details	EPPO GD (2021)
	China	Guangdong	Present, no details	EPPO GD (2021)
	China	Guangxi	Present, no details	García Morales et al. (2016)
	China	Shanxi	Present, no details	EPPO GD (2021)
	China	Xianggang (Hong Kong)	Present, no details	EPPO GD (2021)
	China	Xizhang	Present, no details	EPPO GD (2021)
	China	Yunnan	Present, no details	EPPO GD (2021)
	China	Hong Kong	Present, no details	García Morales et al. (2016)
	India		Present, widespread	EPPO GD (2021)
	India	Andaman and Nicobar Islands	Present, no details	EPPO GD (2021)
	India	Andhra Pradesh	Present, no details	EPPO GD (2021)
	India	Assam	Present, no details	EPPO GD (2021)
	India	Bihar	Present, no details	EPPO GD (2021)
	India	Delhi	Present, no details	EPPO GD (2021)
	India	Gujarat	Present, no details	EPPO GD (2021)
	India	Karnataka	Present, no details	EPPO GD (2021)
	India	Kerala	Present, no details	EPPO GD (2021)
	India	Madhya Pradesh	Present, no details	EPPO GD (2021)
	India	Maharashtra	Present, no details	EPPO GD (2021)
	India	Odisha	Present, no details	EPPO GD (2021)
	India	Punjab	Present, no details	EPPO GD (2021)
	India	Tamil Nadu	Present, no details	EPPO GD (2021)
	India	Telangana	Present, no details	EPPO GD (2021)
	India	Tripura	Present, no details	EPPO GD (2021)
	India	Uttar Pradesh	Present, no details	EPPO GD (2021)
	India	West Bengal	Present, no details	EPPO GD (2021)
	Indonesia		Present, widespread	EPPO GD (2021)
	Indonesia	Flores	Present, no details	García Morales et al. (2016)
	Indonesia	Irian Jaya	Present, no details	EPPO GD (2021)
	Indonesia	Java	Present, no details	EPPO GD (2021)
	Indonesia	Lombok	Present, no details	García Morales et al. (2016)

Region	Country	Sub-national (e.g. State)	Status	Reference
	Indonesia	Nusa Tenggara	Present, no details	EPPO GD (2021)
	Indonesia	Sulawesi	Present, no details	EPPO GD (2021)
	Indonesia	Sumatra	Present, no details	EPPO GD (2021)
	Iran		Present, no details	EPPO GD (2021)
	Israel		Present, few occurrences	EPPO GD (2021)
	Japan		Present, restricted distribution	EPPO GD (2021)
	Japan	Ryukyu Archipelago	Present, no details	EPPO GD (2021)
	Jordan		Present, no details	EPPO GD (2021)
	Laos		Present, no details	EPPO GD (2021)
	Lebanon		Present, no details	EPPO GD (2021)
	Malaya		Present, no details	García Morales et al. (2016)
	Malaysia		Present, no details	EPPO GD (2021)
	Malaysia	West	Present, no details	EPPO GD (2021)
	Maldives		Present, no details	EPPO GD (2021)
	Myanmar		Present, no details	EPPO GD (2021)
	Nepal		Present, no details	EPPO GD (2021)
	Oman		Present, no details	EPPO GD (2021)
	Pakistan		Present, no details	EPPO GD (2021)
	Philippines		Present, no details	EPPO GD (2021)
	Saudi Arabia		Present, no details	EPPO GD (2021)
	Singapore		Present, no details	EPPO GD (2021)
	Sri Lanka		Present, no details	EPPO GD (2021)
	Taiwan		Present, no details	EPPO GD (2021)
	Thailand		Present, no details	EPPO GD (2021)
	Turkey		Present, no details	EPPO GD (2021)
	United Arab Emirates		Present, no details	EPPO GD (2021)
	Vietnam		Present, no details	EPPO GD (2021)
	Yemen		Present, no details	EPPO GD (2021)
Oceania	Australia		Present, no details	EPPO GD (2021)
	Australia	Northern Territory	Present, no details	EPPO GD (2021)
	Australia	Queensland	Present, no details	EPPO GD (2021)
	Australia	South Australia	Present, no details	EPPO GD (2021)
	Australia	Western Australia	Present, no details	EPPO GD (2021)
	Fiji		Present, no details	EPPO GD (2021)
	Guam		Present, no details	EPPO GD (2021)
	Micronesia		Present, restricted distribution	EPPO GD (2021)
	New Caledonia		Present, no details	EPPO GD (2021)
	Northern Mariana Islands		Present, no details	EPPO GD (2021)
	Palau		Present, no details	EPPO GD (2021)
	Papua New Guinea		Present, no details	EPPO GD (2021)
	Samoa		Present, no details	EPPO GD (2021)
	Solomon Islands		Present, no details	EPPO GD (2021)
	Tonga		Present, no details	EPPO GD (2021)
	Tuvalu		Present, no details	EPPO GD (2021)
	USA	Hawaii	Present, no details	EPPO GD (2021)
	Vanuatu		Present, no details	EPPO GD (2021)

## Appendix C – Import data

**Table C.1: Fresh or dried citrus (CN code: 0805) imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 22/9/2021)**

COUNTRY	2016	2017	2018	2019	2020
Australia	3,279.84	1,284.38	644.97	10,645.40	2,733.47
Bangladesh	227.61	229.58	159.67	322.42	1,183.66
Brazil	864,863.09	903,432.95	900,907.24	822,134.46	902,354.68
Burkina Faso	78.14	148.57	103.95	38.95	53.52
Cameroon	10.48	0.20			
China	827,840.57	1,084,857.27	1,024,163.15	1,108,595.22	1,098,691.70
Colombia	44,825.37	79,400.99	123,887.46	136,914.85	172,413.40
Costa Rica	4,700.31	921.32	704.93	231.20	461.60
Cuba	7,165.74	3,863.97	4,438.14	3,422.11	556.03
Dominica	865.67	193.34	57.65	76.50	78.69
Egypt	1,931,586.64	2,246,998.88	2,643,272.02	2,206,932.71	2,850,742.72
Guyana					24.00
Haiti	207.41	176.53	72.10	31.00	248.29
Hong Kong		0.00		2.27	1.00
India	246.80	1.00	449.63	88.51	254.95
Indonesia	566.73	555.70	779.35	836.73	864.54
Iran	1,533.22	1,218.52	1,208.01	2,174.22	1,882.74
Jamaica	3,633.97	3,325.11	675.68	2,409.55	1,646.87
Israel	799,118.49	969,403.62	824,601.66	812,738.57	878,713.15
Jordan	1.17	0.00	3.79	1.40	11.80
Japan	352.58	417.44	270.73	319.24	162.50
Kenya			8.80		34.56
Laos	51.94	2.10			20.23
Lebanon	503.21	1,504.91	7.46	7.28	3.19
Malaysia	4.18	39.02	83.45	7.71	
Mexico	570,402.80	553,818.66	589,021.12	443,743.54	349,628.56
Nepal		1,170.00			
Nigeria			0.03	0.10	200.00
Pakistan			2.45	0.59	
Philippines			0.20	7.71	0.10
Somalia	490.30	193.21	367.52	514.30	342.10
Sudan			2.10		20.58
Taiwan	157.49				0.01
Tanzania	179.90	190.01	144.12	35.95	75.50
Thailand	426.42	1,283.13	659.74	624.93	194.87
Tunisia	175,010.90	172,515.76	125,258.30	133,950.15	75,620.02
Turkey	2,569,671.58	2,026,980.05	3,149,386.85	2,102,077.48	2,573,806.18
United States	301,229.06	231,210.47	185,706.99	177,755.45	148,845.72
Venezuela	744.08	2,216.36	681.07		
Viet Nam	28,649.46	46,738.17	70,934.07	73,964.35	63,730.13

**Table C.2: Cotton linters (CN code: 140420)** imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 22/9/2021)

COUNTRY	2016	2017	2018	2019	2020
Benin	400.00	294.95	608.38	132.94	87.99
Brazil	13,493.54	57,840.63	68,605.72	50,783.56	57,176.03
China	1,530.80	10.00	44.83	102.75	188.29
Egypt		1.47			
India	1,136.10	589.38	487.65	735.71	2,148.17
Indonesia			27.55		5.38
Iran					3.93
Turkey	40,881.83	115,022.78	88,098.66	82,852.55	81,157.09
United States	56,181.45	32,472.85	16,629.25	7,933.06	19,150.08
Viet Nam			0.21	0.34	

**Table C.3: Fresh or dried bananas (CN code: 0803)** imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 22/9/2021)

COUNTRY	2016	2017	2018	2019	2020
Bangladesh	174.66	79.85	72.75	38.05	35.64
Brazil	149,108.03	26,855.08	59,677.31	104,909.74	98,434.39
Cameroon	2,521,882.41	2,341,539.74	1,791,447.01	1,520,648.04	1,579,456.86
China	252.64	188.73	390.56	545.74	854.93
Colombia	10,120,590.13	11,594,479.46	11,282,545.88	11,524,355.75	12,193,049.39
Costa Rica	9,662,138.79	9,663,219.69	10,125,330.57	9,405,488.40	10,342,372.80
Cuba					1.28
Egypt	42.98	0.18	146.87		
India	515.19	445.99	571.13	607.74	1,418.91
Indonesia		0.01	37.27	14.72	64.17
Iran			0.09	2.86	12.33
Israel	2.10				0.75
Kenya	1.90	0.72	6.15	11.23	14.95
Malaysia			8.02		
Mexico	516,367.97	558,896.47	348,905.62	239,173.11	141,492.42
Nigeria	0.72	2.04	2.50	0.84	6.35
Pakistan			2.60	49.70	
Philippines	2,480.90	11,415.47	1,674.92	2,160.35	1,240.80
Saudi Arabia			5.00		
Singapore		0.06	0.12		
Taiwan	0.15				
Tanzania	28.02	11.93	33.68	34.24	34.74
Thailand	550.44	674.34	603.32	526.15	334.58
Turkey	202.06		210.60	0.14	
United States	7.00	6.37	1.54	6.32	10.37
Viet Nam	276.26	178.84	190.96	210.11	142.71
Zambia		0.72			

**Table C.4: Fresh grapes (CN code: 080610)** imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 22/9/2021)

COUNTRY	2016	2017	2018	2019	2020
Australia	2.95	0.50			
Bangladesh	1.05		0.50		
Brazil	194,152.79	249,279.81	271,987.56	196,465.22	228,095.15
China	0.00	6.00	0.03		
Colombia		0.00	381.30	669.12	186.96
Egypt	330,565.57	404,801.23	429,994.87	442,798.85	462,890.07
India	640,933.67	827,467.67	722,802.04	950,910.96	733,881.71
Iran			2,158.50	366.00	399.80
Israel	13,169.16	7,165.09	6,397.33	318.24	1,080.90
Japan	4.84	1.19	1.17	1.15	20.67
Kenya			186.96		
Mexico		358.96		186.71	184.62
Thailand	0.37	0.14	0.16		0.87
Tunisia	657.82		239.62	40.60	192.00
Turkey	298,205.16	375,776.41	227,616.42	272,447.02	287,021.27
United States	1,714.93	8,868.74	4,413.37	1,866.20	1,072.48
Zambia		0.28	0.03		

**Table C.5: Fresh or dried avocados (CN code: 080440)** imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 22/9/2021)

COUNTRY	2016	2017	2018	2019	2020
Brazil	44,357.36	71,040.50	68,697.61	78,673.73	48,183.82
Cameroon	133.50	173.54	221.30	259.38	205.93
China	193.97	35.28		1.23	0.04
Colombia	152,115.55	210,139.60	251,050.33	387,367.23	663,149.95
Costa Rica		21.56	9.98	428.45	686.40
Cuba	109.09	73.94	41.53	131.08	34.33
Egypt	211.20	5.35	4.58	79.92	363.95
India	0.04	2.06	0.52	0.06	
Israel	301,123.91	424,267.97	370,378.23	437,318.01	345,663.97
Kenya	228,426.16	243,947.31	404,593.87	346,231.90	435,309.11
Malaysia	0.03		47.04		
Mexico	503,687.52	445,611.06	463,741.28	767,878.48	716,205.77
Nigeria	1.06	3.15	3.18	0.51	
Tanzania	26,823.05	25,773.58	55,517.16	60,480.96	50,769.74
Thailand	3.68	9.76	9.66	9.06	3.39
Turkey	213.41	477.05	1,530.93	2,172.09	1,864.65
United States	8,819.53	1.19	2,546.86	0.02	4.66
Viet Nam	1.00			0.05	
Zambia			53.68		



**Table C.6: Fresh tamarinds, cashew apples, lychees, jackfruit, sapodillo plums, passion fruit, carambola and pitahaya (CN code: 08109020) imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 22/9/2021)**

COUNTRY	2016	2017	2018	2019	2020
Australia					12.50
Bangladesh	140.15	222.55	291.61	206.12	382.00
Brazil	49.36	147.37	368.88	966.63	1,220.26
Cameroon	41.84	100.53	38.52	92.00	46.11
China	314.75	287.38	1,112.11	1,014.77	823.41
Colombia	69,743.63	72,656.37	83,639.84	89,847.31	90,741.20
Costa Rica	9.11	3.52	0.13	18.62	
Egypt		13.79			39.05
Hong Kong		9.66			
India	324.19	621.75	1,095.12	1,168.69	754.33
Indonesia	103.20	333.37	297.72	246.67	463.60
Iran	6.25		1.75	0.50	3.88
Israel	2,943.37	2,919.30	1,061.09	1,125.92	594.86
Kenya	714.44	221.45	603.11	481.00	697.14
Malaysia	15,348.23	14,205.33	13,879.92	14,235.96	7,849.69
Mexico	543.90	212.78	1,295.08	669.87	2,331.91
Nigeria		0.00		1.91	3.09
Pakistan	2.22	3.34	8.17		
Philippines	9.78	14.26		0.88	
Singapore	9.00		8.48		
Taiwan	11.92		10.59	25.97	8.97
Tanzania	0.35		1.27	8.77	4.52
Thailand	9,774.93	10,279.68	12,461.38	14,900.21	10,138.74
Turkey			8.61	18.92	23.40
United States	3.97	3.00	0.07		0.02
Viet Nam	33,078.82	38,428.61	44,070.83	52,846.33	45,652.75
Zambia		631.60	4,568.50	3,526.04	3,087.70

**Table C.7: Fresh or dried pineapples (CN code: 08043000) imported in 100 kg into the EU (27) from regions where *Maconellicoccus hirsutus* is known to occur (Source: Eurostat accessed on 19/11/2021)**

COUNTRY	2016	2017	2018	2019	2020
Algeria		0.00		0.01	
Aruba		0.00			
Australia		0.00		0.00	0.01
Bahamas		0.00			
Belize		0.00			
Benin	29,484.88	9,456.56	8,065.08	7,481.67	12,849.58
Brazil	1,522.02	1,272.34	484.83	639.05	280.66
Burkina Faso			145.92	19.68	3.57
Cameroon	38,878.76	39,301.85	30,633.74	23,825.83	13,811.36
China	69.90	25.05	9.91	62.65	42.74
Colombia	64,893.82	123,462.45	91,067.04	53,663.49	42,136.78
Congo		0.00		2.87	3.40
Congo, Democratic Republic of	0.78	2.56	0.85		0.07

<b>COUNTRY</b>	<b>2016</b>	<b>2017</b>	<b>2018</b>	<b>2019</b>	<b>2020</b>
Costa Rica	6,095,312.66	6,832,249.09	7,693,551.48	7,543,050.71	6,650,975.31
Côte d'Ivoire (Ivory Coast)	202,205.93	255,038.72	220,581.56	244,175.93	203,552.53
Cuba	10,645.21	4,382.57	3,838.50	1,998.42	976.85
Dominican Republic	29,667.00	15,582.31	19,723.37	20,566.35	20,525.91
Egypt		201.60			28.16
Fiji		0.00			
Gabon		0.00			
Grenada		0.00			
Guatemala		229.74	40.08	64.03	282.50
Guinea	17.35	98.34	83.45	72.90	19.95
Guyana		0.00			22.00
India	186.71	17.99	75.85	11.52	1.00
Indonesia	0.24	543.77	0.09		2.50
Iran, Islamic Republic of		0.00		0.01	0.00
Israel		2.81	0.20	0.01	
Jamaica		0.00			
Japan	0.02	0.00			
Jordan		0.00		36.00	
Kenya		761.13	745.19	2,147.97	23,799.06
Lao People's Democratic Republic (Laos)		0.00			
Lebanon	0.16	0.00		5.05	
Libya		0.00			
Malaysia	13.60	5.00		2.40	
Maldives		0.00			
Mexico	1,268.22	2,957.94	773.74	142.42	174.97
Nicaragua		0.00			
Nigeria	0.54	0.95	0.13	0.24	0.01
Oman		0.00			
Pakistan		0.00			
Palau		0.00			
Philippines	93.71	114.23	183.83	86.03	566.04
Saudi Arabia		0.00		0.45	0.17
Singapore	0.20	0.00		0.29	
Sri Lanka	1,774.66	5,755.44	4,125.57	2,675.19	2,636.02
Suriname		0.00			
Taiwan		0.00		0.07	0.05
Thailand	10,183.30	11,093.21	9,505.48	8,056.49	8,828.72
Trinidad and Tobago		0.00			
Tunisia	0.05	0.00		0.01	0.03
Turkey		0.00		25.20	0.04
United Arab Emirates		0.00			0.02
United States	69.72	56.66	22.03	28.28	57.29
Venezuela, Bolivarian Republic of	0.15	0.00	0.19	0.04	
Viet Nam	91.31	65.87	9.88	20.20	2.18
Virgin Islands, British		0.00			
Virgin Islands, United States		0.00			
Zambia		0.00			