This is the peer reviewd version of the followng article:

INTEGRATING SHALLOW BENTHIC AND CALCAREOUS NANNOFOSSIL ZONES: THE LOWER EOCENE OF THE MONTE POSTALE SECTION (NORTHERN ITALY) / Papazzoni, Cesare Andrea; Fornaciari, Eliana; Giusberti, Luca; Vescogni, Alessandro; Fornaciari, Beatrice. - In: PALAIOS. - ISSN 0883-1351. - STAMPA. - 32:1(2017), pp. 6-17. [10.2110/palo.2016.014]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

30/04/2024 22:08

- 1 RRH: SB AND CN BIOZONATION IN THE EOCENE OF MONTE POSTALE, ITALY
- 2 LLH: C.A. PAPAZZONI ET AL.
- 3 Research Article
- 4 DOI: http://dx.doi.org/10.2110/palo.2016.014
- 5 INTEGRATING SHALLOW BENTHIC AND CALCAREOUS NANNOFOSSIL ZONES: THE
- 6 LOWER EOCENE OF THE MONTE POSTALE SECTION (NORTHERN ITALY)
- 7 CESARE ANDREA PAPAZZONI,<sup>1</sup> ELIANA FORNACIARI,<sup>2</sup> LUCA GIUSBERTI,<sup>2</sup>
- 8 ALESSANDRO VESCOGNI,<sup>1</sup> and BEATRICE FORNACIARI<sup>1</sup>
- 9 <sup>1</sup>University of Modena and Reggio Emilia, Department of Chemical and Geological Sciences, Via
- 10 Campi 103, 41125 Modena, Italy
- <sup>11</sup> <sup>2</sup>University of Padova, Department of Geosciences, Via Gradenigo 6, 35131 Padova, Italy
- 12 email: papazzoni@unimore.it
- 13 ABSTRACT: The Monte Postale section (Bolca, northern Italy), one of the most famous
- 14 Lagerstätten of the Eocene, has been investigated to reconstruct the sedimentary succession and to
- 15 determine both the larger foraminiferal and the calcareous nannofossil biozones. The results
- 16 allowed us to ascribe the Monte Postale limestones to the Shallow Benthic Zone 11 and to the
- 17 calcareous nannofossil Zone CNE 5-?6 (= NP 13-?14a). The direct correlation of the SB and CNE
- 18 Zones is consistent with the current biostratigraphic schemes and allows assignment of the
- 19 deposition of the succession to the interval between 50.7 and 48.9 Ma, in the late Ypresian (early
- 20 Eocene). According to the available biostratigraphic data, the uppermost portion of the Monte
- 21 Postale section should correlate with the Pesciara limestones.

22 INTRODUCTION

- 23 The Eocene Lagerstätten of Bolca (northern Italy) are known worldwide for the exceptional
- 24 preservation and diversity of their faunas, including soft-bodied organisms (e.g., Sorbini 1972
- 25 1999), and especially for their rich ichthyofauna, represented by more than 240 taxa (Carnevale et

al. 2014). Often referred to as "Monte Bolca", the locality is actually represented by two sites: the
Pesciara di Bolca and the Monte Postale (Fig. 1). These sites are geographically close each other,
but no direct, physical correlation is possible because they are separated by a narrow valley,

volcanic rocks, and possibly a fault. Moreover, the thickness of sedimentary rocks exposed varies
considerably, being less than 20 m for the Pesciara limestone (Papazzoni and Trevisani 2006) and
more than 130 m for the Eocene limestones in the Monte Postale (this work).

32 The Pesciara limestones have been ascribed by Medizza (1975) to the Discoaster sublodoensis

Zone (= NP 14 Zone of Martini 1971) based on calcareous nannofossils. More recently, a study of
the *Alveolina* assemblages by Papazzoni and Trevisani (2006) identified the SBZ 11 (Serra- Kiel et
al. 1998; middle Cuisian or upper part of the Ypresian). Accordingly, the Pesciara succession was
restricted to the uppermost part of the SBZ 11.

37 The age of the Monte Postale section has long been debated. Munier-Chalmas (1891) recognized in 38 the Monte Postale the "Tufs et calcaire de Spilecco", attributed to the lower Eocene, lying on the 39 Cretaceous Scaglia Rossa and overlain by the volcanic breccia with "Lithothamnium Bolcense" 40 (nomen nudum) representing the base of the middle Eocene. Fabiani (1914, 1915) later reported the 41 same age for the main body of limestones that constitutes the Monte Postale. Malaroda (1954) provided the same age, but in a footnote (p. 6) remarked that Arni (1939) determined the 42 43 nummulites from the nearby Brusaferri locality were Cuisian age (= late Ypresian), suggesting that the alveolines from the Monte Postale could also be from the Ypresian. Hottinger (1960) recognized 44 45 larger foraminiferal assemblages from the Bolca area not younger than the "middle Cuisian", 46 therefore definitely older than the Lutetian. Unfortunately, in the comprehensive geological and 47 stratigraphic survey of the Bolca area by Barbieri and Medizza (1969) the Monte Postale was not 48 taken into consideration. Trevisani (2015), based on preliminary data by Papazzoni and Trevisani 49 (2009), ascribed the 60 m of "Nummulitic limestone" of his M. Postale 1 succession to the SBZ 10-50 11. However, the study does not report the distribution of fossils in the section, and determined the

51 SBZ 10 on the presence of *Nummulites partschi*, which indeed spans through the SBZ 10 and 11
52 (see Serra-Kiel et al. 1998).

53 With the aim of obtaining a robust biostratigraphic framework for the Monte Postale, possibly with 54 a direct correlation between shallow-water zonation and planktonic faunas, a detailed revision of the Monte Postale section was completed. A series of detailed field surveys were integrated with the 55 56 study of a core sample drilled in 2000 and of field data collected during the quarrying campaigns 57 directed by the Verona Museum of Natural History (2000–2004). More than 80 samples spanning the 135-m thick succession were examined for larger foraminifera (especially alveolines) and 58 calcareous nannofossils (Figs. 2, 3). This dataset allows both the correlation between the Shallow 59 60 Benthic (SB) Zones and the Calcareous Nannofossil Eocene (CNE) Zones and precise dating, thus clarifying the relationships between the laminites with fish and plants of the Monte Postale and 61 those of the Pesciara. 62

#### 63 GEOLOGICAL AND STRATIGRAPHICAL SETTING

64 The Monte Postale section is renowned not only for its vertebrate and mollusk fossil faunas (e.g., Sorbini 1972; Papazzoni et al. 2014a), but also because it is one of the few shallow-water 65 66 successions showing the establishment of the first nuclei of what later became the Lessini Shelf (Bosellini 1989). The Lessini Shelf was one of the main paleogeographic features of Southern Alps 67 68 during the Eocene (e.g., Bosellini 1989; Luciani 1989), representing at that time the northernmost margin of the Adriatic Plate (Carminati et al. 2012). It was the result of uplift of a pre-existing 69 70 structural high, the "Trento Plateau" (Middle Jurassic-Paleocene), which formed with the drowning of the Early Jurassic Trento Platform (Bosellini et al. 1981; Winterer and Bosellini 1981). The 71 72 Paleogene uplift was not uniform, due to the Alpine tectonics acting in this area with block-faulting 73 (Márton et al. 2011). The tectonic activity resulted in deposition of volcanic and volcaniclastic 74 material, mainly basaltic in composition. Beginning in the early Eocene, in shallower areas in the 75 photic zone, these volcanic and volcaniclastic deposits are commonly intercalated with limestones

76 and marls derived from thriving larger foraminifera, mollusks, corals, algae, and other biotic carbonate producers (Doglioni and Bosellini 1987; Bosellini 1989; Luciani 1989; Bassi et al. 2008). 77 78 The main structural element influencing the Lessini area is the Castelvero Fault (Barbieri 1972), 79 which runs a few kilometers west of the Monte Postale. This fault is on the border between two 80 sectors. In particular, the eastern area was characterized by a graben (or semi-graben) known as 81 Alpone-Chiampo graben (Barbieri et al. 1982) or Alpone-Agno half-graben (Barbieri et al. 1991), or 82 Alpone-Agno graben (Zampieri 1995), where subsidence and volcanic activity were more 83 pronounced than in the western part. Barbieri et al. (1991) recognized six volcanic stages from the late Paleocene up to the Bartonian (see also Barbieri and Zampieri 1992; Zampieri 1995). The 84 85 volcanic rocks of the Monte Postale have traditionally been ascribed to the third stage, but this 86 attribution was not supported by radiometric dating. 87 The lithostratigraphy of the Lessini area suffers from a lack of formal names and from the poor biostratigraphic constrains of the most commonly used units. For example, in the past the name 88 89 "Spileccian" or "Calcari di Spilecco" (Spilecco limestones) was used as a synonym of the rocks 90 ascribed to the lower Eocene (e.g., Malaroda 1967a; Ungaro 2001). Currently, it is assumed that 91 only the uppermost part of the Thanetian and the lowermost part of the Ypresian are represented in 92 Spilecco (NP 9-10 calcareous nannofossil Zones; Barbieri and Medizza 1969; Papazzoni et al. 93 2014b), whereas the lower Eocene deposits are generally ascribed to the Chiusole Formation (sensu Luciani 1989). This formation was deposited in a basinal-slope setting, heteropically passing to the 94 95 Malcesine Limestones, deposited on the ramp margins (Luciani 1989), and to the neritic facies. The 96 latter has been usually named "Calcari nummulitici" (Nummulitic limestones) (Malaroda 1967b; 97 Carraro et al. 1969; De Zanche et al. 1977; Sarti 1980; Ungaro 2001), but the litho-98 chronostratigraphy of this unit in the Lessinian area still needs revision (e.g., Papazzoni et al. 99 2014c).

100 The Paleogene limestones at the Monte Postale are usually ascribed to the "Calcari nummulitici"

101 (e.g., Malaroda 1967), even if nummulites in them are very rare. More recently, Dal Degan and
102 Barbieri (2005) ascribed the entire succession to the informal "Monte Postale formation" (also
103 described in the text as "Formazione del Monte Postale-Pesciara") and dated it as early-middle

104 Eocene.

105 The alleged "Monte Postale fault" theory reported by Trevisani (2015) is discarded herein (see

106 below) based on field observations (see Vescogni et al. 2016). Therefore, even if the volcanic neck

107 and dykes are cutting the sedimentary succession, we can affirm that the Monte Postale succession

108 is relatively undisturbed and represents a quite continuous stratigraphic record.

# 109 MATERIALS AND METHODS

110 Fieldwork and Sampling

111 Monte Postale Hill was the object of intensive field activity (beginning 2003, focused 2013–2015)

112 aimed to detect outcrops, reconstruct the stratigraphic succession, and collect a large number of

113 samples for paleoenvironmental and biostratigraphic purposes.

114 Unfortunately, outcrops in the study area are typically small and discontinuous due to dense 115 vegetative cover. Moreover, the current strike and dip of the strata does not necessarily reflect 116 original depositional geometries as volcanic intrusions caused localized displacements of the sedimentary beds. Despite this, eight partial stratigraphic sections (Fig. 2) were measured and 117 118 sampled; all are located on the eastern and northern sides of the hill that, unlike the western side, are absent volcanic intrusion. The relatively short distances among the outcrops usually allows 119 120 straightforward correlations. The resulting stratigraphic framework was developed further by correlation with a 20-m core drilled in 2000 by the Natural History Museum of Verona (samples 121 122 CMP on the column in Fig. 3). The core site (shown on Fig. 2) was below active quarry operations 123 for the years 1999–2011.

124 A synthesis of the stratigraphic succession of the Monte Postale including the positions of the 85

samples analyzed is provided (Fig. 3). The total thickness of the sampled sections reaches about 135

126 m, with some uncertainties due to the cover intermissions. Some tens of meters below the base of

127 section E (Fig. 2), the contact between the Paleogene limestones and the underlying Cretaceous

128 Scaglia Rossa can be observed. However, this portion of the succession was not investigated, as the

129 relatively few outcrops did not allow a reliable reconstruction of the stratigraphic architecture.

130 Larger Foraminifera

131 Forty-four samples were studied for their larger foraminiferal content with particular attention given

132 to the species of the genus *Alveolina*, a genus with high biostratigraphic potential and the most

133 abundant in the Monte Postale limestones.

134 On random thin sections  $(4.5 \times 6.0 \text{ cm})$  true axial sections of *Alveolina* were selected for taxonomic

135 identification, then digitally photographed under optical microscope. All the features useful for

136 species identification were measured using the image analysis software JMicroVision 1.2.7. The

137 species of Alveolina were determined following mainly Hottinger (1960, 1974), Drobne (1977), and

138 Scotto di Carlo (1966).

139 The Shallow Benthic Zones (SBZ) were determined according to the species ranges reported in140 Serra-Kiel et al. (1998).

141 Calcareous Nannofossils

Forty-five samples were collected and prepared from unprocessed material as smear slides and 142 143 examined using a light microscope at 1250× magnification. Because most samples contain rare to few calcareous nannofossils a semi-quantitative counting of the number of specimens in a prefixed 144 area of about 8–7 mm<sup>2</sup> (roughly equivalent to three vertical traverses; modified after Gardin and 145 Monechi 1998) was performed. This kind of count allowed an evaluation of the presence or absence 146 147 of index species. The counting of biostratigraphically useful species within a prefixed number of 148 taxonomically related forms (e.g., 20-40 Discoaster; Rio et al. 1990) was performed in the samples where the number of the selected taxon was statistically significant (at least 25–30 specimens). 149 150 The taxonomy here adopted was outlined by Perch-Nielsen (1985). Taxonomic remarks and the list

151 of the calcareous nannofossil considered are reported in the Appendix. The main marker species are

152 represented in Figure 4.

153 THE MONTE POSTALE SECTION

154 Description and Facies Distribution

155 The section begins at one of the major hairpins turns of the Monte Postale trail (Fig. 2, section E),

156 where massive limestones with coral rubble and encrusting biota crop out. Stratification is clearly

157 visible ascending the sidehill (across the woods) for the first 25 m of thickness, then vegetation

158 covers the next  $\sim$  15 m, with a few small outcrops allowing confirmation that the bedding continues

upwards with the same orientation. The outcrop is visible for another  $\sim 8$  m and is then obscured by

160 vegetative cover for another 10 m (to the road). The strata are mostly Alveolina

161 grainstones/packstones that are alternated with laminated limestones with fish and plants and very162 few, thin marly levels.

163 Upwards, again on the road, there are laminated, fine-grained limestones followed by coarser

164 Alveolina-bearing strata alternated with thin marly beds. A basaltic dyke interrupts the exposure, but

165 the observation of some key levels in a core drilled in 2000 (samples CMP on the column, section

166 "core"; Natural History Museum of Verona) allowed us to correlate the latter with the underlying

167 strata, and to verify that, despite the dyke, the vertical displacement of the strata is negligible.

168 The situation in the uppermost part of the section is more complex, because there are several

169 outcrops (sections A, A1, A2, B, C, D, D1) showing significant lateral facies changes from quite

170 evenly bedded fine-grained limestones, locally laminated and bearing fish and plants, to massive

171 limestones rich in corals and algae, representing a bioconstructed facies (sections A1, A, B, C, D1,

172 D) interpreted as the raised threshold separating the open sea from a protected area ("lagoon")

173 (Vescogni et al. 2016).

The contact between these lateral facies is locally sharp and was interpreted as a probable tectonic
contact by Munier Chalmas (1891) and Trevisani (2015). During field observations we did not find

176 any evidence of tectonic activity in this portion of the succession; instead, we observed a gradual,

177 continuous transition between the bedded limestones and the massive limestones some tens of

178 meters far. Therefore, we are inclined to interpret the sharp contact as a local sedimentary onlap.

179 The *Alveolina* Assemblages

180 The faunal composition of the larger foraminiferal (LF) assemblages at Monte Postale is fairly

181 constant (Fig. 3) and, according to the SB zonation of Serra-Kiel et al. (1998), it allows us to ascribe

182 the entire section to the SB 11 Zone (middle Cuisian, or upper part of the Ypresian) since Alveolina

183 cremae (Fig. 5A), A. dainellii, and A. decastroi (Fig. 5B) are limited to this biozone. Alveolina

184 distefanoi, A. fornasinii, and A. rutimeyeri (SBZ 10-11), and A. rugosa (upper SBZ 10-lower 12) are

also fully consistent with the SBZ 11.

186 Some elements of larger foraminiferal assemblages deserve further discussion. *Alveolina* aff.

187 *croatica*, previously recorded from the Pesciara limestones by Papazzoni and Trevisani (2006),

188 differs from *Alveolina croatica* Drobne 1977 for the size of the proloculus, closer to the one of *A*.

189 *levantina*, and for its smaller overall size, with a lower number of whorls. A comprehensive

190 description of this taxon goes beyond the aims of this paper, therefore we simply suggest that it

191 could be a transitional form between A. levantina (upper SBZ 11-SBZ 12) and A. croatica (upper

192 SBZ 12-SBZ 13). According to data presented herein, A. aff. croatica ranges through the whole

193 SBZ 11.

194 Alveolina frumentiformis has been observed only at the top of the investigated section (sample PST

195 1318). This species is characteristic/indicative of the SBZ 12, but the co-occurrence of A. cremae,

196 A. decastroi, and A. distefanoi shall instead confirm the attribution to the SBZ 11, extending its

197 range downward to include the upper part of the SBZ 11.

198 On the other hand, the sample MPO 0301 at base section (Fig. 5D) contains A. cf. schwageri, which

199 together with *A. rugosa* should indicate the SBZ 10. However, this sample was taken from a slump,

so it could contain reworked material. We consider this A. cf. schwageri as an indication that the

201 lower portion of Mt. Postale section probably was deposited close to the base of SBZ 11.

In summary, the Monte Postale section is totally included in the SBZ 11 and probably spans throughmost of it.

204 The *Alveolina* assemblage contains species that were recorded from the same biozone throughout

205 the Paleogene Adriatic Carbonate Platform (including southern Italy; see Hottinger 1960; Scotto di

206 Carlo 1966) as well as in Spain, France, Greece, and Turkey (Drobne et al. 2011).

207 The Calcareous Nannofossil Assemblages

208 Analytical calcareous nannofossil data are reported in Table 1. The results are reported in Figure 3

209 together with the biostratigraphic classification according to the "standard" zonation of Martini

210 (1971) and the zonal scheme of Agnini et al. (2014). Out of 45 samples studied, only 24 of them

211 contain a calcareous nannofossil assemblage useful for biostratigraphic interpretation while 18 are

barren or almost barren (< 10 specimens per 8–7mm<sup>2</sup>). Calcareous nannofossils are generally rare or

scarce in the investigated material, with the exception of four samples (> 100 specimens per 8–7

214 mm<sup>2</sup>) containing common nannofossils. Preservation of calcareous nannofossil assemblage is

215 moderate to poor. Assemblages are usually dominated by placoliths (mainly *Coccolithus pelagicus*,

216 Dictyococcites, and Reticulofenestra) and sphenoliths (manly Sphenolithus moriformis group and

217 Sphenolithus radians). Among placoliths, Cyclicargolithus floridanus and Ericsonia are generally

218 scarce while *Toweius* is rare. The genus *Discoaster* is poorly represented, except for the samples

219 CMP 1422, CMP 800, PST 15105, PST 15102, and PST 1594 (Table 1).

220 The first 60 m of the Monte Postale composite section is characterized by a limestone facies not

suitable for calcareous nannofossil studies, therefore, the nannofossil content of only three marly

samples from the interval between 10–20 m has been analyzed, and it provided poor assemblages.

223 However, the presence of rare Noelaerhabdaceae (*Reticulofenestra*, *Dictyococcites*, and *C*.

224 *floridanus*) and the absence of *Tribrachiatus orthostylus* and *Discoaster sublodoensis* allow the

ascription of this part of the section to the Biozone CNE 5 of Agnini et al. (2014; = Zone NP 13 of

Martini 1971; Fig. 3, Table 1). In particular, the presence of very rare specimens belonging to genus *Toweius* (Prinsiaceae) and *Coccolithus crassus* suggests that this part of the investigated section
could be assigned to the lower part of the Zone CNE 5 of Agnini et al. (2014; Fig. 3, Table 1). In
fact, according to Agnini et al. (2006, 2014), in the uppermost Zone CNE 4 the Prinsiaceae show a
sharp decrease in abundance concomitantly with the first and sporadic entry of members of the
Noelaerhabdaceae family and *C. crassus* has its first occurrence (Bukry 1971; Agnini et al. 2006;
Shamrock and Watkins 2012).

The high-resolution sampling from approximately 60 to 102 m allows a sound biostratigraphic 233 constraint of this part of the M. Postale composite section. In particular, the segment between 60 234 235 and 87 m is characterized by the presence of *Reticulofenestra*, *Dictvococcites*, and *C. crassus*, while the discoasterids which provide important biostratigraphic biohorizons are very rare except in three 236 samples (Fig. 3, Table 1). The count within the genus Discoaster highlights the presence of 237 238 Discoaster lodoensis (10-30%) and of rare specimens of 6-rayed Discoaster (6-rayed Discoaster cf. 239 sublodoensis in Table 1; 0–3%) with intermediate morphologies between D. sublodoensis and D. 240 lodoensis. In order to overcome this taxonomic problem, we considered as D. sublodoensis only the 241 5-rayed morphotypes, in agreement with Agnini et al (2006, 2014). The sporadic presence of T. orthostylus has been attributed to reworking. In fact, the species becomes extinct in correspondence 242 243 of the base common occurrence of Noelaerhabdaceae (Agnini et al. 2006, 2014) which, in the investigated section, are common from the base. The concomitant occurrence of specimens 244 belonging to Noelaerhabdaceae, the virtual absence of Tribrachiatus orthostylus and the absence of 245 246 Discoaster sublodoensis allow to ascribe the segment to the Biozone CNE 5 of Agnini et al. (2014; 247 = Zone NP 13 of Martini 1971; Fig. 3). The segment between 87–102 m contains a few nannofossils 248 except in the sample at the top of this interval (PST 1594), where we registered the sporadic 249 presence of ambiguous specimens of D. sublodoensis (11%; Table 1) which could represents the 250 first rare and discontinuous occurrence of the species in the upper part of Zone CNE 5 as observed

by Agnini et al. (2006, 2014). On this basis, we suggests to assign the top of this interval to the

252 uppermost part of the calcareous nannofossil Zone CNE 5, according to Agnini et al. (2014; Fig. 3).

253 The segment from 102 m to the top of the M. Postale composite section is represented by lithologies

254 not suitable for calcareous nannofossil analysis (massive limestone, biocalciruditic limestone, etc.).

255 The investigated samples are barren/almost barren and so not useful for biostratigraphic

256 interpretation. However, we suggest that this segment could be ascribed to the transition CNE

257 5/CNE 6 Zones of Agnini et al. (2014) (NP13/NP14 Zones of Martini 1971) or to the lowermost

258 part of the CNE 6 Zone based on its stratigraphic position (Fig. 3). The biostratigraphic assignment

259 of the entire M. Postale composite section (lower CNE 5- uppermost CNE 5 or basal CN6) indicates

260 the age of the section is to be late Ypresian, spanning the interval between  $\sim 50.5$  and 48.96 Ma

according to Vandenberghe et al. (2012) and Agnini et al. (2014; Fig. 3).

### 262 INTEGRATED BIOZONATION (SB-CNE ZONES)

263 To date, a firm and integrated calcareous nannofossil and larger foraminifera biostratigraphic

scheme for the Paleogene is not established, even if some correlation schemes (Serra-Kiel et al.

265 1998, fig. 1; Vandenberghe et al. 2012, fig. 28.1) are published and widely used. The different envi-

ronmental settings (basinal and shallow-water, respectively) of these groups often hinder their direct
correlation, there are therefore relatively few successions suitable for a straightforward correlation

268 between calcareous nannofossils and larger foraminifera biozones.

269 The presence of calcareous nannofossils in the Monte Postale section offers the opportunity to test

270 the reliability of Serra Kiel et al. (1998) correlation scheme in the Ypresian. In this section the

271 limestones with *Alveolina* are intercalated with marly beds containing a number of calcareous

272 nannofossils sufficient for biostratigraphic interpretation.

273 Up to now, the larger foraminiferal assemblages from the Monte Postale were studied either as

single samples from unresolved stratigraphic levels (e.g., Hottinger, 1960) or as the bulk fossil

275 content of the stratigraphic succession (Trevisani 2015). Regarding calcareous nannofossils, no

studies have concerned Monte Postale before now. In the present paper we show for the first time
the complete distribution of both alveolines and calcareous nannofossils throughout the succession
(Fig. 3).

The integrated biostratigraphic study performed on the Monte Postale composite section suggests
that the section encompasses almost all the SBZ 11, up to its top; in terms of calcareous
nannofossils we recognize the Zone CNE 5 (= NP 13) and the possible transition to the nannofossil
Zone CNE 6 in the uppermost portion of the section. These results indicate the age of the entire
section is late Ypresian (middle Cuisian) as also suggested by Hottinger (1960). It is worth pointing
out that these results seem consistent with the correlation proposed by Serra-Kiel et al. (1998) and
by Vandenberghe et al. (2012) (Fig. 6).

286 Our findings are in good agreement with the results from the Agost section (Alicante, southeast

287 Spain), where the top of the NP13 (CNE5) Zone occurs within the SBZ11 (Larrasoaña et al. 2008).

288 On the contrary, at Gorrondatxe (Biscay Province, Basque Country, Spain), the top of the NP 13

289 (CN5) Zone is apparently located within the SBZ 12 (Bernaola et al. 2006; Payros et al. 2007;

290 Molina et al. 2011), but the raw data show that only one sample contains larger foraminifera

attributed to the SBZ 12 (see Bernaola et al. 2006, fig. 11; Molina et al. 2011, fig. 7), and this

sample correlates with the CP 12a (= NP 14a) (see Bernaola et al. 2006, fig. 4; Molina et al. 2011,

fig. 5). No samples with larger foraminifera are reported from below, therefore the extension of the

SBZ 12 to the upper part of the NP 13 is not justified by data.

295 The comparison with the neighboring section of Solane (Giusberti et al. 2014) provides additional

296 information about the lower part of the Monte Postale section. At Solane the SBZ 11 was

recognized in the lower part of the section correlated with the CNE 4 (= NP 12) Zone, in good

agreement with the schemes by Serra-Kiel et al. (1998) and Vandenberghe et al. (2012). At Monte

299 Postale, the first sample with nannofossils (PST 1435), approximately 10 m from the base section,

300 positively indicates the lower CNE 5 Zone, therefore we can assume that probably the SBZ 11

should continue below the section measured. Hence, the *A*. cf. *schwageri* identified at the very base
of the section is most probably reworked.

303 One of the still unresolved questions about the Bolca sites is the relative stratigraphic position of the 304 laminated limestones containing the famous "Monte Bolca" ichthyofauna. This name groups 305 together the Pesciara and the Monte Postale laminites, which are apparently very similar and so 306 presumably correlate to each other. Trevisani (2015) concluded that "the Pesciara stratigraphic 307 section is synchronous with the M. Postale 2 section." This assumption is to some extent vague and 308 leaves uncertainties about the correlation of the 15-m thick Pesciara section with the 90-m thick M. Postale 2 section. Since the SBZ 11 is a long-lasting zone, the simple identity of the larger 309 310 foraminiferal assemblage is not sufficient to allow a precise correlation. Medizza (1975), based on a single sample from the uppermost part of the Pesciara (Level L4 in 311 Papazzoni and Trevisani 2006) ascribed this part of the section to the Discoaster sublodoensis (NP 312 14) calcareous nannofossil Zone. This, together with the recognition of the SBZ 11, allowed 313 314 Papazzoni and Trevisani (2006) to restrict the age of the Pesciara laminites to the uppermost part of 315 the SBZ 11 and lowermost NP 14 (= CNE 6) based on the correlation given by Serra-Kiel et al. 316 (1998). The results of this study allow us to ascribe confidently most of Monte Postale succession to the SBZ 11 and to the CNE 5 Zone. However, we cannot exclude, for stratigraphic position, the 317 318 presence of the CNE 6 Zone in the uppermost part of the section (over 105 m; Fig. 3) where the calcareous nannofossil assemblage does not contain biostratigraphically significant taxa. The larger 319 320 for a miniferal assemblages indicate for this segment the upper part of the SBZ 11. This portion of the M. Postale composite section, which is above the "M. Postale 2" section of Trevisani (2015), 321 322 still contains alveolines of the SBZ 11 and for the aforementioned reasons is possibly synchronous 323 with the Pesciara section. Consequently, we suggest that the laminites of the Monte Postale could be slightly older than the ones of the Pesciara. 324

325 According to the calcareous nannofossils biochronology by Agnini et al. (2014), we can estimate

the interval of deposition of the Monte Postale succession between about 50.7 and 48.9 Ma, with a mean depositional rate of at least 6 cm/kyr. This average rate results essentially from two different depositional processes: one very slow and even, accounting for the deposition of the laminated and non-laminated, fine-grained limestones, the other one quite rapid and abrupt, giving rise to the thick coarse-grained *Alveolina* limestones. Given the difficulties in determining quantitatively the relative extent of the two processes, we cannot estimate with confidence their relative importance, therefore the calculated depositional rate has to be considered as purely indicative.

## 333 CONCLUSIONS

334 In this study, we report for the first time the calcareous nannofossil assemblages from the Monte

335 Postale, together with the distribution of the alveolines throughout the section. These data allowed

336 us to obtain a direct correlation of the Shallow Benthic with the Calcareous Nannofossil Zones and

therefore to confirm the general validity of the correlation schemes (Serra-Kiel et al. 1998;

338 Vandenberghe et al. 2012) in this time interval. The comparison of the Monte Postale section with

339 other sections in Italy and Spain reinforces the correlation between the SBZ 11 and the nannofossil

340 zones CNE 4 (upper part), 5 and 6 (lower part) (Fig. 6).

341 These new data are of major importance for the correlation with the Pesciara limestone. For the

342 latter, the attribution to the SBZ 11 and NP 14a (= CNE 6) (Papazzoni and Trevisani 2006) suggests

343 a correlation with the uppermost part of the Monte Postale section.

344 The famous mollusk faunas described by Malaroda (1954) come from the uppermost levels of the

345 Monte Postale; unfortunately, we do not have nannofossil data from here, but based on the larger

346 for aminiferal assemblages we suggest that they belong to the uppermost part of the SBZ 11 or to the

347 SBZ 12, ruling out the Lutetian age.

### 348 ACKNOWLEDGMENTS

349 Many thanks to R. Zorzin (Verona Museum of Natural History) for his assistance in the field and for

350 giving us access to the Monte Postale core samples and to the field data regarding the quarrying

351	campaigns. Thanks to G. Roghi (CNR Padova) for his contribution to the fieldwork and sampling of
352	the Monte Postale section, and to M. Plessi (UniMORE) for his help during work on the Monte
353	Postale core. Warm thanks to the Cerato family for giving us their kind assistance on the field and
354	precious informations about the site. Many thanks to K. Drobne (Ljubljana) and to an anonymous
355	reviewer for their careful reviews which significantly improved the manuscript. This study was
356	funded by the Italian Ministry of Education and Research (MIUR) funds PRIN 2010-2011 (prot.
357	2010X3PP8J_002 and prot. 2010X3PP8J_003) project "Past ExcessCO2 worlds: biota responses to
358	extreme warmth and ocean acidification".
359	REFERENCES
360	Agnini, C., Fornaciari, E., Raffi, I., Catanzariti, R., Pälike, H., Backman, J., and Rio, D.,

2014, Biozonation and biochronology of Paleogene calcareous nannofossils from low and

- 362 middle latitudes: Newsletters on Stratigraphy, v. 47, p. 131–181, doi: 10.1127/0078363 0421/2014/0042.
- 364 AGNINI, C., MUTTONI, G., KENT, D.V., AND RIO, D., 2006, Eocene biostratigraphy and magnetic
- 365 stratigraphy from Possagno, Italy: the calcareous nannofossil response to climate variability:
- 366 Earth and Planetary Science Letters, v. 241, p. 815–830, doi:10.1016/j.epsl.2005.11.005.
- 367 ARNI, P., 1939, Über die Nummuliten und die Gliederung des Untereocaens: Eclogae Geologicae
  368 Helvetiae, v. 32, p. 113–159.
- BARBIERI, G., 1972, Sul significato geologico della faglia di Castelvero (Lessini veronesi): Atti
  dell'Accademia Patavina di Scienze, Lettere e Arti, v. 84, p. 297–302.
- 371 BARBIERI, G., DE ZANCHE, V., MEDIZZA, F., AND SEDEA, R., 1982, Considerazioni sul vulcanesimo
- 372 terziario del Veneto occidentale e del Trentino meridionale: Rendiconti della Società
- 373 Geologica Italiana, v. 4, p. 267–270.

- 374 BARBIERI, G., DE ZANCHE, V., AND SEDEA, R., 1991, Vulcanismo paleogenico ed evoluzione del
- 375 semigraben Alpone-Agno (Monti Lessini): Rendiconti della Società Geologica Italiana, v. 14,

- 376 p. 5–12.
- BARBIERI, G. AND MEDIZZA, F., 1969, Contributo alla conoscenza geologica della regione di Bolca
  (Monti Lessini): Memorie degli Istituti di Geologia e Mineralogia dell'Università di Padova,
  v. 27, p. 1–36.
- BARBIERI, G. AND ZAMPIERI, D., 1992, Deformazioni sinsedimentarie eoceniche con stile a domino
  e relativo campo di paleostress (Monti Lessini): Atti Ticinensi di Scienze della Terra, v. 35, p.

382 25–31.

- BASSI, D., BIANCHINI, G., MIETTO, P., AND NEBELSICK, J.H., 2008, Southern Alps in Italy: Venetian
  Pre-Alps, *in* McCann, T. (ed.), The Geology of Central Europe: The Geological Society of
  London, v. 2, p. 1087–1092.
- 386 BERNAOLA, G., ORUE-ETXEBARRIA, X., PAYROS, A., DINARÈS-TURELL, J., TOSQUELLA, J.,
- 387 APELLANIZ, E., AND CABALLERO, F., 2006, Biomagnetostratigraphic analysis of the
- 388 Gorrondatxe section (Basque Country, Western Pyrenees): its significance for the definition of
- 389the Ypresian/Lutetian boundary stratotype: Neues Jahrbuch für Geologie und Paläontologie

390 Abhandlungen, v. 241, p. 67–109.

- BLACK, M., 1967, New names for some coccolith taxa: Proceedings of the Geological Society of
  London, v. 1640, p. 139–145.
- 393 BOSELLINI, A., 1989, Dynamics of Tethyan carbonate platforms, in Crevello, P.D., Wilson, J.L.,
- Sarg, J.F., and Read, J.R. (eds.), Controls on Carbonate Platform and Basin Platform: SEPM
  Special Publication 44, p. 3–13.
- 396 BOSELLINI, A., MASETTI, D., AND SARTI, M., 1981, A Jurassic "Tongue of Ocean" infilled with
- 397 oolitic sands: the Belluno Trough, Venetian Alps, Italy: Marine Geology, v. 44, p. 59–95.
- BRAMLETTE, M.N. AND MARTINI, E., 1964, The great change in calcareous nannoplankton fossils
  between the Maestrichtian and Danian: Micropaleontology, v. 10, 291–322.
- 400 BRAMLETTE, M.N. AND RIEDEL, W.R., 1954, Stratigraphic value of discoasters and some other

- 401 microfossils related to Recent coccolithophores: Journal of Paleontology, v. 28, p. 385–403.
- BRAMLETTE, M.N. AND SULLIVAN, F.R., 1961, Coccolithophorids and related nannoplankton of the
  Early Tertiary in California: Micropaleontology, v. 7, p. 129–188.
- 404 BRAMLETTE, M.N. AND WILCOXON, J.A., 1967, Middle Tertiary calcareous nannoplankton of the
- 405 Cipero section, Trinidad, W.I:. Tulane Studies in Geology and Paleontology, v. 5, p. 93–131.
- 406 BRÖNNIMANN, P. AND STRADNER, H., 1960, Die Foraminiferen- und Discoasteriden-zonen von
- 407 Kuba und ihre interkontinentale Korrelation: Erdoel-Zeitschrift für Bohr- und Foerdertechnik,
  408 v. 76, p. 364–369.
- BUKRY, D., 1971, Cenozoic calcareous nannofossils from the Pacific Ocean: Transactions of the
  San Diego Society of Natural History, v. 16, p. 303–327.
- 411 BUKRY, D., 1973, Coccolith stratigraphy, eastern equatorial Pacific, Leg 16, Deep Sea Drilling
- 412 Project, *in* Van Andel, T.H., Heath, G.R., et al. (eds.), Initial Reports of the DSDP: Deep Sea
  413 Drilling Project, Washington (U.S. Government Printing Office), v. 16, p. 653–711.
- 414 doi:10.2973/dsdp.proc.16.126.1973
- 415 CANDE, S.C. AND KENT, D.V., 1995, Revised calibration of the geomagnetic polarity time scale for
- 416 the Late Cretaceous and Cenozoic: Journal of Geophysical Research, v. 100, p. 6093–6096.
- 417 CARMINATI, E., LUSTRINO, M., AND DOGLIONI, C., 2012, Geodynamic evolution of the central and
- 418 western Mediterranean: tectonics vs. igneous petrology constraints: Tectonophysics, v. 579, p.
  419 173–192.
- 420 CARNEVALE, G., BANNIKOV, A.F., MARRAMÀ, G., TYLER, J.C., AND ZORZIN, R., 2014, The
- 421 Pesciara-Monte Postale Fossil-Lagerstätte: 2. Fishes and other vertebrates, *in* Papazzoni,
- 422 C.A., Giusberti, L., Carnevale, G., Roghi, G., Bassi, D., and Zorzin, R. (eds.), The Bolca
- 423 Fossil-Lagerstätten: A Window into the Eocene World: Rendiconti della Società
- 424 Paleontologica Italiana, v. 4, p. 37–63.
- 425 CARRARO, F., MALARODA, R., PICCOLI, G., STURANI, C., AND VENZO, S., 1969, Note Illustrative

- 426 della Carta Geologica D'Italia, Scala 1:100.000: Foglio 48, Peschiera del Garda, p. 1–95.
- 427 CHECCHIA-RISPOLI, G., 1905, Sopra alcune alveoline eoceniche della Sicilia: Palaeontographia
  428 Italica, v. 11, p. 147–167.
- 429 CHECCHIA-RISPOLI, G., 1909, Nuova contribuzione alla conoscenza delle alveoline eoceniche della
  430 Sicilia: Palaeontographia Italica, v. 15, p. 59–70.
- 431 DAL DEGAN, D. AND BARBIERI, S., 2005, Rilievo geologico dell'area di Bolca (Monti Lessini
- 432 orientali): Bollettino del Museo Civico di Storia Naturale di Verona, v. 27, p. 3–10.
- 433 DEFLANDRE, G., 1947, Braarudosphaera nov. gen., type d'une famille nouvelle de
- 434 Coccolithophorides actuels a elements composites: Comptes Rendus Hebdomadaires des
- 435 Séances de l'Académie des Sciences. Paris, v. 225, p. 439–441.
- 436 DEFLANDRE, G., 1952, Classe des Coccolithophoridés. (Coccolithophoridae. Lohmann, 1902, *in*
- 437 Grassé, P.P. (ed.), Traite de Zoologie: Masson, Paris, p. 439–470.
- 438 DEFLANDRE, G., 1959, Sur les nannofossiles calcaires et leur systématique: Revue de
- 439 Micropaléontologie, v. 2, p. 127–152.
- DEFLANDRE, G. AND FERT, C., 1954, Observations sur les coccolithophoridés actuels et fossiles en
  microscopie ordinaire et électronique: Annales de Paléontologie, v. 40, p. 115–176.
- 442 DE ZANCHE, V., SORBINI, L., AND SPAGNA, V., 1977, Geologia del territorio del Comune di Verona:
- 443 Memorie del Museo Civico di Storia Naturale di Verona, v. 1, p. 1–52.
- 444 DOGLIONI, C. AND BOSELLINI, A., 1987, Eoalpine and mesoalpine tectonics in the Southern Alps:
  445 Geologische Rundschau, v. 77, 734–754.
- 446 DROBNE, K., 1977, Alvéolines paléogènes de la Slovénie et de l'Istrie: Schweizerische
- 447 Paläontologische Abhandlungen, v. 99, p. 1–175.
- 448 DROBNE, K., ĆOSOVIĆ, V., MORO, A., AND BUCKOVIĆ, D., 2011, The role of the Palaeogene
- 449 Adriatic Carbonate Platform in the Spatial Distribution of Alveolinids: Turkish Journal of
- 450 Earth Sciences, v. 20, p. 721–751.

- FABIANI, R., 1914, La serie stratigrafica del Monte Bolca e dei suoi dintorni: Memorie dell'Istituto
  Geologico della regia Università di Padova, v. 2, p. 223–235.
- 453 FABIANI, R., 1915, Il Paleogene del Veneto: Memorie dell'Istituto Geologico della regia Università
  454 di Padova, v. 3, p. 1–336.
- GARDIN, S. AND MONECHI, S., 1998, Palaeoecological change in middle to low latitude calcareous
  nannoplankton at the Cretaceous/Tertiary boundary: Bulletin de la Société géologique de
  France, v. 169, p. 709–723.
- 458 GIUSBERTI, L., BANNIKOV, A., BOSCOLO GALAZZO, F., FORNACIARI, E., FRIELING, J., LUCIANI, V.,
- 459 PAPAZZONI, C.A., ROGHI, G., SCHOUTEN, S., SLUIJS, A., BOSELLINI, F.R., AND ZORZIN, R.,
- 460 2014, A new *Fossil-Lagerstätte* from the lower Eocene of Lessini Mountains (northern Italy):
- 461 a multidisciplinary approach: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 403, p.
  462 1–15.
- GRAN, H.H. AND BRAARUD, T., 1935, A quantitative study of the phytoplankton in the Bay of
  Fundy and the Gulf of Maine (including observations on hydrography, chemistry and
  turbidity): Journal of the Biological Board of Canada, v. 1, p. 279–467.
- 466 HAQ, B.U., 1971, Paleogene calcareous nannoflora, part IV. Paleogene nanno-plankton
- 467 biostratigraphy and evolutionary rates in Cenozoic calcareous nannoplankton: Stockholm
  468 Contribution in Geology, v. 25, p. 129–158.
- HAY, W.W. AND MOHLER, H.P., 1967, Calcareous nannoplankton from Early Tertiary rocks at Point
  Labau, France and Paleocene-Early Eocene correlations: Journal of Paleontology, v. 41, p.
  1505–1541.
- 472 HAY, W.W., MOHLER, H.P.; ROTH, P.H., SCHMIDT, R.R., AND BOUDREAUX, J.E., 1967, Calcareous
- 473 nannoplankton zonation of the Cenozoic of the Gulf Coast and Caribbean-Antillean area, and
- 474 transoceanic correlation: Transactions of the Gulf Coast Association of Geological Societies,
- 475 v. 17, p. 428–480.

- 476 HAY, W.W., MOHLER, H.P., AND WADE, M.E., 1966, Calcareous nannofossils from Nal'chik
- 477 (northwest Caucasus): Eclogae Geologicae Helvetiae, v. 59, p. 379–399.
- HAY, W.W. AND TOWE, K.M., 1962, Electron microscope examination of some coccoliths from
  Donzacq (France): Eclogae Geologicae Helvetiae, v. 55, p. 497–517.
- 480 HOTTINGER, L., 1960, Recherches sur les Alvéolines du Paléocène et de l'Eocène: Schweizerische
- 481 Paläontologische Abhandlungen, v. 75-76 Text (I), p. 1–243, atlas (II), 18 pls.
- 482 HOTTINGER, L., 1974, Alveolinids, Cretaceous–Tertiary larger Foraminifera: Esso Production
- 483 Research-European Laboratories, p. 1–84.
- 484 KAMPTNER, E., 1954, Untersuchungen über den Feinbau der Coccolithen. Anzeiger:
- 485 Österreichische Akademie der Wissenschaften. Mathematische-Naturwissenschafliche Klasse,
  486 v. 87, p. 152–158.
- 487 KAMPTNER, E., 1963, Coccolithineen-Skelettreste aus Tiefseeablagerungen des Pazifischen Ozeans:
  488 Annalen des Naturhistorischen Museums in Wien, v. 66, p. 139–204.
- 489 LARRASOAÑA, J.C., GONZALVO, C., MOLINA, E., MONECHI, S., ORTIZ, S., TORI, F., AND
- 490 TOSQUELLA, J., 2008, Integrated magnetobiochronology of the early/middle Eocene transition
- 491 at Agost (Spain): implications for defining the Ypresian/Lutetian boundary stratotype: Lethaia,
- 492 v. 41, p. 395–415, doi: 10.1111/j.1502-3931.2008.00096.x.
- 493 LOCKER, S., 1968, Biostratigraphiedes des Alttertiars von Norddeutschland mit Coccolithophoriden:
  494 Monatsberichte der Deutschen Akademie der Wissenschaften zu Berlin, v. 10, p. 220–229.
- 495 LUCIANI, V., 1989, Stratigrafia sequenziale del Terziario nella Catena del Monte Baldo (provincie di
- 496 Verona e Trento): Memorie di Scienze Geologiche, v. 41, p. 263–351.
- 497 MALARODA, R., 1954, Il Luteziano di Monte Postale (Lessini medî): Memorie degli Istituti di
- 498 Geologia e Mineralogia dell'Università di Padova, v. 19 (1955-1956), p. 3–107.
- 499 MALARODA, R., 1967a, Calcari di Spilecco, in Bosellini, A., Carraro, F., Corsi, M., De Vecchi, G.P.,
- 500 Gatto, G.O., Malaroda, R., Sturani, C., Ungaro, S., and Zanettin, B. (eds.), Note illustrative

- della carta geologica d'Italia scala 1: 100.000, foglio 49 Verona: Servizio Geologico d'Italia,
  Nuova Tecnica Grafica, Roma, p. 19–21.
- 503 MALARODA, R., 1967b, Calcari nummulitici, in Bosellini, A., Carraro, F., Corsi, M., De Vecchi,
- 504 G.P., Gatto, G.O., Malaroda, R., Sturani, C., Ungaro, S., and Zanettin, B. (eds.), Note
- 505 illustrative della carta geologica d'Italia scala 1: 100.000, foglio 49 Verona: Servizio
- 506 Geologico d'Italia, Nuova Tecnica Grafica, Roma, p. 21–22.
- 507 MARTINI, E., 1958, Discoasteriden und verwandte Formen im NW-deutschen Eozän
- 508 (Coccolithophorida). 1. Taxonomische Untersuchungen: Senckenbergiana lethaea, v. 39, p.
  509 353–388.
- 510 MARTINI, E., 1971, Standard Tertiary and Quaternary calcareous nannoplankton zonation, in
- 511 Farinacci, A. (ed.), Proceedings 2nd International Conference Planktonic Microfossils Roma:
  512 Rome, v. 2, p. 739–785.
- 513 MÁRTON, E., ZAMPIERI, D., KÁZMÉR, M., DUNKL, I., AND FRISCH, W., 2011, New Paleocene-
- Eocene paleomagnetic results from the foreland of the Southern Alps confirm decoupling of
- 515 stable Adria from the African plate: Tectonophysics, v. 504, p. 89–99.
- MEDIZZA, F., 1975, Il nannoplancton calcareo della Pesciara di Bolca (Monti Lessini): Studi e
  Ricerche sui Giacimenti Terziari di Bolca, v. 2, p. 433–444.
- 518 MOLINA, E., ALEGRET, L., APELLANIZ, E., BERNAOLA, G., CABALLERO, F., DINARÈS-TURELL, J.,
- 519 HARDENBOL, J., HEILMANN-CLAUSEN, C., LARRASOAÑA J.C., LUTERBACHER, H., MONECHI,
- 520 S., ORTIZ, S., ORUE- ETXEBARRIA, X., PAYROS, A., PUJALTE, V., RODRÍGUEZ-TOVAR F.J.,
- 521 TORI, F., TOSQUELLA, J., AND UCHMAN, A., 2011, The Global Stratotype Section and Point
- 522 (GSSP) for the base of the Lutetian Stage at the Gorrondatxe section, Spain: Episodes, v. 34,
  523 p. 86–108.
- MUNIER-CHALMAS, M., 1891, Étude du Tithonique, du Crétacé et du Tertiaire du Vicentin: Série
  Stratigraphique vol. 1, Savy, Paris, 213 p.

526	PAPAZZONI, C.A., BASSI, D., FORNACIARI, E., GIUSBERTI, L., LUCIANI, V., MIETTO, P., ROGHI, G.
527	AND TREVISANI, E., 2014c, 3. Geological and stratigraphical setting of the Bolca area, in
528	Papazzoni, C.A., Giusberti, L., Carnevale, G., Roghi, G., Bassi, D., and Zorzin, R. (eds.), The
529	Bolca Fossil-Lagerstätten: A window into the Eocene World: Rendiconti della Società
530	Paleontologica Italiana, v. 4, p. 19–28.
531	PAPAZZONI, C.A., GIUSBERTI, L., CARNEVALE, G., ROGHI, G., BASSI, D., AND ZORZIN, R. (eds.),
532	2014a, The Bolca Fossil-Lagerstätten: A window into the Eocene world: Rendiconti della
533	Società Paleontologica Italiana, v. 4, p. 19–28.
534	PAPAZZONI, C.A., GIUSBERTI, L., AND TREVISANI, E., 2014b, 10. The Spilecco site, in Papazzoni,
535	C.A., Giusberti, L., Carnevale, G., Roghi, G., Bassi, D., and Zorzin, R. (eds.), The Bolca
536	Fossil-Lagerstätten: A window into the Eocene World: Rendiconti della Società
537	Paleontologica Italiana, v. 4, p. 105–110.
538	PAPAZZONI, C.A. AND TREVISANI, E., 2006, Facies analysis, palaeoenvironmental reconstruction,
539	and biostratigraphy of the "Pesciara di Bolca" (Verona, northern Italy): an early Eocene
540	Fossil-Lagerstätte: Palaeogeography, Palaeoclimatology, Palaeoecology, v. 242, p. 21-35,
541	doi: 10.1016/j.palaeo.2006.05.01.
542	PAPAZZONI, C.A. AND TREVISANI, E., 2009, Relationships between the Pesciara di Bolca and the
543	Monte Postale Fossil-Lagerstätten (Lessini Mts., northern Italy), in Billon-Bruyat, J.P., Marty,
544	D., Costeur, L., Meyer, C.A., and Thuring, B. (eds.), Fifth International Symposium on
545	Lithographic Limestone and Plattenkalk, Naturhistorisches Museum Basel, Switzerland:
546	Actes 2009 bis de la Société jurassienne d'Emulation (Porrentruy), Basel, p. 65-66.
547	PAYROS, A., BERNAOLA, G., ORUE-ETXEBARRIA, X., DINARÈS-TURELL, J., TOSQUELLA, J., AND
548	APELLANIZ, E., 2007, Reassessment of the early-middle Eocene biomagnetochronology based
549	on evidence from the Gorrondatxe section (Basque Country, western Pyrenees): Lethaia, v.
550	40, p. 183–195, doi: 10.1111/j.1502-3931.2007.00016.x.

551 PERCH-NIELSEN, K., 1969, Die Coccolithen einiger dänischer Maastrichtien- und Danien

552 Lokalitäten: Meddelelser fra Dansk Geologisk Forening, v. 19, p. 51–68.

553 PERCH-NIELSEN, K., 1985. Cenozoic calcareous nannofossils, in Bolli, H.M., Saunders, J.B, and

- 554 Perch-Nielsen, K. (eds.), Plankton Stratigraphy: Cambridge University Press, Cambridge, p.
  555 427–554.
- RENZ, O., 1936, Stratigraphische und mikropalaeontologische Untersuchung der Scaglia (Obere
  Kreide-Tertiär) im zentralen Apennin: Eclogae Geologicae Helvetiae, v. 29, p. 1–149.

558 RIO, D., RAFFI, I., AND VILLA, G., 1990, Pliocene–Pleistocene calcareous nannofossil distribution

patterns in the western Mediterranean, in Kastens, K.A., Mascle, J., et al., Proceedings ODP,

- 560 Scientific Results 107 (Ocean Drilling Program, College Station, TX), p. 513–533, doi:
- 561 10.2973/ odp.proc.sr.107.164.1990.
- 562 SARTI, M., 1980, Frane sottomarine e debris flow in una successione carbonatico torbiditica
- 563 eocenica (Val d'Avesa, Verona): Annali dell'Università di Ferrara, Sez. IX Scienze Geologiche
  564 e Paleontologiche, v. 7, p. 65–89.
- 565 SCHILLER, J., 1930, Coccolithineae, *in* Rabenhorst, L., (ed.), Kryptogamen-Flora von Deutschland,

566Österreich und der Schweiz: Akademische Verlagsgesellschaft, Leipzig, p. 89–267.

- 567 SCOTTO DI CARLO, B., 1966, Le Alveoline del Gargano nord-orientale: Palaeontographia Italica, v.
  568 61, p. 65–73.
- 569 SERRA-KIEL, J., HOTTINGER, L., CAUS, E., DROBNE, K., FERRÀNDEZ, C., JAUHRI, A.K., LESS, G.,

570 PAVLOVEC, R., PIGNATTI, J., SAMSÓ, J.M., SCHAUB, H., SIREL, E., STROUGO, A.,

- 571 TAMBAREAU, Y., TOSQUELLA, J., AND ZAKREVSKAYA, E., 1998, Larger foraminiferal
- 572 biostratigraphy of the Tethyan Paleocene and Eocene: Bulletin de la Société Géologique de
- 573 France, v. 169, p. 281–299.
- 574 SHAMRAI, I.A., 1963, Nekotorye formy verkhnemelovykh i paleogenovykh kokkolitov i
- 575 diskoasterov na yuge russkol platformy [Certain forms of Upper Cretaceous and Paleogene

- 576 coccoliths and discoasters from the southern Russian Platform]: Izvestiya Vysshikh Uchebnykh
- 577 Zavedenii, Geologiya i Razvedka, v. 6, p. 27–40.
- SHAMROCK, J.L. AND WATKINS, D.K., 2012, Eocene calcareous nannofossil biostratigraphy and
   community structure from Exmouth Plateau, Eastern Indian Ocean (ODP Site 762):
- 580 Stratigraphy, v. 9, p. 1–54.
- 581 SORBINI, L., 1972, I fossili di Bolca: Edizioni Corev, Verona, 134 p.
- 582 SORBINI, L., 1999, I giacimenti di Bolca, in Pinna, G. (ed.), Alle radici della storia naturale
- 583 d'Europa. Seicento milioni di anni attraverso i grandi giacimenti paleontologici: Jaca Book,
  584 Milano, p. 172–176.
- 585 STRADNER, H., 1959, Die fossilen Discoasteriden Osterreichs. II: Erdöl-Zeitschrift, v. 75, p. 472–
  586 488.
- 587 STRADNER, H., 1961, Vorkommen von Nannofossilien im Mesozoikum und Alttertiär: Erdöl588 Zeitschrift für Bohr-und Fördertechnik, v. 77, p. 77–88.
- 589 SULLIVAN, F.R., 1965, Lower Tertiary nannoplankton from the California Coast Ranges. II. Eocene:
- 590 University of California Publications in Geological Sciences, v. 53 p. 1–74
- 591 TAN, S.H., 1927, Over de samenstelling en het onstaan van krijt en mergelgesteenten van de
- 592 Molukken: Jaarboek van het Mijnwezen in Nederlandsch-Indie, v. 55, p. 111–122.
- 593 TREVISANI, E., 2015, Upper Cretaceous–lower Eocene succession of the Monte Postale and its
- relationship with the "Pesciara di Bolca" (Lessini Mountains, northern Italy): deposition of a
- 595 fossil-fish *lagerstätte*: Facies, v. 61, p. 1–17, doi: 10.1007/s10347-015-0431-y.
- 596
- 597 UNGARO, S., 2001, Le biofacies paleoceniche ed eoceniche dei Monti Lessini (Veneto, Italia):
  598 Annali dell'Università di Ferrara, Sez. Scienze della Terra, v. 9, p. 1–40.
- 599 VANDENBERGHE, N., HILGEN, F.J., AND SPEIJER, R.P., 2012, The Paleogene Period, in Gradstein,
- 600 F.M., Ogg, J.G., Smith, A.G., and Ogg, G.M. (eds.), The Geologic Time Scale 2012: Elsevier,

- 601 Amsterdam, p. 855–921.
- 602 VAROL, O., 1989, Eocene calcareous nannofossils from Sile (northwest Turkey): Revista Española
  603 de Micropaleontología, v. 21, p. 273–320.
- 604 VESCOGNI, A., BOSELLINI, F.R., PAPAZZONI, C.A., GIUSBERTI, L., ROGHI, G., FORNACIARI, E.,
- 605 DOMINICI, S., AND ZORZIN, R., 2016, Coralgal buildups associated with the Bolca Fossil-
- 606 *Lagerstätten*: new evidences from the Ypresian of Monte Postale (NE Italy): Facies, v. 62: 21,
- 607 p. 1–20. doi: 10.1007/s10347-016-0472-x.
- WALLICH, G.C., 1877, Observations on the coccosphere: Annals and Magazine of Natural History,
  v. 19, p. 342–350.
- 610 WINTERER, E.L. AND BOSELLINI, A., 1981, Subsidence and sedimentation on Jurassic passive
- 611 continental margin, Southern Alps, Italy: AAPG Bulletin, v. 65, p. 394–421.
- 612 ZAMPIERI, D., 1995, Tertiary extension in the southern Trento Platform, Southern Alps, Italy:
  613 Tectonics, v. 14, p. 645–657.
- 614 Received 1 February 2016; accepted 15 July 2016.
- 615 FIGURE CAPTIONS
- 616 FIG. 1.—Location map of the Monte Postale section.
- 617 FIG. 2.—Position of the sections surveyed and sampled, with the relative stratigraphic columns.
- 618 FIG. 3.—Larger foraminifera and calcareous nannofossil biostratigraphy distribution along the
- 619 composite section. The range of selected calcareous nannofossil and larger foraminifera
- 620 (alveolinids) taxa is reported. Abundances of calcareous nannofossils are expressed as number of
- 621 specimens per 7-8 mm<sup>2</sup>, corresponding to three vertical traverses. Light gray areas represent barren
- 622 or almost barren samples for nannofossil assemblages. The presence of larger foraminifera is
- 623 represented by the black rectangles; the gray triangles represent reworked taxa. The star represents
- 624 the presence of D. cf. sublodoensis (11%) within the Discoaster. Columns: A = Thickness (m); B =
- 625 Chronostratigraphy; C = Larger foraminiferal SB Zones (Serra-Kiel et al. 1998); D = Calcareous

- 626 nannofossil zones (Martini 1971); E = Calcareous nannofossil zones (Agnini et al., 2014); F =
- 627 Simplified lithologies and relative position of the samples.
- 628 FIG. 4.—Photomicrographs of selected calcareous nannofossil taxa from the Monte Postale section.
- 629 Scale bar 5 μm. A–Db) Coccolithus crassus Bramlette and Sullivan 1961. Crossed nicols. A)
- 630 Sample CMP 1422. B) Sample MPO P1. C) Sample PST 1594. Db) Sample CMP 1422. Da)
- 631 Coccolithus pelagicus (Wallich 1877) Schiller 1930. Crossed nicols. Sample CMP 1422. E)
- 632 Ericsonia Black 1964. Crossed nicols. Sample PST 15105. F, G) Reticulofenestra Hay et al. 1966.
- 633 F) Sample MPO P1. G) Sample PST 15105. H, I) Dictyococcites Black 1967. Sample PST 15105.
- 634 J) Ericsonia formosa (Kamptner 1963) Haq 1971. Crossed nicols. Sample PST 15105. K)
- 635 *Cyclicargolithus floridanus* (Roth and Hay *in* Hay et al. 1967) Bukry 1971. Sample MPO P1. L)
- 636 *Girgisia gammation* (Bramlette and Sullivan 1961) Varol 1989. Crossed nicols. Sample PST 15105.
- 637 M) Zygrhablithus bijugatus (Deflandre in Deflandre and Fert 1954) Deflandre 1959. Crossed
- 638 nicols. Sample PST 15105. N, O) Sphenolithus radians Deflandre in Grassé 1952. N) Crossed
- 639 nicols 0°. O) nicols 45°. Sample PST 15105. P–R) *Discoaster lodoensis* Bramlette and Riedel 1954.
- 640 Parallel light. P, Q) Sample CMP 1422, same specimen different focus. R) Sample CMP 1422. S)
- 641 *Discoaster* cf. *sublodoensis* Bramlette and Sullivan 1961. Parallel light. Sample PST 1594. T)
- 642 Discoaster kuepperi Stradner 1959. Parallel light. Sample PST 15105. U) Discoaster nodifer
- 643 (Bramlette and Riedel 1954) Bukry 1973. Parallel light. Sample PST 15105. V) Discoaster
- 644 *barbadiensis* Tan 1927. Parallel light. Sample PST 15105. W) *Braarudosphaera* Deflandre 1947.
- 645 Crossed nicols. Sample PST 15105. X, Y) Blackites Hay and Towe 1962. Crossed nicols. Y) Base
- 646 of *Blackites*. Sample PST 15105.
- 647 FIG. 5.—Photomicrographs of selected species of alveolines from the Monte Postale section. A)
- 648 Alveolina cremae Checchia-Rispoli 1905. Sample CMP 1970. B) Alveolina decastroi Scotto di
- 649 Carlo 1966. Sample CMP 1970. C) Alveolina fornasinii Checchia-Rispoli 1909. Sample MPO
- 650 0306. D) Alveolina cf. schwageri Checchia-Rispoli 1905. Sample MPO 0301. E) Glomalveolina

- 651 minutula (Reichel in Renz 1936). Sample PST 1434. F) Alveolina rugosa Hottinger 1960. Sample
- 652 MPO 0307. Scale bar = 1 mm.
- 653 FIG. 6.—Biomagnetostratigraphy of the Ypresian–Lutetian interval. Geomagnetic polarity scale
- after Cande and Kent (1995). Tethyan zonation of larger benthic foraminifera from Serra-Kiel et al.
- (1998). Calcareous nannofossil zonations from Martini (1971; NP Zones) and Agnini et al (2014;
- 656 CNE zones). The stratigraphic positions of Monte Postale section is indicated by the gray bar.
- 657 TABLE CAPTION
- 658 TABLE 1.—Distribution of the calcareous nannofossils in the Monte Postale composite section.