

This is a pre print version of the following article:

The posterior cingulate BOLD duty cycle during meditation predicts attentional skills / Pagnoni, Giuseppe.  
- (2011), pp. 39-39.

*Terms of use:*

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

31/12/2025 14:21

# The posterior cingulate BOLD duty cycle during meditation predicts attentional skills

## Submission No:

4233

## Authors:

Giuseppe Pagnoni<sup>1</sup>

## Institutions:

<sup>1</sup>University of Modena and Reggio Emilia, Modena, Italy

## Introduction:

Recent findings about the human brain's wakeful resting state suggest a link between the spontaneous generation of task-unrelated thoughts (mind-wandering)<sup>1,2</sup>, and the slow fluctuations of activity in the 'default mode network' (DMN)<sup>3</sup>, a consistent set of brain regions with a major hub in the posterior cingulate cortex (PCC)<sup>4</sup>. Meta-awareness and regulation of mind-wandering are core cognitive components of many meditation practices, and to study their relationship to the DMN activity we collected fMRI data from a cohort of experienced Zen meditators and meditation-naive controls engaging in a simple meditative protocol, along with their performance on a computerized sustained attention task. By introducing a simple duty cycle measure for the fMRI BOLD signal from the PCC, we hypothesized that this could be taken as an endophenotype<sup>5</sup> of the individual capacity to regulate mind-wandering and would thus correlate with performance in the sustained attention task.

## Methods:

**Subjects:** 12 Zen meditators with > 3 years of daily practice and 12 control subjects, matched for age and education level.

**Meditative task:** subjects were asked to maintain their attention on their breathing throughout the scan and gently redirect attention to breathing every time they found themselves distracted or mind-wandering.

**MRI acquisition:** a single series of gradient-echo EPI volumes (200 scans, TR=2.35s, 35 3x3x3mm axial slices) and a T1-weighted hi-res volume (MPRAGE, 176 1x1x1 mm sagittal slices) were acquired on a 3.0 Tesla Siemens Magnetom Trio scanner.

**Imaging analysis:** The EPI series were corrected for slice-timing and head motion, and warped to MNI space using the T1-weighted image to estimate the warping parameters. The time series were then band-pass filtered (0.01-0.1 Hz), along with least-square removal of the global signal and estimated motion parameters. The average processed BOLD time courses from a PCC region of interest (center-of-mass MNI coordinates: -6, -56, 22; size=82 voxels), identified by a previous study on the same subject sample using a conceptual processing task<sup>6</sup>, were extracted and a duty cycle measure was computed as the ratio of the cumulative time that the signal lay above its temporal mean and the total scanning time.

**Sustained attention task:** each subject completed the Rapid Visual Information Processing (RVIP) task from the CANTAB neuropsychological computerized battery<sup>7</sup>. Performance was assessed in terms of reaction times (RT) and A-prime, a nonparametric sensitivity index ranging from 0 to 1 based on the number of hits and false alarms (1=perfect performance).

**Correlation analysis:** Pearson's correlation tests were computed between individual scores in the RVIP task (RT and A-prime) and the PCC duty cycle values.

Image processing and statistics were performed with the software packages AFNI, SPM5, and R.


## Results:

We observed a significant correlation between the values of the PCC BOLD duty cycle during the meditative procedure and the individual scores in the RVIP task (A-prime:  $r=-0.64$ ,  $p=0.0007$ ; RT:  $r=+0.58$ ,  $p=0.003$ ): subjects with a lower PCC duty cycle tended to respond faster and more accurately on the sustained attention task (see Figure).

## Conclusions:

Although these findings need to be confirmed by a larger study, they support the notion that individual capacity for top-down attentional control and resistance to mind-wandering is inversely related to the relative abundance of high levels of activity of the central DMN node, the PCC, when such activity is mobilized by the processes of meta-awareness and regulation of spontaneous

mentation that characterize the meditative exercise. The simple time-domain measure of the PCC BOLD activity proposed here, i.e. its duty cycle while subjects attempt to keep their attention on their breathing, appears as a promising candidate for an endophenotype of individual attentional skills.



## Higher Cognitive Functions

### Executive Function

### Abstract Information

### References

1. Mason M.F. (2007) 'Wandering minds: the default network and stimulus-independent thought', *Science*, no. 315, pp. 393-395.
2. Christoff K. (2009) 'Experience sampling during fMRI reveals default network and executive system contributions to mind wandering', *Proceedings of the National Academy of Sciences USA*, no. 21, pp. 8719-8724.
3. Raichle M.E. (1991) 'A default mode of brain function', *Proceedings of the National Academy of Sciences USA*, no. 98, pp. 676-82.
4. Hagmann P. (2008) 'Mapping the Structural Core of Human Cerebral Cortex', *PLoS Biology*, no. 6, pp. e159.
5. Castellanos F.X. (2002) 'Neuroscience of attention-deficit/hyperactivity disorder: the search for endophenotypes', *Nature Reviews Neuroscience*, no. 3, pp. 617-28.
6. Pagnoni G. (2008) '"Thinking about not-thinking": neural correlates of conceptual processing during Zen meditation', *PLoS ONE*, no. 3, pp. e3083.
7. Sahakian B. J. (1992) 'Computerized assessment in neuropsychiatry using CANTAB: discussion paper', *Journal of the Royal Society of Medicine*, no. 85, pp. 399-402.