

This is the peer reviewed version of the following article:

Analytical expression of genuine tripartite quantum discord for symmetrical X-states / Beggi, Andrea; Buscemi, Fabrizio; Bordone, Paolo. - In: QUANTUM INFORMATION PROCESSING. - ISSN 1570-0755. - STAMPA. - 14:2(2015), pp. 573-592. [10.1007/s11128-014-0882-z]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

22/09/2024 06:21

(Article begins on next page)

Analytical Expression of Genuine Tripartite Quantum Discord for Symmetrical X-states

Andrea Beggi · Fabrizio Buscemi · Paolo Bordone

Received: date / Accepted: date

Abstract The study of classical and quantum correlations in bipartite and multipartite systems is crucial for the development of quantum information theory. Among the quantifiers adopted in tripartite systems, the genuine tripartite quantum discord (GTQD), estimating the amount of quantum correlations shared among all the subsystems, plays a key role since it represents the natural extension of quantum discord used in bipartite systems. In this paper, we derive an analytical expression of GTQD for three-qubit systems characterized by a subclass of symmetrical X-states. Our approach has been tested on both GHZ and maximally mixed states reproducing the expected results. Furthermore, we believe that the procedure here developed constitutes a valid guideline to investigate quantum correlations in form of discord in more general multipartite systems.

Keywords Quantum Discord · Analytic expressions · Genuine correlations · X states · Tripartite systems

PACS 03.67.-a · 03.65.Ud · 03.67.Mn

1 INTRODUCTION

Quantum correlations are assuming increasing relevance, since they can be exploited to improve our ability to perform many informational and computational tasks [1, 2, 3, 4]. Therefore, the problem of their characterization and quantification has become a significant topic of research. Traditionally, the most used form of quantum correlation is *entanglement*, and the development of quantum information theory is fundamentally due to its implementation in information and communication protocols [5, 6].

A form of quantum correlation other than entanglement is *quantum discord* (QD) [7, 8, 9], which can be expressed in terms of the difference between the total and the classical correlations for a system when one of its subsystems is subject to an unobserved measure process. Such a quantity, however, significantly depends upon both the subsystem chosen and the measurement performed on it: in particular, if the measurement is carefully selected, we can minimize its “disturbing effect” on the system [10]. This choice corresponds to the minimization of QD firstly over a set of possible measurements (on a fixed subsystem), typically *projective von Neumann measurements* [10, 7], and secondly over all possible subsystem on which the local measurement can be performed. Recent efforts in the study of the optimization processes have led to analytical expression for quantum discord in some particular [9] and more general states [11, 12] in systems composed of two qubits.

Both entanglement and QD have widely been analyzed and used in bipartite systems, while their extension to multipartite systems is still discussed and tackled with different approaches [13, 14, 15, 16, 17]. For instance, *Vinjanampathy et al.* [18] proposed a method to evaluate analytically quantum discord for a n -partite system of qubits

Andrea Beggi · Fabrizio Buscemi · Paolo Bordone
Dipartimento di Scienze Fisiche, Informatiche e Matematiche,
Università di Modena e Reggio Emilia,
Via Campi 213/A, I-41125 Modena, Italy,
E-mail: andrea.beggi@unimore.it

Paolo Bordone
Centro S3, CNR - Istituto Nanoscienze,
Via Campi 213/A, I-41125 Modena, Italy,

in some special cases, but they treated the whole system as a bipartite one (each subparty containing 1 or $n - 1$ qubits, respectively). On the other hand, *Giorgi et al.* [17] defined, for a n -partite system, *genuine n -partite correlations*, which can be divided into total, classical or quantum. These kinds of correlations are shared between all the n parties which form the system, i.e. they cannot be accounted for considering any of the possible subsystems. The quantum part of genuine correlations is quantified by *genuine n -partite quantum discord*. The approach of Ref. [17] represents a natural extension of the concept of QD as introduced for bipartite systems, and this is the reason why we will follow it in the present work. However, it requires massive numerical optimization procedures over a number of parameters, thus making the calculations very demanding [19]. Therefore, the application of such a criterion is not easily amenable.

This justifies the scarce number of works investigating the time evolution of quantum correlations in multipartite system coupled to noisy environments. Specifically only few cases have been considered: two level systems undergoing random telegraph noise [20, 21] and quantum phase transitions in spin systems [22, 23, 24].

The purpose of this paper is to derive an analytical expression for the *genuine tripartite quantum discord* (GTQD) for a class of three qubits systems. In detail, we will focus on those systems described by X-states, which play a relevant role in a large number of physical systems and allow for easy calculations of certain entanglement measures [25, 26]. X-states have been widely investigated also in bipartite systems, where an analytical expression for QD has been proposed in [27]. However, this approach has been questioned, since it is not always providing the correct result [28, 29, 30].

The paper is organized as follows. In Sec. 2 we introduce the genuine quantifiers for correlations in multipartite quantum systems. In Sec. 3 we introduce the expression for a symmetrical tripartite X-state and derive some constraints on its defining parameters. In Sec. 4 we estimate all the quantities required to compute GTQD, and in particular we describe the optimization procedures (both numerical and analytical) appearing in the expression of GTQD. Sec. 5 concerns the comparison between our results on GTQD and others already present in the literature and, finally, in Sec. 6 we draw conclusions.

2 QUANTIFIERS FOR GENUINE TRIPARTITE CORRELATIONS

Here we illustrate the correlation measures adopted in this work to quantify tripartite quantum discord and entanglement.

2.1 Tripartite Quantum Discord

In a tripartite system, described by a state $\rho = \rho_{A,B,C}$, the tripartite quantum mutual information is obtained as a generalization of the quantum mutual information for bipartite systems [31, 32, 17, 33]:

$$T(\rho) = S(\rho_A) + S(\rho_B) + S(\rho_C) - S(\rho), \quad (1)$$

and represents the total amount of correlations encoded in this system¹. Here $S(\rho) = -\text{Tr}[\rho \log_2(\rho)]$ is the von Neumann entropy, and ρ_i ($i = A, B, C$) is the reduced density matrix for the subsystem i . Following the same procedure used in the literature for bipartite systems [7], *Giorgi et al.* [17] define the *tripartite classical correlations* in the system as the quantum version (of a classical analogue) of the mutual information derived from the Bayes' rule:

$$J(\rho) = \max_{i,j,k \in \{A,B,C\}} [S(\rho_i) - S(\rho_{i|j}) + S(\rho_k) - S(\rho_{k|ij})], \quad (2)$$

which has been optimized over the indices i, j, k in the set of all the possible permutation of subsystems $\{A, B, C\}$. Here $S(\rho_{i|j})$ and $S(\rho_{k|ij})$ are relative entropies and $\rho_{i|j}$ and $\rho_{k|ij}$ are the density matrices after a measurement on the subsystem i or after a measurement on both subsystems i and j , respectively [20]. We refer the reader to the Appendix A for a detailed definition of the relative entropies and their optimization. Like for bipartite systems, the tripartite quantum discord is given by the difference between total and classical correlations:

$$D(\rho) = T(\rho) - J(\rho). \quad (3)$$

¹ It can be shown that this quantity measures, in terms of relative entropy, the distance between the state ρ and the nearest classical state with no correlations $\rho^A \otimes \rho^B \otimes \rho^C$. Indeed, by the definition of relative entropy, we get $S(\rho || \rho^A \otimes \rho^B \otimes \rho^C) = -\text{Tr}[\rho \log_2(\rho^A \otimes \rho^B \otimes \rho^C)] - S(\rho)$, then using the linearity of trace and the additivity of logarithm - remember that ρ^i are the marginals of ρ - we get $-\text{Tr}[\rho \log_2(\rho^A \otimes \rho^B \otimes \rho^C)] = -\text{Tr}[\rho \log_2(\rho^A) \otimes I \otimes I] + \dots = S(\rho^A) + S(\rho^B) + S(\rho^C)$ [32, 17].

However, among the correlations included in $T(\rho)$, a subset is shared by all of the three subsystems (*genuine tripartite mutual information*), and can be estimated as:

$$T^{(3)}(\rho) = T(\rho) - T^{(2)}(\rho), \quad (4)$$

where $T^{(2)}(\rho)$ is the maximum amount of mutual information shared by any couple of subsystems:

$$T^{(2)}(\rho) = \max[I(\rho_{A,B}), I(\rho_{A,C}), I(\rho_{B,C})], \quad (5)$$

where $I(\rho_{AB}) = S(\rho_A) + S(\rho_B) - S(\rho_{AB})$. Since all the correlations that cannot be accounted for by $T^{(2)}(\rho)$ must be shared between all of the three subsystems, we can conclude that $T^{(3)}(\rho)$ measures the distance between ρ and the closest product state along any bipartite cut of the system. Indeed it can be shown that $T^{(3)}(\rho) = \min[I(\rho_{AB,C}), I(\rho_{AC,B}), I(\rho_{BC,A})]$ (see Ref. [17]).

Analogously, the genuine tripartite classical correlations reads:

$$J^{(3)}(\rho) = J(\rho) - J^{(2)}(\rho), \quad (6)$$

and GTQD:

$$D^{(3)}(\rho) = D(\rho) - D^{(2)}(\rho), \quad (7)$$

where²:

$$J^{(2)}(\rho) = \max[J(\rho_{A,B}), J(\rho_{A,C}), J(\rho_{B,C})], \quad (8)$$

$$D^{(2)}(\rho) = \min[D(\rho_{A,B}), D(\rho_{A,C}), D(\rho_{B,C})]. \quad (9)$$

Eqs. (4), (6) and (7) can be significantly simplified for the case of a state ρ symmetrical under any exchange of its subsystems. Indeed it can be shown that [20]:

$$T^{(3)}(\rho) = I(\rho_{A,BC}) = S(\rho_A) + S(\rho_{A,B}) - S(\rho), \quad (10)$$

$$D^{(3)}(\rho) = S(\rho_{A|BC}) + S(\rho_{A,B}) - S(\rho), \quad (11)$$

$$J^{(3)}(\rho) = S(\rho_A) - S(\rho_{A|BC}). \quad (12)$$

2.2 Tripartite Negativity

In tripartite systems, represented by a state ρ , we can detect the presence of entanglement between subsystems by using the *negativity* N , which is defined as follows [34]:

$$N(\rho^{tC}) = \sum_i |\lambda_i(\rho^{tC})| - 1. \quad (13)$$

In the previous expression, ρ^{tC} is the partial transpose of ρ with respect to the subsystem C , and $\lambda_i(\rho^{tC})$ are the eigenvalues of ρ^{tC} . The negativity can be equivalently interpreted as the sum of the absolute values of the negative eigenvalues of ρ^{tC} [34], and it depends upon the subsystem on which we make the partial transpose of ρ .

When negativity is higher than zero, we can conclude that there is an entanglement between the subsystem C and the compound subsystem $A - B$, but the converse is not necessarily true. Starting from this point, we can define the *tripartite negativity* as follows [35]:

$$N^{(3)}(\rho) = \sqrt[3]{N(\rho^{tA})N(\rho^{tB})N(\rho^{tC})}, \quad (14)$$

and this quantifier will be different from zero only when the entanglement is shared among all of the three subsystems, i.e. it is a “full” tripartite entanglement [35]. However, apart from pure states, a null negativity could indeed not imply the absence of entanglement. Moreover, we must notice that tripartite negativity cannot distinguish the entanglement of a genuine tripartite entangled state from that of a biseparable state in a generalized sense [35,

² In Eqs. (8) and (9) we used the bipartite quantifiers $J(\rho_{A,B}) = \max[S(\rho_{A,B}) - S(\rho_{A|B}), S(\rho_{A,B}) - S(\rho_{B|A})]$ and $D(\rho_{A,B}) = I(\rho_{A,B}) - J(\rho_{A,B})$ as they are usually defined in literature for 2-qubits systems [17, 20, 9].

23]. For tripartite systems that are symmetrical under any exchange of their qubits, as in our case of study, the tripartite negativity and the negativity always coincide:

$$N^{(3)}(\rho) = N(\rho^{tA}) = N(\rho^{tB}) = N(\rho^{tC}). \quad (15)$$

Another possible quantifier for tripartite entanglement is the *three-tangle* [36], but in this work we use negativity since the three-tangle is not able to detect tripartite entanglement for all states, e.g. W states [37]. However, it should be noticed that $N^{(3)}$ in this work is used simply to provide a further comparison with the outcomes of GTQD, and it is not used to quantify genuine entanglement.

3 THREE-QUBITS SYMMETRICAL X-STATES

Here, we focus on three qubits X-states [38] which, for the particular features of their quantum correlations, have been investigated in the literature, both for bipartite [18, 29, 27] and tripartite systems [26, 25, 20]. A generic tripartite X-state can be written in the form [26]:

$$\rho = \begin{pmatrix} a_1 & 0 & 0 & 0 & 0 & 0 & 0 & c_1 \\ 0 & a_2 & 0 & 0 & 0 & 0 & c_2 & 0 \\ 0 & 0 & a_3 & 0 & 0 & c_3 & 0 & 0 \\ 0 & 0 & 0 & a_4 & c_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & c_4^* & b_4 & 0 & 0 & 0 \\ 0 & 0 & c_3^* & 0 & 0 & b_3 & 0 & 0 \\ 0 & c_2^* & 0 & 0 & 0 & 0 & b_2 & 0 \\ c_1^* & 0 & 0 & 0 & 0 & 0 & 0 & b_1 \end{pmatrix}. \quad (16)$$

In order to simplify the derivation of an analytical expression for GTQD, we limit ourselves to X-states which are symmetrical under any exchange of their subsystems, and invariant under the flip of all of their qubits. This means that ρ can be written in the form:

$$\rho = \frac{1}{8} \begin{pmatrix} 1 - a_1 & 0 & 0 & 0 & 0 & 0 & 0 & c_1 \\ 0 & \alpha_1 & 0 & 0 & 0 & 0 & c_2 & 0 \\ 0 & 0 & \alpha_1 & 0 & 0 & c_2 & 0 & 0 \\ 0 & 0 & 0 & \alpha_1 & c_2 & 0 & 0 & 0 \\ 0 & 0 & 0 & c_2 & \alpha_1 & 0 & 0 & 0 \\ 0 & 0 & c_2 & 0 & 0 & \alpha_1 & 0 & 0 \\ 0 & c_2 & 0 & 0 & 0 & 0 & \alpha_1 & 0 \\ c_1 & 0 & 0 & 0 & 0 & 0 & 0 & 1 - a_1 \end{pmatrix}, \quad (17)$$

where $\alpha_1 = 1 + \frac{a_1}{3}$ (we used the property $\text{Tr}[\rho] = 1$ to express a_2 in terms of a_1 , and then we made the substitutions $a_1 \rightarrow \frac{1-a_1}{8}$, $c_i \rightarrow \frac{c_i}{8}$ to get a simpler expression). Now ρ depends only on the parameters (a_1, c_1, c_2) which, from now on, are assumed to be real due to the qubit-flip invariance. Recently, symmetry features of mixed entangled states have been also exploited in Ref. [39] to evaluate analytically both nonlocality and global quantum discord in multipartite systems.

From the requirement $0 \leq \lambda_i \leq 1 \forall i$, where $\lambda_{1,2} = \frac{1}{8}(1 - a_1 \mp c_1)$ and $\lambda_{3-4-5,6-7-8} = \frac{1}{24}(3 + a_1 \mp 3c_2)$ are the eigenvalues of ρ , we obtain the following constraints for the parameters:

$$\begin{aligned} a_1 &\in [-3, 1], \\ c_1 &\in [a_1 - 1, 1 - a_1] \\ c_2 &\in \left[-1 - \frac{a_1}{3}, 1 + \frac{a_1}{3}\right]. \end{aligned} \quad (18)$$

4 ESTIMATION OF GENUINE TRIPARTITE QUANTUM DISCORD

4.1 von Neumann Entropies for ρ and $\rho_{A,B}$

Now, in order to give an analytical estimation of $D^{(3)}(\rho)$ for the state ρ described by Eq. (17), we calculate the von Neumann entropies for ρ and for the marginal $\rho_{A,B} = \text{Tr}_C[\rho]$, which appears in the expression of GTQD given by Eq. (11).

From the definition of von Neumann entropy, it follows that:

$$S(\rho) = 3 + \frac{1}{8} [2(3 + a_1) \log_2(3) - (1 - a_1 - c_1) \log_2(1 - a_1 - c_1) - (1 - a_1 + c_1) \log_2(1 - a_1 + c_1) - (3 + a_1 - 3c_2) \log_2(3 + a_1 - 3c_2) - (3 + a_1 + 3c_2) \log_2(3 + a_1 + 3c_2)]. \quad (19)$$

From Eq. (17) we obtain:

$$\rho_{A,B} = \begin{pmatrix} \frac{3-a_1}{12} & 0 & 0 & 0 \\ 0 & \frac{3+a_1}{12} & 0 & 0 \\ 0 & 0 & \frac{3+a_1}{12} & 0 \\ 0 & 0 & 0 & \frac{3-a_1}{12} \end{pmatrix}, \quad (20)$$

and after straightforward calculations we find:

$$S(\rho_{A,B}) = -\frac{1}{6} (3 - a_1) \log_2(3 - a_1) - \frac{1}{6} (3 + a_1) \log_2(3 + a_1) + 2 + \log_2(3). \quad (21)$$

4.2 Relative entropy minimization

In order to finally evaluate $D^{(3)}(\rho)$ we need to calculate the relative entropy $S(\rho_{A|BC})$. Following the derivation procedure given in the Appendix A, $S(\rho_{A|BC})$ can be written as:

$$S(\rho_{A|BC}) = \min_{\theta_i, \phi_i} S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2) = \min_{\theta_i, \phi_i} \left\{ 1 + \frac{1}{6} [\lambda_A \log_2 \lambda_A + \lambda_B \log_2 \lambda_B] - \frac{1}{12} \sum_{i=1}^4 \lambda_i \log_2 \lambda_i \right\}, \quad (22)$$

where θ_i and ϕ_i are optimization parameters (the angles defining the basis vectors: see again Appendix A), and:

$$\begin{aligned} \lambda_A &= 3 + a_1 \cos(2\theta_1) \cos(2\theta_2), \\ \lambda_B &= 3 - a_1 \cos(2\theta_1) \cos(2\theta_2), \\ \lambda_C &= \frac{9}{16} \sin^2(2\theta_1) \sin^2(2\theta_2) \left[(c_1 - c_2)^2 + 4c_2 (\cos(\phi_1) + \cos(\phi_2)) (c_2 \cos(\phi_1) + c_1 \cos(\phi_2)) \right], \\ \lambda_{1,2} &= \lambda_B \pm \sqrt{a_1^2 (\cos(2\theta_1) + \cos(2\theta_2))^2 + \lambda_C}, \\ \lambda_{3,4} &= \lambda_A \pm \sqrt{a_1^2 (\cos(2\theta_1) - \cos(2\theta_2))^2 + \lambda_C}. \end{aligned} \quad (23)$$

The optimization of S_{rel} is an hard task, and cannot be performed fully analytically in a simple way. Indeed, it has been proven that in a bipartite system the optimization of the relative entropy (for a general density matrix) involves the solution of equations containing logarithms of nonlinear quantities, that cannot be obtained analytically (see for instance [11, 12]). This is the reason why we developed a numerical approach to the minimization, whose results have been used as guidelines to give an analytical expression for S_{rel} . A similar method has already been adopted independently to estimate the quantum discord of two-qutrit Werner states in Ref. [40].

First, in our procedure, we generate randomly a suitable number of triplets (a_1, c_1, c_2) (obeying to the constraints of Eq. (18)), and then we minimize numerically the corresponding expression of $S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2)$ over a grid of points in the 4D-space $\mathbb{U} = R_{\theta_1} \times R_{\theta_2} \times R_{\phi_1} \times R_{\phi_2}$, where R_{θ_i} and R_{ϕ_i} are the intervals $[0; \pi)$ and $[0; 2\pi)$ respectively, given the periodicity of the functions in Eqs. (23). The optimization procedure, which has been shown to be an NP-complete problem [19], was performed using exhaustive enumeration (i.e. brute force search) over a grid in the \mathbb{U} space, to be sure to find the true absolute minima of S_{rel} . Our calculations indicate that the function S_{rel} exhibits many equivalent absolute minima, and that the ‘‘first’’ one (i.e. the one with the lowest values of its coordinates) is always reached for $\theta_1 = \theta_2 = \theta$ and $\phi_1 = 0$. Specifically, it is found alternatively in one of these three points

$(\theta_1, \theta_2, \phi_1, \phi_2)$ of \mathbb{U} : $(0, 0, 0, 0)$, $(\frac{\pi}{4}, \frac{\pi}{4}, 0, 0)$ or $(\frac{\pi}{4}, \frac{\pi}{4}, 0, \bar{\phi}_2)$, where $\bar{\phi}_2$ depends upon (a_1, c_1, c_2) ³. This means that the minimal relative entropy $S(\rho_{A|BC})$ can take only three possible analytical forms (provided that one can find an analytical expression for $\bar{\phi}_2$).

Starting from these numerical results, we performed an analytical study on the specific case of $S_{rel}(\theta, \theta, \phi_1, \phi_2)$, which confirmed that this function has two extrema in $\theta = 0$ and $\theta = \frac{\pi}{4}$. Moreover, our analytical approach showed that the function $S_{rel}(\frac{\pi}{4}, \frac{\pi}{4}, 0, \phi_2)$ attains its minimum value for $\sin(\phi_2) = 0$ or $\cos(\phi_2) = \left(-\frac{c_1+c_2}{2c_1}\right)$, (which holds only if certain conditions are satisfied - see Eq. (60) in the Appendix B). This is consistent with numerical calculations, which give as minimum $\phi_2 = 0$ or $\phi_2 = \bar{\phi}_2 = \arccos\left(-\frac{c_1+c_2}{2c_1}\right)$. Further details are given in the Appendix B.

Our derivation leads to the following expressions for the minimum values of S_{rel} :

$$\begin{aligned} S_1 &= S_{rel}(0, 0, 0, 0) = 1 - \frac{1}{12}\gamma(a_1), \\ S_2 &= S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, 0\right) = 1 - \frac{1}{2}\varepsilon\left(\frac{3c_2+c_1}{4}\right), \\ S_3 &= S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, \bar{\phi}_2\right) = 1 - \frac{1}{2}\varepsilon\left(\frac{1}{4}\sqrt{\frac{(c_1-c_2)^3}{c_1}}\right), \end{aligned} \quad (24)$$

where

$$\gamma(x) = (3+x)\log_2(3+x) + (3-3x)\log_2(3-3x) - 2(3-x)\log_2(3-x), \quad (25)$$

$$\varepsilon(x) = (1+x)\log_2(1+x) + (1-x)\log_2(1-x). \quad (26)$$

When both S_2 and S_3 are well defined expressions, we found with additional analytical calculations that $S_3 < S_2$ if $c_1 \cdot c_2 < 0$ (see Appendix C). This implies that the relative entropy takes the form:

$$S(\rho_{A|BC}) = \begin{cases} \min\{S_1, S_3\} & |3c_1| \geq |c_2| \text{ and } c_1 \cdot c_2 < 0 \\ \min\{S_1, S_2\} & \text{otherwise} \end{cases}, \quad (27)$$

where the minimization is required only if both entropy expressions are well defined (considering the constraint imposed on a_1 , we can say that the expression of S_1 is always well defined, at least in the limit given by Eq. 18). In our simulations over a set of 6000 triplets of values (a_1, c_1, c_2) randomly generated, we observe that the minimum of $S(\rho_{A|BC})$ occurs in S_1 in the 52% of the cases, in S_2 in the 31% of the cases and in S_3 in the remaining 17% of the cases.

Finally, by using Eqs. (11), (19), (21) and (27), we can write the expression for GTQD:

$$\begin{aligned} D^{(3)}(\rho) &= S(\rho_{A|BC}) - \frac{1}{6}(3-a_1)\log_2(3-a_1) - \frac{1}{6}(3+a_1)\log_2(3+a_1) + 2 + \log_2(3) \\ &\quad - \left\{3 + \frac{1}{8}\left[2(3+a_1)\log_2(3) - (1-a_1-c_1)\log_2(1-a_1-c_1) - (1-a_1+c_1)\log_2(1-a_1+c_1)\right.\right. \\ &\quad \left.\left. - (3+a_1-3c_2)\log_2(3+a_1-3c_2) - (3+a_1+3c_2)\log_2(3+a_1+3c_2)\right]\right\}. \end{aligned} \quad (28)$$

5 RESULTS AND DISCUSSION

To validate our approach, we apply the above expression to two prototypical cases of study. In particular, the well known result $D^{(3)}(\rho_{GHZ}) = 1$ for a pure GHZ state $\rho_{GHZ} = |GHZ\rangle\langle GHZ| = \frac{1}{2}(|000\rangle + |111\rangle)(\langle 000| + \langle 111|)$, is obtained by setting $a_1 = -3$, $c_1 = \pm 4$ and $c_2 = 0$ in Eq. (17).

Analogously, it can be shown that for a maximally mixed state with $c_1 = c_2 = 0$ we find $D^{(3)}(\rho) = 0$ (and $S(\rho_{A|BC}) = S(\rho) - S(\rho_{A,B}) = S_1$, since in this case $S_2 = S_3 = 1$) whatever the value of a_1 is, as expected since all correlations are classical.

Moving towards a more general case, we can set $c_1 = c_2 = c$ and plot the values of $D^{(3)}(\rho)$ with respect to a_1 and c . As we see from Figure 1, along the line $c = 0$ (maximally mixed states) the genuine tripartite discord vanishes - as explained above. Moreover, $D^{(3)}(\rho)$ is zero also along the line $a_1 = 0$, which does not corresponds to mixed states, but to a case where again we have $S(\rho_{A|BC}) = S(\rho) - S(\rho_{A,B}) = S_2$ (since here $S_1 = S_3 = 1$). Maximum values of $D^{(3)}(\rho)$ are achieved when $a_1 \simeq 1.38$ and $c \simeq 0.54$.

³ Notice that when $\theta_i = 0$ other equivalent minima can be found for $\theta_i = \frac{\pi}{2}$ or $\theta_i = \pi$, and when $\theta_i = \frac{\pi}{4}$ other equivalent minima can be found for $\theta_i = \frac{3\pi}{4}$, but we will focus only on the cases $\theta = 0$ or $\theta = \frac{\pi}{4}$, which are the simpler ones.




Fig. 1 Genuine Tripartite Quantum Discord for $c_1 = c_2 = c$ (notice that the maximum value of z axis is set to 0.45 and not to 1.0 in order to make the graph more readable).




Fig. 2 Genuine Tripartite Quantum Discord for $a_1 = 0$ (notice that the maximum value of z axis is set to 0.6 and not to 1.0 in order to make the graph more readable).

A further analysis of the states with $a_1 = 0$ is performed by investigating the behavior of GTQD as a function of c_1 and c_2 (see Figure 2). When $c_1 \neq c_2$ the GTQD never goes to zero, and it reaches its maximum value in $(c_1, c_2) = (1, -1)$ or $(-1, 1)$, where $D^{(3)}(\rho) = 1 - \frac{1}{2}\varepsilon\left(\frac{1}{\sqrt{2}}\right)$. In detail, the density matrix ρ' obtained by setting $(c_1, c_2) = (1, -1)$ in Eq. (17) is a linear combination of density matrices of pure GHZ states of the type:

$$\rho_{GHZ}^{\pm}(k) = \frac{1}{2} (|k\rangle \pm |\bar{k}\rangle) (\langle k| \pm \langle \bar{k}|), \quad (29)$$

where k is a three bit binary number (from 0 to 7) and \bar{k} is the result of flipping each bit of k [26]. Indeed:

$$\rho' = \frac{1}{4} (\rho_{GHZ}^+(0) + \rho_{GHZ}^-(1) + \rho_{GHZ}^-(2) + \rho_{GHZ}^-(3)), \quad (30)$$

and a similar expression can be found for ρ when $(c_1, c_2) = (-1, 1)$. Unlike pure GHZ states, this mixed state ρ' is not a maximally entangled one (indeed its negativity is zero, as we will see in the following), but it shows a GTQD different from zero. Moreover, also the state ρ'' obtained by setting $(c_1, c_2) = (1, 1)$ is a linear combination of pure GHZ states:

$$\rho'' = \frac{1}{4} (\rho_{GHZ}^+(0) + \rho_{GHZ}^+(1) + \rho_{GHZ}^+(2) + \rho_{GHZ}^+(3)), \quad (31)$$

but this state is characterized by zero GTQD. A similar expression can be found for ρ when $(c_1, c_2) = (-1, -1)$, and the value of GTQD is again zero. Therefore, we conclude that a linear combinations of GHZ states is characterized by zero discord when all the states are of kind $\rho_{GHZ}^+(k)$ (or $\rho_{GHZ}^-(k)$), as it occurs for bipartite systems when we combine linearly Bell states with the same sign. Otherwise, if we combine GHZ states of kind $\rho_{GHZ}^+(k)$ and $\rho_{GHZ}^-(k)$ together, the GTQD can be different from zero.

Finally, we study $D^{(3)}(\rho)$ by setting $c_2 = 0$. We see in Figure 3 that the GTQD vanishes along the line $c_1 = 0$ and reaches its absolute maximum value (as expected) for the maximally entangled GHZ states $(a_1, c_1) = (3, \pm 4)$.

Now we compare GTQD and tripartite entanglement, where the latter is quantified by means of tripartite negativity $N^{(3)}(\rho)$, given in Eq. (15). For the state of Eq. (17) we get:

$$\begin{aligned} N^{(3)}(\rho) = & \frac{1}{24} (|3 + a_1 - 3c_1| + |3 + a_1 + 3c_1| + \\ & + 2|3 + a_1 - 3c_2| + 2|3 + a_1 + 3c_2| + 3|1 - a_1 - c_2| + 3|1 - a_1 + c_2|) - 1. \end{aligned} \quad (32)$$

By evaluating this expression for some special values of the parameters (a_1, c_1, c_2) , we see that there are regions in which the tripartite entanglement cannot be detected (i.e. the negativity is zero) but on the other hand the GTQD is different from zero. In particular, for $c_1 = c_2$ or $a_1 = 0$ the tripartite negativity is zero everywhere. On the contrary, for $c_2 = 0$ (see Figure 4) there are regions where GTQD can be both smaller or larger than the negativity. This result is not surprising. Indeed, in bipartite systems for some cases entanglement has been found to be larger than quantum discord [10, 27], since the latter cannot simply be considered as the sum of the entanglement and other forms of nonclassical correlations [27]. However, in our case we can explain this result also on the grounds




Fig. 3 Genuine Tripartite Quantum Discord for $c_2 = 0$.




Fig. 4 Tripartite Negativity for $c_2 = 0$.

that tripartite negativity quantifies a tripartite entanglement not necessarily genuine [35, 23], so it could detect in principle a larger amount of quantum correlations with respect to a genuine quantifier, such as GTQD.

6 CONCLUSIONS

In this paper, we have developed a hybrid analytical-numerical approach to find the analytical expression for GTQD $D^{(3)}(\rho)$, specifically for a subclass of X-states, symmetrical under exchange and flip of all qubits, which are defined by three parameters (a_1 , c_1 and c_2). The expression of $D^{(3)}(\rho)$ depends on the relative entropy $S(\rho_{A|BC})$, whose estimation requires the minimization of the function $S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2)$ depending on 4 angular variables. Numerical calculations show that S_{rel} possesses only three different minimum points, which should correspond to three distinct analytical expressions for $S(\rho_{A|BC})$. Further analytical studies performed over a simplified form of $S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2)$ allowed us to find the exact analytical expressions for $S(\rho_{A|BC})$ and the conditions under which they can be used. These analytical findings have been compared with some thousand of numerical simulations and have been proven always right. Moreover, they are able to reproduce the known results for GTQD in some simple systems, namely GHZ and maximally mixed states. When confronted with tripartite negativity, the calculations show that there are regions in the space of the parameters (a_1 , c_1 and c_2) where entanglement cannot be detected, while genuine quantum correlations (evaluated in terms of GTDQ) differ from zero.

Further possible development of this work include the analytical study of the time evolution of genuine quantum correlations (as accounted by GTQD) and the extension of the hybrid approach here developed to more general cases.

A Appendix: Relative Entropies Definition

Following *Zhao et al.* [41], we can define the *relative entropy* $S(\rho_{A|BC})$ for tripartite systems as:

$$S(\rho_{A|BC}) = \min_{\{E_{ij}^{BC}\}} \sum_{ij} p_{ij} S(\rho_{A|E_{ij}^{BC}}), \quad (33)$$

where:

$$\rho_{A|E_{ij}^{BC}} = \frac{\tilde{\rho}_{A|E_{ij}^{BC}}}{p_{ij}} = \frac{1}{p_{ij}} \text{Tr}_{B,C} \left[\left(I^A \otimes E_{ij}^{BC} \right) \rho \right], \quad (34)$$

$$p_{i,j} = \text{Tr}_{A,B,C} \left[\left(I^A \otimes E_{ij}^{BC} \right) \rho \right]. \quad (35)$$

In the previous expressions, the operators E_{ij}^{BC} are *positive-operator-valued measures* (POVMs) that act on parties B and C (i.e. in the Hilbert space $\mathcal{H}_{BC} = \mathcal{H}_B \otimes \mathcal{H}_C$), and whose outcomes are labeled with two indices (i, j) . For sake of simplicity, we will replace the global POVM E_{ij}^{BC} with the external product of two local POVMs, acting separately on parties B and C , using the same procedure given in [11]. Moreover, following the convention in literature [8, 7, 42, 17, 41]), we use orthogonal *projection-valued measures* (PVMs)

to optimize entropy in Eq. (33), since they are easier to implement in the numerical minimization process⁴. Then, the measurement operators are:

$$E_{ij}^{BC} \rightarrow \Pi_i^B \otimes \Pi_j^C = |\beta_i\rangle\langle\beta_i| \otimes |\gamma_j\rangle\langle\gamma_j|, \quad (36)$$

where $|\beta_i\rangle$ and $|\gamma_j\rangle$ are orthogonal normalized basis states of the Hilbert spaces \mathcal{H}_B and \mathcal{H}_C , respectively.

A possible parametrization of the basis vectors $|\beta_i\rangle$ and $|\gamma_j\rangle$ with respect to the standard basis $\{|0\rangle, |1\rangle\}$ can be found in literature (see [20]; for a full derivation of the basis vectors see [40]):

$$|\beta_1\rangle = \cos\theta_1 |0_B\rangle + e^{+i\phi_1} \sin\theta_1 |1_B\rangle, \quad (37)$$

$$|\beta_2\rangle = \sin\theta_1 |0_B\rangle - e^{+i\phi_1} \cos\theta_1 |1_B\rangle, \quad (38)$$

$$|\gamma_1\rangle = \cos\theta_2 |0_C\rangle + e^{+i\phi_2} \sin\theta_2 |1_C\rangle, \quad (39)$$

$$|\gamma_2\rangle = \sin\theta_2 |0_C\rangle - e^{+i\phi_2} \cos\theta_2 |1_C\rangle, \quad (40)$$

where the angles θ_i and ϕ_i belong to the interval $[0; 2\pi)$.

Since we are studying a system whose state is symmetrical under any permutation of its subsystems, any subscript or superscript referring to a particular subsystem in the relative entropy expression (33) can be dropped. Now, recalling the sum rule $\sum_{i=1}^2 \tilde{\lambda}_i^{(ij)} = p_{ij}$ for the eigenvalues $\tilde{\lambda}_i^{(ij)}$ of $\tilde{\rho}_{ij} = \tilde{\rho}_{A|E_{ij}^{BC}}$ (crf. Eqs. (34) and (35)), we can simplify Eq. (33) as follows:

$$S(\rho_{A|BC}) = \min_{\{E_{ij}^{BC}\}} \left[-H(p) + \sum_{i,j} S(\tilde{\rho}_{ij}) \right], \quad (41)$$

where $H(p) = -\sum_{i,j} p_{ij} \log_2(p_{ij})$ is the Shannon Entropy of the probability ensemble $\{p_{ij}\}$.

Now, using Eqs. (37)-(40) to write the PVMs - together with the change of variables ($\phi_1 - \phi_2 \rightarrow \phi_1$, $\phi_1 + \phi_2 \rightarrow \phi_2$), which simplifies our calculations - the relative entropy in (41) can be written as a function of four angular variables:

$$S(\rho_{A|BC}) = \min_{\theta_i, \phi_i} S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2). \quad (42)$$

The final expression for $S(\rho_{A|BC})$, with all terms written explicitly, is given in Section 4.2.

B Appendix: Analytical study of $S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2)$

The relative entropy $S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2)$ of Eq. (22)

$$S_{rel}(\theta_1, \theta_2, \phi_1, \phi_2) = 1 + \frac{1}{6} [\lambda_A \log_2 \lambda_A + \lambda_B \log_2 \lambda_B] - \frac{1}{12} \sum_{i=1}^4 \lambda_i \log_2 \lambda_i \quad (43)$$

can be studied in a simplified form setting $\theta_1 = \theta_2 = \theta$. Under this condition, the λ_j of Eqs. (23) become:

$$\begin{aligned} \lambda_A &= 3 + a_1 \cos^2(2\theta), \\ \lambda_B &= 3 - a_1 \cos^2(2\theta), \\ \lambda_C &= \frac{9}{16} \sin^4(2\theta) f(\phi_1, \phi_2), \\ f(\phi_1, \phi_2) &= \left[(c_1 - c_2)^2 + 4c_2 (\cos(\phi_1) + \cos(\phi_2)) (c_2 \cos(\phi_1) + c_1 \cos(\phi_2)) \right] \\ \lambda_{1,2} &= \lambda_B \pm \sqrt{4a_1^2 \cos^2(2\theta) + \lambda_C}, \\ \lambda_{3,4} &= \lambda_A \pm \sqrt{\lambda_C}. \end{aligned} \quad (44)$$

The minima of S_{rel} must satisfy the equation:

$$\frac{\partial S_{rel}(\theta, \theta, \phi_1, \phi_2)}{\partial \theta} = 0, \quad (45)$$

which can be rewritten as follows:

$$\frac{\partial S_{rel}}{\partial \lambda_A} \frac{\partial \lambda_A}{\partial \theta} + \frac{\partial S_{rel}}{\partial \lambda_B} \frac{\partial \lambda_B}{\partial \theta} + \sum_{i=1}^4 \frac{\partial S_{rel}}{\partial \lambda_i} \frac{\partial \lambda_i}{\partial \theta} = 0. \quad (46)$$

⁴ This approach has been recently questioned by *Zhao et al.*: in their paper [41], they show that the product POVM $E_i^B \otimes E_j^C$ may not be the optimal POVM E_{ij}^{BC} that minimizes genuine tripartite discord. However, we must notice that the qualitative behaviors of $D^{(3)}(\rho)$ are not changed by this approach (except for the overestimation of $D^{(3)}(\rho)$), i.e. both approaches are able to record the presence of GTQD and its increasing (or decreasing) trend, according to the variations of the parameters which define the density operator ρ .

The derivatives of the λ_j appearing in Eq. (46) are given by:

$$\begin{aligned}
\frac{\partial \lambda_A}{\partial \theta} &= -4a_1 \cos(2\theta) \sin(2\theta) \\
\frac{\partial \lambda_B}{\partial \theta} &= +4a_1 \cos(2\theta) \sin(2\theta) \\
\frac{\partial \lambda_C}{\partial \theta} &= \frac{9}{2} \sin^3(2\theta) \cos(2\theta) f(\phi_1, \phi_2) \\
\frac{\partial \lambda_{1,2}}{\partial \theta} &= \frac{\partial \lambda_B}{\partial \theta} \pm \frac{16a_1^2 \cos(2\theta) \sin(2\theta) + \frac{\partial \lambda_C}{\partial \theta}}{2\sqrt{4a_1^2 \cos^2(2\theta) + \lambda_C}}, \\
\frac{\partial \lambda_{3,4}}{\partial \theta} &= \frac{\partial \lambda_A}{\partial \theta} \pm \frac{\frac{\partial \lambda_C}{\partial \theta}}{2\sqrt{\lambda_C}},
\end{aligned} \tag{47}$$

and furthermore:

$$\frac{\partial S_{rel}}{\partial \lambda_{A,B}} = +\frac{1}{6 \ln 2} (\ln \lambda_{A,B} + 1) \tag{48}$$

$$\frac{\partial S_{rel}}{\partial \lambda_i} = -\frac{1}{12 \ln 2} (\ln \lambda_i + 1) \quad (i = 1, 2, 3, 4) \tag{49}$$

Eq. (46) then becomes:

$$\begin{aligned}
\cos(2\theta) \sin(2\theta) \left[-4a_1 \frac{\partial S_{rel}}{\partial \lambda_A} + 4a_1 \frac{\partial S_{rel}}{\partial \lambda_B} + \right. \\
\left. \left(4a_1 + \frac{16a_1^2 + \frac{9}{2} \sin^2(2\theta) f(\phi_1, \phi_2)}{2\sqrt{4a_1^2 \cos^2(2\theta) + \lambda_C}} \right) \frac{\partial S_{rel}}{\partial \lambda_1} + \left(4a_1 - \frac{16a_1^2 + \frac{9}{2} \sin^2(2\theta) f(\phi_1, \phi_2)}{2\sqrt{4a_1^2 \cos^2(2\theta) + \lambda_C}} \right) \frac{\partial S_{rel}}{\partial \lambda_2} + \right. \\
\left. \left(-4a_1 + \frac{\frac{9}{2} \sin^2(2\theta) f(\phi_1, \phi_2)}{2\sqrt{\lambda_C}} \right) \frac{\partial S_{rel}}{\partial \lambda_3} + \left(-4a_1 - \frac{\frac{9}{2} \sin^2(2\theta) f(\phi_1, \phi_2)}{2\sqrt{\lambda_C}} \right) \frac{\partial S_{rel}}{\partial \lambda_4} \right] = 0 \tag{50}
\end{aligned}$$

This expression shows that $\frac{\partial S_{rel}}{\partial \theta} = 0$ when $\sin(2\theta) = 0$ or $\cos(2\theta) = 0$, that is the function can attain its minimum value for a value in the set $\theta = 0 + k\frac{\pi}{2}$ or $\theta = \frac{\pi}{4} + k\frac{\pi}{2}$, where $k \in \mathbb{Z}$. Indeed, it can be shown that the whole l.h.s. of Eq. (50) goes to zero when θ approaches in the limit the values listed before.

When we make the further assumption that $\theta = \frac{\pi}{4}$ and $\phi_1 = 0$ (as suggested by numerical calculations), we get:

$$\begin{aligned}
\lambda_A &= 3, \\
\lambda_B &= 3, \\
\lambda_C &= \frac{9}{16} f(0, \phi_2), \\
f(0, \phi_2) &= \left[(c_1 - c_2)^2 + 4c_2 (1 + \cos(\phi_2)) (c_2 + c_1 \cos(\phi_2)) \right], \\
\lambda_{1,2} &= \lambda_{3,4} = 3 \pm \sqrt{\lambda_C},
\end{aligned} \tag{51}$$

and

$$S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, \phi_2\right) = 1 + \log_2 3 - \frac{1}{6} \sum_{i=1}^2 \lambda_i \log_2 \lambda_i. \tag{52}$$

Therefore, the minimum is reached when

$$\frac{\partial S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, \phi_2\right)}{\partial \phi_2} = 0, \tag{53}$$

that is

$$\frac{\partial}{\partial \phi_2} (\lambda_1 \log_2 \lambda_1 + \lambda_2 \log_2 \lambda_2) = 0. \tag{54}$$

With further simplifications we get:

$$\left[\ln \left(\frac{3 + \sqrt{\lambda_C}}{3 - \sqrt{\lambda_C}} \right) \frac{1}{2 \ln 2 \sqrt{\lambda_C}} \right] \frac{\partial \lambda_C}{\partial \phi_2} = 0. \tag{55}$$

The expression in the square brackets is always greater than 0, since the argument of the logarithm is always greater than 1 if $\lambda_C > 0$, and when $\lambda_C \rightarrow 0$ the limit is finite, positive and different from zero. Therefore the extremum can be found only for:

$$\frac{\partial \lambda_C}{\partial \phi_2} = 0 \iff \frac{\partial f(0, \phi_2)}{\partial \phi_2} = 0, \tag{56}$$

which leads to the final equation:

$$\sin(\phi_2) (c_2 + c_1 + 2c_1 \cos(\phi_2)) = 0. \tag{57}$$

The solutions are:

$$\sin(\phi_2) = 0 \quad \text{or} \quad \cos(\phi_2) = \left(-\frac{c_1 + c_2}{2c_1}\right), \quad (58)$$

that is

$$\phi_2 = 0 + k\pi \quad \text{or} \quad \phi_2 = \pm \arccos\left(-\frac{c_1 + c_2}{2c_1}\right) + 2k\pi, \quad (59)$$

where $k \in \mathbb{Z}$. Clearly, the second set of extrema exists only if:

$$-1 \leq -\frac{c_1 + c_2}{2c_1} \leq +1 \quad \Rightarrow \quad \begin{cases} c_1 > 0 & \text{and} & -3c_1 \leq c_2 \leq c_1 \\ c_1 < 0 & \text{and} & c_1 \leq c_2 \leq -3c_1 \end{cases}. \quad (60)$$

C Appendix: Comparison between S_2 and S_3

When $\theta = \frac{\pi}{4}$ and $\phi_1 = 0$, the expression for S_{rel} can be written as:

$$\begin{aligned} S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, \phi_2\right) &= 1 + \log_2 3 - \frac{1}{6} \left[\left(3 + \sqrt{\lambda_C}\right) \log_2 \left(3 + \sqrt{\lambda_C}\right) + \left(3 - \sqrt{\lambda_C}\right) \log_2 \left(3 - \sqrt{\lambda_C}\right) \right] \\ &= 1 - \frac{1}{2} \left[\left(1 + \frac{1}{3}\sqrt{\lambda_C}\right) \log_2 \left(1 + \frac{1}{3}\sqrt{\lambda_C}\right) + \left(1 - \frac{1}{3}\sqrt{\lambda_C}\right) \log_2 \left(1 - \frac{1}{3}\sqrt{\lambda_C}\right) \right] \\ &= 1 - \frac{1}{2}\varepsilon\left(\frac{1}{3}\sqrt{\lambda_C}\right), \end{aligned} \quad (61)$$

where $\varepsilon(x)$ is given by (26). The function $S(x) = 1 - \frac{1}{2}\varepsilon(x)$ is known in the literature as an estimator of correlations and relative entropies in bipartite systems [9], and its expression holds only for $-1 \leq x \leq 1$ (in our case it is always $x > 0$). Due to its symmetry properties, $S(x)$ has its maximum value for $x = 0$, and decreases monotonically as x approaches 1 (or -1). Therefore we conclude that:

$$S(x_1) < S(x_2) \iff x_1 > x_2 \quad \forall x_1, x_2 \geq 0 \quad (62)$$

If $\phi_2 = 0$, then λ_C takes the following value:

$$\lambda_C = \frac{9}{16} f(0, 0) = \frac{9}{16} (3c_2 + c_1)^2, \quad (63)$$

and the corresponding expression for S_{rel} is:

$$S_2 = S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, 0\right) = 1 - \frac{1}{2}\varepsilon\left(\frac{|3c_2 + c_1|}{4}\right) = 1 - \frac{1}{2}\varepsilon\left(\frac{3c_2 + c_1}{4}\right) \quad (64)$$

The λ_C expression for $\phi_2 = \bar{\phi}_2 = \arccos\left(-\frac{c_1 + c_2}{2c_1}\right)$ is the following one:

$$\lambda_C = \frac{9}{16} f(0, \bar{\phi}_2) = \frac{9}{16} \frac{(c_1 - c_2)^3}{c_1}, \quad (65)$$

which appears under a square root (for a real eigenvalue), and therefore is acceptable only if:

$$\frac{(c_1 - c_2)}{c_1} \geq 0 \quad \Rightarrow \quad \begin{cases} c_1 \geq c_2 & \text{if } c_1 > 0 \\ c_1 \leq c_2 & \text{if } c_1 < 0 \end{cases}. \quad (66)$$

The corresponding expression for S_{rel} becomes:

$$S_3 = S_{rel}\left(\frac{\pi}{4}, \frac{\pi}{4}, 0, \bar{\phi}_2\right) = 1 - \frac{1}{2}\varepsilon\left(\frac{1}{4}\sqrt{\frac{(c_1 - c_2)^3}{c_1}}\right) \quad (67)$$

When both S_2 and S_3 expressions are well defined (see Eq. (24)), then the absolute minimum of S_{rel} can occur only in the lowest of these two values, and according to Eq. (62) we conclude that

$$S_3 < S_2 \iff \frac{(c_1 - c_2)^3}{c_1} > (3c_2 + c_1)^2 \iff c_1 \cdot c_2 < 0. \quad (68)$$

Now, collecting together the last Eq. (68) and the existence conditions for S_3 (Eqs. (60) and (66)) we conclude that:

$$S_3 < S_2 \iff c_1 \cdot c_2 < 0 \quad \text{and} \quad |c_2| \leq |3c_1|. \quad (69)$$

References

1. A. Datta, S.T. Flammia, C.M. Caves, Phys. Rev. A **72**, 042316 (2005). DOI 10.1103/PhysRevA.72.042316. URL <http://link.aps.org/doi/10.1103/PhysRevA.72.042316>
2. A. Datta, G. Vidal, Phys. Rev. A **75**, 042310 (2007). DOI 10.1103/PhysRevA.75.042310. URL <http://link.aps.org/doi/10.1103/PhysRevA.75.042310>
3. A. Datta, A. Shaji, C.M. Caves, Phys. Rev. Lett. **100**, 050502 (2008). DOI 10.1103/PhysRevLett.100.050502. URL <http://link.aps.org/doi/10.1103/PhysRevLett.100.050502>
4. B.P. Lanyon, M. Barbieri, M.P. Almeida, A.G. White, Phys. Rev. Lett. **101**, 200501 (2008). DOI 10.1103/PhysRevLett.101.200501. URL <http://link.aps.org/doi/10.1103/PhysRevLett.101.200501>
5. M.A. Nielsen, I.L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000)
6. R. Horodecki, P. Horodecki, M. Horodecki, K. Horodecki, Rev. Mod. Phys. **81**, 865 (2009). DOI 10.1103/RevModPhys.81.865. URL <http://link.aps.org/doi/10.1103/RevModPhys.81.865>
7. H. Ollivier, W.H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001). DOI 10.1103/PhysRevLett.88.017901. URL <http://link.aps.org/doi/10.1103/PhysRevLett.88.017901>
8. L. Henderson, V. Vedral, Journal of Physics A: Mathematical and General **34**(35), 6899 (2001). URL <http://stacks.iop.org/0305-4470/34/i=35/a=315>
9. S. Luo, Phys. Rev. A **77**, 042303 (2008). DOI 10.1103/PhysRevA.77.042303. URL <http://link.aps.org/doi/10.1103/PhysRevA.77.042303>
10. S. Luo, Phys. Rev. A **77**, 022301 (2008). DOI 10.1103/PhysRevA.77.022301. URL <http://link.aps.org/doi/10.1103/PhysRevA.77.022301>
11. D. Girolami, G. Adesso, Phys. Rev. A **83**, 052108 (2011). DOI 10.1103/PhysRevA.83.052108. URL <http://link.aps.org/doi/10.1103/PhysRevA.83.052108>
12. S. Javad Akhtarshenas, H. Mohammadi, F.S. Mousavi, V. Nassajpour, arXiv:1304.3914 (2013). URL <http://arxiv.org/abs/1304.3914>
13. C.H. Bennett, A. Grudka, M. Horodecki, P. Horodecki, R. Horodecki, Phys. Rev. A **83**, 012312 (2011). DOI 10.1103/PhysRevA.83.012312. URL <http://link.aps.org/doi/10.1103/PhysRevA.83.012312>
14. C.C. Rulli, M.S. Sarandy, Phys. Rev. A **84**, 042109 (2011). DOI 10.1103/PhysRevA.84.042109. URL <http://link.aps.org/doi/10.1103/PhysRevA.84.042109>
15. J. Xu, Physics Letters A **377**(3&A54), 238 (2013). DOI <http://dx.doi.org/10.1016/j.physleta.2012.11.054>. URL <http://www.sciencedirect.com/science/article/pii/S0375960112012339>
16. I. Chakrabarty, P. Agrawal, A. Pati, The European Physical Journal D **65**(3), 605 (2011). DOI 10.1140/epjd/e2011-20543-y. URL <http://dx.doi.org/10.1140/epjd/e2011-20543-y>
17. G.L. Giorgi, B. Bellomo, F. Galve, R. Zambrini, Phys. Rev. Lett. **107**, 190501 (2011). DOI 10.1103/PhysRevLett.107.190501. URL <http://link.aps.org/doi/10.1103/PhysRevLett.107.190501>
18. S. Vinjanampathy, A.R.P. Rau, Journal of Physics A: Mathematical and Theoretical **45**(9), 095303 (2012). URL <http://stacks.iop.org/1751-8121/45/i=9/a=095303>
19. Y. Huang, New Journal of Physics **16**(3), 033027 (2014). URL <http://stacks.iop.org/1367-2630/16/i=3/a=033027>
20. F. Buscemi, P. Bordone, Phys. Rev. A **87**, 042310 (2013). DOI 10.1103/PhysRevA.87.042310. URL <http://link.aps.org/doi/10.1103/PhysRevA.87.042310>
21. J. Maziero, F.M. Zimmer, Phys. Rev. A **86**, 042121 (2012). DOI 10.1103/PhysRevA.86.042121. URL <http://link.aps.org/doi/10.1103/PhysRevA.86.042121>
22. A.L. Grimsmo, S. Parkins, B.S.K. Skagerstam, Phys. Rev. A **86**, 022310 (2012). DOI 10.1103/PhysRevA.86.022310. URL <http://link.aps.org/doi/10.1103/PhysRevA.86.022310>
23. J.T. Cai, A. Abliz, Physica A: Statistical Mechanics and its Applications **392**(10), 2607 (2013). DOI <http://dx.doi.org/10.1016/j.physa.2013.01.041>. URL <http://www.sciencedirect.com/science/article/pii/S0378437113000939>
24. L. Qiu, G. Tang, X. qing Yang, A. min Wang, EPL (Europhysics Letters) **105**(3), 30005 (2014). URL <http://stacks.iop.org/0295-5075/105/i=3/a=30005>
25. Y.S. Weinstein, Phys. Rev. A **79**, 012318 (2009). DOI 10.1103/PhysRevA.79.012318. URL <http://link.aps.org/doi/10.1103/PhysRevA.79.012318>
26. Y.S. Weinstein, Phys. Rev. A **82**, 032326 (2010). DOI 10.1103/PhysRevA.82.032326. URL <http://link.aps.org/doi/10.1103/PhysRevA.82.032326>
27. M. Ali, A.R.P. Rau, G. Alber, Phys. Rev. A **81**, 042105 (2010). DOI 10.1103/PhysRevA.81.042105. URL <http://link.aps.org/doi/10.1103/PhysRevA.81.042105>
28. X.M. Lu, J. Ma, Z. Xi, X. Wang, Phys. Rev. A **83**, 012327 (2011). DOI 10.1103/PhysRevA.83.012327. URL <http://link.aps.org/doi/10.1103/PhysRevA.83.012327>
29. Q. Chen, C. Zhang, S. Yu, X.X. Yi, C.H. Oh, Phys. Rev. A **84**, 042313 (2011). DOI 10.1103/PhysRevA.84.042313. URL <http://link.aps.org/doi/10.1103/PhysRevA.84.042313>
30. Y. Huang, Phys. Rev. A **88**, 014302 (2013). DOI 10.1103/PhysRevA.88.014302. URL <http://link.aps.org/doi/10.1103/PhysRevA.88.014302>
31. V. Vedral, Rev. Mod. Phys. **74**, 197 (2002). DOI 10.1103/RevModPhys.74.197. URL <http://link.aps.org/doi/10.1103/RevModPhys.74.197>
32. K. Modi, T. Paterek, W. Son, V. Vedral, M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010). DOI 10.1103/PhysRevLett.104.080501. URL <http://link.aps.org/doi/10.1103/PhysRevLett.104.080501>
33. M. Okrasa, Z. Walczak, Europhys. Lett. **96**(6), 60003 (2011). URL <http://stacks.iop.org/0295-5075/96/i=6/a=60003>
34. G. Vidal, R.F. Werner, Phys. Rev. A **65**, 032314 (2002). DOI 10.1103/PhysRevA.65.032314. URL <http://link.aps.org/doi/10.1103/PhysRevA.65.032314>
35. C. Sabín, G. García-Alcaine, The European Physical Journal D **48**(3), 435 (2008). DOI 10.1140/epjd/e2008-00112-5. URL <http://dx.doi.org/10.1140/epjd/e2008-00112-5>
36. V. Coffman, J. Kundu, W.K. Wootters, Phys. Rev. A **61**, 052306 (2000). DOI 10.1103/PhysRevA.61.052306. URL <http://link.aps.org/doi/10.1103/PhysRevA.61.052306>

37. W. Dür, G. Vidal, J.I. Cirac, Phys. Rev. A **62**, 062314 (2000). DOI 10.1103/PhysRevA.62.062314. URL <http://link.aps.org/doi/10.1103/PhysRevA.62.062314>
38. T. Yu, J.H. Eberly, Phys. Rev. Lett. **93**, 140404 (2004). DOI 10.1103/PhysRevLett.93.140404. URL <http://link.aps.org/doi/10.1103/PhysRevLett.93.140404>
39. G.L. Giorgi, S. Campbell, arXiv:1409.1021 (2014)
40. B. Ye, Y. Liu, J. Chen, X. Liu, Z. Zhang, Quantum Information Processing **12**(7), 2355 (2013). DOI 10.1007/s11128-013-0531-y. URL <http://dx.doi.org/10.1007/s11128-013-0531-y>
41. L. Zhao, X. Hu, R.H. Yue, H. Fan, Quantum Information Processing pp. 1–13 (2013). DOI 10.1007/s11128-013-0525-9. URL <http://dx.doi.org/10.1007/s11128-013-0525-9>
42. S. Hamieh, R. Kobes, H. Zaraket, Phys. Rev. A **70**, 052325 (2004). DOI 10.1103/PhysRevA.70.052325. URL <http://link.aps.org/doi/10.1103/PhysRevA.70.052325>