

ACTA BIOMEDICA SUPPLEMENT

ATENEI PARMENSIS | FOUNDED 1887

*Official Journal of the Society of Medicine and Natural Sciences of Parma
and Centre on health systems' organization, quality and sustainability, Parma, Italy*

*The Acta Biomedica is indexed by Index Medicus / Medline Excerpta Medica (EMBASE),
the Elsevier BioBASE*

HEALTH PROFESSIONS (2-2017)

Free on-line www.actabiomedica.it

MATTIOLI 1885



ACTA BIO MEDICA

ATENEI PARMENSIS

FOUNDED 1887

OFFICIAL JOURNAL OF THE SOCIETY OF MEDICINE AND NATURAL SCIENCES OF PARMA
AND CENTRE ON HEALTH SYSTEM'S ORGANIZATION, QUALITY AND SUSTAINABILITY, PARMA, ITALY

free on-line: www.actabiomedica.it

HONORARY EDITOR

Loris Borghi - Parma, Italy

EDITOR IN CHIEF

Maurizio Vanelli - Parma, Italy

ASSOCIATE EDITORS

Antonio Mutti - Parma, Italy
Carlo Signorelli - Parma, Italy
Marco Vitale - Parma, Italy

SECTION EDITORS

Gianfranco Cervellini - Parma, Italy
Vincenzo De Sanctis - Ferrara, Italy
Carlo Signorelli - Parma, Italy

DEPUTY EDITOR FOR HEALTH PROFESSIONS EDITION

Leopoldo Sarli - Parma, Italy

DEPUTY EDITOR FOR SERTOT EDITION

Francesco Pogliacomì - Parma, Italy

EDITORIAL BOARD

Franco Aversa - Parma, Italy
Alberto Bacchi Modena - Parma, Italy
Cesare Beghi - Varese, Italy
Roberto Berretta - Parma, Italy
Corrado Betterle - Padova, Italy
Riccardo Bonadonna - Parma, Italy
Mauro Bonanini - Parma, Italy
Loris Borghi - Parma, Italy
David A. Bushinsky - Rochester, NY, USA
Ovidio Bussolati - Parma, Italy
Ardeville Cabassi - Parma, Italy
Paolo Caffarra - Parma, Italy
Duran Canatan - Antalya, Turkey
Fausto Catena - Parma, Italy
Francesco Ceccarelli - Parma, Italy
Rossana Cecchi - Parma, Italy
Stefano Cecchini - Parma, Italy
Gian Paolo Ceda - Parma, Italy
Graziano Ceresini - Parma, Italy
Gianfranco Cervellini - Parma, Italy
Alfredo Antonio Chetta - Parma, Italy
Marco Colonna - St. Louis, MO, USA
Paolo Coruzzi - Parma, Italy
Lucio Guido Maria Costa - Parma, Italy
Cosimo Costantino - Parma, Italy
Domenico Cucinotta - Bologna, Italy
Alessandro De Fanti - Reggio Emilia, Italy
Massimo De Filippo - Parma, Italy

Filippo De Luca - Messina, Italy
Vincenzo De Sanctis - Ferrara, Italy
Giuseppe Fabrizi - Parma, Italy
Valentina Fainardi - Parma, Italy
Claudio Feliciani - Parma, Italy
Nicola Florindo - Parma, Italy
Lorella Franzoni - Parma, Italy
Antonio Freyrie - Parma, Italy
Vittorio Gallese - Parma, Italy
Livio Garattini - Milano, Italy
Matteo Goldoni - Parma, Italy
Donald J. Hagler - Rochester, MINN, USA
Rick Hippakka - Chicago, IL, USA
Andrew R. Hoffman - Stanford, CA, USA
Joachim Klosterkoetter - Colonia, Germany
Ronald M. Lechan - Boston, MA, USA
Annarosa Leri - Harvard, Boston, MA, USA
Giuseppe Lippi - Verona, Italy
Nicola Longo - Salt Lake City, UT, USA
Wanyun Ma - Beijing, China
Umberto Vittorio Maestroni - Parma, Italy
Marcello Giuseppe Maggio - Parma, Italy
Norman Maitland - York, United Kingdom
Federico Marchesi - Parma, Italy
Carla Mastroilli - Parma, Italy
James A. McCubrey - Greenville, NC, USA
Tiziana Meschi - Parma, Italy
Giuseppe Nuzzi - Parma, Italy

Jose Luis Navia - Cleveland, OH, USA
Anna Odone - Parma, Italy
Donald Orlic - Bethesda, MD, USA
Silvia Pizzi - Parma, Italy
Francesco Pogliacomì - Parma, Italy
Federico Quaini - Parma, Italy
Edoardo Raposio - Parma, Italy
Stephen M. Rao - Cleveland, OH, USA
Luigi Roncoroni - Parma, Italy
Shaukat Sadikot - Mumbai, India
Simone Cherchi Sanna - New York, NY, USA
Leopoldo Sarli - Parma, Italy
Robert S. Schwartz - Denver, Colorado, USA
Anthony Seaton - Edinburgh,
United Kingdom
Ashraf Tawfic Mohamed Soliman - Doha, Qatar
Mario Strazzabosco - New Haven, CT, USA
Nicola Sverzellati - Parma, Italy
Maria Luisa Tanzi - Parma, Italy
Roberto Toni - Parma, Italy
Frederik H. Van Der Veen - Maastricht,
The Netherlands
Vincenzo Vincenti - Parma, Italy
Vincenzo Violi - Parma, Italy
Richard Wallensten - Solna, Sweden
Francesco Ziglioli - Reggio Emilia, Italy

LINGUISTIC ADVISOR

Rossana Di Marzio
Parma, Italy

EDITORIAL OFFICE MANAGER

Anna Scotti
Mattioli 1885 srl - Casa Editrice
Strada di Lodesana 649/sx, Loc. Vaio
43036 Fidenza (PR), Italy
Tel. ++39 0524 530383
Fax ++39 0524 82537
contact@actabiomedica.it

Francesco Covino
Società di Medicina e Scienze Naturali
Azienda Ospedaliero-Universitaria
di Parma - Cattani Building, 2nd floor
Via Gramsci, 14 - Parma, Italy
Tel./Fax ++39 0521 033730
francesco.covino@unipr.it

PUBLISHER

Mattioli 1885 srl Casa Editrice
Strada di Lodesana, 649/sx, Loc. Vaio
43036 Fidenza (PR), Italy
Tel. ++39 0524 530383
Fax ++39 0524 82537
E-mail: edit@mattioli1885.com

HEALTH PROFESSIONS

ACTA BIO MEDICA

EDITOR EXECUTIVE

Leopoldo Sarli - Parma, Italy

DEPUTY EDITORS

Giovanna Artioli - Parma, Italy (nursing topics)

Enrico Bergamaschi - Parma, Italy (prevention topics)

Tiziana Mancini - Parma, Italy (psychosocial topics)

Angelo Mastrillo - Bologna, Italy (diagnostic and rehabilitative topics)

EDITORIAL BOARD

Rodolfo Brianti - Parma, Italy

Rachele La Sala - Parma, Italy

Nicola Parenti - Imola, Italy

Bui Vu Binh - Hanoi, Vietnam

Roberto Lusardi - Bergamo, Italy

Enrico Pasanisi - Parma, Italy

Adriana Calderaro - Parma, Italy

Claudio Macaluso - Parma, Italy

Giovanni Pavesi - Parma, Italy

Luca Caricati - Parma, Italy

Tiziana Mancini - Parma, Italy

Vincenza Pellegrino - Parma, Italy

Franco Carnevale - Montreal, Canada

Sergio Manghi - Parma, Italy

Diletta Priami - Bologna, Italy

Matteo Castaldo - Parma, Italy

Gemma Mantovani - Parma, Italy

Cristina Rossi - Parma, Italy

Luigi Cavanna - Piacenza, Italy

Ardigò Martino - Bologna, Italy

Annavittoria Sarli - Milano, Italy

Francesco Chiampo - Parma, Italy

Giuliana Masera - Piacenza, Italy

Loredana Sasso - Genova, Italy

Cosimo Costantino - Parma, Italy

Maria Messerli Ernst - Berna,

Chiara Scivoletto - Parma, Italy

Renato Costi - Parma, Italy

Switzerland

Alberto Spisni - Parma, Italy

Pham Huy Dung - Hanoi, Vietnam

Nadia Monacelli - Parma, Italy

Angelo Stefanini - Bologna, Italy

Paola Ferri - Modena, Italy

Federico Monaco - Bergamo, Italy

Laura Tibaldi - Piacenza, Italy

Laura Fieschi - Parma, Italy

Maria Mongardi - Bologna, Italy

Stefano Tomelleri - Bergamo, Italy

Chiara Foà - Parma, Italy

Cecilia Morelli - Parma, Italy

Annalisa Tonarelli - Parma, Italy

Laura Fruggeri - Parma, Italy

Mamadou Ndiaye - Dakar, Senegal

Giancarlo Torre - Genova, Italy

LINGUISTIC ADVISOR

Rossana Di Marzio
Parma, Italy

EDITORIAL OFFICE MANAGER

Anna Scotti
Mattioli 1885 srl - Casa Editrice
Strada di Lodesana 649/sx, Loc. Vaio
43036 Fidenza (PR), Italy
Tel. ++39 0524 530383
Fax ++39 0524 82537
E-mail: contact@actabiomedica.it

Francesco Covino
Società di Medicina e
Scienze Naturali
Office of the Faculty of Medicine
Via Gramsci, 14 - Parma, Italy
Tel./Fax ++39 0521 033730

PUBLISHER

Mattioli 1885 srl Casa Editrice
Strada di Lodesana, 649/sx, Loc. Vaio
43036 Fidenza (PR), Italy
Tel. ++39 0524 530383
Fax ++39 0524 82537
E-mail: edit@mattioli1885.com



MATTIOLI 1885

srl- Strada di Lodesana 649/sx
43036 Fidenza (Parma)
tel 0524/530383
fax 0524/82537
www.mattioli1885.com

Direttore Generale
Paolo Cioni
Direttore Scientifico
Federico Cioni

Direttore Commerciale
Marco Spina

Formazione/ECM
Simone Agnello

Project Manager
Natalie Cerioli
Massimo Radaelli

Editing Manager
Anna Scotti

Editing
Valeria Ceci

Foreign Rights
Nausicaa Cerioli

Distribuzione
Massimiliano Franzoni



EXECUTIVE COMMITTEE OF THE SOCIETY OF MEDICINE AND NATURAL SCIENCES OF PARMA

Honorary President
Loris Borghi
President

Maurizio Vanelli
Past-President
Almerico Novarini
General Secretary
Maria Luisa Tanzi

Treasurer
Riccardo Volpi
Members

O. Bussolati	A. Mutti
G. Ceda	P. Muzzetto
G. Cervellin	P. Salcuni
G. Ceresini	L. Sarli
N. Florindo	V. Vincenti
G. Luppino	V. Violi
A. Melpignano	M. Vitale

INDEX

Volume 88 / Suppl. 3-2017

July 2017

Health Professions

Special Issue (2-2017)

- ORIGINAL ARTICLE: ON THE PATIENT'S SIDE: FOCUS ON LEUKAEMIA**
- 5 *Stefano Botti, Gianpaolo Gargiulo, Felice Bombaci, Giovanna Artioli, Chiara Cosentino, Adriana Concetta Pignatelli, Daniela Torino, Maria Marcella Lionetti, Emanuela Samarani, Lorella Cappucciati, Paola Bordiga, Antonella Diodati, Cristiana Caffarri, Irene Rosini, Fabrizio Pane*
Clinical pathway for patients with Chronic Myeloid Leukaemia: The Euriclea Project
- 13 *Giovanna Artioli, Chiara Foà, Chiara Cosentino, Alfonso Sollami, Chiara Taffurelli*
Integrated narrative assessment exemplification: a leukaemia case history
- ORIGINAL ARTICLE: NURSES: INSIGHTS ON NURSING STUDENTS**
- 22 *Paola Ferri, Sergio Rovesti, Nunzio Panzera, Luigi Marcheselli, Alessia Bari, Rosaria Di Lorenzo*
Empathic attitudes among nursing students: a preliminary study
- 31 *Chiara Taffurelli, Alfonso Sollami, Carmen Camera, Francesca Federa, Annise Grandi, Marcella Marino, Tiziano Marrosu, Leopoldo Sarli*
Healthcare associated infection: good practices, knowledge and the locus of control in healthcare professionals
- 37 *Giuseppe Marletta, Leopoldo Sarli, Luca Caricati, Tiziana Mancini*
Intergroup contact and team functioning among nursing students: the mediation role of intergroup anxiety
- ORIGINAL ARTICLE: ANALYSIS OF THE ORGANIZATIONAL ENVIRONMENT**
- 43 *Maura Galletta, Igor Portoghese, Cesar Ivan Aviles Gonzales, Paola Melis, Gabriele Marcias, Marcello Campagna, Luigi Minerba, Claudia Sardu*
Lack of respect, role uncertainty and satisfaction with clinical practice among nursing students: the moderating role of supportive staff
- 51 *Emanuela Alfieri, Alessia Alebbi, M. Giovanna Bedini, Laura Boni, Chiara Foà*
Mapping the nursing competences in neonatology: a qualitative research
- 59 *Chiara Cosentino, Morena Bettuzzi, Giulia Campioli, Valeria Di Marco, Giulia Giacomuzzi, Ivana Marinoni, Lorenzo Orlandini, Angelo Palermo, Siria Pattacini, Giovanna Artioli*
Individual and social variables and their effect on Case/Care Manager Job Satisfaction: an exploratory study
- 67 *Emanuela Alfieri, Marina Mori, Valentina Barbuì, Leopoldo Sarli*
Advanced competencies mapping of critical care nursing: a qualitative research in two Intensive Care Units

Empathic attitudes among nursing students: a preliminary study

Paola Ferri¹, Sergio Rovesti¹, Nunzio Panzera¹, Luigi Marcheselli¹, Alessia Bari¹,
Rosaria Di Lorenzo²

¹Department of Diagnostic, Clinical and Public Health Medicine, University of Modena and Reggio Emilia; ²Department of Mental Health, Local Health Unit (AUSL), Modena, Italy

Abstract. *Background and aim:* An empathic approach is fundamental for therapeutic relationship between nurse and patient. According to some researchers, female nursing students show higher empathic attitude in comparison with males, but both show a decline in empathy level as their studies progress. This preliminary study evaluated the self-reported emotional empathy level among undergraduate students at first and second year of nursing 3-year course. *Method:* To assess empathy level, the Balanced Emotional Empathy Scale (BEES) was administered to all students enrolled in the 2015/16 academic year (N=142), at the beginning of first year (T0) and at mid-point of second year (T1) of nursing course. Data were statistically analyzed. *Results:* 118 nursing students participated in the first and 99 in the second survey. The BEES global mean score for the longitudinal group (n=99) slightly decreased from T0 (mean=37.1±19.5 SD) to T1 (mean=33.5±22.6 SD) (t=1.20, p=0.23; t-test for paired data). Female students reported a statistically significant higher mean BEES score compared to male students in both surveys. *Conclusions:* Our preliminary data suggest a slight decline in empathy level among nursing students with the progress of study course, in accordance with previous studies. In particular, our study shows higher levels of empathy in female students and lower levels in male students, compared to other studies. Further surveys aimed at investigating the empathy attitude at the end of nursing course could confirm the decline tendency reported by this preliminary study. Other research focusing on the causes of empathy decline are necessary to explain this phenomenon.

Key words: preliminary study, empathy, undergraduate nursing students, gender differences

Introduction

Empathy and nursing

If empathy is defined as the ability to “put oneself in others’ shoes and [to] make their experiences one’s own”, it represents an individual’s ability to understand and share another person’s thoughts and feelings (1-6). Empathy influences how we interact with others. It is a process, whereby an empathic listener is able to put his or her own way of perceiving the world into the background in order to better feel and understand

the experiences and perceptions of the speaker (6). A growing number of authors view empathy as a multi-dimensional construct, composed of affective, moral, cognitive, relationship-related and behavioural dimensions (7-12). Although the scientific community recognises the presence of cognitive, affective and behavioural components in the description and measurement of empathy, the emotional aspect remains its cornerstone (4). The emotional empathy, in particular, is a reflection of positive interpersonal relationships and social skills, a pro-social orientation, and engagement in altruistic behaviour (13).

In recent years, the nursing profession has been impacted by growth in scientific knowledge and technological advances. Nonetheless, the essence of the profession is and continues to be a helping relationship with the person receiving care (14). An empathic approach is fundamental to the development of the therapeutic relationship between nurse and patient (15, 16) and, of all of the attitudes that are most desirable in a nurse, empathy is the most important (17). The objective of the nurse-patient relationship is to ensure that the needs of the patient are identified and that the necessary therapies and care are provided. In order to identify these needs, nurses should be capable of fully understanding their patients' feelings, opinions and conditions. All of these things require a high capacity for empathy (18).

A good empathetic capacity in nurses has been linked to greater patient well-being and satisfaction, better patient compliance, and a decrease in errors and complications, not to mention better health outcomes (19-26). It may be that an empathetic approach can result in a shorter treatment period or a reduced demand on resources (27). A lack of empathy, on the other hand, may interfere negatively with the diagnosis, treatment and care processes (14). According to Doyle et al. (17) a review of cases presented to the nurses' disciplinary tribunal in New South Wales (Australia) suggests that the majority of complaints against nurses in this jurisdiction are the result of callousness or lack of empathy. In addition, empathy is associated with lower levels of burnout among nurses and nursing students (28-30) and higher professional satisfaction (14, 31).

Empathic attitudes of undergraduate nursing students

The ability to empathize can be influenced by many factors: gender, age, job training and experience (32). Many studies have shown that women demonstrate greater empathic tendencies in comparison to men (27, 33-35). An Italian longitudinal study, which assessed the efficacy of a specific training course for improving empathy skills in nursing students, highlighted that the training was more effective for the female students than for their male counterparts (36). Another Italian study revealed that the impact of gender on empathetic tendency increased during nursing training,

as demonstrated by the higher Balanced Emotional Empathy Scale (BEES) scores of female students with respect to male students (37). Although empathic skill is a mutually beneficial element in the relationship between nurse and patient (20, 21), according to some researchers nursing students nonetheless demonstrate low to moderate levels of it (21, 30, 35, 37-39). As far as concerns the development of the empathic tendency during the course of nursing studies, the literature offers conflicting results. The cross-sectional studies conducted in Australia by Williams et al. (27) and by McKenna et al. (40) recorded no statistically significant differences between the mean levels of empathy among students from different course years. Three transversal studies, however, which compared a cohort of students at the beginning of their studies with another about to conclude them (34, 41, 42), revealed a "decline" in empathy. Ward also observed, at the Jefferson School of Nursing, a decrease in empathy within the same cohort, at the end of one course year in comparison with its beginning (43). Similar findings were also recorded among medical students and students of other health professions (41, 44-46). Lastly, a recent cross-temporal meta-analysis, conducted by Konrath et al. on 72 samples of American college students, revealed a decline in perspective taking and empathic concern, accompanied, on the other hand, by a rise in narcissistic and individualistic tendencies (47).

Unfortunately, the literature that examines empathy levels in nursing students is often inconsistent, because it is characterised by cross-sectional research and small samples of participants. Longitudinal studies should therefore be encouraged, in order to measure the development of empathy over the course of their studies and analyse the effect of communication skills training on the empathy of nursing students (30, 34). We believe that such an assessment could help us not only to recognise the changing trend, but also to better understand when it becomes necessary to create learning activities that can help students apply empathy or halt its decline.

Aim

The aim of this study is to evaluate the self-reported emotional empathy level among undergraduate students at first and second year of nursing 3-year course.

Methods

Study design

We have carried out the first stage of a three-year longitudinal study whose objective is to assess the development of emotional empathy among nursing students enrolled in the Modena Nursing Degree Programme at the University of Modena and Reggio Emilia, Italy.

Procedures

In order to assess the empathy level among nursing students, two subsequent surveys were scheduled: one at the beginning of their course of study (T0 at 30 October 2015) and one at the mid-point (T1 at 1 March 2017). The empathy level evaluation of this preliminary study will extend to the third year of nursing course (T2 at 30 October 2018), among the same student group, in order to obtain more complete results.

Participants

From a sample composed of all students enrolled in the 2015/16 academic year of nursing course (n=142), 118 students, who participated in the first evaluation (T0), represented our initial group, of whom only 99 participated in the second assessment (T1). This second group (n=99) represented our longitudinal sample, whose BEES scores at T0 and T1 were pairing compared.

Measures

The instrument that we have chosen to assess levels of empathy, intended as a susceptibility to becoming vicariously involved in others' emotional feelings and the tendency to develop positive interpersonal relationships, is the BEES, which has already been used in studies conducted on samples of Italian nursing students (30, 36, 37). BEES is a self-report instrument validated in the Italian language (48, 49). It includes 30 items about which participants express their level of agreement/disagreement on a seven-point Likert scale, with negative and positive answers, designed to avoid

social desirability in the responses. The Italian version of the BEES validated five dimensions that deal with the following areas of emotional empathy (50):

- D1 'Impermeability to the emotional feelings of others', "7 items referring to situations in which the respondent is unwilling to become emotionally involved in another person's feelings (e.g., "I am not affected easily by the strong emotions of people around me"; Cronbach's $\alpha=0.74$ ");
- D2 'Susceptibility to the emotional feelings of others', "6 items tapping the respondent's willingness to become involved in others' feelings and share their suffering (e.g., "I get a strong urge to help when I see someone in distress"; Cronbach's $\alpha=0.66$ ");
- D3 'Emotional spread responsiveness', "7 items referring to the respondent's tendency to identify with characters in films, plays, stories, etc. (e.g., "I don't get caught up easily in the emotions generated by a crowd"; Cronbach's $\alpha=0.68$ ");
- D4 'Susceptibility to emotional involvement with people nearby', "6 items tapping the respondent's feelings shared in the presence of others who are suffering (e.g., "It upsets me to see someone being mistreated"; Cronbach's $\alpha=0.62$ ");
- D5 'Tendency to avoid emotional involvement with fragile people', "4 items reflecting the respondent's tendency to avoid becoming emotionally involved with fragile or vulnerable people like children or the elderly (e.g., "Helpless old people don't have much of an emotional effect on me"; Cronbach's $\alpha=0.55$ ").

The Cronbach's α coefficient for all 30 items ranges from 0.83 to 0.87, according to most authors (49, 50). High scores at D1, D3 and D5 indicate scarce capacity to empathize, while low scores at D2 and D4 indicate a good empathetic tendency. The total BEES score indicates high levels of empathy if it is greater than the mean value of 32 ± 18 (Standard Deviation).

The questionnaire also asked students their gender and age, whilst the date of the questionnaire's completion and course year were filled in each form. The questionnaires were distributed in the classroom and the students were given the time necessary to be able to fill them out in full. All of the information collected

during the study was kept anonymous. However, each student was asked to insert an identification code that only he/she would recognise in order to allow the pairing of first (T0) with the second (T1) questionnaire.

Ethical considerations

The study's purpose and methods were described to the students by the principal investigator. Participants' anonymity and confidentiality were assured and their decision to participate voluntarily in this study was respected. The students were assured that neither the information obtained through administration of the scale nor a failure to participate in the study would have any impact on their course of study. The study was authorized by the Director of the Nursing Degree Programme.

Statistical analyses

The statistical analysis was performed using the software Stata (v12 StataCorp LP, College Station, TX). Continuous variables were reported as arithmetic means and standard deviations (SD).

We computed a total BEES score (reflecting emotional empathy) and five subscale scores (dimensions), in accordance with the indications of the authors who adapted and validated the BEES Italian version (48, 49). The t-test for paired and unpaired data was used to compare the two mean scores of BEES at T0 and T1; the Wilcoxon rank test for paired data was used for the medians. We considered statistical significance to have been attained if $p < 0.05$.

Results

The students who agreed to participate in this study and completed the questionnaire at T0 were 83% of our sample (118/142 students). Of these 118 student participants, 79% were female and 21% male (the imbalance between females and males reflects the distribution of participants). The mean age of students at T0 was 20.20 ± 2.58 (SD) years. 99 of the 118 students who had previously completed the questionnaire at T0, participated in the second BEES administration (the other 19 students were not present at T1 because they were not attending the second year of Nursing course). The mean age of students at T1 was 21.29 ± 2.82 (SD) years.

The empathic tendency

At T0, the nursing students in the initial group ($n=118$) demonstrated a total mean empathy score of 37.0 ± 19.5 SD and mean empathy scores for each dimension, as shown in Table 1. At the second administration (T1), the total mean score recorded in the longitudinal group ($n=99$) was 33.5 ± 22.6 SD.

If we compare the total empathy at T0 with that at T1 for the initial group ($n=118$) and the longitudinal group ($n=99$), we can see that the empathy in the 2nd year has decreased by 3.5 on average. The comparison between T0 and T1 did not evidence any statistically significant difference ($t=1.220$, $p=0.224$; t-test for unpaired data). The mean scores of total and each dimension BEES, registered in the longitudinal group ($n=99$) at T0 and T1, are shown in Table 2.

Table 1. Total empathy and empathy by dimension for the initial and longitudinal groups

BEES Scores	Initial group n=118 T0 Mean (SD)	Longitudinal group n=99 T1 Mean (SD)
D1: Impermeability to the emotional feelings of others	-7.5 (6.0)	-6.0 (7.0)
D2: Susceptibility to the emotional feelings of others	10.7 (4.0)	10.6 (4.7)
D3: Emotional spread responsiveness	-0.60 (5.3)	-1.5 (8.9)
D4: Susceptibility to emotional involvement with people nearby	10.9 (4.9)	10.9 (4.9)
D5: Tendency to avoid emotional involvement with fragile people	-3.3 (2.6)	-2.4 (3.0)
Total	37.0 (19.5)	33.5 (22.6)

The decrease in empathy in the 2nd year of the course was also confirmed by a comparison of the longitudinal group's scores at T0 and T1 (Table 2), although the difference between the two mean values was not statistically significant ($t=1.202$, $p=0.232$; t -test for paired data). The median values of BEES score decreased at T1 in comparison with T0, showing a decline in empathetic tendency (Table 3). Application of the Wilcoxon rank test for paired data did not evidence any statistically significant difference.

The gender difference

The analysis of the first survey (T0) highlighted lower empathy mean scores among males in all BEES dimensions in comparison with females (Table 4). This difference was confirmed by the total BEES empathy score, where males were found to have a mean

score of 17.4 ± 17.5 SD and females one of 42.7 ± 15.9 SD, with a statistically significant difference ($t=6.775$, $p<0.0001$; t -test).

The data from the second administration at T1 confirmed the statistically significant difference between the two genders in total and dimension BEES scores ($t= 5.426$, $p<0.0001$; t -test) (Table 5).

Discussion

This research analysed the empathic tendency among nursing students and its change during the progress of the nursing course. In our study, the mean score of empathetic tendency was found to be slightly higher (37.1 ± 19.5 SD) among the students at the beginning (T0) of their nursing course if compared to the standardized samples identified by the Italian BEES

Table 2. Total empathy and empathy by dimension for the longitudinal group, at T0 and T1

BEES Scores	Longitudinal group	Longitudinal group
	n=99 T0 Mean (SD)	n=99 T1 Mean (SD)
D1: Impermeability to the emotional feelings of others	-7.4 (5.9)	-6.0 (7.0)
D2: Susceptibility to the emotional feelings of others	10.8 (4.1)	10.6 (4.7)
D3: Emotional spread responsiveness	-0.69 (5.3)	-1.45 (8.9)
D4: Susceptibility to emotional involvement with people nearby	11.0 (4.9)	10.9 (4.9)
D5: Tendency to avoid emotional involvement with fragile people	-3.1 (2.5)	-2.4 (3.0)
Total	37.1 (19.5)	33.5 (22.6)

Table 3. Median empathy values for the longitudinal group, at T0 and T1

N=99	Median (2.5-97.5 percentile)		p-value
	T0	T1	
D1: Impermeability to the emotional feelings of others	-7.5 (-20; 3)	-6 (-18; 9)	0.188
D2: Susceptibility to the emotional feelings of others	11 (3; 18)	11 (-1.5; 18)	0.790
D3: Emotional spread responsiveness	-1 (-10; 11)	-2 (-20; 17)	0.470
D4: Susceptibility to emotional involvement with people nearby	11 (-3; 18)	11 (1; 18)	0.976
D5: Tendency to avoid emotional involvement with fragile people	-3 (-8; 2)	-3 (-7.5; 5)	0.102
Total	39 (-15; 71)	34 (-15; 71)	0.325

Table 4. Mean empathy scores at T0 for the initial group, separated by gender

Score	Gender (n=118)		p-value
	Male	Female	
D1: Impermeability to the emotional feelings of others	-3.0 (5.3)	-8.7 (5.5)	<0.0010
D2: Susceptibility to the emotional feelings of others	7.4 (3.6)	11.6 (3.7)	<0.0010
D3: Emotional spread responsiveness	3.2 (4.5)	-1.7 (5.0)	<0.0010
D4: Susceptibility to emotional involvement with people nearby	8.0 (5.5)	11.8 (4.6)	0.0070
D5: Tendency to avoid emotional involvement with fragile people	-1.9 (2.7)	-3.7 (2.4)	0.0019
Total	17.4 (17.5)	42.7 (15.9)	<0.0001

Table 5. Mean empathy scores at T1 for the longitudinal group, separated by gender

Score	Gender (n=99)		p-value
	Male	Female	
D1: Impermeability to the emotional feelings of others	-2.3 (6.4)	-7.2 (6.8)	0.0018
D2: Susceptibility to the emotional feelings of others	7.6 (5.8)	11.6 (3.8)	0.0002
D3: Emotional spread responsiveness	3.1 (5.5)	-1.2 (5.9)	0.0018
D4: Susceptibility to emotional involvement with people nearby	8.3 (4.2)	11.8 (4.8)	0.0011
D5: Tendency to avoid emotional involvement with fragile people	-1.5 (3.7)	-2.7 (2.6)	0.0073
Total	15.0 (20.3)	40.0 (20.0)	<0.0001

study validation: adults between 20 and 24 years of age (25.5 ± 21.0 SD) and nurses (32.0 ± 18.0 SD) (49). Our result indicates an higher level of empathy if compared to other recent Italian studies conducted among nursing students (36, 37). In this regard, the literature put in evidence that the undergraduate nursing students generally show a significantly higher mean score of empathy than the students attending other undergraduate courses (26, 41, 51). This could be explained by the fact that students who choose to attend nursing course probably have a particular aptitude or motivation for helping relationship, which represents a key aspect of the nursing profession (26, 52, 53).

At the time of our second survey (T1), the mean empathy score appeared slightly decreased in comparison with the beginning of study course, although no statistically significant difference came to light. There are no longitudinal studies of empathy among Italian nursing students with which we can compare this result. The comparison with the international studies, which have analysed this issue, is not reliable because they were

conducted using different evaluation instruments and in different educational contexts. With these precautions, we highlight that our findings are in line with those of other studies of health-science and nursing courses that have reported a decline in empathy over the course of study (34, 41-46). Nunes et al., when comparing mean empathy scores at the beginning and at the end of the first academic year, found that students in all five health-science disciplines showed a decrease in empathy scores from the beginning to the end of school, with a statistically significant decline for dental, nursing, and medical students (41). In addition, also a cross-sectional study conducted within the university nursing programme in Seville revealed a progressive decline in empathy as the course of study progressed (34). This is in contrast with the studies conducted by Williams et al. (27, 51), and McKenna et al. (40) in Australian nursing schools, which found no decrease in empathy.

Nunes et al. suggested that the decline in empathy scores during the first year of training is in part due to a 'settling in' effect. Whereas, on entry, the

students are eager to show that they have the positive attributes of a compassionate healthcare provider, their idealism becomes less evident as the year progresses (41). Cunico et al., meanwhile, hypothesized that, since Evidence Based Practice focuses heavily on scientific, technological, diagnostic, and therapeutic approaches to healthcare, it therefore leads students to lose their human and empathetic perspective on the patient-provider relationship (36). Finally, in a longitudinal study conducted on 214 undergraduate nursing students who were exposed more than others to patient encounters during the study period, Ward et al. observed a statistically significant decline in empathy (43). Our findings are consistent with most studies which highlighted that medical and nursing students, who have to frequently encounter patients, can show a decline in empathy tendency due to their empathic engagement in the patients' suffering (44, 46).

Our study suggested a gender difference in empathy attitude and tendency, showing that female students reported a statistically significant higher mean BEES empathy score in comparison with male students. This gender difference can be found also in the standard samples: adults between the ages of 20 and 24 (Females 34.0 ± 20.0 SD, Males 16.5 ± 18.5 SD) and nurses (Females 35.0 ± 17.5 SD, Males 19.5 ± 15.5 SD) (49). However, in our sample, this difference was more pronounced (Females 40.0 ± 20.0 SD, Males 15.0 ± 20.3 SD). These results are consistent with those of other studies (27, 30, 33-37, 51), in which women consistently recorded higher empathy scores, probably due to greater emotional resonance to others' feelings and more sensitivity to interpersonal stimuli probably due to biological and social conditioning (30). According to Williams et al., the traditional and evolutionary role of women as caregivers may also explain the variation in empathy level between males and females, indicating that females are more perceptive to emotions and males take a more rational rather than emotive approach with others (27).

Conclusions

Our preliminary data suggest a slight decline in empathy tendency among nursing students between

the beginning and the mid-point of their undergraduate education, confirming other previous studies. Further study extended to the longitudinal study of nursing course is essential to confirm the empathic decline tendency. If a decline in empathy scores is confirmed, further research would be necessary in order to analyse the causes and verify whether this tendency is related to the clinical training, which exposes students to an overwhelming experience with patient suffering.

Limits and advantages

This study has many limits. It was conducted in only one Italian university, so its results cannot be generalised. The observation period should be extended to the total period of nursing course, as our research planned project. Although the scale evaluates a self-report measure of empathy, it is easily and time efficient to administer. Nevertheless, our research study is one of few longitudinal studies, and certainly the only one Italian study, on student empathy. Our longitudinal findings provide important information that could help us to better understand the capacity of students to maintain and develop, during the nursing course, an empathic attitude towards patients, fundamental condition for any therapeutic approach.

References

1. Eisenberg N, Fabes RA, Murphy B, et al. The relations of emotionality and regulation to dispositional and situational empathy-related responding. *J Pers Soc Psychol* 1994; 66: 776-97.
2. Richendoller NR, Weaver JB. Exploring the links between personality and empathic response style. *Pers Individ Dif* 1994; 17: 303-11.
3. Hoffman ML. *Empathy and moral development: Implications for caring and justice*. New York: Cambridge University Press; 2000.
4. Bonino S. Contagio, empatia e comportamento pro-sociale. [Contagiousness, empathy and pro-social behavior]. In Bonino S, Caprara GV. *Il comportamento prosociale. Aspetti individuali, familiari e sociali*. [Prosocial behavior. Individual, family and social aspects]. Trento: Erickson; 2006.
5. Meneghini AM. *Sentire e condividere. Componenti psicologiche e correlati biologici dell'empatia*. [Feel and share. Psychological and biological components of empathy]. Firenze: SEID; 2010.

6. Depalmas C, Ferro Allodola V. L'“inganno” dell'empatia in ambito sanitario. Dall'empatia all'exotopia: dicotomia del senso attraverso l'uso riflessivo dei film. [The “deception” of empathy in healthcare. From empathy to exotopia: dichotomy of meaning through the reflective use of the films]. *Tutor* 2013; 13: 22-31.
7. Reid-Ponte P. Distress in cancer patients and primary nurses' empathy skills. *Cancer Nurs* 1992; 15: 283-92.
8. Olsen DP. Empathetic maturity: theory of moral point of view in clinical relations. *ANS Adv Nurs Sci* 2001; 24: 36-46.
9. Cliffordson C. The hierarchical structure of empathy: dimensional organization and relations to social functioning. *Scand J Psychol* 2002; 43: 49-59.
10. Stepien KA, Baernstein A. Educating for empathy. A review. *J Gen Intern Med* 2006; 21: 524-30.
11. Devoldre I, Davis MH, Verhofstadt LL, Buysse A. Empathy and social support provision in couples: social support and the need to study the underlying processes. *J Psychol* 2010; 144: 259-84.
12. Lobchuk M, Halas G, West C, Harder N, Tursunova Z, Ramraj C. Development of a novel empathy-related video-feedback intervention to improve empathic accuracy of nursing students: A pilot study. *Nurse Educ Today* 2016; 46: 86-93.
13. Mehrabian A, Epstein N. A measure of emotional empathy. *J Person* 1972; 40: 525-43.
14. Ozcan CT, Oflaz F, Sutcu Cicek H. Empathy: the effects of undergraduate nursing education in Turkey. *Int Nurs Rev* 2010; 57: 493-9.
15. Reynolds WJ, Scott B, Jessiman WC. Empathy has not been measured in clients' terms or effectively taught: a review of the literature. *J Adv Nurs* 1999; 30: 1177-85.
16. McCabe C. Nurse-patient communication: an exploration of patients' experiences. *J Clin Nurs* 2004; 13: 41-9.
17. Doyle K, Hungerford C, Cruickshank M. Reviewing Tribunal cases and nurse behaviour: putting empathy back into nurse education with Bloom's taxonomy. *Nurse Educ Today* 2014; 34: 1069-73.
18. Hemsley B, Sigafos J, Balandin S, et al. Nursing the patient with severe communication impairment. *J Adv Nurs* 2001; 35: 827-35.
19. La Monica EL, Wolf RM, Madea AR, Oberst MT. Empathy and nursing care outcomes. *Sch Inq Nurs Pract* 1987; 1: 197-213.
20. Allgood MR. Empathy: The importance of recognizing two types. *J Psychosoc Nurs Ment Health Serv* 1992; 30: 14-7.
21. Reynolds WJ, Scott B. Do nurses and other professional helpers normally display much empathy? *J Adv Nurs* 2000; 31: 226-34.
22. Yu J, Kirk M. Measurement of empathy in nursing research: systematic review. *J Adv Nurs* 2008; 64: 440-54.
23. Del Canale S, Louis DZ, Maio V, et al. The relationship between physician empathy and disease complications: an empirical study of primary care physicians and their diabetic patients in Parma, Italy. *Acad Med* 2012; 87: 1243-9.
24. Gleichgerrcht E, Decety J. Empathy in clinical practice: how individual dispositions, gender, and experience moderate empathic concern, burnout, and emotional distress in physicians. *PLoS One* 2013; 8: e61526.
25. Hojat M. Empathy in patient care: Antecedents, development, measurement, and outcomes. New York: Springer; 2016.
26. Petrucci C, La Cerra C, Aloisio F, Montanari P, Lancia L. Empathy in health professional students: A comparative cross-sectional study. *Nurse Educ Today* 2016; 41: 1-5.
27. Williams B, Brown T, McKenna L, et al. Empathy levels among health professional students: a cross-sectional study at two universities in Australia. *Adv Med Educ Pract* 2014; 5: 107-13.
28. Aström S, Nilsson M, Norberg A, Winblad B. Empathy, experience of burnout and attitudes towards demented patients among nursing staff in geriatric care. *J Adv Nurs* 1990; 15: 1236-44.
29. Wilczek-Rużyczka E. Empathy vs. professional burnout in health care professionals. *J US China Med Sci* 2011; 8: 526-32.
30. Ferri P, Guerra E, Marcheselli L, Cunico L, Di Lorenzo R. Empathy and burnout: an analytic cross-sectional study among nurses and nursing students. *Acta Biomed* 2015; 86 Suppl 2: 104-15.
31. Norfolk T, Birdi K, Walsh D. The role of empathy in establishing rapport in the consultation: a new model. *Med Educ* 2007; 41: 690-7.
32. Adriaansen M, Van Achterberg T, Borm G. The usefulness of the Staff-Patient Interaction Response Scale for palliative care nursing for measuring the empathetic capacity of nursing students. *J Prof Nurs* 2008; 24: 315-23.
33. Jolliffe D, Farrington DP. Examining the relationship between low empathy and bullying. *Aggressive Behav* 2006; 32: 540-50.
34. Galán González-Serna JM, Serrano RR, Martín MSM, Alarcón Fernández JM. Descenso de empatía en estudiantes de enfermería y análisis de posibles factores Implicados. *Rev Psicol Educ* 2014; 20: 53-60.
35. Fields SK, Mahan P, Tillman P, Harris J, Maxwell K, Hojat M. Measuring empathy in healthcare profession students using the Jefferson Scale of Physician Empathy: health provider - student version. *J Interprof Care* 2011; 25: 287-93.
36. Cunico L, Sartori R, Marognoli O, Meneghini AM. Developing empathy in nursing students: a cohort longitudinal study. *J Clin Nurs* 2012; 21: 2016-25.
37. Bidone S, Beneventi M, Larghero E, Di Pietrantonio C. La tendenza empatica negli studenti del corso di laurea in infermieristica: risultati di uno studio osservazionale. [Empathic tendency in nursing students: results of an observational study]. *Riv Inf* 2014; 51: e8-e13.
38. Ouzouni C, Nakakis K. An exploratory study of student nurses' empathy. *J Health Sci* 2012; 6: 534-52.
39. Kahrman I, Nural N, Arslan U, Topbas M, Can G, Kasim S. The effect of empathy training on the empathic skills of nurses. *Iran Red Crescent Med J* 2016; 18: e24847.

40. McKenna L, Boyle M, Brown T, et al. Levels of empathy in undergraduate nursing students. *Int J Nurs Pract* 2012; 18: 246-51.
41. Nunes P, Williams S, Sa B, Stevenson K. A study of empathy decline in students from five health disciplines during their first year of training. *Int J Med Educ* 2011; 2: 12-17.
42. Wilson SE, Prescott J, Becket G. Empathy Levels in first- and third-year students in health and non-health disciplines. *Am J Pharm Educ* 2012; 76: 24.
43. Ward J, Cody J, Schaal M, Hojat M. The empathy enigma: an empirical study of decline in empathy among undergraduate nursing students. *J Prof Nurs* 2012; 28: 34-40.
44. Sherman JJ, Cramer A. Measurement of changes in empathy during dental school. *J Dent Educ* 2005; 69: 338-45.
45. Hojat M, Vergare MJ, Maxwell K, et al. The devil is in the third year: a longitudinal study of erosion of empathy in medical school. *Acad Med* 2009; 84: 1182-91.
46. Neumann M, Edelhäuser F, Tauschel D, et al. Empathy decline and its reasons: a systematic review of studies with medical students and residents. *Acad Med* 2011; 86: 996-1009.
47. Konrath SH, O'Brien EH, Hsing C. Changes in dispositional empathy in American college students over time: a meta-analysis. *Pers Soc Psychol Rev* 2011; 15: 180-98.
48. Meneghini AM, Sartori R, Cunico L. Adattamento e validazione su campione italiano della Balanced Emotional Empathy Scale di Albert Mehrabian. [Adaptation and validation in an Italian sample of the Balanced Emotional Empathy scale by Albert Mehrabian]. *Ric Psicol* 2006; 29: 123-52.
49. Meneghini AM, Sartori R, Cunico L. Adattamento italiano della Balanced Emotional Empathy Scale (BEES) di Albert Mehrabian. [Italian adaptation of the Balanced Emotional Empathy Scale (BEES) by Albert Mehrabian]. Firenze: Giunti Organizzazioni Speciali; 2016.
50. Meneghini AM, Romaioli D, Nencini A, et al. Validity and reliability of the Caregiving System Scale in the Italian Context. *Swiss J Psychol* 2015; 74: 129-39.
51. Williams B, Brown T, Boyle M, et al. Levels of empathy in undergraduate emergency health, nursing, and midwifery students: a longitudinal study. *Adv Med Educ Pract* 2014; 5: 299-306.
52. Ferri P, Laffi P, Rovesti S, Artioli G, Di Lorenzo R, Magnani D. Motivational factors for choosing the degree course in nursing: a focus group study with nursing students. *Acta Biomed* 2016; 87 Suppl 2: 19-27.
53. Magnani D, Di Lorenzo R, Bari A, Pozzi S, Del Giovane C, Ferri P. The undergraduate nursing student evaluation of clinical learning environment: an Italian survey. *Prof Inferm* 2014; 67: 55-61.

Received: 8 May 2017

Accepted: 7 June 2017

Correspondence:

Paola Ferri

Modena Nursing Degree Programme,

Via del Pozzo, 71 - 41124 Modena

Fax: 059/4222520

E-mail: paola.ferri@unimore.it