

Paolo VENTURA¹
 Valentina BRANCALEONI^{2,a}
 Elena DI PIERRO^{2,a}
 Giovanna GRAZIADEI²
 Annelise MACRÌ³
 Claudio CARMINE GUIDA⁴
 Annamaria NICOLLI⁵
 Maria Teresa ROSSI⁶
 Francesca GRANATA²
 Valeria FIORENTINO²
 Silvia FUSTINONI²
 Raffella SALA⁶
 Piergiacomo Calzavara PINTON⁶
 Andrea TREVISAN⁵
 Stefano MARCHINI¹
 Chiara CUOGHI¹
 Matteo MARCACCI¹
 Elena CORRADINI¹
 Fiammetta SORGE³
 Caterina AURIZI³
 Maria Grazia SAVINO⁴
 Maria Domenica CAPPELLINI²
 Antonello PIETRANGELO¹,
 (Gruppo Italiano Porfiria)

¹ Divisione di Medicina Interna, Centro di riferimento regionale per la diagnosi e la cura delle Porfirie, Policlinico di Modena, Dipartimento di Scienze Medico-Chirurgiche Materno-Infantili e dell'Adulto, Università di Modena e Reggio Emilia

² Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, U.O. di Medicina Generale-Dipartimento di Scienze Cliniche e Comunità, Università degli studi di Milano

³ Centro per le Porfirie, Istituto San Gallicano- IFO IRCCS, Roma

⁴ Centro Interregionale di Riferimento per la Porfiria, U.O.C. Nefrologia e Dialisi-IRCCS Casa Sollievo della Sofferenza, San Giovanni Rotondo (FG)

⁵ Centro per le Malattie Rare del metabolismo-Porfirie, Azienda Ospedaliera-Università degli Studi di Padova

⁶ ASST Spedali Civili di Brescia, UO di Dermatologia, Centro di Fotobiologia e Fotodermatologia

Reprints: Paolo Ventura
 <paoloven@unimore.it>

© The Author(s) 2020

Article accepted on 07/05/2020

Erythropoietic protoporphyria (EPP, OMIM 177000) is a rare inborn metabolic disease characterized by significant and life-long severe acute photosensitivity (often strongly negatively impacting the quality of life of patients) as well as possible haematological and severe liver involvement (2-3% of patients), due to accu-

Clinical and molecular epidemiology of erythropoietic protoporphyria in Italy

Background: Erythropoietic protoporphyria (EPP) is a rare inherited disease associated with heme metabolism, characterized by severe life-long photosensitivity and liver involvement. **Objective:** To provide epidemiological data of EPP in Italy. **Materials & Methods:** Prospective/retrospective data of EPP patients were collected by an Italian network of porphyria specialist centres (Gruppo Italiano Porfiria, GrIP) over a 20-year period (1996-2017). **Results:** In total, 179 patients (79 females) with a clinical and biochemical diagnosis of EPP were assessed, revealing a prevalence of 3.15 cases per million persons and an incidence of 0.13 cases per million persons/year. Incidence significantly increased after 2009 (due to the availability of alfa-melanotide, which effectively limits skin photosensitivity). Mean age at diagnosis was 28 years, with only 22 patients (12.2%) diagnosed ≤ 10 years old. Gene mutations were assessed in 173 (96.6%) patients; most (164; 91.3%) were *FECH* mutations on one allele in association with the hypomorphic variant, c.315-48C, on the other (classic EPP), and nine (5.2%) were *ALAS2* mutations (X-linked EPP). Only one case of autosomal recessive EPP was observed. Of the 42 different *FECH* mutations, 15 are novel, three mutations collectively accounted for 45.9% (75/164) of the mutations (c.215dupT [27.2%], c.901_902delTG [11.5%] and c.67 + 5G > A [7.2%]), and frameshift mutations were prevalent (33.3%). A form of light protection was used by 109/179 (60.8%) patients, and 100 (56%) had at least one α -melanotide implant. Three cases of severe acute liver involvement, requiring OLT, were observed. **Conclusion:** These data define, for the first time, the clinical and molecular epidemiology of EPP in Italy.

Key words: cutaneous porphyrias, erythropoietic protoporphyria, ferrochelatase, photodermatosis, protoporphyria, X-Link EPP

mulation in blood and tissues of protoporphyrin IX (PPIX) [1-4]. The disease affects at least one in 100,000 of the population in Europe and usually presents in early childhood or infancy with severe painful burning and pruritus within minutes of sunlight exposure. Onset of symptoms in adult age is very rare and is mostly a consequence of an acquired somatic mutation within the *ferrochelatase* (*FECH*) gene, secondary to haematological malignancy [1, 2, 4-8]. Four different inheritance patterns of EPP have

doi: 10.1684/ejd.2020.3880




Figure 1. Heme synthesis pathway, enzymes involved and corresponding diseases.

so far been identified. In most cases, the accumulation of PPIX is due to a partial deficiency of FECH, the last enzyme in the heme biosynthesis pathway [1] (figure 1). In the majority, the disease is inherited in a pseudodominant manner (EPP) and photosensitivity is normally overt only in those subjects in whom a deleterious *FECH* mutation (significantly decreasing or abolishing FECH activity) is inherited together with the hypomorphic variant, c.315-48C (also described as FECH IVS3-48C) [9, 10]. Other patients have autosomal recessive EPP (arEPP), which is characterized by the presence of a mutation on both alleles responsible for reduced activity of FECH [11, 12]. In a few cases, PPIX accumulation has been reported to be a consequence of a gain-of-function mutation in an erythroid variant of 5-aminolaevulinic synthase gene (*ALAS2*), the first enzyme of erythroid haem biosynthesis (X-linked Protoporphyrin, XL-EPP; OMIM 300752) [13]. More recently, an increase in activity of ALAS2 protein, responsible for PPIX accumulation and photosensitivity, has been reported to result from a point mutation in the *caseinolytic mitochondrial matrix peptidase chaperone subunit (CLPX)* gene, which encodes for a regulatory protein, responsible for ALAS2 activation [14, 15].

The clinical course of EPP as well as application of an adequate treatment and clinical monitoring justify prompt and accurate diagnosis, requiring specific tests, often supported only by porphyria specialistic centres. As with most rare diseases, healthcare resources and preventative strategies required for support of EPP patients are strictly related to the prevalence of overt cases, however, data concerning the epidemiology of EPP in Italy are so far not available, except for some small studies [16]. The present study aimed to provide clinical and molecular epidemiological data of patients affected by EPP in Italy.

Patients and methods

Patients

A multi-centre cross-sectional national study of EPP in Italy over a 20-year period, 1997-2017, was conducted by Gruppo Italiano Porfiria (GrIP), the Italian Porphyria network including the six reference centres for Porphyrias (Brescia, Milan, Modena, Padova, Roma, and San Giovanni Rotondo). Clinical, biochemical and genetic data of patients with a diagnosis of EPP and XL-EPP were collected. The diagnosis was made in patients with life-long photosensitivity, consistent with EPP, and an increase in total erythrocyte protoporphyrin (usually from 4x to 100x the normal values) with a >50% increase in erythrocyte metal-free protoporphyrin rather than zinc protoporphyrin [1, 2, 17, 18]. In EPP, metal-free protoporphyrin generally represents >85% total porphyrins. In XL-EPP, metal-free protoporphyrin is generally about 50% of total porphyrins. When available, the genetic analysis confirmed the diagnosis. All patients or their parents gave informed consent to be included in this observational survey.

Biochemical analyses

Light-protected collected samples were analysed in the laboratories of porphyria centres for measurement of erythrocyte total PPIX and PPIX fractions that were metal-free and zinc-chelated, as previously described [17, 19, 20].

Molecular analyses

Genomic DNA (gDNA) was extracted from buffy coat by both manual and automated systems. As previously

Table 1. Distribution of EPP patients according to each Italian porphyria centre database.

	Brescia	Milan	Padua	Modena	Rome	San Giovanni Rotondo	Total included	True total*
EPP	24	77	17	12	111	11	252	170
XL-EPP	0	9	0	0	4	0	13	9
Total	24	86	17	12	115	11	265	179

*After excluding multiple referrals (see text for details).

reported [21, 22], for the manual method, gDNA was extracted using the PUREGENE DNA purification system blood kit (Gentra Systems Minneapolis, MN, USA); for the automated systems, the Maxwell 16 DNA Blood Kit with the automatic extractor Maxwell®16 was used (Promega Corp., Madison City, WI). The promoters and the entire coding regions of the human *FECH* and *ALAS2* genes were amplified with specific primers pairs. For the *FECH* and *ALAS2* genes, 11 and 10 primer pairs were used, respectively. gDNA, 100-150 ng, was amplified using BIOTAQ DNA polymerase (Bioline, London, UK) with the specific PCR primer pair for each exon or fragment. Primer sequences and melting temperatures are available from the authors on request. The PCR products were then subjected to automated direct sequencing on an ABI Prism 310 Genetic Analyser (Thermo Fisher Corporation Inc., San Francisco, CA, USA). When no *FECH* mutation was detected by sequencing, multiplex ligation dependent probe amplification (MLPA) technique analysis to identify large deletions and characterization of deletion breakpoints was carried out, as previously described [23]. In the absence of long deletions, a custom enrichment panel for *FECH* gene resequencing was designed; the panel covers 40 kb upstream to 10 kb downstream of the *FECH* gene, including all exons and introns. DNA libraries were produced by a standard Haloplex target enrichment system procedure (Agilent Technologies, Santa Clara, USA) and 150 bp paired-end reads were generated using a MiSeq sequencer (Illumina, San Diego, USA), as already described [24]. Sequence variant nomenclature is according to Human Genome Variation Society (HGVS) recommendations (version 19.01). Nucleotides are numbered in the cDNA sequence of the human *FECH* gene (NCBI reference sequence NM_000140.3) and human *ALAS2* gene (NM_000032.4) with the A of the ATG initiation codon as '+1'.

Statistical methods

Data were analysed using Stata 15.1 statistical software (StataCorp, Texas USA). Measurements were expressed as mean (\pm standard deviation), median and range, or percentage, when appropriate. The significance of differences between quantitative variables was assessed by the Mann-Whitney or Kruskal-Wallis tests, when appropriate. For all statistical analysis, $p < 0.05$ was considered significant.

Results

A total of 179 patients (79 females) among 140 unrelated families were analysed. The cases included in the original

Table 2. General characteristics of studied patients.

Sex	
Male	100 (55.4%)
Female	79 (45.6%)
Prevalence (cases/million inhabitants)	3.15
Incidence (new diagnosis/year/million inhabitants) *	0.13
Mean age (years)	38.2 (\pm 16.1)
Male	38.4 (\pm 15.6)
Female	37.9 (\pm 16.3)
Mean age at diagnosis (years) (min-max)	28.3 (\pm 15.1) (3-71)
Male	28.8 (\pm 15.8)
Female	29.5 (\pm 15.4)
Treatment	
None/unknown	67
Photoprotectors ¹	21
Alfa-melanotide ²	100
Heme arginate	2
Liver transplant	3
Erythrocyte protoporphyrin (μmol/L) median (range)	27.1 (8.3-91.6)
Metal-free protoporphyrin (%) (range)	89 (78-91)

*Mean rate within the considered period (see text for details).¹At least one period during a life-time (see text for details).²At least one subcutaneous implant during a life-time (see text for details).

collected database by the six different Italian porphyria centres participating in the study are presented in table 1. The sum of patients included by each single centre is higher than the true number of patients as a consequence of a "over-sampling bias" (see below in the discussion section for explanation). The mean age at diagnosis was 28 years (with no significant difference between genders) (table 2). No EPP diagnosis was made under three years of age. The frequency of EPP diagnosis was 4/179 (2.2%) and 22/179 (12.2%) under five and under 10 years of age, respectively. Eighteen (10.4%) patients were diagnosed after 50 years of age. None had haematological malignancies. The mean age at diagnosis for patients with XL-EPP ($n = 9$) was older than that of patients with classic EPP (36.7 ± 14.9 vs. 28.6 ± 15.6 , $p = 0.109$) even though the difference was not statically significant. Three patients affected by the classic form of EPP underwent liver transplantation due to severe hepatic complications (in all cases, acute liver failure with significant increase in markers of cholestatic disease). In all cases, liver pathology revealed a diffuse precipitation of protoporphyrin microaggregates into the intrahepatic biliary canaliculi. One patient died during the




Figure 3. Geographical EPP prevalence in Italy.




Figure 4. EPP proband distribution per family.

Discussion

In Italy, most patients with suspected porphyria are referred to expert centres for diagnostic confirmation and clinical management. This study reports, for the first time, data collected by all the leading porphyria expert centres in the country. It should be noted that in Italy some patients consult more than one expert centre, for possible different opinions and clinical management. To reduce this “over-sampling bias”, we performed a deep analysis by comparing each single-centre database and excluding all possible iterative inclusions. For this reason, we believe that the data provided by the present study better reflects the true Italian epidemiology of EPP compared to other previously reported data [7].

The analysis of the overall data shows that in Italy the diagnosis of EPP is made later than in the rest of Europe (on average at the age of 28 rather than 21) [7] and, in partic-

ular, that the rate of EPP diagnosis under 10 years of age is still very low (less than 15% of cases). These data suggest that too many Italian patients with EPP are probably still misdiagnosed with “sun allergy”, this highlighting the need for a greater awareness of the disease among Italian physicians, in particular, paediatricians and dermatologists, and a higher level of clinical suspicion and improvement in diagnosis [7, 25].

Moreover, our data indicate an incidence of about 0.13 new cases/year/million inhabitants, approximately twice as much as that observed in Europe [7]. However, the incidence in Italy was not constant over the considered period, showing a significant increase from 2009 (figure 2). Interestingly, in 2009, alfa-melanotide started to become available and patients taking this medication were enrolled into clinical trials via one Italian centre, in particular [26-28]. One may speculate that the availability of an effective protective drug against PPIX-induced photosensitivity may have favoured the spread awareness of EPP

Table 3. Frequency and type of mutation in the *FECH* gene in the studied population.

Mutation	Number Of Patients	%*	Type	References
Genetic analyses not made	6	–	–	-
None detected (Only Hypomorphic allele)	4	2.4	undefined	-
c.[215dupT]	45	27.4	FS	Wang (1997) J Invest Dermatol 109,688
c.[901_902delTG]	18	11	FS	Schneider-Yin (1994) Hum Genet 93, 711
c.[67+5G>A]	12	7.3	SP	Wang (1999) J Invest Dermatol 113, 87
c.[1-251G>C;194+4350_463+1197del5576]	6	3.6	LD	Di Pierro (2006) Hum Genet 118 776
c.[343C>T]	6	3.6	NS	Henriksson (1996) J Invest Dermatol 106, 346
c.[843delC]	6	3.6	FS	Gouya (1998) J Invest Dermatol 111, 406
c.[599-3C>T]	5	3.0	SP	Aurizi (2007) Mol Genet Metab 90, 402
c.[757_761delAGAAG]	5	3.0	FS	Henriksson (1996) J Invest Dermatol 106, 346
c.[464-1169 A>C]**	4	2.4	SP	Di Pierro (2019) Genet Med 2019
c.[892C>T]	4	2.4	NS	Di Pierro (2004) Hum Genet 114 608
c.[94C>T]	4	2.4	NS	This report
c.[1-251G>C]	3	1.8	RG	Di Pierro (2005) Exp Hematol 33: 584
c.[195-2A>G]	3	1.8	SP	Aurizi (2007) Mol Genet Metab 90, 402
c.[400delA]	3	1.8	FS	Wang (1999) J Invest Dermatol 113, 87
c.[116delG]	2	1.2	FS	This report
c.[286C>T]	2	1.2	NS	Henriksson (1996) J Invest Dermatol 106, 346
c.[315-67G>A]	2	1.2	FS	Aurizi (2007) Mol Genet Metab 90, 402
c.[415C>T]	2	1.2	NS	This report
c.[488_501del]	2	1.2	FS	Di Pierro (2001) Hum Genet 109, 468
c.[706-3C>G]	2	1.2	SP	This report
c.[791C>T]	2	1.2	MS	Martinez (2001) Hum Genet 109, 241
c.[930 G>A]	2	1.2	NS	Di Pierro (2004) Hum Genet 114, 221
c.[942_945delAACA]	2	1.2	FS	This report
c.[1-7887_67+2422del10376bp]	1	.06	LD	Brancaleoni (2007) Hum Genet 121, 646
c.[1-9628_67+2871del12566bp]	1	.06	LD	Brancaleoni (2008) Hum Genet 123, 546
c.[1041_1047delGCTGGACA]	1	.06	FS	This report
c.[1057delT]	1	.06	FS	This report
c.[1080_1081delTG]	1	.06	FS	This report
c.[1232G>A]	1	.06	MS	Balwani (2013) Mol Med 19, 26
c.[163G>T]	1	.06	MS	Lamoril (1991) Biochem Biophys Res Commun 181, 594
c.[195-16A>G] ¹	1	.06	SP	This report
c.[451T>C]	1	.06	MS	Rufenacht (1998) Am J Hum Genet 62, 1341
c.[544delC]	1	.06	FS	This report
c.[680G>A]	1	.06	NS	This report
c.[705+1delG]	1	.06	SP	This report
c.[705+1G>A]	1	.06	SP	This report
c.[727A>G] ²	1	.06	MS	This report
c.[1231T>G] ²	1	.06	MS	Whatley (2010) Br J Dermatol 162, 642
c.[742T>C]	1	.06	MS	This report
c.[782C>T]	1	.06	MS	Brancaleoni (2008) Hum Genet 124, 296
c.[801G>A]	1	.06	MS	Lamoril (1991) Biochem Biophys Res Commun 181, 594
c.598+1G>T(;);599-3C>T ³	1	.06	SP	Frank (1999) J Investig Med 47, 278

FS: frameshift; MS: missense; NS: nonsense; SP: splicing; LD: long deletion.*Percentage calculated from the total 164 subjects.**Presence of IVS in both alleles.¹No IVS analysed.²Mutations found in the same patient (ArEPP).³Two mutations, probably in cis.




Figure 5. *FECH* mutations observed in Italian patients. Upper panel: point mutations; lower panel: long deletions. Previously unreported mutations are shown in black, while previously described mutations are in grey. The two mutations within a square were found in the only autosomal recessive erythropoietic protoporphyria patient.

among clinicians. This may explain the higher EPP incidence observed in Italy, where a large proportion of patients [100/179 (55.8%)] received at least one subcutaneous alfamelanotide implant from 2009 onwards (table 1).

Our study reports a prevalence of 3.15 cases/million/inhabitants, which is lower than that reported for Italy in the porphyria European epidemiological study on porphyria conducted in 2012. It should be noted that in this study, the prevalence of different porphyrias was derived from incidence rates over three consecutive years, assuming that the ratio between incidence and prevalence was stable for many years, however, as presented in figure 2, this was not the case in Italy [7]. Moreover, the Italian incidence data reported by Elder *et al.* [7] may be biased due to an “over-sampling bias”, as mentioned above. Although the data reported in our study, free of this bias, are more likely to be accurate, we cannot exclude that a large number of EPP patients are still undiagnosed.

Moreover, this is supported by the relatively old age at diagnosis, suggesting poor awareness of this disease by clinicians in our country. This hypothesis is also supported by the significant difference in regional distribution of EPP cases (figure 3). In fact, most patients seem to live in regions with an active porphyria reference centre, which probably leads to a greater knowledge of the disease, resulting in a greater number of patients diagnosed.

Table 4. Erythrocyte protoporphyrin concentration in patients with erythropoietic protoporphyria, relative to type of mutation.

	Number of patients	Erythrocyte protoporphyrin ($\mu\text{mol/L}$) median (range)
Total	164	26.7 (8.27-91.6)
Missense mutations	10	18.2 (9.84-54.2)
Splicing mutations	30	19.6 (8.56-91.6)
Frameshift mutations	92	25.8 (8.27-81.9)
Nonsense mutations	20	26.8 (9.82-78.9)
Long-deletion mutations	8	28.9 (9.81-84.5)
Regulatory gene mutations	4	23.6 (10.2-73.9)

In our Italian cohort of EPP patients, 42 different *FECH* gene mutations were detected, confirming the allelic heterogeneity already described in other countries [6, 7, 11, 29-31]. However, in Italy, as well as in other European countries, the level of allelic *FECH* heterogeneity is lower than that observed for autosomal acute porphyria [7], mostly due to the presence of three prevalent mutations accounting for 45.9% of all *FECH* mutations (table 3).

Similar high frequencies of *FECH* mutations among apparently unrelated EPP patients have been reported in other European and non-European countries [6, 7, 29, 31].

Of the 179 patients with EPP, 88.8% had a loss-of-function *FECH* allele and the c.315-48T > C low-expressing variant. This data reinforces the idea that EPP prevalence is strongly influenced by the frequency of the hypomorphic *FECH* IVS3-48C allele [7, 11], which is highly variable in different European countries (ranging from 5-11%) [6, 31]. In Italy, the prevalence is reported to be 3.3% and 3.6%, according to the 1000Genome (<https://www.internationalgenome.org/>) and GnomeAD (<https://gnomad.broadinstitute.org/>) databases, for Tuscany and south European populations, respectively. All but 2.4% of our patients underwent complete molecular characterization, excluding patients for whom it was not possible to analyse DNA. In fact, four patients had only the heterozygous hypomorphic *FECH* allele with classic biochemical and clinical features of EPP. In these patients, it was not possible to assess long deletion by MLPA, nor the presence of *ALAS2* mutation, or deep intronic mutation. Thus, it is conceivable that these patients represent classic EPP patients. We identified only one patient with two *FECH* loss-of-function alleles, accounting for only 0.6% of the total number of EPP patients. This percentage is very low when compared to that reported in other countries (about 5%).

Most of the 42 different mutations identified in the *FECH* gene were small deletions and insertions resulting in: a frameshift with creation of a premature stop codon; splicing defects predicted to cause aberrant mRNA processing; or nonsense variants, introducing premature stop codons. Similarly, among the 15 novel variants detected in our population, most were small deletions, splicing defects or nonsense variants, while only two missense mutations were described. Both missense variants (p.K243L, p.F248L) led to replacement of highly conserved amino acids. The most common mutations in the study population were c.215dupT and c.901_902delTG, which have not been reported with such a high frequency in other EPP populations. Among our cases, patients with XL-EPP accounted for approximately 5% of all EPP patients, which is less than that reported in Europe and the US [6, 32, 33]. In contrast to other reports, our XL-EPP patients carried only one mutation, c.[1706_1709delAGTG]. Male patients with XL-EPP reported early onset of symptoms, while females showed significant heterogeneity of symptoms, ranging from onset of symptoms in childhood to being symptomatic until adulthood at the time of the study. This variability is strongly associated with random X-chromosomal inactivation in females patients, as previously described [20].

In contrast to other studies, we did not identify mutations associated with a milder phenotype [34]. However, several patients were reported to carry the same mutation but with different disease severity, suggesting the existence of as yet undefined factors (modifier genes or acquired or environmental factors) which may influence the phenotype of the disease.

In conclusion, to the best of our knowledge, this study represents the largest collection of data on the prevalence and molecular epidemiology of EPP in Italy available so far. Most patients were found to have a *FECH* gene mutation associated with a low-expressing allele (EPP). The autosomal recessive form of EPP appears to be very rare in

Italy, and one patient was identified in our cohort (0.56%), whereas the recently identified X-linked dominant EPP showed a prevalence of 5% [6, 35]. Considering both the availability of an effective treatment for photosensitivity (which negatively impacts on patients' quality of life) and the importance of the clinical follow-up (especially concerning the possibility of severe liver involvement), a significantly greater clinical awareness and knowledge of EPP is required among clinicians [26, 28, 36-41]. In order to reduce the number of misdiagnosed or/and untreated EPP patients, relevant training for clinicians should be implemented. ■

Acknowledgments and disclosures. *Acknowledgments: the authors are grateful to all patients, who donated samples for this study. We are grateful to Dr. Gigi Mori, PhD (ASST Spedali Civili di Brescia) and Prof. Leonardo Salviati, MD (Università degli studi di Padova) for genetic testing. Financial support: none. Conflicts of interest: Paolo Ventura and Maria Domenica Cappellini received funds for consultation and lecturing by Alnylam Pharmaceuticals and Recordati Rare Disease.*

Open Access. *This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.*

References

1. Cox TM. *Protoporphyrin*. Kadish KM, Guillard R. San Diego: Academic Press, 2003.
2. Thunell S, Harper P, Brun A. Porphyrins, porphyrin metabolism and porphyrias. IV. Pathophysiology of erythropoietic protoporphyria: diagnosis, care and monitoring of the patient. *Scand J Clin Lab Invest* 2000; 60: 581-604.
3. Holme SA, Anstey AV, Finlay AY, Elder GH, Badminton MN. Erythropoietic protoporphyria in the UK: clinical features and effect on quality of life. *Br J Dermatol* 2006; 155: 574-81.
4. Sood G, Anderson KE. *Porphyrias*. Crowther M, Ginsberg J, Schunemann H, et al. Hoboken: Wiley, 2008.
5. Wahlin S, Floderus Y, Stal P, Harper P. Erythropoietic protoporphyria in Sweden: demographic, clinical, biochemical and genetic characteristics. *J Intern Med* 2011; 269: 278-88.
6. Whatley SD, Mason NG, Holme SA, Anstey AV, Elder GH, Badminton MN. Molecular epidemiology of erythropoietic protoporphyria in the U.K. *Br J Dermatol* 2010; 162: 642-6.
7. Elder G, Harper P, Badminton M, Sandberg S, Deybach JC. The incidence of inherited porphyrias in Europe. *J Inherit Metab Dis* 2013; 36: 849-57.

8. Sarkany RP, Ross G, Willis F. Acquired erythropoietic protoporphyria as a result of myelodysplasia causing loss of chromosome 18. *Br J Dermatol* 2006; 155: 464-6.
9. Wiman A, Floderus Y, Harper P. Novel mutations and phenotypic effect of the splice site modulator IVS3-48C in nine Swedish Families with erythropoietic protoporphyria. *J Hum Genet* 2003; 48: 70-6.
10. Gouya L, Puy H, Robreau AM, et al. The penetrance of dominant erythropoietic protoporphyria is modulated by expression of wildtype FECH. *Nat Genet* 2002; 30: 27-8.
11. Gouya L, Martin-Schmitt C, Robreau AM, et al. Contribution of a common single-nucleotide polymorphism to the genetic predisposition for erythropoietic protoporphyria. *Am J Hum Genet* 2006; 78: 2-14.
12. Sarkany RP, Cox TM. Autosomal recessive erythropoietic protoporphyria: a syndrome of severe photosensitivity and liver failure. *QJM* 1995; 88: 541-9.
13. Whatley SD, Ducamp S, Gouya L, et al. C-terminal deletions in the ALAS2 gene lead to gain of function and cause X-linked dominant protoporphyria without anemia or iron overload. *Am J Hum Genet* 2008; 83: 408-14.
14. Yien YY, Ducamp S, van der Vorm LN, et al. Mutation in human CLPX elevates levels of delta-aminolevulinic acid synthase and protoporphyrin IX to promote erythropoietic protoporphyria. *Proc Natl Acad Sci USA* 2017; 114: E8045-52.
15. Whitman JC, Paw BH, Chung J. The role of ClpX in erythropoietic protoporphyria. *Hematol Transfus Cell Ther* 2018; 40: 182-8.
16. Aurizi C, Schneider-Yin X, Sorge F, Macri A, Minder EI, Biolcati G. Heterogeneity of mutations in the ferrochelatase gene in Italian patients with erythropoietic protoporphyria. *Mol Genet Metab* 2007; 90: 402-7.
17. Deacon AC, Elder GH. ACP Best Practice No 165: front line tests for the investigation of suspected porphyria. *J Clin Pathol* 2001; 54: 500-7.
18. Sassaroli M, da Costa R, Vaananen H, Eisinger J, Poh-Fitzpatrick MB. Distribution of erythrocyte free porphyrin content in erythropoietic protoporphyria. *J Lab Clin Med* 1992; 120: 614-23.
19. Sassa S, Granick JL, Granick S, Kappas A, Levere RD. Studies in lead poisoning. I. Microanalysis of erythrocyte protoporphyrin levels by spectrophotometry in the detection of chronic lead intoxication in the subclinical range. *Biochem Med* 1973; 8: 135-48.
20. Brancaleoni V, Balwani M, Granata F, et al. X-chromosomal inactivation directly influences the phenotypic manifestation of X-linked protoporphyria. *Clin Genet* 2016; 89: 20-6.
21. Di Pierro E, Brancaleoni V, Besana V, Ausenda S, Drury S, Cappellini MD. A 10376 bp deletion of FECH gene responsible for erythropoietic protoporphyria. *Blood Cells Mol Dis* 2008; 40: 233-6.
22. Brancaleoni V, Granata F, Colancecco A, Tavazzi D, Cappellini MD, Di Pierro E. Seven novel genetic mutations within the 5'UTR and the housekeeping promoter of HMBS gene responsible for the non-erythroid form of acute intermittent porphyria. *Blood Cells Mol Dis* 2012; 49: 147-51.
23. Di Pierro E, Brancaleoni V, Besana V, Cappellini MD. Multiplex ligation-dependent probe amplification: a novel approach for genetic diagnosis of Porphyria. *J Hum Genet* 2009; 54: 479-87.
24. Chiara M, Primon I, Tarantini L, et al. Targeted resequencing of FECH locus reveals that a novel deep intronic pathogenic variant and eQTLs may cause erythropoietic protoporphyria (EPP) through a methylation-dependent mechanism. *Genet Med* 2020; 22: 35-43.
25. Lala SM, Naik H, Balwani M. Diagnostic delay in erythropoietic protoporphyria. *J Pediatr* 2018; 202: 320-3.e2.
26. Minder EI. Afamelanotide, an agonistic analog of alpha-melanocyte-stimulating hormone, in dermal phototoxicity of erythropoietic protoporphyria. *Expert Opin Investig Drugs* 2010; 19: 1591-602.
27. Luger TA, Bohm M. An alpha-MSH analog in erythropoietic protoporphyria. *J Invest Dermatol* 2015; 135: 929-31.
28. Langendonk JG, Balwani M, Anderson KE, et al. Afamelanotide for erythropoietic protoporphyria. *N Engl J Med* 2015; 373: 48-59.
29. Parker M, Corrigan AV, Hift RJ, Meissner PN. Molecular characterization of erythropoietic protoporphyria in South Africa. *Br J Dermatol* 2008; 159: 182-91.
30. Risheg H, Chen FP, Bloomer JR. Genotypic determinants of phenotype in North American patients with erythropoietic protoporphyria. *Mol Genet Metab* 2003; 80: 196-206.
31. Balwani M. Erythropoietic protoporphyria and X-linked protoporphyria: pathophysiology, genetics, clinical manifestations, and management. *Mol Genet Metab* 2019; 128: 298-303.
32. Balwani M, Doheny D, Bishop DF, et al. Loss-of-function ferrochelatase and gain-of-function erythroid-specific 5-aminolevulinic acid synthase mutations causing erythropoietic protoporphyria and x-linked protoporphyria in North American patients reveal novel mutations and a high prevalence of X-linked protoporphyria. *Mol Med* 2013; 19: 26-35.
33. Balwani M, Naik H, Anderson KE, et al. Clinical, biochemical, and genetic characterization of North American patients with erythropoietic protoporphyria and X-linked protoporphyria. *JAMA Dermatol* 2017; 153: 789-96.
34. Berroeta L, Man I, Goudie DR, Whatley SD, Elder GH, Ibbotson SH. Late presentation of erythropoietic protoporphyria: case report and genetic analysis of family members. *Br J Dermatol* 2007; 157: 1030-1.
35. Schmitt C, Ducamp S, Gouya L, Deybach JC, Puy H. Inheritance in erythropoietic protoporphyria. *Pathol Biol (Paris)* 2010; 58: 372-80.
36. Langendonk JG. Treatment for erythropoietic protoporphyria. *Br J Dermatol* 2015; 172: 1481-2.
37. Singal AK, Parker C, Bowden C, Thapar M, Liu L, McGuire BM. Liver transplantation in the management of porphyria. *Hepatology* 2014; 60: 1082-9.
38. Gross U, Frank M, Doss MO. Hepatic complications of erythropoietic protoporphyria. *Photodermatol Photoimmunol Photomed* 1998; 14: 52-7.
39. Anstey AV, Hift RJ. Liver disease in erythropoietic protoporphyria: insights and implications for management. *Postgrad Med J* 2007; 83: 739-48.
40. McGuire BM, Bonkovsky HL, Carithers RL Jr., et al. Liver transplantation for erythropoietic protoporphyria liver disease. *Liver Transpl* 2005; 11: 1590-6.
41. Rand EB, Bunin N, Cochran W, Ruchelli E, Olthoff KM, Bloomer JR. Sequential liver and bone marrow transplantation for treatment of erythropoietic protoporphyria. *Pediatrics* 2006; 118: e1896-9.