

This is the peer reviewed version of the following article:

On the behavior of periodic solutions of planar autonomous Hamiltonian systems with multivalued periodic perturbations / O., Makarenkov; Malaguti, Luisa; P., Nistri. - In: ZEITSCHRIFT FUR ANALYSIS UND IHRE ANWENDUNGEN. - ISSN 0232-2064. - STAMPA. - 30:2(2011), pp. 129-144. [10.4171/ZAA/1428]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

25/09/2024 07:29

(Article begins on next page)

On the behavior of periodic solutions of planar autonomous Hamiltonian systems with multivalued periodic perturbations

Oleg Makarenkov, Luisa Malaguti and Paolo Nistri *

Abstract. Aim of the paper is to provide a method to analyze the behavior of T -periodic solutions $x_\varepsilon, \varepsilon > 0$, of a perturbed planar Hamiltonian system near a cycle x_0 , of smallest period T , of the unperturbed system. The perturbation is represented by a T -periodic multivalued map which vanishes as $\varepsilon \rightarrow 0$. In several problems from nonsmooth mechanical systems this multivalued perturbation comes from the Filippov regularization of a nonlinear discontinuous T -periodic term. Through the paper, assuming the existence of a T -periodic solution x_ε for $\varepsilon > 0$ small, under the condition that x_0 is a nondegenerate cycle of the linearized unperturbed Hamiltonian system we provide a formula for the distance between any point $x_0(t)$ and the trajectories $x_\varepsilon([0, T])$ along a transversal direction to $x_0(t)$.

Keywords. planar Hamiltonian systems, characteristic multipliers, multivalued periodic perturbations, periodic solutions, topological degree.

Mathematics Subject Classification (2000). 37K05, 34A60, 34C25

1. Introduction

Let x_0 be a T -periodic cycle of the Hamiltonian system

$$\dot{x} = f(x), \quad (1)$$

where $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$. In the recent monographs [2] and [3], on the basis of numerical simulations, was heuristically shown that the subharmonic Melnikov's

*Corresponding author.

Department of Mathematics, Imperial College London, London, SW7 2AZ, UK; e-mail: o.makarenkov@imperial.ac.uk

Dipartimento di Scienze e Metodi dell'Ingegneria, Università di Modena e Reggio Emilia, 42100 Reggio Emilia, Italy; e-mail: luisa.malaguti@unimore.it

Dipartimento di Ingegneria dell'Informazione, Università di Siena, 53100 Siena, Italy; e-mail: pnistri@dii.unisi.it

method ([19], [10, Chapter 4, §6]) correctly predicts the existence of T -periodic solutions x_ε of the differential inclusion

$$\dot{x} \in f(x) + \varepsilon g(t, x, \varepsilon), \quad (2)$$

where $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ is a multivalued map taking the values in the set $K(\mathbb{R}^2)$ of nonempty compact and convex sets of \mathbb{R}^2 . Sufficient conditions for the local and global existence of at least an absolutely continuous solution of (2) starting from any initial condition can be found in ([1, Chapter 2]).

In [2] and [3] the authors have experimentally observed that if θ_0 is a simple zero of the subharmonic Melnikov's bifurcation function then (2) possesses a T -periodic solution x_ε such that

$$x_\varepsilon(t) \rightarrow x_0(t + \theta_0) \quad \text{as } \varepsilon \rightarrow 0, \quad \text{uniformly in } t \in [0, T]. \quad (3)$$

A theoretical justification of this result can be provided along the lines of the papers [7], [11] and [16]. In this paper we do not provide conditions to ensure the existence of T -periodic solutions x_ε , for $\varepsilon > 0$ small, instead we want to evaluate the distance between any point $x_0(t)$ and the curve $x_\varepsilon([0, T])$ providing in this way a tool to study the behavior of the T -periodic solutions of (2) near x_0 . This tool, together with the method based on the Melnikov's bifurcation function mentioned above, permits to perform a complete analysis both for the existence and the behavior near the cycle x_0 of the T -periodic solutions x_ε to (2).

Since in this paper the existence of T -periodic solutions x_ε of (2) is assumed, we only require to the multivalued map g the minimal regularity assumptions needed for our analysis. In fact, through the paper we only assume that the map $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ is measurable or upper semicontinuous.

The interest of considering multivalued perturbation of system (1) is mainly related to the necessity, encountered in the applications, to deal with perturbations, having jump discontinuities, of Hamiltonian autonomous systems. In fact, many physical problems are modeled by ordinary differential equations with discontinuous right hand side whose regularization produces a multivalued map (see for instance [9] and [1]). Among them we like to cite the study of the self-sustained oscillations induced by friction in one-degree of freedom mechanical systems. This problem gives rise to a planar Hamiltonian system perturbed by a periodic perturbation of small amplitude with jump discontinuities, compare e.g. [2, Chapter 15] where the analysis was heuristically performed by means of the Melnikov method.

The paper is organized as follows. In Section 2 assuming that the linearized system

$$\dot{y} = f'(x_0(t))y \quad (4)$$

possesses a not T -periodic solution, in this case following Rhouma-Chicone [21] x_0 is said to be nondegenerate, we show the existence of a family $\{\Delta_\varepsilon\}_{\varepsilon>0}$ of

real numbers with $\Delta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ such that

$$\frac{\|x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)\|}{\varepsilon} \leq \text{const} \quad \text{for any } t \in [0, T] \text{ and any } \varepsilon > 0. \quad (5)$$

This property has been already established by the authors in [16] and [17] in the case when x_0 is an isolated limit cycle and g in (2) is a singlevalued continuous function. In section 3 we employ property (5) together with a suitably defined multivalued function $M^\perp \in C^0(\mathbb{R}, \mathbb{R})$ to obtain

$$x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) \in \varepsilon M^\perp(t)y(t) + \alpha_\varepsilon(t)\dot{x}_0(t) + o(\varepsilon), \quad (6)$$

where y is a not T -periodic solution of the linearized system (4) and $\alpha_\varepsilon(t)$ is a scalar function infinitesimal as $\varepsilon \rightarrow 0$ of order greater or equal to 1. The function $\alpha_\varepsilon(t)$ is given in the formula (46) of the paper. The formula to represent the function M^\perp is provided in Section 3, thus (6) gives an explicit formula for the distance between the trajectories x_0 and x_ε along a transversal direction to x_0 . Finally, in Section 4 we specialize the formula for M^\perp in the case when the Hamiltonian system (1) possesses symmetry properties, as often is the case in the applications.

2. Evaluation of the distance between the periodic solutions of the perturbed system and the cycle of the unperturbed one

In this Section we establish the validity of inequality (5) which is the starting point for (6). This result does not depend on the perturbation term g , indeed the only property we need is the following one.

Definition 2.1. ([21]) We say that the cycle x_0 of autonomous system (1) is nondegenerate if the linearized system (4) has a not T -periodic solution.

If (1) is Hamiltonian then the nondegeneracy of x_0 implies that the period T of x_0 is noncritical (compare [5]).

Definition 2.2. ([15, Definition 2.2.1]) A function $x : [0, T] \rightarrow \mathbb{R}^2$ is said to be a solution of the differential inclusion (2) on $[0, T]$ if x is absolutely continuous and the inclusion in (2) holds for almost all (a.a.) $t \in [0, T]$.

Definition 2.3. ([12, Definition 1.3.1]) For any $\varepsilon > 0$ the multivalued map $g(\cdot, \varepsilon) : \mathbb{R} \times \mathbb{R}^2 \rightarrow K(\mathbb{R}^2)$ is said to be measurable if, for any open $V \subset \mathbb{R}^2$, the set $g^{-1}(V, \varepsilon) := \{(t, x) \in \mathbb{R} \times \mathbb{R}^2 : g(t, x, \varepsilon) \cap V \neq \emptyset\}$ is measurable.

We assume the following condition.

(H) – for any bounded set $B \subset \mathbb{R}^2$ there exists $\mu_B \in L_{loc}^\infty(\mathbb{R})$ such that

$$\|g(t, x, \varepsilon)\| := \sup\{\|y\| : y \in g(t, x, \varepsilon)\} \leq \mu_B(t)$$

for all $t \in \mathbb{R}$, $x \in B$ and $\varepsilon \in [0, 1]$.

Note that the notion of nondegenerate cycles has been used in [16] and [17] in a stronger sense, i.e. x_0 is called nondegenerate if the linearized system (4) has only one characteristic multiplier equal to +1.

In order to introduce the family $\{\Delta_\varepsilon\}_{\varepsilon>0}$, following [17], we define a surface $S \in C(\mathbb{R}, \mathbb{R}^2)$ as follows

$$\begin{aligned} S(v) &= \Omega(T, 0, h(v)), \\ h(v) &= x_0(0) + A_1 v, \end{aligned} \tag{7}$$

where $\Omega(\cdot, t_0, \xi)$ is the solution of (1) satisfying $\Omega(t_0, t_0, \xi) = \xi$ and A_1 is an arbitrary 2×1 vector such that the 2×2 matrix $(\dot{x}_0(0), A_1)$ is nonsingular.

The following result shows that the surface S intersects x_0 transversally.

Lemma 2.4. ([17, Lemma 2.2]) *Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$. Let x_0 be a nondegenerate T -periodic cycle of (1). Then $\dot{x}_0(0) \notin S'(0)(\mathbb{R})$.*

Using the previous Lemma we can prove the following result.

Lemma 2.5. *Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and that $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ is measurable and satisfying (H). Let x_0 be a nondegenerate T -periodic cycle of (1). Let x_ε be a T -periodic solution to perturbed system (2) satisfying*

$$\|x_\varepsilon(t) - x_0(t)\| \rightarrow 0$$

as $\varepsilon \rightarrow 0$ uniformly with respect to $t \in \mathbb{R}$, then there exists $\varepsilon_0 > 0$ and $r_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0]$ the equation $x_\varepsilon(\Delta) = S(v)$ has a unique solution $(\Delta_\varepsilon, v_\varepsilon)$ in $[-r_0, r_0] \times \{v \in \mathbb{R} : |v| \leq r_0\}$. Moreover, the functions $\varepsilon \rightarrow \Delta_\varepsilon$, $\varepsilon \rightarrow v_\varepsilon$ are continuous at $\varepsilon = 0$ with $\Delta_0 = 0$ and $v_0 = 0$.

In the case when g in (2) is singlevalued and continuous Lemma 2.5 is a simple consequence of Lemma 2.4 ([17, Corollary 2.3]). In the present case of g multivalued map we should provide a proof.

Proof. Define the function $F : \mathbb{R}^2 \times [0, 1] \rightarrow \mathbb{R}^2$ as follows

$$F((t, v), \varepsilon) = x_\varepsilon(t) - S(v),$$

then $F((0, 0), 0) = 0$. Our assumptions and definitions (7) guarantee that F is a continuous function at the points $\mathbb{R}^2 \times \{0\}$. Since $F(\cdot, 0)$ is differentiable at $(0, 0)$ and $F'_{(t,v)}((0, 0), 0) = (\dot{x}_0(0), -S'(0))$ is nonsingular by Lemma 1, then there exists $r_0 > 0$ such that

$$d(F(\cdot, 0), [-r_0, r_0] \times [-r_0, r_0], 0) \neq 0,$$

here $d(\Phi, V, 0)$ denotes the topological degree of the map Φ in the set V with respect to 0.

Therefore, there exists $\varepsilon_0 > 0$ such that

$$d(F(\cdot, \varepsilon), [-r_0, r_0] \times [-r_0, r_0], 0) \neq 0, \quad \text{for any } \varepsilon \in [0, \varepsilon_0].$$

This implies that for any $\varepsilon \in [0, \varepsilon_0]$, by the solution property of the topological degree, there exists at least one pair $(\Delta_\varepsilon, v_\varepsilon) \in [-r_0, r_0] \times [-r_0, r_0]$ such that $x_\varepsilon(\Delta_\varepsilon) - S(v_\varepsilon) = 0$.

Let us show that this solution is unique in $[-r_0, r_0] \times [-r_0, r_0]$ provided that $r_0 > 0$ and $\varepsilon_0 > 0$ are sufficiently small. Assume the contrary, hence there exist $\varepsilon_k \rightarrow 0$ as $k \rightarrow \infty$ and $(\tilde{\Delta}_{\varepsilon_k}, \tilde{v}_{\varepsilon_k}) \rightarrow (0, 0)$ as $k \rightarrow \infty$ such that

$$x_{\varepsilon_k}(\tilde{\Delta}_{\varepsilon_k}) - S(\tilde{v}_{\varepsilon_k}) = 0 \quad \text{and} \quad (\tilde{\Delta}_{\varepsilon_k}, \tilde{v}_{\varepsilon_k}) \neq (\Delta_{\varepsilon_k}, v_{\varepsilon_k}), \quad \text{for any } k \in \mathbb{N}.$$

Since $S : [-r_0, r_0] \rightarrow S([-r_0, r_0])$ is invertible then $(\tilde{\Delta}_{\varepsilon_k}, \tilde{v}_{\varepsilon_k}) \neq (\Delta_{\varepsilon_k}, v_{\varepsilon_k})$ implies $\tilde{\Delta}_{\varepsilon_k} \neq \Delta_{\varepsilon_k}$, say $\tilde{\Delta}_{\varepsilon_k} < \Delta_{\varepsilon_k}$. On the other hand $\dot{x}(0) \neq 0$ and so we can assume $\tilde{v}_{\varepsilon_k} \neq v_{\varepsilon_k}$. For any $v_1, v_2 \in \mathbb{R}^2$ we define $\angle(v_1, v_2)$ as follows

$$\angle(v_1, v_2) = \arccos \frac{\langle v_1, v_2 \rangle}{\|v_1\| \cdot \|v_2\|}.$$

Then we have

$$\angle(x_{\varepsilon_k}(\Delta_{\varepsilon_k}) - x_{\varepsilon_k}(\tilde{\Delta}_{\varepsilon_k}), \dot{x}_0(0)) = \angle(S(v_{\varepsilon_k}) - S(\tilde{v}_{\varepsilon_k}), \dot{x}_0(0)).$$

Passing to a subsequence if necessary we have that $\left\{ \frac{v_{\varepsilon_k} - \tilde{v}_{\varepsilon_k}}{\|v_{\varepsilon_k} - \tilde{v}_{\varepsilon_k}\|} \right\}_{k=1}^\infty$ converges.

Denote by $q \in \mathbb{R}$, $|q| = 1$, the limit of this sequence. Then

$$\angle(S(v_{\varepsilon_k}) - S(\tilde{v}_{\varepsilon_k}), \dot{x}_0(0)) \rightarrow \angle(S'(0)q, \dot{x}_0(0)) \quad \text{as } k \rightarrow \infty,$$

with $\angle(S'(0)q, \dot{x}_0(0)) \neq 0$, since, by Lemma 1, $\dot{x}(0) \notin S'(0)(\mathbb{R})$. Therefore, there exists $\alpha > 0$ such that

$$|\angle(x_{\varepsilon_k}(\Delta_{\varepsilon_k}) - x_{\varepsilon_k}(\tilde{\Delta}_{\varepsilon_k}), \dot{x}_0(0))| \geq \alpha > 0, \quad \text{for any } k \in \mathbb{N}. \quad (8)$$

Since $t \rightarrow x_{\varepsilon_k}(t)$ is a solution of (2) then, by Filippov's lemma ([8] or [4, Theorem 1.5.10]), there exists a singlevalued measurable function $h_{\varepsilon_k} : [0, T] \rightarrow \mathbb{R}^2$ such that

$$\begin{aligned}\dot{x}_{\varepsilon_k}(t) &= f(x_{\varepsilon_k}(t)) + \varepsilon_k h_{\varepsilon_k}(t), \quad \text{for a.a. } t \in [0, T], \\ h_{\varepsilon_k}(t) &\in g(t, x_{\varepsilon_k}(t), \varepsilon_k), \quad \text{for a.a. } t \in [0, T].\end{aligned}$$

Therefore

$$x_{\varepsilon_k}(\Delta_{\varepsilon_k}) - x_{\varepsilon_k}(\tilde{\Delta}_{\varepsilon_k}) = \int_{\tilde{\Delta}_{\varepsilon_k}}^{\Delta_{\varepsilon_k}} f(x_{\varepsilon_k}(\tau)) d\tau + \varepsilon_k \int_{\tilde{\Delta}_{\varepsilon_k}}^{\Delta_{\varepsilon_k}} h_{\varepsilon_k}(\tau) d\tau.$$

Due to the uniform convergence of x_{ε} to x_0 as $\varepsilon \rightarrow 0$ we have that

$$\sup_{k \in N} \{ \|x_{\varepsilon_k}(\tau)\| : \tau \in [0, T] \} < \infty$$

thus the assumptions on f and g permit to conclude that

$$\angle(x_{\varepsilon_k}(\Delta_{\varepsilon_k}) - x_{\varepsilon_k}(\tilde{\Delta}_{\varepsilon_k}), \dot{x}_0(0)) \rightarrow \angle(f(x_0(0)), \dot{x}_0(0)) \quad \text{as } k \rightarrow \infty,$$

hence $\angle(f(x_0(0)), \dot{x}_0(0)) = 0$ since $f(x_0(0)) = \dot{x}_0(0)$. This is a contradiction with (8) and so the proof is complete.

We are now in the position to prove inequality (5).

Theorem 2.6. *Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ is measurable and satisfying (H). Let x_{ε} be a T -periodic solution to the perturbed system (2) satisfying*

$$\|x_{\varepsilon}(t) - x_0(t)\| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0 \tag{9}$$

uniformly with respect to $t \in [0, T]$, where x_0 is a nondegenerate T -periodic cycle of the unperturbed system (1). Let $\varepsilon_0 > 0$ and $\{\Delta_{\varepsilon}\}_{\varepsilon \in (0, \varepsilon_0]} \subset \mathbb{R}$ be as in Lemma 2.5. Then there exists $M > 0$ such that

$$\|x_{\varepsilon}(t + \Delta_{\varepsilon}) - x_0(t)\| \leq M\varepsilon \quad \text{for any } t \in [0, T] \text{ and any } \varepsilon \in (0, \varepsilon_0]. \tag{10}$$

Proof. In the sequel $\varepsilon \in (0, \varepsilon_0]$ and $\tau \in [0, T]$. Consider the change of variables $\nu_{\varepsilon}(\tau) = \Omega(0, \tau, x_{\varepsilon}(\tau + \Delta_{\varepsilon}))$ in system (2). Observe that

$$x_{\varepsilon}(\tau + \Delta_{\varepsilon}) = \Omega(\tau, 0, \nu_{\varepsilon}(\tau)). \tag{11}$$

Taking the derivative in (11) with respect to τ we obtain

$$\dot{x}_{\varepsilon}(\tau + \Delta_{\varepsilon}) = f(\Omega(\tau, 0, \nu_{\varepsilon}(\tau))) + \Omega'(\tau, 0, \nu_{\varepsilon}(\tau))\dot{\nu}_{\varepsilon}(\tau). \tag{12}$$

On the other hand from (2) we have

$$\dot{x}_\varepsilon(\tau + \Delta_\varepsilon) \in f(\Omega(\tau, 0, \nu_\varepsilon(\tau))) + \varepsilon g(\tau + \Delta_\varepsilon, \Omega(\tau, 0, \nu_\varepsilon(\tau)), \varepsilon). \quad (13)$$

Since $\Omega'_\xi(\tau, 0, \nu_\varepsilon(\tau))$ is the fundamental matrix of a linear system thus it is invertible, then from (12) and (13) it follows

$$\dot{\nu}_\varepsilon(\tau) \in \varepsilon (\Omega'_\xi(\tau, 0, \nu_\varepsilon(\tau)))^{-1} g(\tau + \Delta_\varepsilon, \Omega(\tau, 0, \nu_\varepsilon(\tau)), \varepsilon),$$

and

$$\nu_\varepsilon(0) = x_\varepsilon(\Delta_\varepsilon) = x_\varepsilon(T + \Delta_\varepsilon) = \Omega(T, 0, \nu_\varepsilon(T)).$$

Since g is measurable then again by Filippov's lemma there exists a measurable singlevalued function $h_\varepsilon : [0, T] \rightarrow \mathbb{R}^2$ such that

$$h_\varepsilon(\tau) \in (\Omega'_\xi(\tau, 0, \nu_\varepsilon(\tau)))^{-1} g(\tau + \Delta_\varepsilon, \Omega(\tau, 0, \nu_\varepsilon(\tau)), \varepsilon), \quad \text{for a.a. } \tau \in [0, T]$$

and

$$\dot{\nu}_\varepsilon(\tau) = \varepsilon h_\varepsilon(\tau), \quad \text{for a.a. } \tau \in [0, T].$$

Therefore, $h_\varepsilon \in L^\infty([0, T], \mathbb{R}^2)$ and

$$\nu_\varepsilon(\tau) = \Omega(T, 0, \nu_\varepsilon(T)) + \varepsilon \int_0^\tau h_\varepsilon(s) ds, \quad \text{for any } \tau \in [0, T]. \quad (14)$$

Since, for any $\tau \geq 0$, $\nu_\varepsilon(\tau) \rightarrow x_0(0)$ as $\varepsilon \rightarrow 0$ we can write $\nu_\varepsilon(\tau)$ in the following form

$$\nu_\varepsilon(\tau) = x_0(0) + \varepsilon \mu_\varepsilon(\tau). \quad (15)$$

We now prove that the functions μ_ε are bounded on $[0, T]$ uniformly with respect to $\varepsilon \in (0, \varepsilon_0]$. For this, we first subtract $x_0(0)$ from both sides of (14), with $\tau = T$, obtaining

$$\varepsilon \mu_\varepsilon(T) = \varepsilon \Omega'_\xi(T, 0, x_0(0)) \mu_\varepsilon(T) + o(\varepsilon \mu_\varepsilon(T)) + \varepsilon \int_0^T h_\varepsilon(s) ds, \quad (16)$$

where, from (15), $\frac{o(\varepsilon \mu_\varepsilon(T))}{\|\varepsilon \mu_\varepsilon(T)\|} \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Since $x_\varepsilon(\Delta_\varepsilon) \in S(\{v \in \mathbb{R} : |v| \leq r_0\})$, then by Lemma 2.5 there exists $v_\varepsilon \in \mathbb{R}$, $|v_\varepsilon| \leq r_0$, such that

$$x_\varepsilon(\Delta_\varepsilon) = \Omega(T, 0, x_0(0) + A_1 v_\varepsilon) \quad (17)$$

and

$$v_\varepsilon \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (18)$$

Now by using (17) we can represent $\varepsilon\mu_\varepsilon(T)$ as follows

$$\begin{aligned}\varepsilon\mu_\varepsilon(T) &= \nu_\varepsilon(T) - x_0(0) = \Omega(0, T, x_\varepsilon(\Delta_\varepsilon)) - x_0(0) = \\ &= \Omega(0, T, \Omega(T, 0, x_0(0) + A_1 v_\varepsilon)) - x_0(0) = A_1 v_\varepsilon.\end{aligned}\quad (19)$$

Therefore (16) can be rewritten as follows

$$A_1 v_\varepsilon = \Omega'_\xi(T, 0, x_0(0)) A_1 v_\varepsilon + o(A_1 v_\varepsilon) + \varepsilon \int_0^T h_\varepsilon(s) ds. \quad (20)$$

Let us show that there exists $M_1 > 0$ such that

$$|v_\varepsilon| \leq \varepsilon M_1, \quad \text{for any } \varepsilon \in (0, \varepsilon_0]. \quad (21)$$

Arguing by contradiction we assume that there exist sequences $\{\varepsilon_k\}_{k \in \mathbb{N}} \subset (0, \varepsilon_0]$, $\varepsilon_k \rightarrow 0$ as $k \rightarrow \infty$, such that $|v_{\varepsilon_k}| = \varepsilon_k c_k$, where $c_k \rightarrow \infty$ as $k \rightarrow \infty$. Let $q_k = \frac{v_{\varepsilon_k}}{|v_{\varepsilon_k}|}$, then from (20) we have

$$A_1 q_k = \Omega'_\xi(T, 0, x_0(0)) A_1 q_k + \frac{o(A_1 v_{\varepsilon_k})}{|v_{\varepsilon_k}|} + \frac{1}{c_k} \int_0^T h_{\varepsilon_k}(s) ds, \quad (22)$$

where $\frac{o(A_1 v_{\varepsilon_k})}{|v_{\varepsilon_k}|} \rightarrow 0$ as $k \rightarrow \infty$, in fact $\frac{o(A_1 v_{\varepsilon_k})}{|v_{\varepsilon_k}|} = \frac{o(A_1 v_{\varepsilon_k})}{\|A_1 v_{\varepsilon_k}\|} \cdot \frac{\|A_1 v_{\varepsilon_k}\|}{|v_{\varepsilon_k}|}$.

Let $B = \{v_\varepsilon(\tau) : \tau \in [0, T], \varepsilon \in [0, 1]\}$. The continuity of Ω and condition (9) imply that B is bounded. Since also $(\Omega'_\xi)^{-1}$ is continuous, we can find $\Lambda > 0$ satisfying $\left\| (\Omega'_\xi(T, 0, v_\varepsilon(\tau))^{-1} \right\| \leq \Lambda$ for any $\tau \in [0, T]$ and any $\varepsilon \in [0, \varepsilon_0]$. Therefore, from assumption (H) we obtain

$$\left\| \int_0^T h_\varepsilon(s) ds \right\| \leq \varepsilon \Lambda \int_0^T \mu_B(s + \Delta_\varepsilon) ds < +\infty, \quad \text{for } \varepsilon \in [0, \varepsilon_0]. \quad (23)$$

Without loss of generality we may assume that the sequence $\{q_k\}_{k \in \mathbb{N}} \subset \mathbb{R}$ converges, let $q_0 = \lim_{k \rightarrow \infty} q_k$ with $|q_0| = 1$. By passing to the limit as $k \rightarrow \infty$ in (22) we have that

$$A_1 q_0 = \Omega'_\xi(T, 0, x_0(0)) A_1 q_0.$$

Therefore $A_1 q_0$ is the initial condition of a T -periodic solution to (4). On the other hand the cycle x_0 is nondegenerate, hence $A_1 q_0$ is linearly dependent with $\dot{x}_0(0)$ contradicting the choice of A_1 . Thus (21) is true for some $M_1 > 0$. From (15) and the fact that $\nu_\varepsilon(0) = x_\varepsilon(\Delta_\varepsilon)$ we have

$$\begin{aligned}\|x_\varepsilon(\Delta_\varepsilon) - x_0(0)\| &= \varepsilon \|\mu_\varepsilon(0)\| \leq \varepsilon \|\mu_\varepsilon(T)\| + \|\varepsilon \mu_\varepsilon(T) - \varepsilon \mu_\varepsilon(0)\| = \\ &= \varepsilon \|\mu_\varepsilon(T)\| + \|\nu_\varepsilon(T) - \nu_\varepsilon(0)\|.\end{aligned}\quad (24)$$

From (14) and (23) we have that there exists $M_2 > 0$ such that

$$\|\nu_\varepsilon(T) - \nu_\varepsilon(0)\| \leq \varepsilon M_2, \quad \text{for any } \varepsilon \in (0, \varepsilon_0]. \quad (25)$$

Therefore combining (19) with (21) and taking into account (25) we have from (24) that

$$\|x_\varepsilon(\Delta_\varepsilon) - x_0(0)\| \leq \varepsilon \|A_1\| M_1 + \varepsilon M_2, \quad \text{for any } \varepsilon \in (0, \varepsilon_0].$$

Since

$$\dot{x}_\varepsilon(t + \Delta_\varepsilon) \in f(x_\varepsilon(t + \Delta_\varepsilon)) + \varepsilon g(t + \Delta_\varepsilon, x_\varepsilon(t + \Delta_\varepsilon), \varepsilon)$$

and g is measurable then Filippov's lemma ensures the existence of a measurable singlevalued function $m_\varepsilon : [0, T] \rightarrow \mathbb{R}^2$ such that

$$\dot{x}_\varepsilon(t + \Delta_\varepsilon) = f(x_\varepsilon(t + \Delta_\varepsilon)) + \varepsilon m_\varepsilon(t), \quad \text{for a.a. } t \in [0, T]$$

and

$$m_\varepsilon(t) \in g(t + \Delta_\varepsilon, x_\varepsilon(t + \Delta_\varepsilon), \varepsilon), \quad \text{for a.a. } t \in [0, T].$$

This allows to conclude that

$$x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) = x_\varepsilon(\Delta_\varepsilon) - x_0(0) + \int_0^t (f(x_\varepsilon(s + \Delta_\varepsilon)) - f(x_0(s))) ds + \varepsilon \int_0^t m_\varepsilon(s) ds.$$

Therefore, there exists a constant $M_3 \geq 0$ such that, for any $\varepsilon \in (0, \varepsilon_0]$, we have

$$\begin{aligned} \|x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)\| &\leq (\varepsilon \|A_1\| M_1 + \varepsilon M_2) + \\ &\quad + M_3 \int_0^t \|x_\varepsilon(s + \Delta_\varepsilon) - x_0(s)\| ds + \varepsilon M_3, \end{aligned} \quad (26)$$

By means of the Gronwall-Bellman lemma, (compare e.g. [6, Chapter II, § 11]), inequality (26) implies

$$\|x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)\| \leq \varepsilon (\|A_1\| M_1 + M_2 + M_3) e^{M_3 T} \quad \text{for any } \varepsilon \in (0, \varepsilon_0].$$

and thus the proof is complete.

Remark 2.7. Observe that Theorem 2.6 does not require that (1) is a Hamiltonian system, indeed the crucial assumption is that the linearized system (4) has a not T -periodic solution.

3. First approximation formula for periodic solutions of the perturbed system

Denote by \tilde{z} a non-trivial T -periodic solution of the adjoint system

$$\dot{z} = -(f'(x_0(t)))^* z. \quad (27)$$

Let $t_* \in [0, T]$ such that

$$\tilde{z}_1(t_*) = 0,$$

hence $\tilde{z}_2(t_*) \neq 0$. We begin the Section by studying the behavior, as $\varepsilon \rightarrow 0$, of the scalar product

$$\left\langle \tilde{z}(t), \frac{x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)}{\varepsilon} \right\rangle \quad (28)$$

which is the starting point for deriving the first approximation formula (6). To this end we denote by $\hat{z} = (\hat{z}_1, \hat{z}_2)$ any solution of (27) defined in $[0, T]$ linearly independent with \tilde{z} and introduce the multivalued map $M^\perp : [0, T] \rightarrow K(\mathbb{R})$ as follows

$$M^\perp(t) = \left\{ \gamma(t_*) \int_{t-T}^t \langle -\hat{z}(\tau), h(\tau) \rangle d\tau : h \in L^\infty([-T, T], \mathbb{R}^2), h(t) \in g(t, x_0(t), 0) \text{ for a.a } t \in [-T, T] \right\}, \quad (29)$$

$$\text{where } \gamma(t_*) = \frac{\tilde{z}_2(t_*)}{\tilde{z}_2(T + t_*) - \tilde{z}_2(t_*)}.$$

We can prove the following result.

Theorem 3.1. *Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ upper semicontinuous and satisfying (H). Let x_ε be a T -periodic solution to the perturbed system (2) such that*

$$\|x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)\| \leq M\varepsilon \quad \text{for any } t \in [0, T] \text{ and any } \varepsilon \in (0, \varepsilon_0], \quad (30)$$

where $\Delta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$, M and ε_0 are positive constants and x_0 is a nondegenerate cycle of the Hamiltonian system (1). Then

$$\lim_{\varepsilon \rightarrow 0} \rho \left(\frac{1}{\varepsilon} \langle \tilde{z}(t), x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) \rangle, \frac{\tilde{z}_2(t_*)}{\tilde{z}_2(T + t_*) - \tilde{z}_2(t_*)} M^\perp(t) \right) = 0 \quad (31)$$

uniformly with respect to $t \in [0, T]$, where for any $v \in \mathbb{R}^n$ and $S \subset \mathbb{R}^n$ the distance $\rho(v, S)$ is defined as $\rho(v, S) = \inf_{s \in S} \|v - s\|$.

To prove Theorem 3.1 we need the following lemma.

Lemma 3.2. *Assume that the T -periodic system*

$$\dot{u} = A(t)u, \quad u \in \mathbb{R}^2 \quad (32)$$

has the characteristic multiplier +1 of algebraic multiplicity 2. Let us denote by $\tilde{u} = (\tilde{u}_1, \tilde{u}_2)$ a T -periodic solution of (32) such that

$$\tilde{u}_1(0) = 0, \quad \tilde{u}_2(0) \neq 0.$$

Denote by $\hat{u} = (\hat{u}_1, \hat{u}_2)$ any solution of (32) satisfying

$$\hat{u}_1(0) \neq 0.$$

Then

$$\hat{u}(t+T) = \hat{u}(t) + \frac{\hat{u}_2(T) - \hat{u}_2(0)}{\tilde{u}_2(0)} \tilde{u}(t), \quad \text{for any } t \in \mathbb{R}.$$

This result has been proved in [18, Lemma 4.2] under the additional assumption $\hat{u}_2(0) = 0$. Though it is immediate to see that avoiding this assumption does not affect the proof of [18, Lemma 4.2] at all we provide here a proof of Lemma 3.2 for a sake completeness.

Proof. Denote by X the fundamental matrix of system (32) such that $X(0) = I$. Since

$$X(T) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix},$$

then $X(T) = \begin{pmatrix} a & 0 \\ b & 1 \end{pmatrix}$ with $a, b \in \mathbb{R}$. By our assumption $X(T)$ has two eigenvalues equal to +1, therefore

$$X(T) = \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix}, \quad b \in \mathbb{R}.$$

We have

$$\begin{aligned} X(t+T)\hat{u}(0) &= X(t)X(T)\hat{u}(0) = X(t) \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix} \hat{u}(0) = \\ &= X(t)\hat{u}(0) + X(t) \begin{pmatrix} 0 \\ b\hat{u}_1(0) \end{pmatrix} = X(t)\hat{u}(0) + \frac{b\hat{u}_1(0)}{\tilde{u}_2(0)} \tilde{u}(t). \end{aligned}$$

On the other hand

$$X(T)\hat{u}(0) = \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix} \hat{u}(0) = \hat{u}(0) + \begin{pmatrix} 0 \\ b\hat{u}_1(0) \end{pmatrix},$$

which implies $b\hat{u}_1(0) = \hat{u}_2(T) - \hat{u}_2(0)$. This completes the proof.

We now prove Theorem 3.1.

Proof. In what follows $\varepsilon \in (0, \varepsilon_0]$, $t, \tau \in [-T, T]$ and \tilde{z}, \hat{z} are the functions introduced at the beginning of this section. Let A be a nonsingular 2×2 matrix such that

$$\hat{z}(0)^* A = (0, 1). \quad (33)$$

Let $Y(t)$ be the fundamental matrix of the linearized system (4) with initial condition $Y(0) = A$. Let

$$Z(t) = (Y(t)^*)^{-1} \quad (34)$$

and define $a_\varepsilon \in C([-T, T], \mathbb{R}^2)$ as follows

$$a_\varepsilon(t) = Z(t)^* \frac{x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)}{\varepsilon}.$$

Then we have

$$x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) = \varepsilon Y(t) a_\varepsilon(t). \quad (35)$$

In what follows by $o(\varepsilon)$, $\varepsilon > 0$, we will denote a function, which may depend also on other variables, having the property that $\frac{o(\varepsilon)}{\varepsilon} \rightarrow 0$ as $\varepsilon \rightarrow 0$ uniformly with respect to these variables when they belong to any bounded set. Since

$$\dot{x}_\varepsilon(t + \Delta_\varepsilon) \in f(x_\varepsilon(t + \Delta_\varepsilon)) + \varepsilon g(t + \Delta_\varepsilon, x_\varepsilon(t + \Delta_\varepsilon), \varepsilon), \quad \text{for a.a. } t \in \mathbb{R}$$

then, again by Filippov's lemma there exists a measurable singlevalued function $h_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}^2$ such that

$$\dot{x}_\varepsilon(t + \Delta_\varepsilon) = f(x_\varepsilon(t + \Delta_\varepsilon)) + \varepsilon h_\varepsilon(t), \quad \text{for a.a. } t \in \mathbb{R} \quad (36)$$

and

$$h_\varepsilon(t) \in g(t + \Delta_\varepsilon, x_\varepsilon(t + \Delta_\varepsilon), \varepsilon), \quad \text{for a.a. } t \in \mathbb{R}.$$

By subtracting (1) where $x(t)$ is replaced by $x_0(t)$ from (36) we obtain

$$\dot{x}_\varepsilon(t + \Delta_\varepsilon) - \dot{x}_0(t) = f'(x_0(t))(x_\varepsilon(t + \Delta_\varepsilon) - x_0(t)) + \varepsilon h_\varepsilon(t) + o_t(\varepsilon), \quad (37)$$

for a.a. $t \in [-T, T]$, here $\varepsilon \rightarrow o_t(\varepsilon)$ is such that $o_{t+T}(\cdot) = o_t(\cdot)$ for any $t \in \mathbb{R}$. By substituting (35) into (37) we have

$$\varepsilon \dot{Y}(t) a_\varepsilon(t) + \varepsilon Y(t) \dot{a}_\varepsilon(t) = \varepsilon f'(x_0(t)) Y(t) a_\varepsilon(t) + \varepsilon h_\varepsilon(t) + o_t(\varepsilon),$$

for a.a. $t \in [-T, T]$. Since $f'(x_0(t)) Y(t) = \dot{Y}(t)$ the last formula can be rewritten as follows

$$\varepsilon Y(t) \dot{a}_\varepsilon(t) = \varepsilon h_\varepsilon(t) + o_t(\varepsilon), \quad \text{for a.a. } t \in [-T, T]. \quad (38)$$

By means of Perron's lemma [20], (see also Demidovich [6, Sec. III, §12]), formula (33) implies that

$$\widehat{z}(t)^* Y(t) = (0, 1) \quad \text{for any } t \in \mathbb{R}. \quad (39)$$

Therefore, applying $\widehat{z}(t)^*$ to both sides of (38) we have

$$\varepsilon(\dot{a}_{\varepsilon,2})(t) = \varepsilon \widehat{z}(t)^* h_{\varepsilon}(t) + \widehat{z}(t)^* o_t(\varepsilon), \quad \text{for a.a. } t \in [-T, T],$$

where $a_{\varepsilon,2}(t)$ is the second component of the vector $a_{\varepsilon}(t)$, and so

$$a_{\varepsilon,2}(t) = a_{\varepsilon,2}(t_0) + \int_{t_0}^t \langle \widehat{z}(\tau), h_{\varepsilon}(\tau) \rangle d\tau + \int_{t_0}^t \left\langle \widehat{z}(\tau), \frac{o_{\tau}(\varepsilon)}{\varepsilon} \right\rangle d\tau, \quad (40)$$

for all $t, t_0 \in [-T, T]$. From (34) we have that $Z(0)^* Y(0) = I$. Therefore

$$([Z(0)]_2)^* A = (0, 1),$$

where $[Z(0)]_2$ denotes the second column of $Z(0)$. Thus $[Z(0)]_2 = \widehat{z}(0)$. Therefore,

$$a_{\varepsilon,2}(t) = \left\langle \widehat{z}(t), \frac{x_{\varepsilon}(t + \Delta_{\varepsilon}) - x_0(t)}{\varepsilon} \right\rangle.$$

Since \widehat{z} is linearly independent with \widetilde{z} then

$$\widehat{z}_1(t_*) \neq 0.$$

Since system (1) is Hamiltonian then the algebraic multiplicity of the characteristic multiplier +1 of linearized system (4) is equal to 2. By lemma 3.2 we have

$$\widehat{z}(t) = \widehat{z}(t - T) + \frac{\widehat{z}_2(T + t_*) - \widehat{z}_2(t_*)}{\widetilde{z}_2(t_*)} \widetilde{z}(t) = \widehat{z}(t - T) + \frac{1}{\gamma(t_*)} \widetilde{z}(t),$$

that implies

$$a_{\varepsilon,2}(t_0) = a_{\varepsilon,2}(t_0 - T) + \frac{1}{\gamma(t_*)} \left\langle \widetilde{z}(t_0), \frac{x_{\varepsilon}(t_0 + \Delta_{\varepsilon}) - x_0(t_0)}{\varepsilon} \right\rangle.$$

Substituting the last formula into (40) we obtain

$$\begin{aligned} \int_{t_0}^{t_0-T} \langle \widehat{z}(\tau), h_{\varepsilon}(\tau) \rangle d\tau &= -\frac{1}{\gamma(t_*)} \left\langle \widetilde{z}(t_0), \frac{x_{\varepsilon}(t_0 + \Delta_{\varepsilon}) - x_0(t_0)}{\varepsilon} \right\rangle - \\ &\quad - \int_{t_0}^{t_0-T} \left\langle \widehat{z}(\tau), \frac{o_{\tau}(\varepsilon)}{\varepsilon} \right\rangle d\tau. \end{aligned} \quad (41)$$

We claim that

$$\lim_{\varepsilon \rightarrow 0} \rho \left(\gamma(t_*) \int_{t_0}^{t_0-T} \langle \widehat{z}(\tau), h_{\varepsilon}(\tau) \rangle d\tau, M^{\perp}(t) \right) = 0, \quad (42)$$

uniformly with respect to $t_0 \in [0, T]$, with h_ε defined as in (36). To prove this we observe that the subset of \mathbb{R} given by

$$\begin{aligned} M : &= \left\{ \gamma(t_*) \int_{-T}^T \langle \hat{z}(\tau), h(\tau) \rangle d\tau : \right. \\ &\quad \left. h \in L^\infty([-T, T], \mathbb{R}^2) \text{ and } h(t) \in g(t, x_0(t), 0) \text{ for a.a. } t \in [-T, T] \right\} \end{aligned}$$

is nonempty and compact; hence, for each $\varepsilon \in (0, \varepsilon_0]$, there exists $k_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ such that $k_\varepsilon(t) \in g(t, x_0(t), 0)$ for a.a. t and

$$\rho \left(\gamma(t_*) \int_{-T}^T \langle \hat{z}(\tau), h_\varepsilon(\tau) \rangle d\tau, M \right) = |\gamma(t_*)| \left| \int_{-T}^T \langle \hat{z}(\tau), h_\varepsilon(\tau) - k_\varepsilon(\tau) \rangle d\tau \right|.$$

The upper semicontinuity of g in the bounded set

$$[-T, T] \times \{x_\varepsilon(t) : t \in [0, T], \varepsilon \in [0, \varepsilon_0]\} \times [0, \varepsilon_0]$$

implies that, given $\delta > 0$, there exists $\varepsilon_1 \in (0, \varepsilon_0]$ such that, for all $\varepsilon \in [0, \varepsilon_1]$, we have that

$$h_\varepsilon(t) \in g(t + \Delta_\varepsilon, x_\varepsilon(t + \Delta_\varepsilon), \varepsilon) \subset B_\delta(g(t, x_0(t), 0)), \quad \text{for a.a. } t \in [-T, T], \quad (43)$$

Fix an arbitrary $\delta > 0$ and let $\varepsilon_1 \in (0, \varepsilon_0]$ satisfying (43). Let $\varepsilon \in [0, \varepsilon_1]$ and $t_0 \in [0, T]$. We obtain

$$\begin{aligned} &\rho \left(\gamma(t_*) \int_{t_0-T}^{t_0} \langle \hat{z}(\tau), h_\varepsilon(\tau) \rangle d\tau, M^\perp(t) \right) \leq \\ &\leq |\gamma(t_*)| \left| \int_{t_0-T}^{t_0} \langle \hat{z}(\tau), h_\varepsilon(\tau) - k_\varepsilon(\tau) \rangle d\tau \right| \leq \\ &\leq |\gamma(t_*)| \int_{-T}^T | \langle \hat{z}(\tau), h_\varepsilon(\tau) - k_\varepsilon(\tau) \rangle | d\tau \leq 2T |\gamma(t_*)| \delta \|\hat{z}\|_C. \end{aligned}$$

which implies our assertion (42). According to (41), the proof is complete.

Remark 3.3. The assumption that (1) is Hamiltonian ensures that the linearized system (4) has a characteristic multiplier $+1$ of algebraic multiplicity 2 and so the assumption of Lemma 3.2. Alternatively, we could directly assume that the algebraic multiplicity of the characteristic multiplier $+1$ of (4) is equal to 2. The latter is a bit more general. The same consideration applies to Theorems 3.6 and 4.2 below.

We have the following result.

Lemma 3.4. *Let x_0 be a nondegenerate T -periodic cycle of the Hamiltonian system (1). Let \tilde{z} be any T -periodic solution of the adjoint system (27). Then*

$$\langle \dot{x}_0(t), \tilde{z}(t) \rangle = 0, \quad \text{for any } t \in \mathbb{R}. \quad (44)$$

Proof. Let $t_* \in [0, T]$ be such that $\tilde{z}_1(t_*) = 0$. Let \hat{z} be any solution of (27) linearly independent with \tilde{z} . Then from Lemma 3.2 we have

$$\langle \dot{x}_0(t), \hat{z}(t+T) \rangle = \langle \dot{x}_0(t), \hat{z}(t) \rangle + \frac{\hat{z}_2(T+t_*)}{\hat{z}_2(t_*)} \langle \dot{x}_0(t), \tilde{z}(t) \rangle, \quad \text{for any } t \in \mathbb{R}.$$

Perron's lemma [20] implies that $\langle \dot{x}_0(t), \hat{z}(t+T) \rangle = \langle \dot{x}_0(t), \hat{z}(t) \rangle$ for any $t \in \mathbb{R}$ and thus (44).

Lemma 3.4 allows the reader to better understand the substantial difference between the situation when the cycle x_0 is isolated, which is studied in [16] and [17] and the present situation when the cycle is non-isolated. In fact, in [16] and [17] it is shown that $\langle \dot{x}_0(t), \tilde{z}(t) \rangle \neq 0$, for any $t \in \mathbb{R}$, which is the contrary of (44).

Remark 3.5. Let \tilde{z} be any T -periodic solution of the adjoint system (27) and \hat{z} any solution of (27) linearly independent with \tilde{z} . Lemma 3.4 ensures that $\langle \dot{x}_0(t), \tilde{z}(t) \rangle = 0$ for any $t \in \mathbb{R}$, moreover from the Perron's Lemma $\langle \dot{x}_0(t), \hat{z}(t) \rangle = \langle \dot{x}_0(0), \hat{z}(0) \rangle \neq 0$ for any $t \in \mathbb{R}$. Without loss of generality we can assume that $\langle \dot{x}_0(0), \hat{z}(0) \rangle = 1$.

Let y be the function defined by

$$y(t)^* = \left(\frac{-\hat{z}_2(t)}{\det(\hat{z}(t), \tilde{z}(t))}, \frac{\hat{z}_1(t)}{\det(\hat{z}(t), \tilde{z}(t))} \right)$$

then

$$(\dot{x}_0(t), y(t)) = \begin{pmatrix} \hat{z}(t)^* \\ \tilde{z}(t)^* \end{pmatrix}^{-1}. \quad (45)$$

is a matrix solution of the linearized system (4) ([6, Chapter III, §12]).

We can now formulate the following result.

Theorem 3.6. *Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ upper semicontinuous and satisfying (H). Let x_ε be a T -periodic solution to perturbed system (2) such that*

$$\|x_\varepsilon(t) - x_0(t)\| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

uniformly with respect to $t \in [0, T]$, where x_0 is a nondegenerate T -periodic cycle of the Hamiltonian system (1). Let \tilde{z}, \hat{z} be as in Remark 3.5 and \dot{x}_0, y as in (45). Then there exists a family $\{\Delta_\varepsilon\}_{\varepsilon>0}$ such that $\Delta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ and

$$\lim_{\varepsilon \rightarrow 0} \rho(x_\varepsilon(t + \Delta_\varepsilon) - x_0(t), \varepsilon M^\perp(t)y(t) + \langle \hat{z}(t), x_\varepsilon(t - \Delta_\varepsilon) - x_0(t) \rangle \dot{x}_0(t)) = 0, \quad (46)$$

uniformly with respect to $t \in [0, T]$.

Proof. The proof of theorem 3.6 follows from the following representation which is a consequence of (45)

$$\begin{aligned} x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) = & \langle \tilde{z}(t), x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) \rangle y(t) + \\ & + \langle \hat{z}(t), x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) \rangle \dot{x}_0(t), \end{aligned}$$

and Theorem 3.1.

4. A symmetric case

In this section we consider the situation when the unperturbed Hamiltonian system (1) possesses the following symmetry properties:

$$f_1(\xi_1, \xi_2) = f_1(-\xi_1, \xi_2), \quad (47)$$

$$f_2(\xi_1, \xi_2) = -f_2(-\xi_1, \xi_2), \quad (48)$$

$$(f_1)'_{(1)}(\xi_1, \xi_2) = -(f_2)'_{(2)}(\xi). \quad (49)$$

where $(h)'_{(i)}$, $i = 1, 2$ denotes the derivative of h with respect to the i -variable. The main consequence of this symmetry assumption is given by the following lemma whose prove is immediate.

Lemma 4.1. ([18, Lemma 4.4]) Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and that properties (47)-(49) hold true. Let x_0 be a nondegenerate cycle of the Hamiltonian system (1) and denote by y the solution of the linearized system (4) satisfying

$$\begin{pmatrix} y_1(0) \\ y_2(0) \end{pmatrix} = \begin{pmatrix} -\dot{x}_{0,2}(0) \\ \dot{x}_{0,1}(0) \end{pmatrix}. \quad (50)$$

Then the functions

$$\hat{z}(\theta) = \begin{pmatrix} y_2(\theta) \\ -y_1(\theta) \end{pmatrix}, \quad \tilde{z}(\theta) = \begin{pmatrix} -\dot{x}_{0,2}(\theta) \\ \dot{x}_{0,1}(\theta) \end{pmatrix}, \quad \theta \in \mathbb{R},$$

where $\dot{x}_0(\theta) = (\dot{x}_{0,1}(\theta), \dot{x}_{0,2}(\theta))$, are linearly independent solutions of the adjoint system (27).

Lemma 4.1 allows us to rewrite the multivalued map $M^\perp : [0, T] \rightarrow K(\mathbb{R})$ defined in (29) as follows

$$\begin{aligned} M^\perp(t) = & \left\{ \frac{\dot{x}_{0,1}(t_*)}{y_1(T + t_*)} \int_{t-T}^t \det(-y(\tau), h(\tau)) d\tau : \right. \\ & \left. h \in L^\infty([-T, T], \mathbb{R}^2) : h(t) \in g(t, x_0(t), 0) \text{ for a.a } t \in [-T, T] \right\}. \end{aligned}$$

where $t_* \in [0, T]$ is such that $\dot{x}_{0,2}(t_*) = 0$. Therefore Theorem 3.6 takes the form of the following Theorem 4.2 when the symmetry assumptions (47)-(49) are satisfied. In particular, observe that the statement of Theorem 4.2 refers only to the linearized system (4) and not to the adjoint system (27).

Theorem 4.2. Assume $f \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and $g : \mathbb{R} \times \mathbb{R}^2 \times [0, 1] \rightarrow K(\mathbb{R}^2)$ upper semicontinuous and satisfying (H). Let x_ε be a T -periodic solution to perturbed system (2) satisfying

$$\|x_\varepsilon(t) - x_0(t)\| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

uniformly with respect to $t \in [0, T]$, where x_0 is a nondegenerate T -periodic cycle of the Hamiltonian system (1). Let y be the solution of the linearized system (4) with the initial condition (50). Then there exists a family $\{\Delta_\varepsilon\}_{\varepsilon>0}$ such that $\Delta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ and

$$x_\varepsilon(t+\Delta_\varepsilon) - x_0(t) \in \varepsilon M^\perp(t)y(t) + \left\langle \begin{pmatrix} y_2(t) \\ -y_1(t) \end{pmatrix}, x_\varepsilon(t + \Delta_\varepsilon) - x_0(t) \right\rangle \dot{x}_0(t) + o(\varepsilon),$$

uniformly with respect to $t \in [0, T]$.

Acknowledgement. The first author was supported by the Grant BF6M10 of Russian Federation Ministry of Education and U.S. CRDF (BRHE), by RFBR Grants 09-01-90407, 09-01-00468, by the President of Russian Federation Young PhD Student grant MK-1620.2008.1 and by "Researchers Mobility in the Field of Scientific and Cultural Cooperation Programmes" of the University of Modena and Reggio Emilia.

The second author was supported by the national research project PRIN "Ordinary Differential Equations and Applications"

Finally, the third author was supported by the national research project PRIN "Control, Optimization and Stability of Nonlinear Systems: Geometric and Topological Methods".

References

- [1] Aubin, J.P. and Cellina, A., *Differential inclusions*, Grundlehren der mathematischen Wissenschaften, 264, Springer-Verlag, 1984.
- [2] Awrejcewicz, J. and Lamarque, C.-H., *Bifurcation and chaos in nonsmooth mechanical systems*, World Scientific Series on Nonlinear Science, Series A-45, World Scientific Publishing Co., 2003.
- [3] Awrejcewicz, J. and Holicke, M.M., *Smooth and nonsmooth high dimensional chaos and the Melnikov-type methods*, World Scientific Series on Nonlinear Science, Series A-60, World Scientific Publishing Co., 2007.
- [4] Borisovich, Yu. G., Gelman, B.D., Myshkis, A.D. and Obukhovskii, V.V., *Introduction to the theory of multivalued mappings*, Voronezhskii Gosudarstvennyi Universitet, Voronezh, 1986.
- [5] Chicone, C. and Jacob, M., Bifurcations of critical periods for plane vector fields, *Trans. Amer. Math. Soc.* 312 (1989), 433–486.

- [6] Demidovich, B.P., *Lectures on the mathematical theory of stability* (in Russian). Izdat. Nauka, Moscow, 1967.
- [7] Fečkan, M., Bifurcation of periodic solutions in differential inclusions, *Appl. Math.* 42 (1997), 369–393.
- [8] Filippov, A. F., On some questions in the theory of optimal regulation: existence of a solution of the problem of optimal regulation in the class of bounded measurable functions (in Russian). *Vestnik Moskov. Univ. Ser. Mat. Meh. Astr. Fiz. Him.* (1959), 25–32.
- [9] Filippov, A. F., *Differential equations with discontinuous righthand side*, Mathematics and its Applications, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1988.
- [10] Guckenheimer, J. and Holmes, P., *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields*, Revised and corrected reprint of the 1983 original. Applied Mathematical Sciences No. 42, Springer-Verlag, New York, 1990.
- [11] Kamenskii, M. Makarenkov, O. and Nistri, P., A continuation principle for a class of periodically perturbed autonomous systems, *Math. Nachr.* 281 (2008), 42–61.
- [12] Kamenskii, M., Obukhovskii, V.V. and Zecca, P., *Condensing Multivalued Maps and Semilinear Differential Inclusions in Banach Space*, W. deGruyter, Berlin, (2001).
- [13] Kolmogorov, A.N. and Fomin, S.V., *Elements of the theory of functions and functional analysis* (in Russian). Fourth edition, revised. Izdat. “Nauka”, Moscow, 1976.
- [14] Kolovskii, M.Z., An application of the small-parameter method for determining discontinuous periodic solutions, in *Proc. Internat. Sympos. Non-linear Vibrations* (in Russian.) Izdat. Akad. Nauk Ukrainsk. SSR, Kiev. I (1961), 264–276.
- [15] Markus, K., *Non-smooth dynamical systems*, Lecture Notes in Mathematics, No. 1744. Springer-Verlag, Berlin, 2000.
- [16] Makarenkov, O. and Nistri, P., Periodic solutions for planar autonomous systems with nonsmooth periodic perturbations, *J. Math. Anal. Appl.* 338 (2008), 1401–1417.
- [17] Makarenkov, O. and Nistri, P., On the rate of convergence of periodic solutions in perturbed autonomous systems as the perturbation vanishes, *Commun. Pure Appl. Anal.* 7 (2008), 49–61.
- [18] Makarenkov, O., Poincaré index and periodic solutions of perturbed autonomous systems, *Trudy Moskov. Mat. Obšč.* 70 (2008), in press.
- [19] Melnikov, V. K. On the stability of a center for time-periodic perturbations (in Russian). *Trudy Moskov. Mat. Obšč.* 12 (1963), 3–52.
- [20] Perron, O., Die Ordnungszahlen der Differentialgleichungssysteme, *Math. Zeitschr.* 31 (1930), 748–766.
- [21] Rhouma, M.B.H. and Chicone, C., On the continuation of periodic orbits, *Methods Appl. Anal.* 7 (2000), 85–104.